## SUPPLEMENTAL LABEL

# Acephate 90% SP

INSECTICIDE

## KEEP OUT OF REACH OF CHILDREN CAUTION

EPA Reg. No. 81964-3

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THE ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

This labeling must be in the possession of the user at the time of the application. Read the entire label affixed to the Acephate 90% SP container for additional precautionary statements. Use of Acephate 90% SP according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for Acephate 90% SP.

#### **SOYBEANS**

#### USE PRECAUTIONS:

- Do not apply more than 1.66 pounds per acre (1.5 lbs. of active ingredient) per season.
- Do not harvest soybeans for at least 14 days after last application.
- Do not graze or cut vines for hay or forage.
- Always read and follow all label directions, restrictions and precautions when using any pesticide alone or in tank mix combinations. The most restrictive labeling applies when using a tank mix.

#### **FOLIAR APPLICATIONS:**

By Air:

Apply using 5 to 10 gallons per acre of spray.

By Ground: Apply using 10 to 50 gallons per acre of spray.

PESTS CONTROLLED	ACEPHATE 90% SP APPLICATION RATE (PER ACRE)	SPECIFIC INSTRUCTIONS
Grasshopper Thrips	0.28 to 0.56 lb. (4.5 to 9 oz.)	To maintain control: For applications of 0.5 lbs. or less active ingredient per acre, repeat applications at 3-day spray intervals as necessary.*
Potato Leafhopper Stink Bugs	0.56 to 1.1 lbs. (9 to 17.6 oz.)	

ACCEPTED 2007 MAR 1 Under the Pederal Insecticide, Pungicide, and Rodenticide Act. as amended, for the pesticide Replaced under EPA Reg. No. 8/9

CIFIC INSTRUCTIONS
tions greater than 0.5 lbs. dient per acre, repeat at 7-day spray intervals as
i

### Manufactured for:

ChemStarr LLC 21 Hubble Irvine, CA 92618