| 82493.2 | | 11 | 5/2005 | | 1 |
|---|--|---|--|--|--|
| Please read instructions on reverse bef | United Stat United Stat Vironmental Prote Washington, DC | ction Agenc | ☐ Regis У ☐ Amen | ved, OMB No. 2070. tration dment - Notification | 0060, Approval expires 05-31-9 OPP Identifier Number 265394 |
| | Applic | cation for Pe | esticide - Sectio | on l | |
| 1. Company/Product Number 82493-2 | | | Product Manager mpkins | | 3. Proposed Classification |
| 4. Company/Product (Name) Glygran® WDG | | PM# 25 | <u></u> | <u>, es</u> | |
| 5. Name and Address of Applicant (Ir Agriventures, Inc. 9775 East Balancing Rock Scottsdale, AZ 85262 NOTE: PLEASE SEND ALL CORRE "CONTACT POINT" LISTED BELOW Check if this is a new a | Rd SPONDENCE TO | (b)(l), n to: EPA R | ny product is simil Reg. No | lar or identical in | th FIFRA Section 3(c)(3) composition and labeling |
| | | Secti | on - II | | |
| | I page(s) if necessa Please refe <u>Notificat</u> ne provisions of PR Notification is provisions of PR Notifications of PR Notification is provisions of PR Notification is provisions of PR Notification is provisions of PR Notifications of PR Notifi | Ary. (For Sect Prence co lon in Accordar lice 98-10 and EP product. I underst in is not consister action and penalti Date: Section No. per t. container (s) Retail Contair | Other - Explain Other - Explain Over letter for New Presence of | n below II.) or details PR 152.46, and no o on of 18 U.S.C. Sec. R Notice 95-2 and 40 and 14 of FIFRA. 2 3 / C Ackaging No. per container 5. Location On Label | 1001 to willfully make any false 0 CFR 152.46, this product may 2. Type of Container Metal Plastic Glass Paper Other (Specify) of Label Directions |
| 6. Manner in Which Label is Affixed | 🔲 Pa | lhograph aper glued rencile <u>d</u> | Other _ | | ······ |
| | | Sectio | on – IV | <u></u> | |
| Contact Point (Complete items di Name Georgia Anastasiou - Lewis 122 C St. NW #740, Washing I certify that the statements I have r acknowledge that any knowingly fall under applicable law. | & Harrison, LLC, ton DC 20001 Certific nade on this form and a | Title Agent cation | t for Agriventure | s, Inc. | this application) elephone No. (Include Area Co 202-393-3903 ext. 19 6. Date Application Received (Stamperi) |
| 2. Signature | | | nt for Agriventur | res, Inc. | |
| 4. Typed Name Georgia Anastasiou - Lewis a | & Harrison, LLC | 5. Date | October 28, 20 | 05 | |

| EPA Form 8570-1 (Rev. 8-94) Previous editions are obsole | te |
|--|----|
|--|----|

NOTIFICATION

Glygran[®] WDG

- A Non-selective and Systemic Post Emergent Herbicide for Weed Control on Commercial, Industrial & Residential Sites.
- Provides Systemic Control of Listed Broadleaf Weeds, Grasses, Woody Brush and Trees.

ACTIVE INGREDIENT:

| Glyphosate*: N-(phosphonomethyl) glycine | 80.0% |
|--|--------------|
| INERT INGREDIENTS: | <u>20.0%</u> |
| TOTAL: | 100.0% |

*The glyphosate in this product is manufactured in Brazil.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si Usted no entiende la etiqueta, busque a alguien para que se la explique a Usted en detalle. (If you do not understand this label, find someone to explain it to you in detail)

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND COMPLETE DIRECTIONS FOR USE

NET CONTENTS

Agriventures, Inc. 9775 E. Balancing Rock Road Scottsdale, AZ 85262 USA EPA Reg. No. 82493-2 EPA Est. No. 11773 IA-01

Glygran[®] WDG ts a registered trademark of Agriventures, Inc.

NOV 1 5 2005

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing.

FIRST AID

| If on skin | Take off contaminated clothing. |
|----------------|--|
| or clothing: | Rinse skin immediately with plenty of water for 15-20 minutes. |
| | Call a poison control center or doctor for treatment advice. |
| if in eyes: | Hold eye open and rinse slowly and gently with water for 15-20 minutes. |
| | • Remove contact lenses if present, after the first 5 minutes, then continue rinsing. |
| | Call a poison control center or doctor for treatment advice. |
| if | Call a poison control center or doctor immediately for treatment |
| swallowed: | advice. |
| | Have person sip a glass of water if able to swallow. |
| | • Do not induce vomiting unless told to do so by a poison control center or doctor. |
| | Do not give anything by mouth to an unconscious person. |
| If inhaled: | Move person to fresh air. |
| | • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. |
| | Call a poison control center or doctor for further treatment advice |
| Have the pr | oduct container or label with you when calling a poison control |
| | center or doctor or going for treatment. |
| | ician: Probable mucosal damage may contraindicate the use of |
| gastric lavage |). |

[The following language is only for use with labels containing fallow uses]

Personal Protective Equipment (PPE)

Some of the materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear: long sleeved shirts, long pants, shoes plus socks, and chemical resistant gloves made of any waterproof material such as polyethytene or polyvinytchloride. Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements: When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Úser Safety Recommendations: Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

Spray solutions containing this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers. DO NOT mix, store or apply this product or spray solutions of this product in galvanized steel or unlined steel (except stainless steel) containers or spray tanks. This product or spray solutions containing this product react with such containers and tanks to produce hydrogen gas, which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, lighted cigarette, welder's torch or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ AND FOLLOW ALL DIRECTIONS AND USE PRECAUTIONS ON THIS LABEL PRIOR TO APPLICATION. NOT FOR USE ON PLANTS BEING GROWN FOR SALE, OR OTHER COMMERCIAL USE ON SOD, OR FOR COMMERCIAL SEED PRODUCTION, OR FOR RESEARCH PURPOSES.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Do not apply this product through any type of irrigation system. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

[The following language is for use only with labels including fallow uses]

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, Chemicalresistant gloves made of any waterproof material, and, shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Turf, Landscape, and other Non-crop Uses: Do not enter or allow entry into treated areas until sprays have dried unless wearing the personal protective equipment listed on this label.

STORAGE AND DISPOSAL

Prohibitions: Do not contaminate water, food or feed by storage or disposal. Do not store under conditions which might adversely affect the container or its ability to function properly. Such conditions include, but are not limited to, positioning of the container in storage, storage temperature, potential for crushing or damage due to stacking, and penetration of moisture.

Pesticide Storage: Store in a safe manner. Store in original container only. Store in a cool, dry place. Reduce stacking height where local conditions, such as humidity or pallet overhang can affect package strength.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal¹: Completely empty contents of bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Container Disposal²: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.

NOTICE: Apply this product in a manner to avoid drift. Extreme care must be used when applying this product to prevent injury to desirable plants. This product is a non-selective herbicide and can injure or destroy all vegetation contacted. Avoid spray contact with the foliage of desirable turf, ornamentals, shrubs or trees. Avoid spray contact with green bark or exposed non-woody roots of desirable trees and shrubs. If spray contacts desirable plants, they should be rinsed immediately with water. Do not apply during gusty or windy conditions. Use pressure and nozzle types that will not result in splatter or fine particles (mist) which are likely to drift.

General Information: This product mixes readily with water and should be used as a foliar spray to control listed plants. This product may be applied through equipment providing a broadcast spray or hand-held and high-volume spray equipment such as knapsack and backpack sprayers, pump-up pressure sprayers, handguns, handwands and other hand-held and motorized spray equipment to direct spray onto weed foliage. Refer to the appropriate sections below for specific equipment use instructions and rates of application.

GLYGRAN WDG is recommended for noncrop or nonplanted areas such as right-ofways, including railroads, dividers and medians; pipelines, public utility lines, including pumping stations, transformer stations and substations; around electric yards, fence lines and parkways. Also, around ornamental plantings and trees,

GLYGRAN WDG\ revised 10-27-05 Page 2 of 2

¹ For product packaged in plastic bags

² For product packaged in plastic containers/jugs/pails

walkways, patios; along driveways and around golf courses; parking areas, industrial sites, parks, playgrounds, petroleum farms, fence posts, schools, signposts, and storage areas.

Apply this product when plants are actively growing and sufficient foliage is available. Unless otherwise directed, allow 7 days or more after application before tillage, mowing or removal of treated weeds. Watering or irrigation should be delayed for 24 to 48 hours following treatment. Repeat treatments may be necessary to control plants regenerating from underground plant parts or seed. Application to undesirable deciduous species containing some autumn color development is acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatment is made following a severe frost.

GLYGRAN WDG is a non-selective herbicide that, upon foliar contact, is translocated throughout the plant system into the roots. Visible effects on most annual weeds will be observed within 2 to 4 days. For perennial weeds, visible effects may not be observed for 7 days or more. Cool or overcast conditions following application may slow activity of this product, resulting in a delay of visible effects. Visible effects include wilting and yellowing or browning of above ground plant growth and detorioration of underground plant parts.

Rainfall or irrigation shortly after application may reduce the effectiveness of this product and a repeat application may be needed for desirable control.

This product does not provide residual weed control. For extended weed control, reapplications of this product will be necessary or tank mixture use of a residual herbicide will be needed such as, Oust XP™, Karmex IWC™, Krovar I DF™, Arsenal™*, Escort XP™*, and Oust Extra™* †. (*not in California)

PREPARATION OF SPRAY SOLUTION

Broadcast Applications: When applied as directed and under conditions described, GLYGRAN WDG will control weeds listed below. Apply in sufficient volume of water to obtain uniform coverage. See the "Weeds Controlled" section of this label for specific rates. As density of weed growth increases, spray volume should be increased within the recommended range to ensure adequate coverage. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

To prepare the spray solution, completely fill the mixing or spray tank with the required amount of water and start agitation. Add the recommended amount of this product and surfactant if used, and continue agitation until product is completely dissolved and or suspended. If foaming occurs, use an approved anti-foam or defoaming agent. Continue agitation during application.

Hand-Held and High-Volume Equipment Applications: Fill spray tank with clean water and add the recommended amount GLYGRAN WDG. Agitate the mixture until product is thoroughly mixed. Apply on a spray-to-wet basis. Spray coverage should be uniform and complete. Do not spray to the point of run-off.

For control of annual weeds listed on this label, apply 1/2 oz. GLYGRAN WDG per gallon of desired spray volume for weeds under 6 inches in height. Apply prior to seedhead formation in grass or bud formation in broadleaf weeds. For

annual weeds over 6 inches tall, apply 2/3 oz. GLYGRAN WDG per gallon of desired spray volume.

For control of perennials, woody brush and trees, apply 1 oz. GLYGRAN WDG per gallon of desired spray volume. Agriculturally approved colorants and marking dyes may be added to this product as desired. Follow manufacturer's recommendations when using colorants or dyes.

Nonlonic surfactants which are labeled for use with herbicides may be used with this product to increase effectiveness for hard to control weeds. Follow manufacturers labeled recommendations for appropriate use rates. Do not reduce the rate of this product when adding additional surfactant.

Equipment Maintenance and Cleaning: Triple-rinse sprayer and flush all sprayer components with water to remove residues of this product. Check sprayer nozzles and seals prior to and after each use to remove foreign debris and confirm integrity of seals. After thorough cleaning, spray equipment may be used to apply other chemical products.

Commercial/Residential Site Treatment: This product is inactive in the soil and following removal of weeds, any ornamental species, bedding plant, turfgrass, tree or shrub species may be planted. Where treated vegetation will not be disturbed, planting may be done 1 day following treatment.

Lawn/Turf Area Renovation: Seedbed preparation and seeding, sodding or sprigging of turfgrass species may take place 7 days following treatment of undesirable weeds or turfgrass. Where treated vegetation will not be disturbed by seedbed preparation (tillage, aeration, power raking, etc.), seeding may take place 1 day following treatment.

Post-Directed Treatments: This product can be used for spot-treatment of weeds and undesirable vegetation existing in shrub beds, around ornamental plantings, and ground covers.

Avoid spray contact with desirable plants. To avoid injury on desirable plants, avoid spray or drift contact with foliage, stems, bark or exposed non-woody roots. Do not spray into, or over the top of, desirable plants, trees or shrubs.

Edge Treatments: GLYGRAN WDG can be used to control weeds for trimming and edging around many structures, including buildings, sidewalks, roadways, park benches, golf course markers, trees, light posts, signs, fence posts, patios, fence rows, gravel and bark mulch beds, etc.

WEEDS CONTROLLED

Annual Weeds:

Apply to actively growing annual grass and broadleaf weeds. Allow at least 3 days following treatment before disturbing treated vegetation. After 3 days, treated weeds and grass may be removed.

For broadcast applications, use 0.9 lbs. GLYGRAN WDG per acre on weeds smaller

[†] Oust XP™, Karmex IWC™, Krovar I DF™, Escort XP™*, and Oust Extra™* are registered trademarks of E.I. DuPont. Arsenal™ is a registered trademark of BASF Ltd.

than 6"-and not growing under stressed conditions. Use higher rates from 1.35 to 3.75 lbs. per acre on weeds larger than 6" or if weeds are growing under stressed conditions.

When applied as directed, this product will control the following annual weeds:

Foxtails

Barley: little, wild Barnvardgrass Bassia, fivehook Bedstraw, catchweed Blueorass, annual Bluegrass, bulbous Bromes Buttercup Cheat Chervil Chickweed: jagged. mouseear Clovers Cocklebur Crabgrass Dwarfdandelion Falseflax, smallseed Fiddleneck Flaxleaf fleabane Foxtail, giant

Geranium, Carolina Goosegrass Groundsel, common Hemp sesbania Henbit Horseweed/Marestail Johnsongrass, seedling Kochia Lambsquarters, common Lettuce, prickly Momingolory (ipomoea spp.) Mustards Oats Panicum Pennycress, field Pigweeds Purslane, common Ragweed, common

Ragweed, glant Rye Ryegrass, Italian Sandbur, field Shattercane Shepherdspurse Signalgrass, broadleaf Smartweed, Pennsylvania Sowthistle, annual Spanishneedles Speedwell, com Spurry, umbrella Stinkorasa Sunflower Thistle, Russian Velvetleaf Wheat Witchorass

When applied as directed, this product will control the following biennial/perennial weeds:

| | Rate - Ibs./acre | | Rate - Ibs./acre |
|---------------------|------------------|--------------------------|------------------|
| Alfalfa* | 0.9 | Knapweed | 3.6 |
| Alligator weed* | 3.6 | Kudzu | 2.7 - 3.6 |
| Bahiagrass | 2.7 - 4.5 | Lantana | 2.7 - 3.6 |
| Bentgrass* | 1.35 | Lespedeza | 2.7 - 4.5 |
| Bermudagrass | 4.5 | Milkweed | 2.7 - 3.6 |
| Bindweed, field | 3.6 - 4.5 | Muhly, wirestern | 1.8 |
| Bluegrass, Kentucky | 1.8 | Mullein, common | 2.7 |
| Bluestern, silver | 0.9 - 2.7 | Naplergrass | 2.7 - 4.5 |
| Blueweed, Texas | 3.6 - 4.5 | Nightshade, silver leaf | 1.8 |
| Brackenfern | 2.7 - 3.6 | Nutsedge: purple, yellow | 2.7 |
| Bromegrass, smooth | 1.8 | Orchardgrass | 1.8 |
| Broomsedge | 1.8 - 2.7 | Pampasgrass | 2.7 - 4.5 |
| Canarygrass, reed | 1.8 - 2.7 | Paragrass | 2.7 - 4.5 |
| Cattail | 2.7 - 4.5 | Phragmites* | 2.7 - 4.5 |
| Clovers | 2.7 - 4.5 | Puncturevine | 0.9 - 2.7 |
| Cogongrass | 2.7 - 4.5 | Quackgrass | 1.8 - 2.7 |
| Dallisgrass | 2.7 - 4.5 | Raspberry | 3.6 - 4.5 |
| Dandelion | 2.7 - 4.5 | Redvine* | 1.8 |
| Dewberry | 0.9 - 2.7 | Reed, giant | 3.6 - 4.5 |
| Dock, curty | 2.7 - 4.5 | Ryegrass, perennial | 1.8 - 2.7 |
| Dogbane, hemp | 3.6 | Smartweed, swamp | 2.7 - 4.5 |
| Fescues | 0.9 - 2.7 | Starthistle, yellow | 1.8 - 2.7 |
| Fescue, tall | 0.9 - 2.7 | Thistle, artichoke | 1.8 - 2.7 |
| Guineagrass | 2.7 | Thistle, Canada | 1.8 - 2.7 |
| Honeysuckle | 2.7 - 3.6 | Timothy | 1.8 - 2.7 |
| Horsenettle | 2.7 - 4.5 | Torpedograss* | 3.6 - 4.5 |
| Horseradish | 3.6 | Trumpet creeper* | 3.6 - 4.5 |
| Ice Plant | 1.8 | Vaseygrass | 2.7 - 4.5 |
| Johnsongrass | 1.8 - 2.7 | Velvetgrass | 2.7 - 4.5 |
| Kikuyugrass | 1.8 - 2.7 | Wheatgrass, western | 1.8 - 2.7 |

Biennial/Perennial Weeds:

Apply to actively growing weeds. Allow at least 7 days following treatment before disturbing treated vegetation. After 7 days, treated weeds may be mowed, tilled or burned. Re-treatments may be needed to control weeds regenerating from underground plant parts or seeds.

For broadcast applications, apply at the rate indicated below. Use the higher recommended rate for larger plants and/or dense areas of growth.

*Partial control

Woody Brush and Trees:

For broadcast applications, apply at the rate indicated below. Use higher recommended rates for larger plants and/or dense growth. Ensure thorough coverage when using spray-to-wet treatments.

When applied as directed, this product will control or partially control the following woody brush, plants and trees:

| | Rate - Ibs./acre | | Rate - Ibs./acre |
|----------------------------|------------------|-----------------------|------------------|
| Alder | 3.6 - 3.9 | Oak, southern red | 2.7 - 3.6 |
| Ash | 1.8 - 4.5 | Oak, white* | 2.7 - 4.5 |
| Aspen, guaking | 1.8 - 2.7 | Persimmon* | 1.8 - 4.5 |
| Bearmat (Bearclover)* | 1.8 - 4.5 | Poison Ivy | 3.6 - 4.5 |
| Birch | 1.8 | Polson Oak | 3.6 - 4.5 |
| Blackberry | 2.7 - 3.6 | Poplar, yellow* | 1.8 - 4.5 |
| Cascara* | 1.8 - 4.5 | Redbud, eastern | 2.7 - 4,5 |
| Ceanothus* | 1.8 - 4.5 | Rose, multiflora | 1.8 |
| Chamise* | 1.8 - 4.5 | Russian-olive | 1.8 - 4.5 |
| Cherry: bitter, black, pin | 1.8 - 2.7 | Sage, Black | 2.7 - 4.5 |
| Elderberry | 1.8 | Sagebrush, California | 2.7 - 4.5 |
| Elm* | 1.8 - 4.5 | Salmonberry | 1.8 |
| Hawthorn | 1.8 - 2.7 | Sassafras* | 1.8 - 4.5 |
| Eucalyptus | 3.6 - 4.5 | Sourwood* | 1.8 - 4.5 |
| Hazel | 1.8 | Sumac* | 1.8 - 3.6 |
| Locust, Black* | 1.8 - 3.6 | Sweetgum | 1.8 - 2.7 |
| Manzanita | 1.8 - 4.5 | Swordfern* | 1.8 - 4.5 |
| Oak | 1.8 - 3.6 | Sycamore | 2.7 - 4.5 |
| Oak, northern | 3.6 - 4.5 | Thimbleberry | 1.8 |
| Oak, pin | 3.6 - 4.5 | Waxmyrtle, southern* | 1.8 - 4.5 |
| Oak, post | 4.5 | Willow | 2.7 |

Partial Control

Time of Application

For best results, in arid climates make treatments in the spring or early summer when brush species are at a high moisture content. In areas of spring and summer rainfall, best results are obtained when application is made in late summer or early fall after fruit formation.

Fallow Systems:

This product may be applied during the fallow period prior to planting.

TYPES OF APPLICATIONS: Chemical Failow, Preplant Failow Beds, Aid-to-Tillage.

Chemical Fallow:

USE INSTRUCTIONS: This product may be applied during the fallow period prior to planting. This product may be used as a substitute for tillage to control annual weeds in fallow fields. Also, broadcast or spot treatments will control or suppress many perennial weeds in fallow fields. Ground or aerial application equipment may be used. Tank mixtures with 2,4-D and dicamba may be used. Applications up to 1.9 pounds per acre may be made by aerial application in fallow sites where there is sufficient buffer to prevent injury due to drift onto adjacent crops.

PRECAUTIONS, RESTRICTIONS: Applications must be made at least 30 days prior to planting. Do not apply dicamba mixtures by air in California.

Refer to specific product labels for crop rotation restrictions and cautionary statements of all products used in tank mixtures. Some crop injury may occur if dicamba is applied within 45 days of planting.

Preplant Fallow Beds

USE INSTRUCTIONS: This product must be applied at least 30 days prior to planting when being applied to fallow beds. This product will control weeds listed in the "ANNUAL WEEDS", "PERENNIAL WEEDS", and "WOOD BRUSH AND TREES RATE TABLES" sections of this label.

TANK MIXTURES: In addition, 7 ounces of this product plus 2 to 3 fluid ounces of Goal™ 2XL per acre will control the following weeds with the maximum height or length indicated: 3 inches – common cheeseweed, chickweed, groundsel; 6 inches – London rocket, shepherd's purse.

9.5 ounces of this product plus 2 to 3 fluid ounces of Goal™ 2XL per acre will control the following weeds with the maximum height or length indicated: 6 inches – common cheeseweed, groundsel, marestail (*Conyza Canadensis*); 12 inches – chickweed, London rocket, shepherd's purse.

Aid-to-Tillage

USE INSTRUCTIONS: This product may be used in conjunction with tillage practices in fallow systems to control downy brome, cheat, volunteer wheat, tansy mustard and foxtail. Apply 7 ounces of this product in 3 to 10 gallons of water per acre. Make applications before weeds are 6 inches in height. Application must be followed by conventional tillage practices no later than 15 days after treatment and before regrowth occurs. Allow at least 1 day after application before tillage.

PRECAUTIONS, RESTRICTIONS: Tank mixtures with residual herbicides may result in reduced performance.

• •

CONDITION OF SALE

The label instructions and directions for use on this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, soll composition, type of vegetation, presence of other materials, or the manner of use or application, all of which are beyond the control of AGRIVENTURES, INC. [AGRIVENTURES].

ALL SUCH RISKS ARE ASSUMED BY THE BUYER.

LIMITED WARRANTY AND DISCLAIMER

AGRIVENTURES warrants that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use. AGRIVENUTRES makes NO OTHER warranty, expressed or implied, concerning the use of this product other than as indicated on the label. Buyer assumes all risk of use and/or handling that is contrary to label instructions.

The warranty does not extend to use of this product contrary to the label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to AGRIVENTURES, and the buyer assumes the risk of any such use.

READ THE LABEL BEFORE USING

Neither AGRIVENTURES nor any selfer of this product makes, and their employees and agents are not permitted to make, any representations or warranties, express or implied by law, not specifically set forth herein.

Buyer assumes the risk to persons or property arising from the use of handling this product not in strict accordance with the directions given herewith, and ACCEPTS THE PRODUCT ON THAT CONDITION.

USES WITH OTHER PRODUCTS (TANK MIXES)

Glygran WDG does not provide residual vegetation control. Weeds germinating from seed after treatment will not be controlled. For residual vegetation control, follow a label approved herbicide program. When tank mixing Glygran WDG with other herbicides, all applicable directions, restrictions, and precautions for the additional herbicides must be followed. Use the most restrictive limitations stated on the product labels. Before initiating full scale tank mix application, the physical compatibility of the proposed mixture should be evaluated on a small scale of recommended spray mixture concentrations. Also, such mixtures should be evaluated for vegetation control before expanding commercial use.

LIMITATIONS OF LIABILITY AND DAMAGES

The liability of AGRIVENTURES, and any other seller for any loss or injury, wheth based on negligence, breach of warranty, strict liability in tort, or any other cause action, shall be limited to the purchase price of the product. Prompt notice of a claim must be given by the buyer to AGRIVENTURES within ninety days after th application of the product or ninety days after the first planting, whichever is late Notification to AGRIVENTURES, should be at this address: 9775 East Balancia Rock Rd., Scottsdale, AZ 85262.

AGRIVENTURES and any other seller of the product shall not be liable, and any a all claims against AGRIVENTURES, and any other seller are waived, for speci indirect, incidental, or consequential damages or expenses of any nature, includir but not limited to, loss of profits or income, and crop or property loss or dama resulting from use of the product not in strict accordance with the directions, wheth or not based on negligence, breach of warranty, strict liability in tort or any oth cause of action.

AGRIVENTURES OFFERS THIS PRODUCT AND THE BUYER AND USE ACCEPT IT SUBJECT TO THE FOREGOING CONDITIONS OF SALE AN LIMITATIONS OR WARRANTY, LIABILITY AND DAMAGES, WHICH MAY E VARIED ONLY BY AGREEMENT IN WRITING SIGNED BY A DULY AUTHORIZE REPRESENTATIVE OF AGRIVENTURES.



122 C Street, N.W., Suite 740 Washington, D.C. 20001 telephone 202.393.3903 fax 202.393.3906

Consultants In Government Affairs

October 28, 2005

Office of Pesticide Programs Document Processing Desk [NOTIF] U. S. Environmental Protection Agency (7504C) Room 266A, Crystal Mall 2 1921 Jefferson Davis Highway Arlington, VA 22202

ATTENTION: Jim Tompkins Product Manager, Team 25

SUBJECT: Agriventures, Inc. GLYGRAN® WDG (EPA Reg. No. 82493-2) Notification per PR Notice 98-10

Dear Mr. Tompkins:

On behalf of Agriventures, Inc., I am submitting a notification pursuant to PR Notice 98-10 for their product Glygran® WDG (EPA Reg. No. 82493-2). Below I have listed the purpose of this notification:

- 1) Addition of the statement "other non-cop uses" in the Non-Agricultural Use Requirements section. Please note this addition is consistent with the labeling.
- 2) Revised the "Preparation of Spray Solution" (mixing) directions on page 3 in order to clarify that applicators thoroughly mix the solution.
- 3) Revised the DuPont residual herbicide product trade names listed on page 3.
- 4) Removed the partial control language for the annual weeds listed on page 4. Please note this was a typographical error on the original label.

In support of this Notification, enclosed please find the following documents:

- 1. Application for Pesticide Notification (OPP Id. No. 265394);
- 2. One (1) copy of the master label with changes highlighted; and,
- 3. Two (2) copies of the master label

Insofar as Lewis & Harrison serves as the "Company Contact" and "Company Agent" for Agriventures, Inc. please relay all correspondence regarding this submission directly to us. If you have any questions, please contact me at (202) 393-3903 x.19 or e-mail me at georgia@lewisharrison.com.

I thank you in advance for your cooperation.

Sincerely,

Georgia Anastasiou

Agent for Agriventures, Inc.

Enclosures

cc: Nina Reynolds (Agriventures, Inc.)