

82542-31

06/14/2011

1/27



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Chemical Safety and Pollution Prevention
 Office of Pesticide Programs
 Registration Division (7504P)
 1200 Pennsylvania Ave., N.W.
 Washington, DC 20460

EPA Reg. Number:
82542-31

Date of Issuance:
JUN 14 2011

Term of Issuance:
 Unconditional

Name of Pesticide Product:
Tebuconazole 3.6F Ag Fungicide

NOTICE OF PESTICIDE:

Registration
 Reregistration
 (under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):
Source Dynamics
10039 E. Troon North Drive
Scottsdale, AZ 85262

Mailed to:
 Robert Hawk
 Registration Manager

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA sec 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

Signature of Approving Official:

 Mary L. Waller, Product Manager (21)
 Fungicide Branch/Registration Division/OPP/OCSP (7504P)

Date:
 6/14/2011

2. Make the following changes to the label:
 - a. Change the product registration number to "EPA Reg. No. 82542-31"
 - b. On page 12 in the first column of the table for **BULB VEGETABLES** change "Welch onion" to "Welsh onion"
 - c. On page 19 change the header "**Animal Feeding and Grazing Directions**" to "**Animal Feeding and Grazing Restrictions**".
 - d. On page 23 change the header "**Animal Feeding and Grazing Directions**" to "**Animal Feeding and Grazing Restrictions**".
3. Submit one copy of the revised final printed label for the record before the product is released for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A copy of the label stamped "Accepted with Comments" is enclosed for your records.



Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7504P)

Enclosure:
Label stamped "Accepted with Comments"

Source Dynamics LLC proposed label; revised 5/2/2011

TEBUCONAZOLE 3.6F Ag Fungicide

For control of listed diseases on asparagus, barley, beans, corn, cotton, cucurbit vegetables, garlic, grasses grown for seed, hops, leafy Brassica greens, garden beets, lychee, okra, onion, peanuts, pecan, soybeans, sunflower, turnip and wheat.

ACTIVE INGREDIENT:

tebuconazole, α -[2-(4-chlorophenyl)ethyl]- α -(1,1-dimethylethyl)-

1H-1,2,4-triazole-1-ethanol 38.7%

OTHER INGREDIENTS: 61.3%

TOTAL..... 100.0%

Contains 3.6 pounds tebuconazole per gallon

KEEP OUT OF REACH OF CHILDREN

CAUTION

| FIRST AID | |
|---|--|
| If swallowed | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person. |
| If on skin or clothing | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15- 20 minutes. • Call a poison control center or doctor for treatment advice. |
| If in eyes | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| If inhaled | <ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice. |
| Note to Physician: No specific antidote. Treat symptomatically. The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation. | |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For Emergency Medical Assistance, call the National Pesticide Information Center 1-800-858-7378. | |

For chemical emergency: spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300.

Manufactured for:
Source Dynamics LLC

EPA Registration No. 82542-GR

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

6/14/2011

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 82542-31

10039 E. Troon North Drive
Scottsdale, AZ 85262
tel. 480-502-9289

Est. No. _____
Net Contents 2.5 Gal

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or Viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean

high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory: Tebuconazole is known to leach through soil into ground water under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) specified in the application directions for the treated crop.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Do not store this product near fertilizers, seeds or other pesticides. Store in original containers only. Store in a cool, dry, locked facility and avoid excess heat. Carefully open containers. Reclose all partially used containers by thoroughly tightening screw cap. Do not put concentrate or dilute material in food or drink containers. Keep containers closed when not in use.

In case of spill, confine spill by surrounding area with sand, cat litter or commercial clay and dispose as directed below.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full of water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

PRODUCT INFORMATION

Read the entire Directions for Use and Conditions of Sale before using this product.


Chemigation: Do not apply this product through any type of irrigation system.

For the most effective results, equipment calibration should be checked regularly. When using lower spray volumes, be sure to maintain uniform application and full crop coverage so as to ensure effective control. Increase spray volume to ensure proper application, if required.

NOTE: FOLLOW THESE DIRECTIONS WHEN MAKING APPLICATIONS NEAR AQUATIC AREAS (ESTUARIES, LAKES, MARSHES, NATURAL PONDS, PERMANENT STREAMS, RESERVOIRS AND RIVERS).

See Aerial Drift Reduction Advisory Sections in following pages.

- Ground and aerial application with 100 feet of aquatic areas listed above is prohibited.
- Application to fields next to aquatic areas may only be made every other year.
- To prevent unwanted exposure to bodies of water maintain a 10 foot wide non-cultivated vegetative strip filter.
- See Spray Drift Management section for further information.



Mixing: Continuous agitation is required during mixing. When mixing this product and water, add the labeled amount of Tebuconazole 3.6F Ag Fungicide. Before combining any other substances with the mixture, ensure that the Tebuconazole 3.6F Ag Fungicide is complete dispersed in the mixture.

Compatibility Test for Mix Components:

Before mixing components, always perform a compatibility jar test. For 20 gallons per acre spray volume, use 3.3 cups (800 ml) of water in a clear, clean mixing jar. For other spray volumes adjust accordingly. Only use water from the intended source at the source temperature. Add components in the sequence indicated below in Mixing Order using 2 teaspoons for each pound of dry product or 1 teaspoon for each pint of liquid product of recommended label rate per acre. Always cap the jar and invert 10 cycles between component additions.

When the components have all been added to the jar and fully mixed, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent and use the compatibility agent as directed on its label.

Mixing Order:

1. Water. Begin by agitating a thoroughly clean sprayer tank three-quarters full of clean water.
2. Agitation. Maintain constant agitation throughout mixing and application.
3. Inductor. If an inductor is used, rinse it thoroughly after each component has been added.
4. Products in PVA Bags. Place any product contained in water soluble PVA bags into the mixing tank. Wait until all water soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5. Water Dispersible Products. Including dry flowables (DF), wettable powders (WP), suspension concentrates (SC) or suspo-emulsions (SE).
6. Water-soluble products.
7. Emulsifiable concentrates (such as oil concentrates when applicable).
8. Water soluble additives (such as AMS or UAN when applicable).
9. Remaining quantity of water.

Maintain constant agitation during application.

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, and relative humidity) and method of application can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Apply only as a medium or coarser spray (ASAE Standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

For aerial applications, the boom should be mounted on the aircraft so as to minimize

drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter.

Use the largest droplet size consistent with pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure. Apply in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment.

Spray should be released at the lowest possible height consistent with good pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided. When applications are made with a crosswind, the swath must be displaced downward. The applicator must compensate for this displacement at the up and downwind edge of the application area by adjusting the path of the aircraft upwind.

Make aerial or ground application when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperatures.

Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

ROTATIONAL CROPS:

Treated areas may be replanted with any crop specified on this label as soon as practical after last application. Any crop not specified on this label may be planted into treated areas 120 days after last application.

ASPARAGUS

APPLICATION DIRECTIONS

| CROP | TARGET DISEASES | PER ACRE PRODUCT USE RATE PER APPLICATION | FOLLOW-UP APPLICATION TIMING | MAXIMUM USE RATE OF PRODUCT/ACRE/CROP PER SEASON | PHI |
|-----------|------------------------------|---|------------------------------|--|---|
| Asparagus | Rust (<i>Puccinia</i> spp.) | 4 – 6 fl. oz. | 14 days | 18 fl oz. | 100 days in California 180 days in all other states. |

Application Directions

Apply Tebuconazole 3.6F Ag Fungicide as a foliar spray to the developing ferns after

harvest of spears is completed. For optimum control apply at the earliest sign of rust pustules or when weather conditions are conducive for rust development. Under conditions of severe rust pressure, use the higher rate.

Apply in alternation with another effective fungicide. Tebuconazole 3.6F Ag Fungicide is a sterol demethylation inhibitor (DMI) fungicide (Group 3). Alternating Tebuconazole 3.6F Ag Fungicide with other DMI fungicides may lead to resistance.

Tebuconazole 3.6F Ag Fungicide must have 2 - 4 hours of drying time on the plant foliage for the active ingredient to move systematically in to the plant tissue. After 2 - 4 hours the Tebuconazole 3.6F Ag Fungicide will be resistant to weathering.

Spray Volume:

Ground Application

Apply Tebuconazole 3.6F Ag Fungicide in no less than 10 gallons of spray solution per acre.

Aerial Application

Apply Tebuconazole 3.6F Ag Fungicide in no less than 5 gallons of spray solution per acre. A 50 foot spray drift buffer zone is required for all aerial applications.

Do not make more than 3 applications per season (18 fl. oz./acre or 0.51 lb. a.i./acre). Do not apply to harvestable spears.

Restricted-entry interval (REI) = 12 hours.

BARLEY

APPLICATION DIRECTIONS

| CROP | TARGET DISEASES | PER ACRE PRODUCT USE RATE PER APPLICATION | FOLLOW-UP APPLICATION TIMING | MAXIMUM USE RATE OF PRODUCT/ACRE/CROP PER SEASON | PHI |
|--------|---|---|------------------------------|--|---------|
| Barley | Rust (<i>Puccinia</i> spp.) | 4 fluid ounces | Not Allowed | 4 fluid ounces | 30 days |
| | Head Blight (<i>Fusarium</i> spp.)- Supression | | | | |

Application Directions

For optimum control, barley fields should be kept under observation for early disease symptoms. This is particularly important when conditions favoring disease development are favorable or when varieties susceptible to disease are planted.

For Rusts, apply the Tebuconazole 3.6F Ag Fungicide at the earliest sign of rust pustules on foliage.

For Fusarium head blight apply Tebuconazole 3.6F Ag Fungicide when the main stem heads have fully emerged (Feekes 10.5) on 50% of the plants for optimum suppression.

For optimum disease control, the lowest labeled rate of a spray surfactant may be tank mixed with Tebuconazole 3.6F Ag Fungicide.

Tebuconazole 3.6F Ag Fungicide must have 2 - 4 hours of drying time on the plant foliage for the active ingredient to move systematically in to the plant tissue. After 2 - 4 hours the Tebuconazole 3.6F Ag Fungicide will be resistant to weathering.

Ground Application

Apply Tebuconazole 3.6F Ag Fungicide in no less than 10 gallons of spray solution per acre.

Aerial Application

Apply Tebuconazole 3.6F Ag Fungicide in no less than 5 gallons of spray solution per acre.

Animal Feeding and Grazing Directions:

Following application of this product, do not permit animals to graze or forage in the treated areas for at least 6 days. Straw cut after harvest may be used for feed or bedding.

Restricted-entry interval (REI) = 12 hours.

BEANS

APPLICATION DIRECTIONS


| CROP | TARGET DISEASES | PER ACRE PRODUCT USE RATE PER APPLICATION | FOLLOW-UP APPLICATION TIMING | MAXIMUM USE RATE OF PRODUCT/ACRE/CROP PER SEASON | PHI |
|--|---|---|------------------------------|--|---------|
| Beans (fresh & dry except succulent shelled. | Rust (<i>Uromyces appendiculatus</i>) | 4 - 6 fl. oz. | 14 days | Beans, fresh: 24 fl. oz. | 7 days |
| | | | | Beans, dry: 12 fl. oz. | 14 days |

Application Directions

Apply Tebuconazole 3.6F Ag Fungicide in a protective spray schedule or when weather conditions are favorable for rust development.

For optimum disease control, the lowest labeled rate of a spray surfactant may be tank mixed with Tebuconazole 3.6F Ag Fungicide .

Tebuconazole 3.6F Ag Fungicide must have 2 - 4 hours of drying time on bean foliage



for the active ingredient to move systematically in to the plant tissue before rain or irrigation occurs. After 2 - 4 hours the Tebuconazole 3.6F Ag Fungicide will be resistant to weathering.

Restricted-entry interval (REI) = 12 hours.

CORN

APPLICATION DIRECTIONS

| CROP | TARGET DISEASES | PER ACRE PRODUCT USE RATE PER APPLICATION | FOLLOW-UP APPLICATION TIMING | MAXIMUM USE RATE OF PRODUCT/ACRE/ CROP PER SEASON | PHI |
|---|---|---|------------------------------|---|--|
| Corn (sweet corn, field corn, field corn grown for seed, and popcorn) | Rust (<i>Puccinia</i> spp.) | 4 - 6 fl. oz. | 7 - 14 days | 24 fl oz. | Sweet Corn: 7 days before harvest of ears or forage, 49 days before harvest of fodder. Field, seed or popcorn: 21 days before harvest of forage, 36 days before harvest of grain or fodder. |
| | Northern leaf blight (<i>Helminthosporium turcicum</i>) | | | | |
| | Southern leaf blight (<i>Helminthosporium maydis</i>) | | | | |
| | Northern leaf spot (<i>Helminthosporium carbonum</i>) | | | | |
| | Gray leaf spot (<i>Cercospora zeae-maydis</i>) | | | | |

Application Directions


Apply Tebuconazole 3.6F Ag Fungicide in a protective spray schedule or when weather conditions are favorable for disease development.

For optimum disease control, the lowest labeled rate of a spray surfactant may be tank mixed with Tebuconazole 3.6F Ag Fungicide.

Tebuconazole 3.6F Ag Fungicide must have 2 - 4 hours of drying time on bean foliage for the active ingredient to move systematically in to the plant tissue before rain or irrigation occurs. After 2 - 4 hours the Tebuconazole 3.6F Ag Fungicide will be resistant to weathering.

Restricted-entry interval (REI) for sweet corn = 19 days.

Restricted-entry interval (REI) for all corn except sweet corn = 12 hours.



COTTON

APPLICATION DIRECTIONS

| CROP | TARGET DISEASES | PER ACRE PRODUCT USE RATE PER APPLICATION | FOLLOW-UP APPLICATION TIMING | MAXIMUM USE RATE OF PRODUCT/ACRE/CROP PER SEASON | PHI |
|-------------|---|--|-------------------------------------|---|------------|
| Cotton | Southwestern cotton rust (<i>Puccinia cacabata</i>) | 6 - 8 fl. oz. | 7 - 14 days | 24 fl. oz. | 30 days |


Application Directions

Apply Tebuconazole 3.6F Ag Fungicide in a protective spray schedule or when weather conditions are favorable for rust development.

For optimum disease control, the lowest labeled rate of a spray surfactant may be tank mixed with Tebuconazole 3.6F Ag Fungicide.

Tebuconazole 3.6F Ag Fungicide must have 2 - 4 hours of drying time on cotton foliage for the active ingredient to move systematically in to the plant tissue before rain or irrigation occurs. After 2 - 4 hours the Tebuconazole 3.6F Ag Fungicide will be resistant to weathering.

Restricted-entry interval (REI) = 12 hours.



CUCURBIT VEGETABLES

APPLICATION DIRECTIONS

| CROP | TARGET DISEASES | PER ACRE PRODUCT USE RATE PER APPLICATION | FOLLOW-UP APPLICATION TIMING | MAXIMUM USE RATE OF PRODUCT/ACRE CROP PER SEASON | PHI |
|--|--|---|------------------------------|--|--------|
| Cucurbit Vegetables Group Chayote Chinese waxgourd Citron melon Cucumber Gherkin Edible gourd (includes hyotan, cucuzza, hechima and Chinese okra) Momordica spp. (includes balsam apple, balsam pear, bitter melon and Chinese cucumber) Muskmelon (includes cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon and snake melon) Pumpkin Summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow and zucchini) Winter squash (includes butternut squash, calabaza, hubbard squash, acorn squash and spaghetti squash) Watermelon | Powdery mildew (<i>Sphaerotheca fuliginea</i> / <i>Podosphaera xanthii</i>) (<i>Erysiphe cichoracearum</i>) | 4 – 6 fl. oz.. | 10 – 14 days | 24 fl. oz. | 7 days |
| | Gummy stem blight – suppression (<i>Didymella bryonae</i>) (watermelon, squash, pumpkin and melons only) | 8 fl. oz. | | | |

Application Directions

Apply Tebuconazole 3.6F Ag Fungicide to foliage and fruit in a protective spray

schedule.

For optimum disease control, the lowest labeled rate of a spray surfactant may be tank mixed with Tebuconazole 3.6F Ag Fungicide.

Tebuconazole 3.6F Ag Fungicide must have 2 - 4 hours of drying time on cotton foliage for the active ingredient to move systematically in to the plant tissue before rain or irrigation occurs. After 2 - 4 hours the Tebuconazole 3.6F Ag Fungicide will be resistant to weathering.

Restricted-entry interval (REI) = 12 hours.

BULB VEGETABLES

APPLICATION DIRECTIONS

| CROP | TARGET DISEASES | PER ACRE PRODUCT USE RATE PER APPLICATION | FOLLOW-UP APPLICATION TIMING | MAXIMUM USE RATE OF PRODUCT/ACRE/CROP PER SEASON | PHI |
|--|--|---|---|--|--------|
| Dry bulb onion Garlic Great-headed (elephant) garlic Welch onion Shallot | White rot (<i>Sclerotium cepivorum</i>) | 20.5 fl oz in a 4 to 6 inch band over/into each furrow at the time of planting. | Two foliar applications at 4 - 6 fl oz/acre may be used to obtain additional control. | 32.5 fl oz. for in-furrow treatment. | 7 days |
| | Rust (<i>Puccinia allii</i> , <i>Puccinia porri</i>) | 4 - 6 fl. oz. | 10 - 14 days | 12 fl oz. as a foliar spray | |
| | Purple blotch (<i>Alternaria porii</i>) | | | | |

Application Directions

Apply Tebuconazole 3.6F Ag Fungicide as a preventative treatment for optimum results. Begin applications as soon as crop and/or environmental conditions become favorable for disease development.

For optimum disease control, the lowest labeled rate of a spray surfactant may be tank mixed with Tebuconazole 3.6F Ag Fungicide.

Tebuconazole 3.6F Ag Fungicide must have 2 - 4 hours of drying time on plant foliage for the active ingredient to move systematically in to the plant tissue before rain or irrigation occurs. After 2 - 4 hours the Tebuconazole 3.6F Ag Fungicide will be resistant to weathering.

Restricted-entry interval (REI) = 12 hours.

GRASSES GROWN FOR SEED

APPLICATION DIRECTIONS

| CROP | TARGET DISEASES | PER ACRE PRODUCT USE RATE PER APPLICATION | FOLLOW-UP APPLICATION TIMING | MAXIMUM USE RATE OF PRODUCT/ACRE/CROP PER SEASON | PHI |
|------------------------|---|---|------------------------------|--|--------|
| Grasses grown for seed | Rust (<i>Puccinia</i> spp.) | 4 – 8 fluid ounces | 14 – 16 days | 16 fluid ounces | 4 days |
| | Powdery Mildew (<i>Erysiphe graminis</i>) | 4 – 8 fluid ounces | | | |

Application Directions:

For optimal disease control, begin applications of Tebuconazole 3.6F Ag Fungicide prior to disease development, as favorable weather conditions for disease development are noted in the crop area.

Use the higher rate and shorter intervals when disease pressure is high or if disease is present prior to fungicide application. For best results, use a minimum rate of recommended spray adjuvant when mixing this product for application. Uniform and complete distribution of applied spray is critical for best disease control.

Ground Application:

Use specified rate of Tebuconazole 3.6F Ag Fungicide in no less than 20 gallons of water per acre.

Aerial Application:

Use the specified rate of Tebuconazole 3.6F Ag Fungicide in to less than 10 gallons of water per acre.

Animal Feeding and Grazing Directions:

Following the application of this product, do not permit animals to graze or forage in the treated areas for at least 17 days. While straw, chaff and screenings from the treated area may be used for feed, do not use seed for animal feed purposes. Do not forage or cut green crop.

Restricted-entry interval = 12 hours

GREEN ONIONS

APPLICATION DIRECTIONS

| CROP | TARGET DISEASES | PER ACRE PRODUCT USE RATE PER APPLICATION | FOLLOW-UP APPLICATION TIMING | MAXIMUM USE RATE OF PRODUCT/ACRE/ CROP PER SEASON | PHI |
|--|--|---|------------------------------|---|--------|
| Green onion Leek Spring onion Scallion Japanese bunching onion Green Shallot Green Eschalots | White rot (<i>Sclerotium cepivorum</i>) Suppression only Rust (<i>Puccinia allii</i> , <i>Puccinia porri</i>) Purple blotch (<i>Alternaria porii</i>) | 4 – 6 fl. ounces | 10 – 14 days | 24 fl. ounces | 7 days |

Application Directions

Apply Tebuconazole 3.6F Ag Fungicide as a preventative treatment in a protective spray schedule. Begin applications as soon as crop and/or environmental conditions become favorable for disease development.

For optimum disease control, the lowest labeled rate of a spray surfactant may be tank mixed with Tebuconazole 3.6F Ag Fungicide.

Tebuconazole 3.6F Ag Fungicide must have 2 - 4 hours of drying time on plant foliage for the active ingredient to move systematically in to the plant tissue before rain or irrigation occurs. After 2 - 4 hours the Tebuconazole 3.6F Ag Fungicide will be resistant to weathering.

Restricted-entry interval (REI) = 12 hours.

HOPS

APPLICATION DIRECTIONS

| CROP | TARGET DISEASES | PER ACRE PRODUCT USE RATE PER APPLICATION | FOLLOW-UP APPLICATION TIMING | MAXIMUM USE RATE OF PRODUCT/ACRE/ CROP PER SEASON | PHI |
|------|---|---|------------------------------|---|---------|
| Hops | Powdery mildew (<i>Sphaerotheca humuli</i> / <i>Sphaerotheca macularis</i>) | 4 – 8 fl. oz. | 10 – 14 days | 32 fl. oz. | 14 days |

Application Directions

Apply the specified dosage of Tebuconazole 3.6F Ag Fungicide in a protective spray schedule to foliage. Increase the spray volume and the application rate as vine growth increases during the season.

For optimum disease control, the lowest labeled rate of a spray surfactant may be tank mixed with Tebuconazole 3.6F Ag Fungicide.

Tebuconazole 3.6F Ag Fungicide must have 2 - 4 hours of drying time on plant foliage for the active ingredient to move systematically in to the plant tissue before rain or irrigation occurs. After 2 - 4 hours the Tebuconazole 3.6F Ag Fungicide will be resistant to weathering.

Restricted-entry interval (REI) = 12 hours.

LEAFY BRASSICA GREENS

APPLICATION DIRECTIONS

| CROP | TARGET DISEASES | PER ACRE PRODUCT USE RATE PER APPLICATION | FOLLOW-UP APPLICATION TIMING | MAXIMUM USE RATE OF PRODUCT/ACRE/CR OP PER SEASON | PHI |
|--|---|---|------------------------------|---|--------|
| Leafy Brassica Greens Broccoli raab Chinese cabbage (bok choy) Collards Kale Mizuna Mustard greens Mustard spinach Rape greens Turnip greens [Application to turnip greens is limited to east of the Rockies.] | Cercospora leaf spot (<i>Cercospora brassicola</i>) Powdery mildew (<i>Erysiphe cruciferarum</i>) Alternaria leaf spot (<i>Alternaria brassicola</i>) | 3 - 4 fl. ounces | 14 days | 16 fl ounces | 7 days |

Application Directions

Apply Tebuconazole 3.6F Ag Fungicide as a preventative treatment for optimum results. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. Do not apply more often than once every 10 days.

For optimum disease control, the lowest labeled rate of a spray surfactant may be tank mixed with Tebuconazole 3.6F Ag Fungicide.

Tebuconazole 3.6F Ag Fungicide must have 2 - 4 hours of time on plant foliage for the active ingredient to move systematically in to the plant tissue before rain or irrigation

occurs. After 2 - 4 hours the Tebuconazole 3.6F Ag Fungicide will be resistant to weathering.

Restricted-entry interval (REI) = 12 hours.

GARDEN BEETS

APPLICATION DIRECTIONS

| CROP | TARGET DISEASES | PER ACRE PRODUCT USE RATE PER APPLICATION | FOLLOW-UP APPLICATION TIMING | MAXIMUM USE RATE OF PRODUCT/ACRE/CROP PER SEASON | PHI |
|-------------------------------------|---|---|------------------------------|--|--------|
| Garden beet roots and tops (leaves) | Cercospora leaf spot (<i>Cercospora beticola</i>) | 3 - 7.2 fl. oz. | 14 days | 28.8 fl. oz. | 7 days |

Application Directions

Apply Tebuconazole 3.6F Ag Fungicide as a preventative treatment for optimum results. Begin applications as soon as crop and/or environmental conditions become favorable for disease development.

For optimum disease control, the lowest labeled rate of a spray surfactant may be tank mixed with Tebuconazole 3.6F Ag Fungicide.

Tebuconazole 3.6F Ag Fungicide must have 2 - 4 hours of time on plant foliage for the active ingredient to move systematically in to the plant tissue before rain or irrigation occurs. After 2 - 4 hours the Tebuconazole 3.6F Ag Fungicide will be resistant to weathering.

Restricted-entry interval (REI) = 12 hours.

LYCHEE

APPLICATION DIRECTIONS

| CROP | TARGET DISEASES | PER ACRE PRODUCT USE RATE PER APPLICATION | FOLLOW-UP APPLICATION TIMING | MAXIMUM USE RATE OF PRODUCT/ACRE/CROP PER SEASON | PHI |
|--------|---|---|------------------------------|--|--------|
| Lychee | Anthracoise (<i>Colletotrichum gloeosporioides</i>) | 4 - 6 fl. oz. | 10 days | 48 fl. oz. | 0 days |

Application Directions

Begin first application of Tebuconazole 3.6F Ag Fungicide as panicle emerges. Apply the specified dosage in a minimum of 50 gallons of spray solution per acre by ground only. Tebuconazole 3.6F Ag Fungicide can be applied up to and including the day of harvest.

For optimum disease control, the lowest labeled rate of a spray surfactant may be tank mixed with Tebuconazole 3.6F Ag Fungicide.

Tebuconazole 3.6F Ag Fungicide must have 2 - 4 hours of drying time on plant foliage for the active ingredient to move systematically in to the plant tissue before rain or irrigation occurs. After 2 - 4 hours the Tebuconazole 3.6F Ag Fungicide will be resistant to weathering.

Restricted-entry interval (REI) = 2 days.

OKRA

APPLICATION DIRECTIONS

| CROP | TARGET DISEASES | PER ACRE PRODUCT USE RATE PER APPLICATION | FOLLOW-UP APPLICATION TIMING | MAXIMUM USE RATE OF PRODUCT/ACRE/CROP PER SEASON | PHI |
|------|---|---|------------------------------|--|--------|
| Okra | Cercospora leaf spot (<i>Cercospora</i> spp.) | 4 - 6 fl. ounces. | 14 days | 24 fl. ounces | 3 days |

Application Directions

Apply specified dosage of Tebuconazole 3.6F Ag Fungicide in a preventative spray program. Use the highest rate when disease conditions are favorable and in areas where high disease pressure is expected. Apply specified dosage as a foliar spray in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air.

For optimum disease control, the lowest labeled rate of a spray surfactant may be tank mixed with Tebuconazole 3.6F Ag Fungicide.

Tebuconazole 3.6F Ag Fungicide must have 2 - 4 hours of drying time on plant foliage for the active ingredient to move systematically in to the plant tissue before rain or irrigation occurs. After 2 - 4 hours the Tebuconazole 3.6F Ag Fungicide will be resistant to weathering.

Restricted-entry interval (REI) = 12 hours.

PEANUTS

APPLICATION DIRECTIONS

| CROP | TARGET DISEASES | PER ACRE PRODUCT USE RATE PER APPLICATION | FOLLOW UP APPLICATION TIMING | MAXIMUM USE RATE OF PRODUCT/ACRE/CROP PER SEASON | PHI |
|-----------------------|---|--|-------------------------------------|---|------------|
| Peanuts Foliar | Early leaf spot (<i>Cercospora arachidicola</i>) | 7.2 fluid ounces | 14 days | 28.8 fluid ounces | 14 days |
| Foliar | Late leaf spot (<i>Cercosporidium personatum</i>) | | | | |
| Foliar | Leaf rust (<i>Puccinia</i> spp.) | | | | |
| Foliar | Pepper spot (<i>Leptosphaerulina crassiasca</i>) | | | | |
| Foliar | Web blotch (<i>Phoma arachidicola</i>) | | | | |
| Peanuts Soil borne | Southern stem rot Southern blight White mold (<i>Sclerotium</i> spp.) | | | | |
| Soil borne | Rhizoctonia limp rot Rhizoctonia pod rot* (<i>Rhizoctonia solanii</i>) | | | | |

Application Directions

*Rhizoctonia pod rot – North Carolina and Virginia only.

General Use Directions For Peanuts:

Ground Application:

Apply Tebuconazole 3.6F Ag Fungicide in no less than 10 gallons of spray solution per acre.

Aerial Application

Apply Tebuconazole 3.6F Ag Fungicide in no less than 5 gallons of spray solution per acre.

Traditional and university proven anti-disease techniques, such as specific crop rotation, along with industry approved best management practices, will contribute to optimum disease control when used with Tebuconazole 3.6F Ag Fungicide.

For optimum disease control, the lowest labeled rate of a spray surfactant may be tank mixed with Tebuconazole 3.6F Ag Fungicide.

Tebuconazole 3.6F Ag Fungicide will be less effective when the area to be treated is subject to drought. Product must be moved into the lower plant area and surrounding soil area by rain and overhead irrigation. Moving the applied product down into the plant structure and surrounding soil is especially important in the control of root, stem and pod diseases.

Mode of Action Information

The active ingredient in tebuconazole 3.6F Ag Fungicide is a member of the DMI (Demethylation Inhibitor) fungicide group and FRAC grouping 3. Its mode of action inhibits synthesis of sterols. The triazole fungicide's actions are protective, curative (when applied early in the fungal pathogen's life cycle) and systemic in nature. The active ingredient is absorbed by root and leaf tissue, and then moves to the growing tissue. (Chlorothalonil is a Substituted Benzene fungicide that slows sporulation and growth rates of fungi and a member of FRAC group Y, Multi Site Action. Its action is protective and makes it a good resistance management partner).

Soilborne Disease Preventative Spray Program:

For best results in controlling White Mold and other Soilborne diseases (such as Sclerotium stem and pod rots or Rhizoctonia limb and pod rots), apply the above specified rate as part of a seven application spray program. Treatments should be initiated as preventative in nature. Chlorothalonil should be used in the beginning treatments (1st and 2nd) and those following four (4) consecutive Tebuconazole 3.6F Ag Fungicide applications (14 day scheduled) to lessen the risks of disease resistance. All treatments after mid August should be tank mixed with chlorothalonil.

Leaf Spot Resistance:

Care should be taken not to alternate or tank mix DMI fungicides in the same application. Non-DMI fungicides should be used in rotation or alternation with Tebuconazole 3.6F Ag Fungicide for disease resistance management. Contact your local extension peanut specialist or crop consultant about management programs proven for your area.

Animal Feeding and Grazing Directions:

Following application of this product, do not permit animals to graze or forage in the treated areas. Hay and harvester thrashings from the treated area may not be used for animal feed.

Restricted entry interval (REI) = 12 hours.

PECAN

APPLICATION DIRECTIONS

| CROP | TARGET DISEASES | PER ACRE PRODUCT USE RATE PER APPLICATION | FOLLOW-UP APPLICATION TIMING | MAXIMUM USE RATE OF PRODUCT/ACRE/ CROP PER SEASON | PHI |
|-------|---|---|------------------------------|---|--|
| Pecan | Brown leaf spot <i>(Sirosporium diffusium)</i> Downy spot <i>(Mycosphaerella caryigena)</i> Liver spot <i>(Gnomonia caryae)</i> Scab <i>(Cladosporium caryigenum)</i> Vein spot <i>(Gnomonia nerviseda)</i> Zonate leaf spot <i>(Grovesinia pyramidalis)</i> | 4 – 8 fl. oz. | 10 – 14 days | 32 fl. oz. | Do not apply Tebuconazole 3.6F Ag Fungicide after shucks begin to split. |

Application Directions

Apply Tebuconazole 3.6F Ag Fungicide in a preventive spray schedule beginning at early bud break (young leaves unfolding), and continue applications through the pollination period. Tebuconazole 3.6F Ag Fungicide should be applied at 4 fl. oz. per acre in a tank-mix with the labeled rate of Super-Tin^R in cover sprays. Follow label directions for the use of Super-Tin. Do not add a surfactant to the spray solution when tank-mixing Tebuconazole 3.6F Ag Fungicide with Super-Tin. Apply Tebuconazole 3.6F Ag Fungicide in a spray volume of 15 gallons or more per acre by air or 50 gallons or more per acre by ground.

Apply 7 to 8 fl. oz. per acre of Tebuconazole 3.6F Ag Fungicide to full-size mature trees and 4 to 6 fl. oz. per acre of Tebuconazole 3.6F Ag Fungicide to smaller trees. Apply the high rate to varieties that are highly susceptible to the indicated diseases, or when severe disease conditions exist.

For optimum disease control, the lowest labeled rate of a spray surfactant may be tank mixed with Tebuconazole 3.6F Ag Fungicide.

Tebuconazole 3.6F Ag Fungicide must have 2 - 4 hours of drying time on plant foliage for the active ingredient to move systematically in to the plant tissue before rain or irrigation occurs. After 2 - 4 hours the Tebuconazole 3.6F Ag Fungicide will be resistant to weathering.

Do not cut cover crops in treated areas for feed or allow livestock to graze treated areas.

Restricted-entry interval (REI) = 12 hours.

SOYBEANS

APPLICATION DIRECTIONS

| CROP | TARGET DISEASES | PER ACRE PRODUCT USE RATE PER APPLICATION | FOLLOW-UP APPLICATION TIMING | MAXIMUM USE RATE OF PRODUCT/ACRE/CROP PER SEASON | PHI |
|---------|---|---|------------------------------|--|---------|
| Soybean | Rust (<i>Phakopsora pachyrhizi</i>) Powdery mildew (<i>Microsphaera diffusa</i>) | 3 – 4 fl. oz. | 10 – 14 days | 12 fl. oz. | 21 days |

Application Directions

Apply specified dosage of Tebuconazole 3.6F Ag Fungicide as a broadcast foliar spray as a preventative or at first visible symptoms of disease. Use the higher rate and shorter spray interval when disease pressure is severe. Apply specified dosage as a foliar spray in a minimum of 10 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air.

For optimum disease control, the lowest labeled rate of a spray surfactant may be tank mixed with Tebuconazole 3.6F Ag Fungicide.

Do not apply more than 3 applications per season.

Restricted-entry interval (REI) = 12 hours.

SUNFLOWER

APPLICATION DIRECTIONS

| CROP | TARGET DISEASES | PER ACRE PRODUCT USE RATE PER APPLICATION | FOLLOW-UP APPLICATION TIMING | MAXIMUM USE RATE OF PRODUCT/ACRE/CROP PER SEASON | PHI |
|-----------|------------------------------------|---|------------------------------|--|---------|
| Sunflower | Rust (<i>Puccinia helianthi</i>) | 4 – 6 fl. oz. | 14 days | 16 fl. oz. | 50 days |

Application Directions

Apply specified dosage of Tebuconazole 3.6F Ag Fungicide at the earliest sign of infection (rust pustules developing) or when weather conditions are favorable for rust development. Apply higher rate to highly susceptible varieties and/or under severe disease conditions. Apply specified dosage as a foliar spray in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air.

For optimum disease control, the lowest labeled rate of a spray surfactant may be tank

mixed with Tebuconazole 3.6F Ag Fungicide.

Tebuconazole 3.6F Ag Fungicide must have 2 - 4 hours of drying time on plant foliage for the active ingredient to move systematically in to the plant tissue before rain or irrigation occurs. After 2 - 4 hours the Tebuconazole 3.6F Ag Fungicide will be resistant to weathering.

Restricted-entry interval (REI) = 12 hours.

TURNIP

APPLICATION DIRECTIONS

| CROP | TARGET DISEASES | PER ACRE PRODUCT USE RATE PER APPLICATION | FOLLOW-UP APPLICATION TIMING | MAXIMUM USE RATE OF PRODUCT/ACRE/ CROP PER SEASON | PHI |
|---|--|---|------------------------------|---|--------|
| Turnip [Application is limited to east the Rockies.] | Cercospora leaf spot (<i>Cercospora brassicicola</i>) | 4 - 7.2 fl. oz. | 12 - 14 days | 28.8 fl. oz. | 7 days |

Application Directions

Apply specified dosage of Tebuconazole 3.6F Ag Fungicide in a protective spray schedule to foliage.

For optimum disease control, the lowest labeled rate of a spray surfactant may be tank mixed with Tebuconazole 3.6F Ag Fungicide.

Tebuconazole 3.6F Ag Fungicide must have 2 - 4 hours of drying time on plant foliage for the active ingredient to move systematically in to the plant tissue before rain or irrigation occurs. After 2 - 4 hours the Tebuconazole 3.6F Ag Fungicide will be resistant to weathering.

Restricted-entry interval (REI) = 12 hours.

WHEAT

APPLICATION DIRECTIONS

| CROP | TARGET DISEASES | PER ACRE PRODUCT USE RATE PER APPLICATION | FOLLOW-UP APPLICATION TIMING | MAXIMUM USE RATE OF PRODUCT/ACRE/ CROP PER SEASON | PHI |
|-------|--|---|------------------------------|---|---------|
| Wheat | Rusts-leaf, stem and stripe (<i>Puccinia</i> spp.) | 4 fl. oz. | Not Allowed | 4 fl. oz. | 30 days |
| | Head Blight (<i>Fusarium</i> spp.)- Suppression | | | | |

Application Directions

For optimum control, wheat fields should be kept under observation for early disease symptoms. This is particularly important when conditions favoring disease development are favorable or when varieties susceptible to disease are planted.

For Rusts, apply the Tebuconazole 3.6F Ag Fungicide at the earliest sign of rust pustules on foliage.

For Fusarium head blight apply Tebuconazole 3.6F Ag Fungicide at the beginning of flowering on the main stem heads (Feekes 10.51) of the plants for optimum suppression.

For optimum disease control, the lowest labeled rate of a spray surfactant may be tank mixed with Tebuconazole 3.6F Ag Fungicide.

Tebuconazole 3.6F Ag Fungicide must have 2 - 4 hours of drying time on the plant foliage for the active ingredient to move systematically in to the plant tissue. After 2 - 4 hours the Tebuconazole 3.6F Ag Fungicide will be resistant to weathering.

Spray Volume:

Ground Application

Apply Tebuconazole 3.6F Ag Fungicide in no less than 10 gallons of spray solution per acre.

Aerial Application

Apply Tebuconazole 3.6F Ag Fungicide in no less than 5 gallons of spray solution per acre.

Animal Feeding and Grazing Directions:

Following application of this product, do not permit animals to graze or forage in the treated areas for at least 6 days. Straw may be used for feed or bedding.

Restricted-entry interval (REI) = 12 hours.

| SEED TREATMENT - Corn (Sweet Corn, Field Corn, Field Corn Grown For Seed, and Popcorn). For control of soilborne and seedborne Fusarium and soilborne and seedborne head smut. | | |
|--|-----------------------|---|
| SEED LABELING: To meet U.S. Federal Seed Act requirements, all seed treated with Tebuconazole 3.6F Ag Fungicide must be labeled: TREATED SEED. DO NOT USE FOR FOOD, FEED OR OIL PURPOSES. Treated with Tebuconazole. | | |
| USE PRECAUTION: When using formulations that do not contain dye, to comply with 40 CFR 153.155, all seed treated with an economic poison must be colored to distinguish and prevent subsequent inadvertent use as a food for man or feed for animals. | | |
| DISEASE | RATE Fl Oz/CWT | DIRECTIONS FOR USE |
| Soilborne and Seedborne <i>Fusarium</i> spp. | 0.071 | Apply as a seed treatment using standard slurry or mist-type seed treatment equipment. Uniform applications of seed is necessary to ensure seed safety and best defense protection. Seed should be sound and well |
| Soilborne and Seedborne Head smut (<i>Sphacelotheca reiliana</i>) | 0.27 - 0.54 | |

| | | |
|--|--|---|
| | | <p>cured prior to treatment. Product should be diluted with sufficient water to ensure complete seed coverage. Consult a seed treatment specialist regarding slurry rates recommended for the crop to be treated with Tebuconazole 3.6F Ag Fungicide. The length of control will vary depending on the rate used.</p> |
|--|--|---|

ROTATIONAL CROPS: Treated areas may be replanted with any crop specified on this label as soon as practical after last application. Any crop not specified on this label may be planted into treated areas 120 days after last application.

**IMPORTANT INFORMATION
READ BEFORE USING PRODUCT**

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded. The Directions for Use of this product must be followed carefully. The Directions for Use of this product reflect the opinion of experts based on field use and tests. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SOURCE DYNAMICS LLC or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of SOURCE DYNAMICS LLC and Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SOURCE DYNAMICS LLC and Seller harmless for any claims relating to such factors.

SOURCE DYNAMICS, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SOURCE DYNAMICS LLC, and to the extent consistent with applicable law Buyer and User assume the risk of any such use. SOURCE DYNAMICS LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, SOURCE DYNAMICS LLC or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SOURCE DYNAMICS LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING**

CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE, AT THE ELECTION OF SOURCE DYNAMICS LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT, OR COMPENSATION LIMITED TO DAMAGES NOT EXCEEDING THE FAIR MARKET PURCHASE PRICE, AND SHALL NOT INCLUDE INCIDENTAL OR CONSEQUENTIAL DAMAGES.

SOURCE DYNAMICS LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of SOURCE DYNAMICS LLC.