| UNITED STATED | U.S. ENVIRONMENTAL PROTECTION AGENCY | EPA Reg. Number: | Date of Issuance: |
|---|---|---|--|
| SAN HOME TAL PROTECTION | Office of Pesticide Programs Registration Division (7505C) 1200 Pennsylvania Ave., N.W. Washington, D.C. 20460 | 82957-2 | FEB 12 2 |
| (under FIFRA, as amen | NOTICE OF PESTICIDE: <u>X</u> Registration Reregistration | Term of Issuance: Conditional Name of Pesticide Product: Prothor WP | |
| | | | |
| Name and Address of | Registrant (include ZIP Code): | | |
| 4110 136'th Str Gig Harbor, WA | A 98332 | | |
| Registration Division pr On the basis of informat and Rodenticide Act. Registration is in no way | ng differing in substance from that accepted in connection with this regi- tion to use of the label in commerce. In any correspondence on this pro- tion furnished by the registrant, the above named pesticide is hereby reg | huct always refer to the above EF istered/reregistered under the Fe by the Agency. In order to prote | PA registration number deral Insecticide, Fung ect health and the |
| Registration Division pr On the basis of informat and Rodenticide Act. Registration is in no way environment, the Admin of any name in connecti- or to its use if it has been | for to use of the label in commerce. In any correspondence on this product tion furnished by the registrant, the above named pesticide is hereby reg y to be construed as an endorsement or recommendation of this product distrator, on his motion, may at any time suspend or cancel the registration on with the registration of a product under this Act is not to be construed | huct always refer to the above EF istered/reregistered under the Fe by the Agency. In order to prote on of a pesticide in accordance w d as giving the registrant a right | PA registration number deral Insecticide, Fung ect health and the with the Act. The accept to exclusive use of the |
| Registration Division pr On the basis of informat and Rodenticide Act. Registration is in no way environment, the Admin of any name in connecti- or to its use if it has been This produ- that you: 1. Subn | tion furnished by the registrant, the above named pesticide is hereby reg y to be construed as an endorsement or recommendation of this product instrator, on his motion, may at any time suspend or cancel the registration on with the registration of a product under this Act is not to be construed in covered by others. | huct always refer to the above EF istered/reregistered under the Fe by the Agency. In order to prote on of a pesticide in accordance w d as giving the registrant a right h FIFRA section 3(c)(eregistration of your p | PA registration number deral Insecticide, Fung ect health and the vith the Act. The accep to exclusive use of the (7)(A) provided |
| Registration Division pr On the basis of informat and Rodenticide Act. Registration is in no way environment, the Admin of any name in connecti- or to its use if it has been This produ- that you: 1. Subm Agen 2. Make a. b. | tion furnished by the registrant, the above named pesticide is hereby registration furnished by the registrant, the above named pesticide is hereby registration on his motion, may at any time suspend or cancel the registration on with the registration of a product under this Act is not to be construed in covered by others. | huct always refer to the above EF istered/reregistered under the Fe by the Agency. In order to prote on of a pesticide in accordance w d as giving the registrant a right h FIFRA section 3(c)(eregistration of your p submit such data. the product for shipm <i>EPA Reg. No. 82957</i> e the following statem <i>y not be necessary to p</i> <i>nces, it may be necess</i> <i>products such as terr</i> | PA registration number deral Insecticide, Fung ect health and the with the Act. The accepto to exclusive use of the (7)(A) provided product when the nent: -2". nents: "All provide adequation cary to supplem nite baits or |
| Registration Division pr On the basis of informat and Rodenticide Act. Registration is in no way environment, the Admin of any name in connecti- or to its use if it has been This produ- that you: 1. Subm Agen 2. Make a. b. | tion furnished by the registrant, the above named pesticide is hereby reg of the construed as an endorsement or recommendation of this product distrator, on his motion, may at any time suspend or cancel the registration on with the registration of a product under this Act is not to be construed in covered by others. The conditionally registered in accordance with the following label changes before you release Revise the EPA Registration Number to read, " Within the "General" section of the label, delet treatment directions contained in this label may protection against termites. In some circumstant the use of PROTHOR WP with other termiticide products approved for direct or injection applica- the property." | huct always refer to the above EF istered/reregistered under the Fe by the Agency. In order to prote on of a pesticide in accordance w d as giving the registrant a right h FIFRA section 3(c)(eregistration of your p submit such data. the product for shipm <i>EPA Reg. No. 82957</i> e the following statem <i>y not be necessary to p</i> <i>nces, it may be necess</i> <i>products such as terr</i> | PA registration number deral Insecticide, Fung ect health and the with the Act. The acce to exclusive use of the (7)(A) provided product when the nent: -2". nents: "All provide adequation cary to supplem nite baits or |

Page 2 EPA Reg. No. 82957-2

c. All references to pre-construction treatments must be deleted from the label. To have pre-construction treatments added to the label at a future date, a thorough "PRE-CONSTRUCTION TREATMENT" section, comparable to the cited label, must be submitted to the Agency.

2110

- d. All references to foam applications must be deleted from the label. To have foam treatments added to the label at a future date, a thorough "FOAM APPLICATION" section, comparable to the cited label, must be submitted to the Agency.
- e. Within the "STRUCTURES CONTAINING CONCRETE SLABS on GROUND" section of the label, include the following statements: "Treat all existing cracks and cold, construction or expansion joints. Also, treat around bath traps, plumbing and utility services which penetrate the slab. Apply 4 gallons of solution per 10 linear feet per foot of depth to provide a uniform treated zone."
- f. Within the "DISCLAIMER OF WARRANTIES" section of the label, begin the first sentence with "*To the extent consistent with applicable law*..."
- g. Within the "DISCLAIMER OF WARRANTIES" section of the label, rephrase the first sentence to read "To the extent consistent with applicable law..."
- h. Within the "CONDITIONS OF SALE" section of the label, rephrase the first sentence to read "*The Directions for Use of this product are believed to be adequate and must be followed carefully*."
- 3. The data requirements for storage stability (830-6317) and corrosion characteristics (830-6320) have not been satisfied, and must be submitted within eighteen months of the date of this letter.
- 4. Submit one copy of the revised final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. If you have any questions regarding this letter, please contact Kable Bo Davis at 703 306-0415.

A stamped copy of the label is enclosed for your records.

John Hebert Product Manager (07) Insecticide-Rodenticide Branch Registration Division (7505P)

Enclosure



PROTHOR WP

For use only by individuals/firms licensed or registered by the state to apply termiticide products. States may have more restrictive requirements regarding qualifications of persons using this product. Consult the structural pest control regulatory agency of your state prior to use of this product.

For prevention or control of subterranean termites in and around residential, commercial, industrial, institutional and public structures and buildings.

| Active Ingredient: | By Wt. |
|--------------------|--------------|
| Imidacloprid | . 75.0% |
| Other Ingredients: | <u>25.0%</u> |
| TOTAL: | 100.0% |
| | |

Keep water soluble packets in outer container until immediately before use. Store in a cool, dry place but not below freezing (32⁰F).

EPA Reg. No. 82957-X EPA Est. 81824-NC-001

STOP – Read the label before use

KEEP OUT OF REACH OF CHILDREN

CAUTION

(PRECAUCION AL USUARIO: Si usted no puede leer o entender ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.)

(TO THE USER: If you cannot read and understand English, do not use this product until the label has been fully explained to you.)

For product use information call 1-866-FOR-THOR (367-8467) or visit www.for-thor.com.

NET WEIGHT: As marked on container

Manufactured by: ENSYSTEX III, Inc. Favetteville, NC 28303

ACCEPTED with COMMENTS In EPA Letter Dated: FEB 1 2 2007

Under the Dedecri Intecticide, Fungicide, and Heferricide Act, as anothered, for the pesticide registered under EFA Reg. No. 82957-2

| FIRST AID | | | | |
|---------------------------|--|--|--|--|
| If swallowed | Call a poison control center or doctor immediately for treatmen advice Have person sip a glass of water if able to swallow Do not induce vomiting unless told to do so by a poison control center or doctor Do not give anything by mouth to an unconscious person | | | |
| lf on skin or clothing | Take off contaminated clothing Rinse skin (mmediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice. | | | |
| If in eyes | Hold eye open and rise slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes then continue rinsing eye Call a poison control center or doctor for treatment advice. | | | |
| | HOTLINE NUMBER | | | |

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-866-367-8467 for emergency medical treatment information.

NOTE TO PHYSICIAN

No specific antidote is available. Treat the patient symptomatically

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Causes 8ye imitation. Avoid contact with skin, eyes or clothing. Avoid breathing dust or vapor. Wash thoroughly with scap and water after handling and before eating, drinking chewing gum, using tobacco or using the toillet. Remove contaminated clothing and wash before reuse. Keep children and pets away from treated area until dry.

Personal Protective Equipment: All pesilicide handlers (mixers, loaders and applicators) must wear long-sleeved shirt and long pants, socks, shoes and water-proof gloves. After the product is diluted in accordance with label directions for use, shirt, pants, socks and water-proof gloves are sufficient protection. All pesticide handlers must wear protective eyewear, such as goggles, faceshield or safety glasses, when working in a non-ventilated space or when applying as a termiticide by rodding or sub-slab injection.

Termite Control Treatment: When freating adjacent to an existing structure, the applicator must check the area to be treated and immediately adjacent areas of the structure for visible and accessible cracks and holes to prevent any leaks or significant exposures to persons occupying the structure. People present or residing in the structure during application must be advised to remove their pets and themselves from the structure if they see any signs of leakage. After application, the applicator is required to check for leaks. All teaks resulting in the deposition of termitoide in locations other than those prescribed on this label must be cleaned up prior to leaving the application site. Do not allow people or pets to contact contaminated areas or to reoccupy contaminated areas of the structure unit the cleanup is completed.

Environmental Hazards

This pesticide is highly toxic to aqualic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label. Extreme care must be taken to avoid runoff. Apply only to soll or other fill substrate that will accept the solution at the specified rate. Do not treat soll that is water-saturated or frozen or in any conditions where run-off or movement from the treated area (site) is likely to occur.

Physical and Chemical Hazards

Do not apply this product or solutions of this product around electrical equipment such as electrical conduits, motor housings, junction boxes, switch boxes, etc. due to the possibility of shock hazard.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Keep out of reach of children and animals Store in original containers only Store in a cool (but not below freezing ($32^{O}F$)), dry place and avoid excess heat Handle and open container in a manner so as to prevent spillage. Do not put concentrate or dilute material into food or drink containers' Preferably store in a locked area

Exposure to moisture or excessive handling of water soluble packets can cause them to break. Store water soluble packets in original container preferably in a locked area. Do not cut the water soluble packets when opening outer container.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site (in the treatment area) or at an approved waste disposal facility

Container Dispossi: [For product packaged in plastic bottle]: Thele nnse (or equivalent). Then ofter for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incrineration, or if allowed by state and local authorities, by burning. If burned stay out of smoke

[For product packaged in WSP]: Do not use carton in connection with food, feed or drinking water The empty foil wrappers may be disposed of in the trash. After removing all PVA packets, the carton may be disposed of in the trash.

in Case of Spill: Confine II, avoid contact, isotate area and keep animals and unprotected persons away. If spill is liquid, form dike around spill area and/or absorb spill with absorbent materials, such as sand, cat litter or clay. If spill is dry powder only sweep material into a suitable container. Place damaged package in a holding container and identify contents. Contact Ensystex til at 1-866-357-8467 or Chemtrec at 1-800-424-9300 for any assistance.

APPLICATION FOR CONTROL OF SUBTERRANEAN TERMITES

General

PROTHOR WP, in the form of a dilute insecticidal solution, prevents and controls subterranean termite infestations in and around structures and other items by creating a continuous chemically treated zone (horizontal and/or vertical as needed) between the wood and other cellulose material in a structure and termite colonies in the soit. In order to establish a zone between the wood in the structure and the termites in the soit, adequately disperse the solution of this product in the soil.

To effectively control subterranean termites with this product, the service technician should be familiar with current subterranean termite control practices including trenching, rodding, sub-slab and void injection, soil surface fan spraying and excavated soil treatment. Correct use of these techniques is necessary to effectively control infestations by subterranean termites such as *Coptotermes*, *Heterotermes* and *Refuciliermes*. The service technician should consider the biology and behavior of the termite specie(s) to be controlled to determine which control practices to use.

Treatment standards and procedures for subternanean termite control may vary due to regulations, water table level, structure design, soil types, construction practices and other factors. For advice concerning current control practices with respect to specific local conditions, consult resources in structural pest control and state cooperative extension and regulatory agencies. Follow all faderal, state and local regulations and treatment standards for protection of a structure from subternaem remities. All treatment directions contained in this table may not be necessary to provide adequate protection against termities. In some circumstances, it may be necessary to supplement the use of PROTHOR WP with other termiticide products such as termite baits or products approved for direct or injection application to wood to adequately protect the property.

Effective termite control may also include mechanical attention of the structure. Elimination of leaks or points of moisture accumulation within or on the exterior of the structure that result in an increase in the moisture content of wooden structural components is advised. Removal of nonessential cellulose containing materials that are in contact with the ground under or around the structure can reduce termite foraging in the area.

PROTHOR WP is labeled for use against subterranean termites as a 0.05% to 0.10% solution in water. Generally, the 0.05% rate is used for typical control situations. When severe or persistent infestations are occurring, a 0.10% solution may be more appropriate. When difficult or problem solls or construction types are encountered, it may be necessary to use 0.10% PROTHOR WP mixed in reduced volumes of water.

Avoid contamination of water supplies due to backflow under reduced water system pressure by using anti-backflow equipment or procedures to prevent sphorning of any solution back into a water supply. Do not contaminate cistems or wells. Do not treat soli that is water saturated or frozen. Do not treat white pracipitation is occurring. Do not apply solution to an area or site if the soil at the area or site is in such a state or condition that runoff or movement of the solution from the treated area or site is likely to occur. Structures that contain wells or cistems within the loundation of the structure can only be treated using the treated backfill method described in the treatment around wells and cistems section of this label. Consult state and local specifications for recommended distances of wells from treated areas, or if such regulations do not exist, refer to Federal Housing Administration Specifications (H.U.D.) for guidance.

Dilution and Mixing of PROTHOR WP

[Note to Reviewer: This paragraph will only appear on label for product packaged in WSP] Each PROTHOR WP outer container contains a number of clear water soluble packets. Do not allow the packets to become wet at any time before putting them into a sprey tank at the time of intended mixing. Handling the packets with wet hands or gloves or in a rough manner can cause them to leak. Reseal the outer container if water soluble packets are left inside after opening.

Use rates for PROTHOR WP are expressed and the solution is mixed according to the percentage (%) concentration. It forms when mixed in water. Use the *Mixing Table for PROTHOR WP* or alternately the formulas below to determine the amount of PROTHOR WP to add to any quantity of water.

To mix, measure out the required amount of PROTHOR WP according to the *Mixing Table for PROTHOR WP* with the PROTHOR WP Mixing Scoop. Pour this amount of PROTHOR WP into the spray tank as it is being filled with water with the agitator operating.

Mix PROTHOR WP to create a use dilution in the following manner:

1 Fill tank 1/4 to 1/3 full

2 Start pump to begin by-pass agitation and place end of treating tool in tank to allow circulation through hose

3. Add appropriate amount of PROTHOR WP

4 Add remaining amount of water.

5. Let pump run and allow recirculation through the hose for 2 to 3 minutes."

PROTHOR WP may also be mixed into full tanks of water, but substantial agitation is required to ensure uniformity of the solution.

[Note to Reviewer: The following two paragraphs will only appear on label for product packaged in WSP]

To mix, open the outer container and remove the required number of packets needed for the gallon amount of solution to be mixed according to the *Mixing Table for PROTHOR WP*. Drop the required number of packets (unopened) into the spray tank as it is being filled with water with the agitator operating. The elapsed time until the packets are fully dissolved depends on the degree of agitation and the temperature of the water with more time needed for lower levels of agitation and lower temporatures. Generally, a few minutes is sufficient time for the packets to completely dissolve.

Avoid using PROTHOR WP packets in tanks also containing products that contain boron or free chlorine. The packet material may react with the boron or chlorine to form a non-water soluble compound instead of dissolving into the water. However chlorinated water use is acceptable and will not cause this compound to form or reaction to occur.

| Mixing Table for PROTHOR WP | | | | |
|--|---|--|--|--|
| Solution Percentage Concentration Desired | Gallons of Finished Solution Desired | Amount of PROTHOR WP to add # of packets or # of level Scoops / Ounces / Grams | | |
| 0.05% | 25 | 1 / 2.25 oz. / 64 grams | | |
| | 50 | 2 / 4.50 oz. / 128 grams | | |
| | 100 | 4 / 9.00 oz. / 255 grams | | |
| 0.10% | 25 | 2 / 4.50 oz. / 128 grams | | |
| | 50 | 4 / 9.00 oz. / 255 grams | | |
| | 100 | 8 18.00 oz. / 511 grams | | |

Calculating an Amount of PROTHOR WP to Mix

To mix any amount of PROTHOR WP determine.

A = Gallons of water into which PROTHOR WP will be mixed. Express any partial gallons as decimal fractions ($1/2 \approx .5$).

Level scoops of PROTHOR WP to add to A gailons for 0.05% = A/25Level scoops of PROTHOR WP to add to A gailons for 0.10% = A/12.5

Ounces of PROTHOR WP to add to A gallons for 0.05% = A x 0.09

Grams of PROTHOR WP to add to A gallons for 0.05% = A x 2.55

Ounces of PROTHOR WP to add to A gations for 0.10% = A x 0.18

Grams of PROTHOR WP to add to A gallons for 0.05% = A x 5.11

Application Volume

To provide maximum control and protection against termite infestation, apply the specified volume of the finished water solution containing the specified amount of PROTHOR WP as set out below or as otherwise directed in this label.

Prescribed Horizontal Barrier Rate: Unless otherwise directed, horizontal barriers are created by applying a 0.05% to 0.10% solution at a rate of one gallon of solution per 10 square feet.

Prescribed Vertical Barrier Rate: Unless otherwise directed, vertical barriers are created by applying a 0.05% to 0.10% solution at a rate of four gallons of solution per 10 linear feet per foot of depth.

Adjustments to Application Volume

If soit will not accept the tabeled application volumes, the volume may be reduced provided there is a corresponding increase in concentration so that the amount of active ingredient applied to the soit remains the same.

Note: Large reductions of application volume reduce the likelihood of obtaining a continuous barrier. Variance is allowed when volume and concentration are consistent with label directed rates and a continuous barrier can still be achieved. When volume is reduced, the spacing of holes created for sub stab injection and soit rodding may need to be reduced to account for decreased dispersion of the solution in the soil.

For example, adjust the amount of solution applied to deliver a horizontal barrier of 1D square feet from 1 gallon to as low as 0.5 gallons and as high as 2 gallons while maintaining the amount of PROTHOR WP applied per 10 square feet.

For example, adjust the amount of solution applied to deliver a vertical barrier 10 feet long by one foct deep from 4 gallons to as low as 2 gallons and as high as 8 gallons while maintaining the amount of PROTHOR WP applied per 10 linear feet.

PRE-CONSTRUCTION TREATMENT

All Structures

Pre-construction treatment: Do not apply at a lower dosage and/or concentration than specified on this label for applications prior to the installation of the finished grade.

Prior to each application, applicators must notify the general contractor, construction superintendent, or similar responsible party, of the intended termiticide application and intended sites of application and instruct the responsible person to notify construction workers and other individuals to leave the area to be treasted during application and until the termiticide is absorbed into the soil.

Effective control of subterranean termities can be accomplished during construction by using a 0.05% solution of PROTHOR WP to establish vertical end/or horizontal barriers between the structure and the soli as directed. To meet current termite proofing requirements, follow the procedures in the latest edition of the Housing and Urban Development Minimum Property Standards.

Horizontal Barriers Under Stabs on Ground Including Basements

Create a horizontal barrier on the entire surface of soil or substrate that will be covered by a slab, including, but not limited to slab floors, garages, carports, basements, porches and entrance platforms by treating the soil or substrate with the solution at the Prescribed Horizontal Barrier Rate.

If the fill under the slab is a coarse material such as washed gravel, make sure that a sufficient enough amount of dilution is applied that the solution reaches the soil beneath the fill.

Apply solution using a coarse spray nozzle. If the stab over the treated area will not be poured on the same day as the application (and there are no foundation wails in place around the treated soil) cover treated soil with a water-proof barrier such as polyethylene sheeting.

Vertical Barriers

Create a vertical barrier along the inside and outside of foundation walls, around piers, plumbing and utility service entrances and other points of possible future termile access and entry by treating the soit at these points at the Prescribed Vertical Barrier Rate. When trenching and rodding into the trench, or trenching alone, it is important that the solution reaches the top of the footing. Rod holes must be spaced so as to achieve a continuous termiticidal barrier, but they should in no case be more than 12 inches spart. Trenches need not be wider than 6 inches. Mix the solution into the soil as it is being replaced in the trench. Care should be taken to avoid washing soil out from around footings thereby undermining the stability of the structure. An inside vertical barrier may not be required for a monolithic stab.

If distance from final grade to top of footing will be tass than four fest, it is permissible to wait until final grade is established to apply the vertical barrier. When treating foundations deeper than 4 fest, apply the termiticide as the backfill is being replaced, or if the construction contractor fails to notify the applicator in time to permit this, treat the foundation to a minimum depth of 4 fest after the backfill has been installed. The applicator must trench and rod into the trench arong the backfill has been installed. The applicator must trench and rod into the trench arong the grade to a minimum depth of 4 fest. When the top of the footing is exposed, the applicator must treat the soil adjacent to the footing to a depth not to exceed the bottom of the footing. However, in no case should a structure be treated below the footing.

Hollow Block Foundations and Volds

Hollow block foundations and voids may be treated at a rate of 2 gallons of solution per 10 linear feet to create a continuous treated zone within the voids at the footing.

POST CONSTRUCTION TREATMENT

All Structures

Do not apply treatment until the identity and location of all wells, radiant heat pipes, water and sever lines, electrical conduits and sub-slab heating and air conditioning ducts is detablished. Caution must be taken to avoid puncturing these elements and/or injecting solution into them. All holes in commonly occupied areas into which material has been applied must be plugged. Plugs must be of a non-cellulose material or covered by an impervious, non-cellulose material.

Vertical Barrier Depth: For applications made after the final grade is installed, the applicator must irench and rod into the trench or trench along the foundation walls and around pillars and other foundation elements and treat at the rate prescribed from grade to the top of the footing. When the footing is more than four (4) feet below grade, the applicator must trench and rod into the trench or trench along the foundation walls and treat at the rate prescribed to a minimum depth of four feet. The actual depth of treatment will vary depending on the soil type, degree of compaction, and location of termite activity. When the top of the footing is exposed, the applicator must treat the soil adjacent to the footing to a depth not to exceed the bottom of the footing. However, in no case should a structure be treated below the footing Structures Containing Concrete Stabs on Ground (Monolithic/Floating/Supported) including Basements

To make an application beneath existing slabs, it may be necessary to drill holes in the slab or adjacent foundation and to apply solution. Holes should be spaced such that when treatment is applied through them, a continuous treated zone is applied beneath the slab.

Vertical Barriers Along Exterior of Foundation Walls: Trench and rod into the trench or trench along the outside of foundation walls and treat at the Prescribed Vertical Barrier Rate to the depth specified under Vertical Barrier Depth. Where physical obstructions such as concrete walkways adjacent to foundation elements or soil type and/or conditions make trenching prohibitive, treatment may be made by rodding alone

Vertical Barriers Along Interior of Foundation Walts: Vertical barriers may be established on the interior side of foundation walls by sub-slab injection of the solution at the Prescribed Vertical Barrier Rate. Injection openings can be drilled either vertically through the slab along the interior of the foundation wall or horizontally from the exterior through the foundation wall or horizontally from the exterior through the foundation wall ow enough on the wall to allow for the deposition of the solution beneath the slab along the interior side of the foundation wall. Onli holes should be spaced so as to achieve a continuous chemical barrier but in no case farther apart than 12 inches. Special care must be taken to distribute the solution evenly. Vertical barriers may also be established beneath the slab along both sides of interior footing-supported walls, one side of interior partitions and along all cracks and expansion joints and utility service entrances and bath traps.

Horizontal Barriers Beneeth Slabs on Ground: Create a horizontal barrier by treating at the Prescribed Horizontal Barrier Rate beneath slabs by either drilling and long rodding from the exterior or by grid pattern drilling and injection vertically through the slab. Long rodding should be used only when grid pattern drilling and injection and horizontal short rodding and injection cannot be used to deliver the sub slab treatment.

Bath Traps: Exposed soil beneath and around areas where plumbing and utility services penetrate the slab should be treated at the rate of 3 gallons of solution per square fool of soil.

Structures Containing Accessible Crawl Spaces

For crawl spaces, including sealed underfloor spaces that serve as heating and sir conditioning pienums, apply vertical termiticide barriers at the rate of 4 gallons of solution per 10 linear feet per foot of depth from grade to the top of the footing, or if the footing is more than 4 feet below grade, to a minimum depth of 4 feet. Apply by trenching and rodding into the trench, or trenching. Treat both sides of foundation and around all piers and pipes. Where physical obstructions such as concrete walkways adjacent to foundation elements prevent trenching, treatment may be made by rodding atome. When soil type and/or conditions make trenching prohibitive, rodding may be used. When not to exceed the bottom of the footing. Read and follow the mixing and use direction section of the label if situations are encountered where the soil will not accept the full application volume.

1. Rod holes and trenches must not extend below the bottom of the footing.

Rod holes must be spaced so as to achieve a continuous termiticide barrier but in no case more than 12 inches apart.

3. Trenches must be a minimum of 6 inches deep or to the bottom of the footing, whichever is less, and need not be wider than 6 inches. When trenching in stoping (lisred) soil, the trench must be stepped to ensure adequate distribution and to prevent termiticide from running off. The solution must be mixed with the soil as it is replaced in the trench.

4. When treating crawl spaces used as plenums, turn off the air disculation system of the structure until application has been completed and all solution has been absorbed by the soil.

Subterranean termites can be prevented from constructing shelter tubes directly between the crawl space soil surface and overhead crawl space wooden members by the application of an overail treatment of the crawl space soil surface at the Prescribed Hortzontal Barrier Rate using a 0.05% to 0.10% solution of PROTHOR WP.

PROTHOR WP can be applied as a general fan spray within crawl speces directly to swarming and exposed worker termites at the Prescribed Horizontal Barrier Rate using a 0.05% to 0.10% solution of PROTHOR WP.

Note: Overalli treatments (treatments where chemical is applied more than 18 inches from the foundation walls, piers and pipes) should not be applied within a crawl space that serves as a plenum

Structures Containing Inaccessible Crawl Spaces

For inaccessible interior areas, such as areas where there is insufficient dearance between floor joists and ground surfaces to allow operator access, excavate, if possible, and treat according to the instructions for accessible crawl spaces. Otherwise, apply one, or a combination of the following two methods.

1. To establish a horizontal barrier, apply to the soil surface, 1 gallon of solution per 10 square feet overall using a nozzle pressure of less than 25 p.s.i. and a coarse application nozzle (e.g., Delavan Type RD Raindrop, RD-7 or Isrger, or Spraying Systems Co. 80160,P TeoJet or comparable nozzle). For an area that cannot be reached with the application wand, use one or more extension roots to make the application to the soil. Do not broadcast or power spray with higher pressures.

2. To establish a horizontal barrier, drill through the foundation wall or through the floor above and treat the soil perimeter at a rate of 1 gallon of solution per 10 square feet. Drill spacing must be at intervals not to exceed 16 inches. Many states have smaller intervals, so check state regulations which may apply.

When treating crawl spaces used as plenums, turn off the air circulation system of the structure until application has been completed and all termiticide has been absorbed by the soil.

Note. Overall treatments (treatments where chemical is applied more than 18 inches from the foundation walls, piers and pipes) should not be applied within a crawl space that serves as a plenum.

Masonry Volds

Drill and treat voids in multiple mesonry elements of the structure extending from the structure to the soil in order to create a continuous treatment barrier in the area to be treated. Apply all the rate of 2 gallons of solution per 10 linear feel of footing using a nozzle pressure of less than 25 p.s.t. When using this treatment access holes must be drilled below the still plate and should be as close as possible to the footing as is practical. Treatment of voids in block or rubble foundation waits must be dosely examined. Applicators must inspect areas of possible runoff as a precaution against application leakage in the treated ereas. Some areas may not be treatable or may require mechanical alteration prior to treatment.

All leaks resulting in the deposition of termiticide in locations other than those prescribed on this label must be cleaned up prior to leaving the application site. Do not allow people or pets to contact contaminated areas or to reoccupy the contaminated areas of the structure until the clean-up is completed

Note: When drilling veneer walls, care should be taken to not drill beyond the depth of the void behind the veneer into another construction layer behind the veneer. It is however permissible to drill through the veneer and into concrete blocks behind the veneer and to treat the veneer and the concrete blocks at the same time.

Note: Not for use in voids insulated with rigid foam.

TREATMENT OF STRUCTURES WITH WELLS AND CISTERNS Do not contaminate wells or cisterns.

Structures with Wells/Cisterns Inside Foundations

Structures that contain wells or cisterns within the foundation of a structure can only be treated

using the following techniques:

Do not treat soil while it is beneath or within the foundation or along the exterior perimeter of a structure that contains a well or cistern. The treated backfill method must be used if soil is removed and treated outside/gway from the foundation. The treated backfill technique is described as follows:

a. Trench and remove soil to be treated onto heavy plastic sheeting or similar material or into a wheelbarrow.

b. Treat the soil at the rate of 4 gallons of dilute solution per 10 linear feet per foot of depth of the trench, or 1 galkon per 1.0 cubic feet of soil. See Mixing Directions for PROTHOR WP for Use as a Termiticide section of the label. Mix thoroughly into the soil taking care to contain the liquid and prevent runoff or spillage.

c. After the treated soil has absorbed the solution, replace the soil into the trench

Structures with Adjacent Wells/Cisterns and/or Other Water Bodies

Applicators must inspect all structures with nearby water sources such as wells, cisterns, surface ponds, streams, and other bodies of water and evaluate, at a minimum, the treatment recommendations listed below prior to making an application.

 Prior to treatment, if feasible, expose the water pipe(s) coming from a well to the structure. If the pipe(s) enter the structure within 3 feet of grade.

2. Prior to treatment, applicators are advised to take precautions to limit the risk of applying the termilicide Into subsurface drains that could emply into any bodies of water. These precautions include evaluating whether application of the termilicide to the top of the footer may result in contamination of the subsurface drain. Factors such as depth to the drain system and soil type and degree of compaction should be taken into account in determining the depth of treatment.

3. When appropriate (for example, on the water side of the structure), the treated backfill technique (described above) can also be used to minimize offsite movement of termiticide

FOAM APPLICATION

PROTHOR WP, in the form of a foam, can be used to deliver PROTHOR WP as a termiticide any time it appears likely this form of delivery will improve the dispersal of PROTHOR WP into and within the intended target area. Foam can be particularly useful to deliver PROTHOR WP where it either cannot be depended upon to be delivered as just a solution or due to a need to reduce the amount of water used in order to avoid water damage to the target or adjacent areas. In some situations, for example under some slabs, a solution cannot be depended upon to disperse as completely as a foam because of deflection of the liquid stream or some other structural obstacle or defect.

Depending on the circumstances, foam applications of PROTHOR WP may be used alone or in combination with liquid solution applications, provided that the cumulative amount of active ingrecient per unit of area applied is equivalent to that which would be contained in a 0.05% to 0.10% solution-only application applied to the same area.

Using foam generating equipment, a solution of PROTHOR WP, ranging in concentration from 0.05% to 0.10%, may be converted into a foam according to the foaming agent and foaming equipment manufacturer's recommendations.

First, form a solution of PROTHOR WP of the appropriate percentage concentration and volume. Then add the recommended volume of a fearning agent. Verify that the fearning agent is compatible with PROTHOR WP.

Applications may be made behind veneers, piers, chimney bases, into rubble foundations, into block volds, structural volds or other similar volds, under stabs, stoops, porches or to the soil in crawlspaces.

RETREATMENT

Relreatments for subtemanean termites can only be performed if there is clear evidence of reinfestation or disruption of the barrier due to construction, excavation or landscaping and/or evidence of the breakdown of the termitide barrier in the soil. These vulnerable or reinfestate areas may be retreated in accordance with application techniques described in this product's labeling. The timing and type of these retreatments will vary, depending on factors such as termite pressure, soil types, soil conditions and other factors which may reduce the effectiveness of the barrier. Retreatment may be made as either a spot or compile treatment.

Releatments in the absence of reinfestation or barrier disruption may be performed five or more years after a complete treatment was last applied to the structure. Such retreatments should be made based on the judgment of the applicator that such retreatment is medessary to ensure the continued protection of the structure from termite attack. In making such judgment, the applicator should take into account the expected useful life of the last treatment administered (based on efficacy testing) and conditions specific to the structure in question that may increase it vulnerability to attack.

Annual retreatment of the structure is prohibited unless there is clear evidence that reinfestation or treated zone disruption has occurred.

APPLICATION IN CONJUNCTION WITH BORATES AND TERMITE BAITS

Spot only applications of PROTHOR WP can be used as a supplement to borate treatments and termite baiting system installations that are labeled for stand alone protection against termite attack. Stand alone product is defined as a product that is labeled for the protection of a structure when applied alone without the use of other termite control products. Spot only applications are defined as the use of PROTHOR WP according to any of the permitted and applications are defined application techniques contained in this label, alone or in combination, to the extent needed or deemed necessary or useful as an adjunct to the application of a standalone product.

APPLICATION TO PROTECT UNDERGROUND ITEMS FROM SUBTERRANEAN TERMITE ATTACK

To protect components Installed underground such as wires, conduits, cables and pipes buried in soil against termite attack, create an envelope of PROTHOR WP treated soil around the components along the entire underground length of the component. First, treat soil through which components will be run with 0.05% to 0.10% solution of PROTHOR WP at a rate of 2 galions of solution per 10 linear feet. Install components, laying them on the treated soli. Cover components with untreated soli and then treat this covering soil using the same percent solution at 2 galions of solution per 10 linear feet.

Underground components to be protected may be located within the foundation of a structure or outside of a structure such as within a utility right of way, for example. Do not treat items that are electrically energized at the time of application. If the soil will not absorb the indicated amount of solution, as little as 1 gallon of 0.10% solution per 10 linear feet can be used. Treat points where services emerge from the ground at a rate of 1 to 2 gallons of solution at the point of emergence.

APPLICATIONS TO PROTECT POLES, POSTS AND OTHER WOODEN ITEMS FROM SUBTERRANEAN TERMITE ATTACK

PRCTHOR WP can be used to protect the below ground portions of wooden structural components from termites. Form a treated zone around components below ground by vertically rodding the soil around their perimeter to a depth of six inches below their maximum depth of placement in the soil and applying a 0.05% to 0.10% solution of PROTHOR WP at a rate of 0.4 gallons of solution per linear foot of perimeter around the component per foot of treated depth. Measure the perimeter of the component ax inches from the outside of the component.

APPLICATIONS TO TERMITE CARTON NESTS LOCATED IN ABOVE GROUND WALL VOIDS

Apply a 0.05% to 0 10% solution of PROTHOR WP directly into above ground lermite carton nests including nests tocated in wall volds using a directional injector. Apply as a solution or foam under pressure to distribute solution thoroughly throughout the nest. It may be necessary to inject solution at one or more points and at varying depths within the nest to adequately distribute solution within the interfor of the nest.

EXTERIOR APPLICATION FOR ANT CONTROL

Apply a 0.05% to 0.10% solution of PROTHOR WP to the exterior of the structure as a general surface, spot, crack and crevice or wall void treatment. Apply at points where nits may enter the structure or crawl and hide including exterior surfaces, around doors and windows, under eaves, attic and foundation vents, utility entrances and cracks in the surface of the structure. Spray solution or foam into voids where ants or their nests are present. Apply a volume of solution sufficient to cover the target surface(s) however avoid excess dripping or runoff from vertical or overhead surfaces.

Treat soil, turf or ground cover (flower, shrub and plant beds) adjacent to the structure where ants are trailing or may find food. Ants tunneling in the soil may be controlled by applying a 0.05% to 0.10% solution of PROTHOR WP as a drench or soil injection along the edge of foundations or other hard surfaces such as driveways. Apply in a volume sufficient to treat or cover the soil or foliage.

Inject a 0.05% to 0.10% solution of PROTHOR WP in the form of a spray or foam into tree cavities or other parts of trees where ant nests are located.

Do not treat more often than once per month. Do not allow residents or pets into the immediate area during application or allow them to make contact with treated areas until spray has dried.

It is recommended to remove or prune away shrubbery, bushes and tree branches touching the structure. Vegetation touching the structure may offer a route of entry for ants into the structure that allow ants to inhabit the structure without coming in contact with the treatment. If nests are found, direct treatment of nests with PROTHOR WP can be made.

Do not use PROTHOR WP against native fire ants, imported fire ants, pharaoh ants or harvester ants. Limit applications for control of carpenter ants to treatment of non-wooden parts or surfaces of structures.

ATTENTION

Do not apply to soil in areas where edible plants may be planted. Do not plant edible plants in soil that has been treated with PROTHOR WP.

IMPORTANT READ BEFORE USE

NOTICE: Read the entire Directions for Use, Conditions of Sale, Discletimer of Warranties and Limitations of Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

CONDITIONS OF SALE: The Directions for Use of this product are believed to be adequate and should be followed carefully. However, because of manner of use and other factors beyond the control of Ensystex III, how, it is impossible for Ensystex III to eliminate all inteks associated with the use of this product such as ineffectiveness or unintended consequences. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Ensystex III formises for any claims relations to such as factors.

DISCLAIMER OF WARRANTIES: Selier warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the Directions for Use under namel conditions of use. ENSYSTEX Itil MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. Any warrantiles, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, under abnormal conditions or under conditions not reasonably foreseeable by (or beyond the control of) selier or Ensystex III, Inc., and buyer assumes the risk of any such use.

LIMITATIONS OF LIABILITY: To the extent permitted by law, Ensystex III shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ENSYSTEX III AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ENSYSTEX III, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

PROTHOR is a registered trademark of Ensystex III, Inc. Revised 2/07