



**UNITED STATES ENVIRONMENTAL PROTECTION AGENCY**  
WASHINGTON, DC 20460

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

Rotam Agrochemical Company Limited  
% Frank E. Sobotka  
IPM Resources LLC  
660 Newtown-Yardley Road, Suite 105  
Newtown, PA 18940

AUG 7 2008

Subject: Rotam Tebuconazole 3.6F  
EPA Reg. No. 83100-1  
Your amendment dated May 13, 2008  
EPA Decision Number 394557

Dear Dr. Sobotka:

The supplemental label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act as amended is acceptable provided the following changes are made:

1. Add the following text to the supplemental label:

**"ENVIRONMENTAL HAZARDS**

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

**Ground Water Advisory:** Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

**Surface Water Advisory:** This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours."

One copy of the supplemental label stamped "Accepted with comments" is enclosed for your records. Please submit one copy of the final printed label that incorporates the required changes before the product is released for shipment.

# Rotam Tebuconazole 3.6F

**FUNGICIDE**

EPA Reg. No. 83100-1

## SUPPLEMENTAL LABEL

For Use on Wheat and Barley

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of application. Read the label affixed to the container for Rotam Tebuconazole 3.6F before applying. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the registered product label for Rotam Tebuconazole 3.6F.

### USE DIRECTIONS FOR WHEAT AND BARLEY

Crop	Disease	Rate of Rotam Tebuconazole 3.6F
Barley	Rusts ( <i>Puccinia spp.</i> ) Head blight ( <i>Fusarium spp.</i> ) Suppression	4 fl. oz. per acre
	<b>APPLICATION DIRECTIONS</b> Apply Rotam Tebuconazole 3.6F in a minimum of 10 gallons of spray solution per acre by ground or in a minimum of 5 gallons of spray solution per acre by air. A maximum of 4 fl. oz. of Rotam Tebuconazole 3.6F may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw cut after harvest may be fed or used for bedding. Grazing livestock or feeding of green forage is permitted 6 or more days after the last application of Rotam Tebuconazole 3.6F. Barley fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development.  <u>Rusts:</u> Apply Rotam Tebuconazole 3.6F at the earliest sign of rust pustules on foliage. <u>Fusarium head blight:</u> Optimal timing of Rotam Tebuconazole 3.6F for Fusarium head blight suppression is when main stem heads have fully emerged (Feekes 10.5) on 50% of the plants.	

**NOTE:** For optimum disease control, the lowest specified rate of a spray surfactant should be tank-mixed with Rotam Tebuconazole 3.6F. Rotam Tebuconazole 3.6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Rotam Tebuconazole 3.6F will be resistant to weathering.

ACCEPTED  
with COMMENTS  
In EPA Letter Dated

AUG 7 2008

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No. 83100-1

Crop	Disease	Rate of Rotam Tebuconazole 3.6F
Wheat	Rusts leaf, stem, and stripe ( <i>Puccinia spp.</i> )  Head blight or scab ( <i>Fusarium spp.</i> ) – Suppression	4 fl. oz. per acre
	<p><b>APPLICATION DIRECTIONS</b> Wheat fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. A maximum of 4 fl. oz. of Rotam Tebuconazole 3.6F may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw may be fed or used for bedding. Do not allow livestock to graze or feed green forage to livestock prior to 6 days after treatment with Rotam Tebuconazole 3.6F. Apply Rotam Tebuconazole 3.6F in a minimum of 10 gallons of spray solution per acre by ground, or in a minimum of 5 gallons of spray solution per acre by air.</p> <p><u>Rusts:</u> Apply Rotam Tebuconazole 3.6F at the earliest sign of rust pustules on foliage. <u>Fusarium head blight:</u> Optimal timing of Rotam Tebuconazole 3.6F for Fusarium head blight suppression is the beginning of flowering on main stem heads (Feekes 10.51).</p>	
<p><b>NOTE:</b> For optimum disease control, the lowest specified rate of a spray surfactant should be tank mixed with Rotam Tebuconazole 3.6F. Rotam Tebuconazole 3.6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Rotam Tebuconazole 3.6F will be resistant to weathering.</p>		

Manufactured by:  
Rotam Agrochemicals Company Limited  
660 Newtown-Yardley Rd.  
Newtown, PA 18940

Registered:

[Directions for use booklet]

# Rotam Tebuconazole 3.6F

## Fungicide

FOR CONTROL OF SPECIFIC DISEASES ON GRASSES GROWN FOR SEED, BARLEY, PEANUTS AND WHEAT.

### ACTIVE INGREDIENT:

Tebuconazole*	38.7%
Other Ingredients:	61.3%
<b>Total:</b>	<b>100.0%</b>

Contains 3.6 pounds tebuconazole per gallon

\*alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-H-1,2,4-triazole-1-ethanol  
(CASRN: 107534-96-3)

EPA Reg. No. 83100 - 1

EPA Est. No.: 069821-CHN-005

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

ACCEPTED  
with COMMENTS  
In EPA Letter Dated

AUG 7 2008

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 83100-1

FIRST AID	
If Swallowed	<ul style="list-style-type: none"> <li>• Call poison control center or doctor for treatment advice</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
If On Skin Or Clothing	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
If In Eyes	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
If Inhaled	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call poison control center or doctor for treatment advice.</li> </ul>
<p align="center"><b>Note to Physician</b></p> <p><b>Symptoms of Poisoning and Recommendations for Medical Treatment:</b> The compound does not cause any definite symptoms that would be diagnostic. No specific antidote. Treat symptomatically.</p>	
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. For Medical Emergency Treatment Call your Local Emergency Response Center.</p>	

Manufactured by:  
Rotam Agrochemical Company, Ltd.  
7/F Cheung Tat Center  
No. 18, Cheung Lee Street  
Chai Wan, Hong Kong

Net Contents:  
1.0 GALLON (3.78 Liters)

# TABLE OF CONTENTS

[TO BE ADDED]

## RECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist.

#### Personal Protective Equipment

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category C on an EPA chemical resistance category selection chart.

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### User Safety Recommendations

##### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to estuarine and marine invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water or rinsate.

## DIRECTIONS FOR USE

**It is a violation of federal law to use this product in a manner inconsistent with its labeling.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements on this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton)
- Shoes plus socks

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

**Storage:** Store in a tightly closed container in a cool, dry place.

**Pesticide Disposal:** Pesticide spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Container Disposal (Nonrefillable container 5 gallons or less):** Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Offer for recycling, if available.

**Residue Removal:** Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call CHEMTREC AT 1-800-424-9300**

## GENERAL DIRECTIONS

Rotam Tebuconazole 3.6F is a soluble concentrate which will control certain pests on the crops listed on this label when applied according to the **Directions for Use**.

**Spray Volume:** Rotam Tebuconazole 3.6F may be applied in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied. If necessary, increase the spray volume per acre for complete crop coverage.



**Chemigation:** Do not apply this product through any type of irrigation system.

**Mixing:** Add recommended amount of Rotam Tebuconazole 3.6F into the spray tank while filling with water to the desired level. Operate the agitator while mixing. If other materials are added to the spray tank, the Rotam Tebuconazole 3.6F should be thoroughly dispersed prior to the addition of other materials.

**Compatibility:** Rotam Tebuconazole 3.6F has been tested for phytotoxicity and has a wide margin of safety on a variety of crops. Rotam Tebuconazole 3.6F has also been shown to be compatible with many commonly used pesticides, crop oils, and nutritional sprays. However, since it is not possible to test a large number of possible mixtures, the user should pre-test to assure the physical compatibility and lack of phytotoxicity effect of any proposed mixtures with Rotam Tebuconazole 3.6F.

To determine the compatibility of Rotam Tebuconazole 3.6F with other products, the following procedure should be followed: Pour the recommended proportions of the products into a suitable container of water, mix thoroughly and allow to stand at least five (5) minutes. If the combination remains mixed or can be re-mixed readily, the mixture is considered physically compatible.

### Application Rate Table

#### GRASSES GROWN FOR SEED

CROP	DISEASE	RATE OF ROTAM TEBUCONAZOLE 3.6F
Grasses Grown for Seed	Rusts ( <i>Puccinia spp.</i> )	4 to 8 fl oz per acre
	Apply the specified rate of Rotam Tebuconazole 3.6F as soon as weather conditions are favorable for rust development or when first rust pustules are present. Repeat applications at 14 to 16 day intervals. Under heavy disease pressure use 6 to 8 fl oz/A and shorter spray intervals.	
	Powdery mildew	4 to 8 fl oz per acre
	Apply specified rate of Rotam Tebuconazole 3.6F when powdery mildew first appears on the leaves. Repeat applications at 14 to 16 day intervals. Under heavy disease pressure use 6 to 8 fl oz/A and shorter intervals.	
	<p>For optimum benefit, the lowest recommended rate of a spray surfactant should be tank mixed with Rotam Tebuconazole 3.6F.</p> <p><b>Use Directions:</b> Apply the specified rate in a minimum of 20 gallons of water per acre with ground sprayers or in a minimum of 10 gallons of water per acre with aircraft. Thorough coverage is important for optimum disease control.</p>	

	<p><b>Restrictions:</b> A maximum of 16 fluid ounces (1 pint) may be applied per acre per crop season.</p> <p><b>Preharvest Interval (PHI):</b> Rotam Tebuconazole 3.6F may be applied up to 4 days before harvest.</p> <p>Chaff, screenings and straw from treated areas may be used for feed purposes; however, do not forage, cut green crop, or use seed for feed purposes. Regrowth may be grazed starting 17 days after last application.</p>
--	---

**BARLEY**

Crop	Disease	Rate of Rotam Tebuconazole 3.6F
Barley	Rusts ( <i>Puccinia spp.</i> ) Head blight ( <i>Fusarium spp.</i> ) Suppression	4 fl. oz. per acre
	<p><b>APPLICATION DIRECTIONS</b> Apply Rotam Tebuconazole 3.6F in a minimum of 10 gallons of spray solution per acre by ground or in a minimum of 5 gallons of spray solution per acre by air. A maximum of 4 fl. oz. of Rotam Tebuconazole 3.6F may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw cut after harvest may be fed or used for bedding. Grazing livestock or feeding of green forage is permitted 6 or more days after the last application of Rotam Tebuconazole 3.6F. Barley fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development.</p> <p><u>Rusts:</u> Apply Rotam Tebuconazole 3.6F at the earliest sign of rust pustules on foliage.</p> <p><u>Fusarium head blight:</u> Optimal timing of Rotam Tebuconazole 3.6F for Fusarium head blight suppression is when main stem heads have fully emerged (Feekes 10.5) on 50% of the plants.</p>	
<p><b>NOTE:</b> For optimum disease control, the lowest specified rate of a spray surfactant should be tank-mixed with Rotam Tebuconazole 3.6F. Rotam Tebuconazole 3.6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Rotam Tebuconazole 3.6F will be resistant to weathering.</p>		

# PEANUT

CROP	DISEASE	RATE OF ROTAM TEBUCONAZOLE 3.6F FUNGICIDE
Peanut	<b>SOIL BORNE:</b> Sclerotium stem and pod rot (white mold, southern blight, southern stem rot) Rhizoctonia limb rot Rhizoctonia pod rot (Virginia and North Carolina only) <b>FOLIAR:</b> Early leaf spot Late leaf spot Leaf rust Web blotch (Phoma) Pepper spot (Leptosphaerulina)	7.2 fl oz per acre
<p><b>FOUR-APPLICATION SPRAY PROGRAM:</b> Apply the specified rate in a preventive spray schedule. (*See table below for proper timing of applications) Applications of chlorothalonil should be made prior to and following applications of Rotam Tebuconazole 3.6F to discourage development of resistant strains of fungi. For optimum control of foliar diseases such as leaf rust, web blotch, and pepper spot, the lowest label recommended rate of a spray surfactant should be tank-mixed with Rotam Tebuconazole 3.6F.</p> <p><b>LEAF SPOT ADVISORY SCHEDULE:</b> For control of soil borne diseases in an advisory schedule, apply Rotam Tebuconazole 3.6F in the first advisory spray in July and continue Rotam Tebuconazole 3.6F applications at 14-day intervals. Applications after August 15 should be tank mixed with chlorothalonil for resistance management purposes.</p>		

Rotam Tebuconazole 3.6F is a sterol demethylation inhibitor (DMI) fungicide. Chlorothalonil may be tank mixed at the rate of 12 ounces of active ingredient with Rotam Tebuconazole 3.6F as a leaf spot resistance management strategy. A spray surfactant is not necessary when Rotam Tebuconazole 3.6F is tank mixed with chlorothalonil. Mixing or alternating Rotam Tebuconazole 3.6F with other DMI fungicides may lead to resistance.

**Use Directions:** For optimum control of the specified soilborne diseases, four consecutive applications of Rotam Tebuconazole 3.6F must be made at 14-day intervals. Rotam Tebuconazole 3.6F must be carried by rainfall or irrigation into the root and pod zone for control of root and pod rots caused by *Sclerotium rolfsii* and *Rhizoctonia solani*. Drought conditions will decrease the effectiveness of Rotam Tebuconazole 3.6F against the root and pod rots. Use Rotam Tebuconazole 3.6F in conjunction with cultural practices that are known to reduce the severity of soil borne diseases, such as proper crop rotation practices.

**Restrictions:** Do not apply more than 28.8 fluid ounces of Rotam Tebuconazole 3.6F per acre per crop season.

**Preharvest Interval (PHI):** Rotam Tebuconazole 3.6F may be applied up to 14 days before harvest.

Do not feed hay or threshings or allow livestock to graze in treated areas.

**\*Timing of Rotam Tebuconazole 3.6F Application for Optimum Control of White Mold and Rhizoctonia Limb and Pod Rot**

Spray Program	Rotam Tebuconazole 3.6F Application No.	Chlorothalonil Application No.
7 applications	3, 4, 5 & 6	1, 2 & 7

**WHEAT**

Crop	Disease	Rate of Rotam Tebuconazole 3.6F
Wheat	<p>Rusts leaf, stem, and stripe (<i>Puccinia spp.</i>)</p> <p>Head blight or scab (<i>Fusarium spp.</i>) – Suppression</p>	4 fl. oz. per acre
<p><b>APPLICATION DIRECTIONS</b></p> <p>Wheat fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. A maximum of 4 fl. oz. of Rotam Tebuconazole 3.6F may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw may be fed or used for bedding. Do not allow livestock to graze or feed green forage to livestock prior to 6 days after treatment with Rotam Tebuconazole 3.6F. Apply Rotam Tebuconazole 3.6F in a minimum of 10 gallons of spray solution per acre by ground, or in a minimum of 5 gallons of spray solution per acre by air.</p> <p><u>Rusts:</u> Apply Rotam Tebuconazole 3.6F at the earliest sign of rust pustules on foliage.</p> <p><u>Fusarium head blight:</u> Optimal timing of Rotam Tebuconazole 3.6F for Fusarium head blight suppression is the beginning of flowering on main stem heads (Feekes 10.51).</p>		
<p><b>NOTE:</b> For optimum disease control, the lowest specified rate of a spray surfactant should be tank mixed with Rotam Tebuconazole 3.6F. Rotam Tebuconazole 3.6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Rotam Tebuconazole 3.6F will be resistant to weathering.</p>		

**OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.**

Apply only during alternate years in fields adjacent to aquatic areas listed above.

Do not apply by ground or air within 100 feet of aquatic areas listed above.

Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip.

Spray Drift Management: For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter.

Use the largest droplet size consistent with pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure. Apply in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment.

Spray should be released at the lowest possible height consistent with good pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

Make aerial or ground applications when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.

Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

**ROTATIONAL CROPS**

Treated areas may be replanted with any crop specified on this label as soon as practical after last application. Any crop not specified on this label may be planted into treated areas 120 days after last application.

15/1

## CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of ROTAM AGROCHEMICAL COMPANY LIMITED or Seller. To the extent allowed by applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold ROTAM AGROCHEMICAL COMPANY LIMITED and Seller harmless for any claims relating to such factors.

ROTAM AGROCHEMICAL COMPANY LIMITED warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or ROTAM AGROCHEMICAL COMPANY LIMITED, and to the extent allowed by applicable law, Buyer and User assume the risk of any such use. **ROTAM USA LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

To the extent allowed by law, ROTAM AGROCHEMICAL COMPANY LIMITED or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT ALLOWED BY LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ROTAM AGROCHEMICAL COMPANY LIMITED AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ROTAM AGROCHEMICAL COMPANY LIMITED OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

ROTAM AGROCHEMICAL COMPANY LIMITED and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of ROTAM AGROCHEMICAL COMPANY LIMITED.

Registered: [Date]

16/18

[Base Container Label]

# Rotam Tebuconazole 3.6F

## **Fungicide**

**FOR CONTROL OF SPECIFIC DISEASES ON GRASSES GROWN FOR SEED, BARLEY, PEANUTS AND WHEAT.**

**ACTIVE INGREDIENT:**

Tebuconazole*	38.7%
Other Ingredients:	61.3%
<b>Total:</b>	<b>100.0%</b>

Contains 3.6 pounds tebuconazole per gallon

\*alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-H-1,2,4-triazole-1-ethanol  
(CASRN: 107534-96-3)

EPA Reg. No. 83100 - 1

EPA Est. No.: 069821-CHN-005

## **KEEP OUT OF REACH OF CHILDREN CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

<b>FIRST AID</b>	
If Swallowed	<ul style="list-style-type: none"> <li>• Call poison control center or doctor for treatment advice</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
If On Skin Or Clothing	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
If In Eyes	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
If Inhaled	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call poison control center or doctor for treatment advice.</li> </ul>
<b>Note to Physician</b>	
<p><b>Symptoms of Poisoning and Recommendations for Medical Treatment:</b> The compound does not cause any definite symptoms that would be diagnostic. No specific antidote. Treat symptomatically.</p>	
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. For Medical Emergency Treatment Call your Local Emergency Response Center.</p>	

Manufactured by:  
Rotam Agrochemical Company, Ltd.  
7/F Cheung Tat Center  
No. 18, Cheung Lee Street  
Chai Wan, Hong Kong

Net Contents:  
1.0 GALLON (3.78 Liters)



## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to labeling under "AGRICULTURAL USE REQUIREMENTS" in the DIRECTIONS FOR USE section for information about this standard.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to estuarine and marine invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water or rinsate.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

**Storage:** Store in a tightly closed container in a cool, dry place.

**Pesticide Disposal:** Pesticide spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Container Disposal (Nonrefillable container 5 gallons or less):** Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Offer for recycling, if available.

**Residue Removal:** Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call CHEMTREC AT 1-800-424-9300**

### DIRECTIONS FOR USE

Refer to attached booklet for complete Precautionary Statements and Directions For Use.