

83100-1

05/05/2009

1/36



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Rotam Agrochemical Company Limited
% Frank E. Sobotka
IPM Resources LLC
660 Newtown-Yardley Road, Suite 105
Newtown, PA 18940

MAY 5 2009

Subject: Rotam Tebuconazole 3.6F
EPA Reg. No. 83100-1
Your amendment dated November 3, 2008
EPA Decision Number 403105

Dear Dr. Sobotka:

The revised amended label referred to above, submitted March 31, 2009 in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act as amended is acceptable provided the following changes are made:

1. On page 4 in the **AGRICULTURAL USE REQUIREMENTS** box change "...interval (REI) of 12 hours." to "...interval (REI). See specific crop directions for applicable REI."
2. On page 6 in the paragraph starting **Mixing** change "Add recommended amount...." to "Add specified amount...."
3. On page 10 move the Pre-harvest Interval instructions for "Beans (Fresh)...7 days before harvest" and "Beans (dry)...14 days before harvest" to the bottom of the table below the header **RESTRICTIONS**.
4. On page 14 in the first column move the word "Welch" to the next line so it reads "Welch Onion"
5. On page 23 under **RESTRICTIONS** add "Restricted-entry Interval (REI) = 12 hours"
6. On page 24 in the first paragraph of the **Applications Directions** change "Rotam Tebuconazole 3.6F should be applied at...." to "Apply Rotam Tebuconazole 3.6F at...."
7. On page 29 in the instructions for **SEED LABELING** put quotation marks around "TREATED SEED. DO...Tebuconazole"
8. On page 29 in the **Application Directions** change "Product should be diluted with sufficient water to ensure complete seed coverage." to "Dilute product with sufficient water to ensure complete seed coverage. Add dye to resulting slurry."

MAY 5 2009


2
36

9. On page 29 in the Rate column for Seed Treatment, change the rate for Fusarium from 0.071 fl oz/CWT to 0.055 fl oz/CWT.

One copy of the label stamped "Accepted with comments" is enclosed for your records. This label supersedes all labels previously accepted for this product. Please submit one copy of the final printed label that incorporates the required changes before the product is released for shipment.

If you have any questions, please contact Robert Westin by phone at (703) 305-5721 or via email at westin.robert@epa.gov.

Sincerely,



Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7505P)

Enclosure

[Directions for use booklet]
Rotam Tebuconazole 3.6F

3
 36

Agricultural Fungicide

FOR CONTROL OF SPECIFIC DISEASES ON VARIOUS CROPS

ACTIVE INGREDIENT:

Tebuconazole*.....	38.7%
Other Ingredients:.....	61.3%
Total:.....	100.0%

Contains 3.6 pounds tebuconazole per gallon

*alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-H-1,2,4-triazole-1-ethanol with **COMMENTS**
 (CASRN: 107534-96-3) **In EPA Letter Dated**

EPA Reg. No. 83100 -1

EPA Est. No.: 069821-CHN-005

ACCEPTED
MAY 5 2009

**Under the Federal Insecticide,
 Fungicide, and Rodenticide Act
 as amended, for the pesticide
 registered under EPA Reg. No.**

**KEEP OUT OF REACH OF CHILDREN
 CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

83100-1

FIRST AID	
If Swallowed	<ul style="list-style-type: none"> • Call poison control center or doctor for treatment advice • Do not induce vomiting unless told to do so by the poison control center or doctor. • Have person sip a glass of water if able to swallow. • Do not give anything by mouth to an unconscious person.
If On Skin Or Clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for further treatment advice.
If In Eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for further treatment advice.
If Inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call poison control center or doctor for treatment advice.
Note to Physician	
<p>Symptoms of Poisoning and Recommendations for Medical Treatment: The compound does not cause any definite symptoms that would be diagnostic. No specific antidote. Treat symptomatically.</p>	
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. For Medical Emergency Treatment Call your Local Emergency Response Center.</p>	

See Additional Precautionary Statements and Directions for Use Inside Booklet.

Manufactured by:
 Rotam Agrochemical Company, Ltd.
 7/F Cheung Tat Center
 No. 18, Cheung Lee Street
 Chai Wan, Hong Kong

Net Contents:
 1.0 GALLON (3.78 Liters)

PECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist.

Personal Protective Equipment

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category C on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

6
36

Ground Water Advisory: Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements on this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton)
- Shoes plus socks

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

Pesticide Storage: Store in a tightly closed container in a cool, dry place.

Pesticide Disposal: Pesticide spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal (Nonrefillable container 5 gallons or less): Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Offer for recycling, if available.

Residue Removal: Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call CHEMTREC AT 1-800-424-9300

GENERAL DIRECTIONS

Rotam Tebuconazole 3.6F is a soluble concentrate which will control certain pests on the crops listed on this label when applied according to the **Directions for Use**.

Spray Volume: Rotam Tebuconazole 3.6F may be applied in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied. If necessary, increase

the spray volume per acre for complete crop coverage.

Chemigation: Apply Rotam Tebuconazole 3.6F through irrigation equipment only to crops and diseases for which the chemigation use is specified. Apply this product only through center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move; or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally dosed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. Pesticide may be applied continuously for the duration of the water application.

Mixing: Add recommended amount of Rotam Tebuconazole 3.6F into the spray tank while filling with water to the desired level. Operate the agitator while mixing. If other materials are added to the spray tank, the Rotam Tebuconazole 3.6F should be thoroughly dispersed prior to the addition of other materials.

Compatibility: Rotam Tebuconazole 3.6F has been tested for phytotoxicity and has a wide margin of safety on a variety of crops. Rotam Tebuconazole 3.6F has also been shown to be compatible with many commonly used pesticides, crop oils, and nutritional sprays. However, since it is not possible to test a large number of possible mixtures, the user should pre-test to assure the physical compatibility and lack of phytotoxicity effect of any proposed mixtures with Rotam Tebuconazole 3.6F.

To determine the compatibility of Rotam Tebuconazole 3.6F with other products, the following procedure should be followed: Pour the recommended proportions of the products into a suitable container of water, mix thoroughly and allow to

stand at least five (5) minutes. If the combination remains mixed or can be re-mixed readily, the mixture is considered physically compatible.

9
—
36

Resistance Management

Rotam Tebuconazole 3.6F is a Group 3 fungicide which exhibits no known cross-resistance to other fungicide groups. However, fungal pathogens are known to develop resistance to products with the same mode of action when used repeatedly. Any fungal population may contain or develop individuals that are resistant to Rotam Tebuconazole 3.6F and other Group 3 fungicides. If Group 3 fungicides are used repeatedly in the same field or in successive years as the primary method of control for targeted diseases, the resistant isolates may eventually dominate the fungal population. Because resistance development cannot be predicted, the use of this product should conform to resistance management strategies established for the crop and use area. Such strategies may include rotation and /or tank mixing with products having different modes of action or limiting the total number of applications per season. Contact your local extension specialist, certified crop advisor, and/or manufacturer for fungicide resistance management and/or integrated disease management recommendations for specific crops and resistant disease populations. Rotam encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label.

Application Rate Table

Crop	Disease	Rate of Rotam Tebuconazole 3.6F
Asparagus	Rust (<i>Puccinia</i> spp.)	4 to 6 fl. oz. per acre
	<p>Application Directions: Apply Rotam Tebuconazole 3.6F as a foliar spray to the developing ferns after harvest of spears is completed. Apply at the earliest sign of rust pustules or when weather conditions are conducive for rust development. Apply 4 to 6 fl oz of Rotam Tebuconazole 3.6F per acre (0.11 lb ai – 0.17 lb ai per acre) in alternation with another effective fungicide. Under conditions of severe rust pressure, use the higher rate. Repeat applications on a 14-day interval as necessary to maintain control of rust.</p> <p>Applications may be made using ground or aerial application equipment. A 50 foot spray drift buffer zone is required for all aerial applications. For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Rotam Tebuconazole 3.6F.</p>	
<p>Restricted-entry interval (REI) = 12 hours.</p> <p>RESTRICTION: Do not apply within 100 days of harvest in California and 180 days in all other states. Do not make more than three foliar applications per season (18 fl oz/acre or 0.51 lb ai/acre). Do not apply to harvestable spears.</p>		

11
36

Crop	Disease	Rate of Rotam Tebuconazole 3.6F
Barley	Rusts (<i>Puccinia spp.</i>) Head blight (<i>Fusarium spp.</i>) - Suppression	4 fl. oz. per acre
	<p>Application Directions: Apply Rotam Tebuconazole 3.6F in a minimum of 10 gallons of spray solution per acre by ground or in a minimum of 5 gallons of spray solution per acre by air. Barley fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development.</p> <p>Application Timing:</p> <p><u>Rusts:</u> Apply Rotam Tebuconazole 3.6F at the earliest sign of rust pustules on foliage.</p> <p><u>Fusarium head blight:</u> Optimal timing of Rotam Tebuconazole 3.6F for Fusarium head blight suppression is when main stem heads have fully emerged (Feekes 10.5) on 50% of the plants.</p> <p>NOTE: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Rotam Tebuconazole 3.6F. Rotam Tebuconazole 3.6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Rotam Tebuconazole 3.6F will be resistant to weathering.</p>	
<p>Restricted-entry interval (REI) = 12 hours</p> <p>RESTRICTIONS: A maximum of 4 fl. oz. of Rotam Tebuconazole 3.6F may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw cut after harvest may be fed or used for bedding. Grazing livestock or feeding of green forage is permitted 6 or more days after the last application of Rotam Tebuconazole 3.6F.</p>		

12
36

Crop	Disease	Rate of Rotam Tebuconazole 3.6F
Beans (fresh and dried except succulent shelled)	Rusts (<i>Uromyces appendiculatus</i>)	4 to 6 fl. oz. per acre
	<p>Application Directions: Apply Rotam Tebuconazole 3.6F in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 14-day intervals, or as necessary to maintain control.</p> <p><u>Beans (fresh):</u> Rotam Tebuconazole 3.6F may be applied up to 7 days before harvest.</p> <p><u>Beans (dry):</u> Rotam Tebuconazole 3.6F may be applied up to 14 days before harvest.</p> <p>For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Rotam Tebuconazole 3.6F. Rotam Tebuconazole 3.6F must have two to four hours of drying time on bean foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Rotam Tebuconazole 3.6F will be resistant to weathering.</p>	
<p>Restricted-entry interval (REI) = 12 hours</p> <p>RESTRICTIONS:</p> <p><u>Beans (fresh):</u> Do not apply more than 24 fl. oz. of Rotam Tebuconazole 3.6F per acre per crop season.</p> <p><u>Beans (dry):</u> Do not apply more than 12 fl. oz. of Rotam Tebuconazole 3.6F per acre per crop season.</p>		

13
36

Crop	Disease	Rate of Rotam Tebuconazole 3.6F
<p>Corn (sweet corn, field corn, field corn grown for seed, and popcorn)</p>	<p>Rust (<i>Puccinia spp.</i>) Northern leaf blight (<i>Helminthosporium turcicum</i>) Southern leaf blight (<i>Helminthosporium maydis</i>) Northern leaf spot (<i>Helminthosporium carbonum</i>) Gray leaf spot (<i>Cercospora zea-maydis</i>)</p>	<p>4 to 6 fl. oz. per acre</p>
<p>Application Directions: Apply Rotam Tebuconazole 3.6F in a protective spray schedule or when weather conditions are favorable for disease development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control.</p> <p><u>Sweet corn:</u> Rotam Tebuconazole 3.6F may be applied up to 7 days before the harvest of ears or forage, and 49 days before the harvest of fodder.</p> <p><u>Field, seed or popcorn:</u> Rotam Tebuconazole 3.6F may be applied up to 21 days before the harvest of forage, and 36 days before the harvest of grain or fodder.</p> <p>For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Rotam Tebuconazole 3.6F. Rotam Tebuconazole 3.6F must have two to four hours of drying time on corn foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Rotam Tebuconazole 3.6F will be resistant to weathering.</p>		
<p>Restricted-entry interval (REI) for sweet corn = 19 days. Restricted-entry interval (REI) for all corn except sweet corn = 12 hours.</p> <p>RESTRICTIONS: Do not apply more than 24 fl. oz. of Rotam Tebuconazole 3.6F per acre per crop season.</p>		

14
38

Crop	Disease	Rate of Rotam Tebuconazole 3.6F
Cotton	Southwestern cotton rust (<i>Puccinia cacabata</i>)	6 to 8 fl. oz. per acre
	<p>Application Directions: Apply Rotam Tebuconazole 3.6F in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. Rotam Tebuconazole 3.6F may be applied up to 30 days before harvest.</p> <p>For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Rotam Tebuconazole 3.6F. Rotam Tebuconazole 3.6F must have two to four hours of drying time on cotton foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Rotam Tebuconazole 3.6F will be resistant to weathering.</p>	
<p>Restricted-entry interval (REI) = 12 hours.</p> <p>RESTRICTIONS: Do not apply more than 24 fl. oz. of Rotam Tebuconazole 3.6F per acre per crop season.</p>		

15
36

Crop	Disease	Rate of Rotam Tebuconazole 3.6F
Cucurbit Vegetables Group Chayote Chinese waxgourd Citron melon Cucumber Gherkin	Powdery mildew (<i>Sphaerotheca fuliginea</i> / <i>Podosphaera xanthii</i>) (<i>Erysiphe cichoracearum</i>)	4 to 6 fl. oz. per acre
Edible gourd, (includes hyotan, cucuzza, hechima and Chinese okra) Momordica spp. (includes balsam apple, balsam pear, bitter melon and Chinese cucumber)	Gummy stem blight - suppression (<i>Didymella toyonae</i>) (watermelon, squash, pumpkin, and melons only)	8 fl. oz. per acre
Muskmelon (includes cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon and snake melon) Pumpkin Summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow and zucchini) Winter squash (includes butternut squash, calabaza, hubbard squash, acorn squash and spaghetti squash) Watermelon	<p>Application Directions: Apply the specified dosage in a protective spray schedule to foliage and fruit. Repeat applications at 10- to 14-day intervals. Rotam Tebuconazole 3.6F may be applied up to 7 days before harvest. Do not apply more than 24 fl. oz. of Rotam Tebuconazole 3.6F per acre per crop season.</p> <p>For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Rotam Tebuconazole 3.6F F. Rotam Tebuconazole 3.6F must have two to four hours of drying time for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Rotam Tebuconazole 3.6F will be resistant to weathering.</p>	
<p>.Restricted-entry interval (REI) = 12 hours.</p>		
<p>RESTRICTIONS: Do not apply more than 24 fl. oz. of Rotam Tebuconazole 3.6F per acre per crop season.</p>		

16
36

Crop	Disease	Rate of Rotam Tebuconazole 3.6F
Dry bulb onion Garlic Great-headed (elephant) garlic Welch onion Shallot	White rot <i>(Sclerotium cepivorum)</i>	White rot: 20.5 fl oz per acre applied in a 4 to 6 inch band over/into each furrow. May be applied by chemigation to control white rot.
	Rust <i>(Puccinia aHii, Puccinia porri)</i> Purple blotch <i>(Alternaria porii)</i>	4 to 6 fl. oz per acre
	<p>White rot: For the control of white rot, make one application in the furrow at the time of planting. The in-furrow application should be made at the rate of 20.5 fl. oz Rotam Tebuconazole 3.6F per acre. Apply the entire per acre rate in a 4 to 6 inch band over/into each furrow. Additional control may be obtained by including two foliar applications at 4 to 6 fl oz/acre.</p> <p>Rust: For the control of rust make foliar applications at the rate of 4 to 6 fl. oz Rotam Tebuconazole 3.6F per acre per application. Repeat at an interval of 10 to 14 days. Apply Rotam Tebuconazole 3.6F in a protective spray schedule or when weather conditions are favorable for rust development.</p> <p>Application Directions: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Rotam Tebuconazole 3.6F. Rotam Tebuconazole 3.6F must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Rotam Tebuconazole 3.6F will be resistant to weathering.</p>	
<p>Restricted-entry interval (REI) = 12 hours.</p> <p>RESTRICTIONS: Do not apply more than 32.5 fl.oz. Rotam Tebuconazole 3.6F per acre per season if an in-furrow treatment is made. If Rotam Tebuconazole 3.6F is not applied as an in-furrow treatment then do not apply more than 12 fl.oz. Rotam Tebuconazole 3.6F per acre per season as a foliar spray. Do not apply within 7 days of harvest (PHI = 7 days).</p>		

17
36

Crop	Disease	Rate of Rotam Tebuconazole 3.6F
Grasses Grown for Seed	Rusts (<i>Puccinia spp.</i>)	4 to 8 fl oz per acre
	<p>Application Directions for Rusts: Apply the specified rate of Rotam Tebuconazole 3.6F as soon as weather conditions are favorable for rust development or when first rust pustules are present. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl oz/A and shorter spray intervals.</p>	
	Powdery mildew	4 to 8 fl oz per acre
	<p>Application Directions for Powdery mildew: Apply specified rate of Rotam Tebuconazole 3.6F when powdery mildew first appears on the leaves. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl oz/A and shorter spray intervals.</p> <p>Apply the specified rate in a minimum of 20 gallons of water per acre with ground sprayers or in a minimum of 10 gallons of water per acre with aircraft. Thorough coverage is important for optimum disease control.</p> <p>For optimum benefit, the lowest labeled rate of a spray surfactant should be tank mixed with Rotam Tebuconazole 3.6F.</p>	
<p>Restricted-entry interval (REI) = 12 hours.</p> <p>RESTRICTIONS: A maximum of 16 fluid ounces (1 pint) may be applied per acre per crop season. Rotam Tebuconazole 3.6F may be applied up to 4 days before harvest. Chaff, screenings and straw from treated areas may be used for feed purposes; however, do not forage, cut green crop, or use seed for feed purposes. Regrowth may be grazed starting 17 days after last application.</p>		

18
36

Crop	Disease	Rate of Rotam Tebuconazole 3.6F
Green onion Leek Spring onion Scallion Japanese bunching onion Green shallots Green eschalots	White rot <i>(Sclerotium cepivorum)</i> suppression only Rust <i>(Puccinia allii, Puccinia porri)</i> Purple blotch <i>(Alternaria porii)</i>	4 to 6 fl. oz per acre
<p>Application Directions: For the control of diseases make foliar applications using an interval of 10 to 14 days. Apply Rotam Tebuconazole 3.6F in a protective spray schedule or when weather conditions are favorable for rust development.</p> <p>For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest labeled rate of a spray surfactant may be tank-mixed with Rotam Tebuconazole 3.6F. Rotam Tebuconazole 3.6F must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Rotam Tebuconazole 3.6F will be resistant to weathering.</p>		
<p>Restricted-entry interval (REI) = 12 hours.</p> <p>RESTRICTIONS: Do not apply more than 24 fl. oz. Rotam Tebuconazole 3.6F per acre per season. Do not apply within 7 days of harvest (PHI = 7 days).</p>		

19
36

Crop	Disease	Rate of Rotam Tebuconazole 3.6F
Hops	Powdery mildew (<i>Sphaerotheca humuli</i> / <i>Sphaerotheca macularis</i>)	4 to 8 fl. oz. per acre
	<p>Application Directions: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Rotam Tebuconazole 3.6F. Rotam Tebuconazole 3.6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Rotam Tebuconazole 3.6F will be resistant to weathering.</p> <p>Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 10- to 14-day intervals. Rotam Tebuconazole 3.6F may be applied up to 14 days before harvest. Increase the spray volume and the application rate as vine growth increases during the season.</p>	
<p>Restricted-entry interval (REI) = 12 hours.</p> <p>RESTRICTIONS: Do not apply more than 32 fl. oz. of Rotam Tebuconazole 3.6F per acre per crop season.</p>		

Crop	Disease	Rate of Rotam Tebuconazole 3.6F
Leafy Brassica Greens Broccoli raab Chinese cabbage (bok choy) Collards Kale	Cercospora leaf spot (<i>Cercospora brassicicola</i>) Powdery mildew (<i>Erysiphe cruciferarum</i>) Alternaria leaf spot (<i>Altemaria brassicicola</i>)	3 to 4 fl. oz per acre
Mizuma Mustard greens Mustard spinach Rape greens Turnip greens	<p>Application Directions: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest labeled rate of a spray surfactant may be tank-mixed with Rotam Tebuconazole 3.6F. Rotam Tebuconazole 3.6F must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Rotam Tebuconazole 3.6F will be resistant to weathering.</p>	
<p>Restricted-entry interval (REI) = 12 hours.</p> <p>RESTRICTION: Do not apply more than 16 fl.oz. Rotam Tebuconazole 3.6F per acre per season. Do not apply within 7 days of harvest (PHI = 7 days).</p> <p>Application to turnip greens is limited to East of the Rockies</p>		

Crop	Disease	Rate of Rotam Tebuconazole 3.6F
Garden beet roots and tops (leaves)	Cercospora leaf spot (<i>Cercospora beticola</i>)	3 to 7.2 fl. oz per acre
	<p>Application Directions: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest labeled rate of a spray surfactant may be tank-mixed with Rotam Tebuconazole 3.6F. Rotam Tebuconazole 3.6F must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Rotam Tebuconazole 3.6F will be resistant to weathering. Make applications on a 14 day intervals.</p>	
<p>Restricted-entry interval (REI) = 12 hours.</p> <p>RESTRICTIONS: Do not apply more than 28.8 fl. oz. Rotam Tebuconazole 3.6F per acre per season. Do not apply within 7 days of harvest (PHI = 7 days).</p>		

22
36

Crop	Disease	Rate of Rotam Tebuconazole 3.6F
Lychee	Anthracnose (<i>Colletotrichum gloeosporioides</i>)	4 to 6 fl. oz. per acre
	<p>Application Directions: For optimum disease control, the lowest labeled rate of a non-ionic spray surfactant should be tank-mixed with Rotam Tebuconazole 3.6F. Rotam Tebuconazole 3.6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Rotam Tebuconazole 3.6F will be resistant to weathering.</p> <p>Begin first application of Rotam Tebuconazole 3.6F as panicle emerges. Spray up to 6 fl. oz. per acre every 10 days thereafter for a total of 8 sprays. Apply specified dosage in a minimum of 50 gallons of spray solution per acre by ground only.</p>	
<p>Restricted-entry interval (REI) = 2 days</p> <p>RESTRICTIONS: Do not apply more than 48 fl. oz. of Rotam Tebuconazole 3.6F per acre per season. Rotam Tebuconazole 3.6F can be applied up to and including the day of harvest (PHI = 0 days).</p>		

23
36

Crop	Disease	Rate of Rotam Tebuconazole 3.6F
Okra	Cercospora leaf spot (<i>Cercospora spp.</i>)	4 to 6 fl. oz. per acre
	<p>Application Directions: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Rotam Tebuconazole 3.6F. Rotam Tebuconazole 3.6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Rotam Tebuconazole 3.6F will be resistant to weathering. Apply specific dosage of Rotam Tebuconazole 3.6F in a preventative spray program. Use the highest rate when disease conditions are favorable and in areas where high disease pressure is expected. Applications may be repeated at 14-day intervals in order to maintain control of the disease. Apply specified dosage as a foliar spray in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air.</p>	
<p>Restricted-entry interval (REI) = 12 hours.</p> <p>RESTRICTIONS: Applications may be made no closer than 3 days before harvest. Do not apply more than 24 fl. oz. of Rotam Tebuconazole 3.6F per acre per season.</p>		

24
36

Crop	Disease	Rate of Rotam Tebuconazole 3.6F
Peanut	<p>SOIL BORNE: Sclerotium stem and pod rot (white mold, southern blight, southern stem rot) Rhizoctonia limb rot Rhizoctonia pod rot (Virginia and North Carolina only)</p> <p>FOLIAR: Early leaf spot Late leaf spot Leaf rust Web blotch (Phoma) Pepper spot (Leptosphaerulina)</p>	7.2 fl oz per acre
<p>FOUR-APPLICATION SPRAY PROGRAM: Apply the specified rate in a preventive spray schedule. (*See table below for proper timing of applications) Applications of chlorothalonil should be made prior to and following applications of Rotam Tebuconazole 3.6F to discourage development of resistant strains of fungi. For optimum control of foliar diseases such as leaf rust, web blotch, and pepper spot, the lowest label recommended rate of a spray surfactant should be tank-mixed with Rotam Tebuconazole 3.6F.</p> <p>LEAF SPOT ADVISORY SCHEDULE: For control of soil borne diseases in an advisory schedule, apply Rotam Tebuconazole 3.6F in the first advisory spray in July and continue Rotam Tebuconazole 3.6F applications at 14-day intervals. Applications after August 15 should be tank mixed with chlorothalonil for resistance management purposes.</p>		

25
36

Rotam Tebuconazole 3.6F is a sterol demethylation inhibitor (DMI) fungicide. Chlorothalonil may be tank mixed at the rate of 12 ounces of active ingredient with Rotam Tebuconazole 3.6F as a leaf spot resistance management strategy. A spray surfactant is not necessary when Rotam Tebuconazole 3.6F is tank mixed with chlorothalonil. Mixing or alternating Rotam Tebuconazole 3.6F with other DMI fungicides may lead to resistance.

Use Directions: For optimum control of the specified soilborne diseases, four consecutive applications of Rotam Tebuconazole 3.6F must be made at 14-day intervals. Rotam Tebuconazole 3.6F must be carried by rainfall or irrigation into the root and pod zone for control of root and pod rots caused by *Sclerotium rolfsii* and *Rhizoctonia solani*. Drought conditions will decrease the effectiveness of Rotam Tebuconazole 3.6F against the root and pod rots. Use Rotam Tebuconazole 3.6F in conjunction with cultural practices that are known to reduce the severity of soil borne diseases, such as proper crop rotation practices.

RESTRICTIONS: Do not apply more than 28.8 fluid ounces of Rotam Tebuconazole 3.6F per acre per crop season. Do not feed hay or threshings or allow livestock to graze in treated areas.

Preharvest Interval (PHI): Rotam Tebuconazole 3.6F may be applied up to 14 days before harvest.

Timing of Rotam Tebuconazole 3.6F Application for Optimum Control of White Mold and Rhizoctonia Limb and Pod Rot

Spray Program	Rotam Tebuconazole 3.6F Application No.	Chlorothalonil Application No.
7 applications	3, 4, 5 & 6	1, 2 & 7

26
36

Crop	Disease	Rate of Rotam Tebuconazole 3.6F
Pecan	Brown leaf spot <i>(Sirosporium diffusum)</i> Downy spot <i>(Mycosphaerella caryigena)</i> Liver spot <i>(Gnomonia caryae)</i> Scab <i>(Cladosporium caryigenum)</i> Vein spot <i>(Gnomonia nerviseda)</i> Zonate leaf spot <i>(Grovesinia pyramidalis)</i>	4 to 8 fl. oz. per acre
<p>Application Directions: Apply Rotam Tebuconazole 3.6F in a preventive spray schedule beginning at early bud break (young leaves unfolding), and continue applications at 10- to 14-day intervals through the pollination period. Rotam Tebuconazole 3.6F should be applied at 4 fl. oz. per acre in a tank-mix with the recommended rate of Super-Tin® in cover sprays. Follow label directions for the use of Super-Tin. Do not add a surfactant to the spray solution when tank-mixing Rotam Tebuconazole 3.6F F with Super-Tin. Apply Rotam Tebuconazole 3.6F in a spray volume of 15 or more gallons per acre by air or 50 or more gallons per acre by ground. Apply 7 to 8 fl. oz. per acre of Rotam Tebuconazole 3.6F to full-size mature trees, and 4 to 6 fl. oz. per acre of Rotam Tebuconazole 3.6F to smaller trees. Apply the high rate to varieties that are highly susceptible to the indicated diseases, or when severe disease conditions exist. The lowest labeled rate of a surfactant may be added to the spray solution for optimum control of the indicated diseases.</p> <p>For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Rotam Tebuconazole 3.6F. Rotam Tebuconazole 3.6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Rotam Tebuconazole 3.6F will be resistant to weathering. Rotam Tebuconazole 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3). It may be applied in a tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy.</p>		

27
36

Restricted-entry interval (REI) = 12 hours.

RESTRICTIONS: Do not apply after shucks begin to split. A maximum of 32 fl. oz. of Rotam Tebuconazole 3.6F may be applied per acre per crop season. Do not cut cover crops in treated areas for feed or allow livestock to graze treated areas.

Crop	Disease	Rate of Rotam Tebuconazole 3.6F
Soybean	Rust (<i>Phakopsora pachyrhizi</i>) Powdery mildew (<i>Microsphaera diffusa</i>)	3 to 4 fl oz per acre

Application Directions: Apply Rotam Tebuconazole 3.6F as a broadcast foliar spray as a preventative spray or at first visible symptoms of disease. Repeat applications on a 10- to 14-day spray interval if environmental conditions are favorable for continued disease development. Use of the higher rates and shorter spray intervals are recommended when disease pressure is severe. The lowest labeled rate of a spray surfactant must be tank-mixed with Rotam Tebuconazole 3.6F. Rotam Tebuconazole 3.6F should be applied in a minimum for 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons per acre by aircraft spray equipment.

Restricted-entry interval (REI) = 12 hours.

RESTRICTIONS: Applications may not be made within 21 days of harvest. Do not apply more than 3 applications per season. Do not apply more than 12 fl. oz/a per use season.

28
36

Crop	Disease	Rate of Rotam Tebuconazole 3.6F
Sunflower	Rust (<i>Puccinia helianthi</i>)	4 to 6 fl. oz. per acre
	<p>Application Directions: Apply specific dosage of Rotam Tebuconazole 3.6F at the earliest sign of infection (rust pustules developing) or when weather conditions are favorable for rust development. Apply higher rate to highly susceptible varieties and/or under severe disease conditions. Application may be repeated at 14 days if necessary to maintain control of the disease. Apply specified dosage in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air.</p> <p>For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Rotam Tebuconazole 3.6F. Contact your state Extension Service or Bayer representative for a list of approved surfactants. Rotam Tebuconazole 3.6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Rotam Tebuconazole 3.6F will be resistant to weathering.</p>	
<p>Restricted-entry interval (REI) = 12 hours.</p> <p>RESTRICTIONS: Do not apply more than 16 fl. oz. of Rotam Tebuconazole 3.6F per acre per season or within 50 days of harvest.</p>		

29
36

Crop	Disease	Rate of Rotam Tebuconazole 3.6F
Turnip (Application is limited to East of the Rockies)	Cercospora leaf spot (<i>Cercospora brassicicola</i>)	4 to 7.2 fl. oz. per acre
	<p>Application Directions: Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 12- to 14-day intervals. Rotam Tebuconazole 3.6F F may be applied up to 7 days before harvest. Do not apply more than 28.8 fl. oz. of Rotam Tebuconazole 3.6F per acre per crop season.</p> <p>For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Rotam Tebuconazole 3.6F. Rotam Tebuconazole 3.6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Rotam Tebuconazole 3.6F will be resistant to weathering.</p>	
<p>Restricted-entry interval (REI) = 12 hours.</p> <p>RESTRICTIONS: Do not apply more than 28.8 fl. oz. of Rotam Tebuconazole 3.6F per acre per crop season.</p>		

30
36

Crop	Disease	Rate of Rotam Tebuconazole 3.6F
Wheat	Rusts leaf, stem, and stripe (<i>Puccinia spp.</i>)	4 fl. oz. per acre
	<p>Head blight or scab (<i>Fusarium spp.</i>) – Suppression</p> <p>Application Directions: Wheat fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. A maximum of 4 fl. oz. of Rotam Tebuconazole 3.6F may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw may be fed or used for bedding. Apply Rotam Tebuconazole 3.6F in a minimum of 10 gallons of spray solution per acre by ground, or in a minimum of 5 gallons of spray solution per acre by air.</p> <p><u>Rusts:</u> Apply Rotam Tebuconazole 3.6F at the earliest sign of rust pustules on foliage.</p> <p><u>Fusarium head blight:</u> Optimal timing of Rotam Tebuconazole 3.6F for Fusarium head blight suppression is the beginning of flowering on main stem heads (Feekes 10.51).</p> <p>For optimum disease control, the lowest labeled rate of a spray surfactant should be tank mixed with Rotam Tebuconazole 3.6F. Rotam Tebuconazole 3.6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Rotam Tebuconazole 3.6F will be resistant to weathering.</p>	
<p>Restricted-entry interval (REI) = 12 hours.</p> <p>RESTRICTIONS: Do not allow livestock to graze or feed green forage to livestock prior to 6 days after treatment with Rotam Tebuconazole 3.6F.</p>		

3)
36

SEED TREATMENT - Corn (Sweet Corn, Field Corn, Field Corn Grown For Seed, and Popcorn) For control of soilborne and seedborne Fusarium and soilborne and seedborne head smut.

SEED LABELING: To meet U.S. Federal Seed Act requirements, all seed treated with Rotam Tebuconazole 3.6F must be labeled: TREATED SEED. DO NOT USE FOR FOOD, FEED OR OIL PURPOSES. Treated with Tebuconazole.

USE PRECAUTION: When using formulations that do not contain dye, to comply with 40 CFR 153.155, all seed treated with an economic poison must be colored to distinguish and prevent subsequent inadvertent use as a food for man or feed for animals.

Disease	Rate Fl Oz/CWT	Application Directions
Soilborne and Seedborne Fusarium	0.071	Apply as a seed treatment using standard slurry or mist-type seed treatment equipment. Uniform application of seed is necessary to ensure seed safety and best disease protection. Seed should be sound and well cured prior to treatment. Product should be diluted with sufficient water to ensure complete seed coverage. Consult a seed treatment specialist regarding slurry rates recommended for the crop to be treated with Rotam Tebuconazole 3.6F. The length of control will vary depending on the rate used.
Soilborne and Seedborne Head smut (<i>Sphacelotheca reiliana</i>)	0.27 – 0.54	

32
36

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.

Apply only during alternate years in fields adjacent to aquatic areas listed above.

Do not apply by ground or air within 100 feet of aquatic areas listed above.

Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip.

Spray Drift Management: For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter.

Use the largest droplet size consistent with pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure. Apply in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment.

Spray should be released at the lowest possible height consistent with good pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

Make aerial or ground applications when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.

Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

ROTATIONAL CROPS

Treated areas may be replanted with any crop specified on this label as soon as practical after last application. Any crop not specified on this label may be planted into treated areas 120 days after last application.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of ROTAM AGROCHEMICAL COMPANY LIMITED or Seller. To the extent allowed by applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold ROTAM AGROCHEMICAL COMPANY LIMITED and Seller harmless for any claims relating to such factors.

ROTAM AGROCHEMICAL COMPANY LIMITED warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or ROTAM AGROCHEMICAL COMPANY LIMITED, and to the extent allowed by applicable law, Buyer and User assume the risk of any such use. **ROTAM USA LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

To the extent allowed by law, ROTAM AGROCHEMICAL COMPANY LIMITED or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT ALLOWED BY LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ROTAM AGROCHEMICAL COMPANY LIMITED AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ROTAM AGROCHEMICAL COMPANY LIMITED OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

ROTAM AGROCHEMICAL COMPANY LIMITED and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of ROTAM AGROCHEMICAL COMPANY LIMITED.

34
36

[Base Container Label/Remains on Container when Label is Removed]

Rotam Tebuconazole 3.6F

Agricultural Fungicide

FOR CONTROL OF SPECIFIC DISEASES ON VARIOUS CROPS

ACTIVE INGREDIENT:

Tebuconazole*	38.7%
Other Ingredients:	61.3%
Total:	100.0%

Contains 3.6 pounds tebuconazole per gallon

*alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-H-1,2,4-triazole-1-ethanol
(CASRN: 107534-96-3)

EPA Reg. No. 83100 – 1

EPA Est. No.: 069821-CHN-005

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If Swallowed	<ul style="list-style-type: none"> • Call poison control center or doctor for treatment advice • Do not induce vomiting unless told to do so by the poison control center or doctor. • Have person sip a glass of water if able to swallow. • Do not give anything by mouth to an unconscious person.
If On Skin Or Clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for further treatment advice.
If In Eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for further treatment advice.
If Inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call poison control center or doctor for treatment advice.
Note to Physician	
<p>Symptoms of Poisoning and Recommendations for Medical Treatment: The compound does not cause any definite symptoms that would be diagnostic. No specific antidote. Treat symptomatically.</p>	
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. For Medical Emergency Treatment Call your Local Emergency Response Center.</p>	

35
36

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to labeling under "AGRICULTURAL USE REQUIREMENTS" in the DIRECTIONS FOR USE section for information about this standard.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory: Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

36
36

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

Pesticide Storage: Store in a tightly closed container in a cool, dry place.

Pesticide Disposal: Pesticide spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal (Nonrefillable container 5 gallons or less): Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Offer for recycling, if available.

Residue Removal: Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call CHEMTREC AT 1-800-424-9300

Refer to Attached Booklet for Complete Precautionary Statements and Directions For Use.

Manufactured by:
Rotam Agrochemical Company, Ltd.
7/F Cheung Tat Center
No. 18, Cheung Lee Street
Chai Wan, Hong Kong

Net Contents:
1.0 GALLON (3.78 Liters)