



## OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

August 22, 2025

Blake Cowen  
Product Registration Manager  
Albaugh, LLC  
1525 NE 36<sup>th</sup> Street  
Ankeny, IA 50021

Subject: Approval of Label Amendment; Only Indicated Changes Reviewed – Add grazing, update hotline section, revise booklet statement, addition of optional marketing graphics, and update company information  
Product Name: Rotam Thifen + Triben 750WG Herbicide  
EPA Registration Number: 83100-24  
Application Date: 03/21/2025  
Case Number: 651834

Dear Blake Cowen:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable. However, EPA reviewed only the label changes highlighted, marked, or otherwise indicated on the submitted label. Any other changes to the previously approved label that were not clearly highlighted, marked, or otherwise indicated in your submission were not reviewed and may form the basis of regulatory and/or enforcement action if later discovered by the Agency. Further, submission of a label amendment application with unidentified changes may be considered a knowing submission of false information to the Agency. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

The label submitted with the application has been stamped “Accepted Only Indicated Revisions Reviewed” and is enclosed for your records.

This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 C.F.R. § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently

approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 C.F.R. § 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website contains any false or misleading statement, design, or graphic, the product may be misbranded and unlawful to sell or distribute under FIFRA Sections 2(q)(1)(A) and 12(a)(1)(E). 40 C.F.R. § 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on the product label, claims made as part of the product's sale or distribution may not substantially differ from those claims approved through the registration process under FIFRA Section 12(a)(1)(B). Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the product will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Hester Dingle at 202-566-2596 or at [dingle.hester@epa.gov](mailto:dingle.hester@epa.gov).

Sincerely,

*Kable Bo Davis*

Kable Bo Davis; Senior Advisor  
Office of Pesticide Programs  
Registration Division; Immediate Office

Enclosure

|                       |   |           |
|-----------------------|---|-----------|
| THIFENSULFURON-METHYL | 2 | HERBICIDE |
| TRIBENURON-METHYL     | 2 | HERBICIDE |

[Master Label]

# Rotam Thifen + Triben 750WG Herbicide

DRY FLOWABLE

FOR USE ON WHEAT (INCLUDING DURUM), BARLEY, OAT, TRITICALE,  
SU TOLERANT CANOLA, AND FALLOW.

## ACTIVE INGREDIENTS

### Thifensulfuron-methyl

Methyl 3-[[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl) amino]carbonyl]amino]  
sulfonyl]-2-thiophenecarboxylate ..... 50%

### Tribenuron-methyl

Methyl 2-[[[[(N-(4-methoxy-6-methyl-1,3,5-triazin-2-yl)  
methylamino]carbonyl]amino]sulfonyl]benzoate..... 25%

OTHER INGREDIENTS..... 25%

TOTAL..... 100%

## KEEP OUT OF REACH OF CHILDREN CAUTION / PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

~~[Optional referral statements when booklets and container labels are used:]~~

~~[See label booklet for [complete] [additional] [First Aid,] [Precautionary Statements,] [Directions For Use,] and [Storage and Disposal.]]~~

EPA REG. NO. 83100-24

EPA EST. NO. -----

NET CONTENTS: \_\_\_\_\_ [Oz./Lbs]

**ACCEPTED**  
ONLY INDICATED  
REVISIONS REVIEWED

08/22/2025

Under the Federal Insecticide, Fungicide and  
Rodenticide Act as amended, for the pesticide  
registered under EPA Reg. No.

83100-24

No label revisions other than those indicated were  
reported to the Agency.

### Manufactured For [By]:

Albaugh, LLC

Ankeny, Iowa 50021 Rotam Agrochemical Co., Ltd.

26/F E-Trade Plaza

24 Lee Chung Street

Chaiwan, Hong Kong



**SELECTIVE HERBICIDE**



**SELECTIVE  
HERBICIDE**

[Table of Contents to be added before the Precautionary Statement]

### FIRST AID

#### IF ON SKIN OR CLOTHING

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 to 20 minutes.
- Call a poison control center or doctor for treatment advice.

#### IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For non-emergency exposure information on this product, call 1-888-347-6732 (7 days/week, 24-hr). For medical emergencies, dial 911. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact your local poison control center for emergency medical treatment information.

[See [inside] booklet for [additional/complete] First Aid, [Precautionary Statements/Directions For Use/Storage and Disposal/and/Conditions of Sale and Warranty].]

For Medical Emergencies, Call (888) 347-6732

For 24-hour chemical spill, leak, fire, exposure or accident response information, call CHEMTREC toll free at 1-800-424-9300.

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION – PRECAUCION

Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are listed below.

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, (including butyl rubber ≥14 mils, natural rubber ≥14 mils, neoprene rubber ≥14 mils, nitrile rubber ≥14 mils, poly-ethylene, polyvinyl Chloride (PVC) ≥14 mils and Viton ≥14 mils).
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### USER SAFETY RECOMMENDATIONS

#### Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

Thifensulfuron-methyl has properties and characteristics associated with chemicals detected in groundwater. Thifensulfuron-methyl may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Thifensulfuron-methyl and tribenuron-methyl may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for days after application. A level, well-maintained vegetative buffer strip

between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of thifensulfuron-methyl from runoff water and sediment. Runoff of this product will be greatly reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

### Non-Target Organism Advisory

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated area. Protect the forage and habitat of non-target organisms by minimizing spray drift. For further guidance and instructions on how to minimize spray drift, refer to the Spray Drift Management section of this label.

### PESTICIDE HANDLING

- Calibrate sprayers only with clean water away from the well site.
- Make scheduled checks of spray equipment.
- Ensure that all operation employees accurately measure pesticides.
- Mix only enough product for the job at hand.
- Avoid over-filling of spray tank.
- Do not discharge excess material on the soil at a single spot in the field/grove or mixing/loading station.
- Dilute and agitate excess solution and apply at labeled rates/uses.
- Avoid storage of pesticides near well sites.
- When triple rinsing the pesticide container, be sure to add the rinsate to the spray mix.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical resistant gloves (including butyl rubber ≥14 mils, natural rubber ≥14 mils, neoprene rubber ≥14 mils, nitrile rubber ≥14 mils, poly-ethylene, polyvinyl Chloride (PVC) ≥14 mils and Viton ≥14 mils)
- Shoes plus socks

### PRODUCT INFORMATION

This product is a dry flowable granule that is used for selective post-emergence weed control in wheat (including durum), barley, oat, triticale and fallow. The best control is obtained when this product is applied to young, actively growing weeds. The use rate will depend on weed spectrum and size of weed at time of application. The degree and duration of control may depend on the following:

- weed spectrum and infestation intensity
- weed size at application
- environmental conditions at and following treatment

This product is noncorrosive, nonflammable, nonvolatile, and does not freeze. This product should be mixed in water

and applied as a uniform broadcast spray.

### **PRECAUTIONS**

- Prevent drift of spray to desirable plants. Take all necessary precautions to avoid all direct or indirect contact (such as spray drift) with non-target plants or areas.
- Carefully observe all sprayer cleanup instructions both prior to and after using this product, as spray tank residue may damage crops other than wheat, barley, triticale, oat or SU Tolerant Canola.
- Varieties of wheat (including durum), barley and triticale may differ in their response to various herbicides. **Rotam Albaugh LLC** recommends that you first consult your state experiment station, university, or extension agent as to sensitivity to any herbicide. If no information is available, limit the initial use to a small area.
- Under certain conditions such as heavy rainfall, prolonged cold weather, or wide fluctuations in day/night temperatures prior to or soon after product application, temporary discoloration and/or crop injury may occur. To reduce the potential of crop injury, tank mix this product with 2,4-D (ester formulations perform best—see Tank Mixtures) and apply after the crop is in the tillering stage of growth.
- Dry, dusty field conditions may result in reduced control in wheel track areas.

### **RESTRICTIONS**

- Do not use on any other crop other than wheat, barley, oat, triticale, fallow and SU Tolerant Canola.
- Do not apply, drain or flush equipment on or near desirable trees or other plants or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Do not use on lawns, walks, driveways, tennis courts, or similar areas.
- Do not be apply to wheat, barley, triticale or oat that is stressed by severe weather conditions, drought, low fertility, water-saturated soil, disease, or insect damage, as crop injury may result. Risk of injury is greatest when crop is in the 2- to 5-leaf stage.
- Do not apply when severe winter stress, drought, disease, or insect damage is evident.
- Do not apply to wheat, barley triticale or oat crops underseeded with another crop.
- Do not harvest sooner than 45 days after the last application of this product.

### **BIOLOGICAL ACTIVITY AND ENVIRONMENTAL CONDITIONS**

This product is absorbed primarily through the foliage of plants, rapidly inhibiting the growth of susceptible weeds. 1 to 3 weeks after application to weeds (2 to 5 weeks for wild garlic), leaves of susceptible plants appear chlorotic, and the growing point subsequently dies.

This product provides the best control in vigorously growing crops that shade competitive weeds. Weed control in areas of thin crop stand or seeding skips may not be as satisfactory. However, a crop canopy that is too dense at application can intercept spray and reduce weed control.

The herbicidal action of this product may be affected in crops stressed from adverse environmental conditions (such as extreme temperatures or moisture), abnormal soil conditions, cultural practices, or variations in crop variety. In warm, moist conditions, the expression of herbicide symptoms is accelerated; in cold, dry conditions, expression of herbicide symptoms is delayed. In addition, weeds hardened-off by drought stress are less susceptible to this product.

### **WINDBLOWN SOIL PARTICLES**

Rotam Thifen + Triben 750WG Herbicide has the potential to move off-site due to wind erosion. Soils that are subject to wind erosion usually have a high silt and/or fine to very fine sand fractions and low organic matter content. Other factors which can affects the movement of windblown soil include the intensity and direction of prevailing winds, vegetative cover, site slope, rainfall, and drainage patterns. Avoid applying Rotam Thifen + Triben 750WG Herbicide if prevailing local conditions may be expected to result in off-site movement.

### **SPRAY DRIFT MANAGEMENT**

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

#### **Aerial Applications:**

- Do not release spray at a height greater than 10 ft. above the vegetative canopy, unless a greater application height is necessary for pilot safety.

- For applications prior to the emergence of crops and target weeds, applicators are required to use a Coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a Medium or coarser droplet size (ASABE S572.1).
- The boom length must not exceed 65% of the wingspan for airplanes or 75% of the rotor blade diameter for helicopters.
- Applicators must use ½ swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

#### **Ground Boom Applications:**

- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy unless making a turf, pasture, or rangeland application, in which case applicators may apply with a nozzle height no more than 4 feet above the ground.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a Coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a Medium or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

### **SPRAY DRIFT ADVISORIES**

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

### **IMPORTANCE OF DROPLET SIZE**

#### **Controlling Droplet Size – Ground Boom**

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

#### **Controlling Droplet Size – Aircraft**

- Adjust Nozzles - Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

### **BOOM HEIGHT – Ground Boom**

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.

### **RELEASE HEIGHT - Aircraft**

Higher release heights increase the potential for spray drift. When applying aurally to crops, do not release spray at a height greater than 10 ft. above the crop canopy, unless a greater application height is necessary for pilot safety.

### **SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

### **TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

### **TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an



inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

## WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

## AIR ASSISTED (AIR BLAST) FIELD CROP SPRAYERS

Air assisted field crop sprayers carry droplets to the target via a downward directed air stream. Some may reduce the potential for drift, but if a sprayer is unsuitable for the application and/or set up improperly, high drift potential can result. It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application, is configured properly, and that drift is not occurring.

**Note:** Air assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration. Consult the spray equipment section of this label to determine if use of an air assist sprayer is recommended.

## WEED RESISTANCE MANAGEMENT

Rotam Thifen + Triben 750WG Herbicide contains two active ingredients with the same mode of action. Thifensulfuron-methyl and tribenuron-methyl are classified as a Group 2 herbicide - an Acetolactate Synthase (ALS) or Acetohydroxy Acid Synthase (AHAS) inhibitor.

Herbicide resistance is defined as the inherited ability of a plant to survive and reproduce following exposure to a dose of herbicide normally lethal to the wild type. In a plant, resistance may be naturally occurring or induced by such techniques as genetic engineering or selection of variants produced by tissue culture or mutagenesis. Any weed population may contain or develop plants that are naturally resistant to Rotam Thifen + Triben 750WG Herbicide and other Group 2 herbicides. Weed species with acquired resistance to Group 2 herbicides may eventually dominate the weed population if Group 2 herbicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by Rotam Thifen + Triben 750WG Herbicide or other Group 2 herbicides.

Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed. If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.

To delay herbicide resistance, consider:

- Avoiding the consecutive use of Rotam Thifen + Triben 750WG Herbicide or other target site of action Group 2 herbicides that have a similar target site of action, on the same weed species.
- Using tank mixtures or premixes with herbicides from different target site of action Groups as long as the involved products are all registered for the same use, have different sites of action, and are both effective at the tank mix or prepack rate on the weed(s) of concern.
- Basing herbicide use on a comprehensive Integrated Pest Management (IPM) program.
- Monitoring treated weed populations for loss of field efficacy.
- Users should scout before and after application. Users should report lack of performance to registrant or their representative.

Contact your local extension specialist, certified crop advisors, and/or manufacturer for herbicide resistance management and/or integrated weed management recommendations for specific crops and resistant weed biotypes.

## INTEGRATED PEST MANAGEMENT

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and



genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

Injury to or loss of adjacent sensitive crops, desirable trees or vegetation may result from failure to observe the Precautions and Restrictions listed below:

| WEEDS CONTROLLED - ALL USES             |                          |                             |
|---|--------------------------|-----------------------------|
| Annual knawel                           | Curly dock               | Redroot pigweed             |
| Annual sowthistle                       | False chamomile          | Russian thistle*†           |
| Black mustard                           | Field chickweed          | Scentless chamomile/mayweed |
| Blue/Purple mustard                     | Field pennycress         | Shepherdspurse              |
| Broadleaf dock                          | Filaree (redstem, Texas) | Slimleaf lambsquarters      |
| Bur buttercup                           | Flixweed                 | Smallflower buttercup       |
| Bushy wallflower/Treacle mustard        | Green smartweed          | Smallseed false lax         |
| Clasping pepperweed                     | Kochia**                 | Stinking chickweed          |
| Coast fiddleneck                        | Ladysthumb               | Stinking mayweed /Dogfennel |
| Common buckwheat                        | Lanceleaf sage*          | Swinecress                  |
| Common chickweed                        | London rocket            | Tansymustard                |
| Common groundsel                        | Mayweed chamomile        | Tarweed fiddleneck          |
| Common lambsquarters                    | Miners lettuce           | Tumble/Jim Hill mustard     |
| Common radish                           | Narrowleaf lambsquarters | Volunteer lentils           |
| Common ragweed*                         | Nightflowering catchfly  | Volunteer peas              |
| Common sunflower                        | Pennsylvania smartweed   | Volunteer sunflower         |
| Corn chamomile                          | Pineappleweed            | Wild buckwheat*             |
| Corn gromwell*                          | Prickly lettuce*†        | Wild chamomile              |
| Corn spurry                             | Prostrate knotweed       | Wild garlic*                |
| Cowcockle                               | Prostrate pigweed        | Wild mustard                |
| Cress (mouse-ear)                       | Redmaids                 | Wild radish*                |
| WEEDS PARTIALLY CONTROLLED** - ALL USES |                          |                             |
| Canada thistle*                         | Cutleaf evening primrose | Nightshade (cutleaf, hairy) |
| Carolina geranium                       | Mallow (common, little)  | Vetch*                      |
| Catchweed bedstraw                      |                          |                             |

\* See SPECIFIC WEED PROBLEMS in the CEREALS section for more information.

\*\*Partial control: A visual reduction of weed population as well as a significant loss of vigor for individual weed plants. For better results, use 0.5 or 0.6 ounce of this product per acre and include a tank mix partner such as 2,4-D, bromoxynil, MCPA and/or dicamba. Refer to the TANK MIXTURES section of this label.

†Naturally occurring resistant biotypes of kochia, prickly lettuce and Russian thistle are known to occur. See the TANK MIXTURES and SPECIFIC WEED PROBLEMS sections of this label for additional details.

## **FALLOW**

### **APPLICATION TIMING IN FALLOW**

Apply this product as a fallow treatment, in the spring or fall when the majority of weeds have emerged and are actively growing.

### **USE RATES IN FALLOW**

Apply this product at 0.3 oz. per acre (0.0047 lbs. a.i. tribenuron-methyl/acre and 0.0094 lbs a.i. thifensulfuron-methyl/acre) to 0.6 oz. per acre (0.0094 lbs. a.i. tribenuron-methyl/acre and 0.0188 lbs a.i. thifensulfuron-methyl/acre) to fallow. Two applications of this product may be made provided the total amount applied does not exceed 1.0 oz. per acre (0.0156 lbs. a.i. tribenuron-methyl/acre and 0.0313 lbs a.i. thifensulfuron-methyl/acre) per year.

This product should be applied in combination with other suitable registered fallow herbicides including glyphosate, dicamba, glyphosate plus 2,4-D (ester formulations work best), glyphosate plus dicamba.

### **TANK MIXTURES IN FALLOW**

This product, when used as a fallow treatment, should be tank mixed with other herbicides that are registered for use in fallow. Read and follow all manufacturer label instructions for the companion herbicide. If those instructions conflict with this label, do not tank mix the herbicide with this product.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

### PREPLANT BURNDOWN

#### USE RATES IN PREPLANT BURNDOWN

**Wheat (including durum), Barley, Triticale and Oat:** Apply this product at 0.3 oz. per acre (0.0047 lbs. a.i. tribenuron-methyl/acre and 0.0094 lbs a.i. thifensulfuron-methyl/acre) to 0.6 oz. per acre (0.0094 lbs. a.i. tribenuron-methyl/acre and 0.0188 lbs a.i. thifensulfuron-methyl/acre) as a burndown treatment to wheat (including durum), barley, triticale, and oat to control emerged weeds prior to, or shortly after planting (prior to emergence). Make applications when the majority of weeds have emerged and are actively growing.

**Cotton:** This product may be applied for burndown of emerged weeds prior to the emergence of cotton. This product may be used as part of a preplant burndown treatment, in combination with other suitable registered preplant herbicides.

Read and follow all manufacturers' label instructions for the companion herbicide. If those instructions conflict with this label, do not tank mix the herbicide with this product.

Apply this product at 0.3 oz per acre (0.0047 lbs. a.i. tribenuron-methyl/acre and 0.0094 lbs a.i. thifensulfuron-methyl/acre) to 0.5 oz. per acre (0.0078 lbs. a.i. tribenuron-methyl/acre and 0.0156 lbs a.i. thifensulfuron-methyl/acre) for control or partial control of the weeds listed on this label. Allow at least 14 days between application of this product and planting of cotton. Include a non-ionic surfactant, petroleum-based crop oil concentrate, or vegetable-seed oil-based product (methylated seed oils are considered a vegetable seed-based oil).

If another herbicide is tank mixed with this product to increase the broadleaf weed spectrum, select adjuvants based on the adjuvant limitations of the companion herbicide.

#### SPRAY ADJUVANTS IN PREPLANT BURNDOWN

**Nonionic Surfactant (NIS):** Apply at a rate (concentration) of 0.25-0.5% v/v (1-2 qts. per 100 gals. spray solution). Use the higher rate in hot and dry conditions to enhance control.

**Crop Oil Concentrate:** Under dry conditions or during cool weather, a petroleum based crop oil concentrate, or vegetable-seed oil-based product may be used in place of a nonionic surfactant at 1-2 gallon/100 gals. of spray solution (1-2% v/v) to enhance weed control. Use a petroleum-based crop oil concentrate with at least 14% emulsifiers/surfactant and 80% oil.

**Ammonium Nitrogen Fertilizer:** An ammonium nitrogen fertilizer can be added to a surfactant or a crop oil concentrate to enhance control. Alternatively, a high-quality, sprayable grade of ammonium sulfate (21-0-0) may be used.

#### IMPORTANT PRECAUTIONS IN PREPLANT BURNDOWN

Seedling disease, nematodes, cold weather, deep planting (more than 2"), excessive moisture, high salt concentration, and/or drought may weaken cotton seedlings and increases the possibility of crop injury. Cotton resumes normal growth once favorable growing conditions return.

**RESTRICTION:** DO NOT apply later than 14 days before planting cotton.

**Cotton:** Apply this product at 0.3 oz. per acre (0.0047 lbs. a.i. tribenuron-methyl/acre and 0.0094 lbs a.i. thifensulfuron-methyl/acre) to 0.5 oz. per acre (0.0078 lbs. a.i. tribenuron-methyl/acre and 0.0156 lbs a.i. thifensulfuron-methyl/acre) as a burndown treatment to cotton. Allow at least 14 days between application of this product and planting of cotton. Include a nonionic surfactant, petroleum based crop oil concentrate, or vegetable-seed oil-based product (methylated seed oils are considered a vegetable seed-based oil). If another herbicide is tank mixed with this product to increase the broadleaf weed spectrum, select adjuvants based on the adjuvant limitations of the companion herbicide.

**Sugarbeets, Winter Rape and Canola:** Apply this product at 0.3 oz. per acre (0.0047 lbs. a.i. tribenuron-methyl/acre and 0.0094 lbs a.i. thifensulfuron-methyl/acre) to 0.6 oz. per acre (0.0094 lbs. a.i. tribenuron-methyl/acre and 0.0188 lbs a.i. thifensulfuron-methyl/acre) as a burndown treatment to sugarbeets, winter rape and canola. Allow at least 60 days between application of this product and planting of sugarbeets, winter rape and canola.

**Any other crop (such as corn, rice, grain sorghum or soybeans):** Apply this product at 0.3 oz. per acre (0.0047 lbs. a.i. tribenuron-methyl/acre and 0.0094 lbs a.i. thifensulfuron-methyl/acre) to 0.6 oz. per acre (0.0094 lbs. a.i. tribenuron-methyl/acre and 0.0188 lbs a.i. thifensulfuron-methyl/acre) as a burndown treatment to any other crop (such as corn, rice, grain sorghum or soybeans).

Sequential treatments of this product may also be made provided the total amount of product applied during one fallow/preplant year does not exceed 1.0 oz. per acre (0.0156 lbs. a.i. tribenuron-methyl/acre and 0.0313 lbs a.i. thifensulfuron-methyl/acre); for example, 0.5 oz. (0.0078 lbs. a.i. tribenuron-methyl/acre and 0.0156 lbs a.i. thifensulfuron-methyl/acre) in the fall followed by 0.5 oz. (0.0078 lbs. a.i. tribenuron-methyl/acre and 0.0156 lbs a.i. thifensulfuron-methyl/acre) in the spring. Do not exceed more than 0.5 oz. (0.0078 lbs. a.i. tribenuron-methyl/acre and 0.0156 lbs a.i. thifensulfuron-methyl/acre) per acre when making two applications per year.

Use the 0.6 oz. per acre (0.0094 lbs. a.i. tribenuron-methyl/acre and 0.0188 lbs a.i. thifensulfuron-methyl/acre) rate when weed infestation is heavy and predominantly consists of those weeds listed under WEEDS PARTIALLY CONTROLLED, or when application timing and environmental conditions are marginal. (See APPLICATION TIMING in this section for restriction on planting intervals.)

**PREPLANT APPLICATION FOR BROADLEAF WEED AND WILD GARLIC CONTROL PRIOR TO PLANTING CORN, COTTON, RICE, GRAIN SORGHUM OR SOYBEANS**

In the states of AL, AR, DE, GA, IA, IN, IL, KY, LA, MD, MO, MS, NC, OH, PA, SC, TN, TX, and VA, this product may be applied prior to planting corn, cotton, rice, grain sorghum or soybeans for the control of certain broadleaf weeds and wild garlic. This product may be applied from late fall through early spring.

Do not use this product through any type of irrigation equipment.

**GENERAL INFORMATION**

This product provides postemergence activity for short term control or partial control of labeled weeds. Degree of control and duration of effect are dependent on rate used, sensitivity and size of target weed, environmental conditions at the time of and following application, and spray coverage. Best results are obtained when this product is applied to young, actively growing weeds. The use rate will depend on weed spectrum and size of weed at time of application. However, typical symptoms of dying weeds (discoloration) may not be noticeable for 1-3 weeks after application (2-5 weeks for wild garlic) depending on the environmental conditions and weed susceptibility. Warm, moist conditions following treatment promote the activity of this product while cold, dry conditions delay the activity. Weeds hardened off by cold weather, drought stress or too wet conditions will be less susceptible.

Some naturally occurring weed biotypes\* resistant to this product are known to exist. If weeds listed on this label are not satisfactorily controlled, respray problem areas in a timely and effective manner using a broadleaf herbicide having a different mode of action.

\*Biotypes are naturally occurring individuals of the species, which have a slightly different make-up. Resistant biotypes may look exactly the same as susceptible biotypes. Herbicide-resistant biotypes are able to survive a use rate several times higher than needed to control susceptible biotypes.

**APPLICATION TIMING**

Annual broadleaf weeds should be past the cotyledon stage, actively growing, but less than 4 inches tall or across. See SPECIFIC WEED PROBLEMS for information on vetch (hairy, common), wild garlic and wild radish.

**WEED CONTROL APPLICATION RATES**

Apply this product at 0.5 oz. per acre (0.0078 lbs. a.i. tribenuron-methyl/acre and 0.0156 lbs a.i. thifensulfuron-methyl/acre) to 0.6 oz. per acre (0.0094 lbs. a.i. tribenuron-methyl/acre and 0.0188 lbs a.i. thifensulfuron-methyl/acre) to control emerged weeds. Use 0.6 oz. per acre (0.0094 lbs. a.i. tribenuron-methyl/acre and 0.0188 lbs a.i.

thifensulfuron-methyl/acre) rate when weed infestation is heavy and predominantly consists of those weeds listed under WEEDS PARTIALLY CONTROLLED or when application timing and environmental conditions are marginal. Reduce the application rate to 0.5 ounces (0.0078 lbs. a.i. tribenuron-methyl/acre and 0.0156 lbs a.i. thifensulfuron-methyl/acre) for fields to be planted to cotton.

Sequential treatments of this product to control emerged weeds may also be made provided the total amount of this product applied during one fallow cropland season does not exceed 1.0 oz. per acre (0.0156 lbs. a.i. tribenuron-methyl/acre and 0.0313 lbs a.i. thifensulfuron-methyl/acre); for example, 0.5 oz. (0.0078 lbs. a.i. tribenuron-methyl/acre and 0.0156 lbs a.i. thifensulfuron-methyl/acre) in the fall followed by 0.5 oz. (0.0078 lbs. a.i. tribenuron-methyl/acre and 0.0156 lbs a.i. thifensulfuron-methyl/acre) in the spring. At least one of the sequential treatments should include a tank mix partner herbicide having a different mode of action than this product.

Foliar absorption is the primary means of this product uptake by plants; therefore, thorough coverage of all target weeds is essential.

Conditions which are conducive to healthy, actively growing plants optimize this product's weed control performance. Ideal conditions include warm temperatures and adequate soil moisture before, during and immediately after application.

### TANK MIXTURES IN PREPLANT BURNDOWN

This product may be used as a preplant burndown treatment alone or tank mixed with other herbicides that are registered for use as a preplant burndown product, including glyphosate, dicamba, 2,4-D (ester formulations work best) or glyphosate plus dicamba.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Read and follow all manufacturers' label instructions for the companion herbicide. If those instructions conflict with this label, follow the most restrictive labeling (such as planting interval after application), or do not tank mix the herbicide with this product.

### Precaution

- **For cotton:** Seedling disease, nematodes, cold weather, deep planting (more than 2 inches), excessive moisture, high salt concentration, and/or drought may weaken cotton seedlings and increases the possibility of crop injury. Cotton resumes normal growth once favorable growing conditions return.

### Restrictions

- Sequential treatments of this product may be made provided the total amount of product applied does not exceed 1.0 ounce (0.0156 lbs. a.i. tribenuron-methyl/acre and 0.0313 lbs a.i. thifensulfuron-methyl/acre) per acre or two applications per year. Allow at least 7 days between applications.
- **For Preplant or At-Planting Burndown – Cotton, Field Corn, Grain Sorghum, Rice and Soybeans**
  - Do not apply after planting field corn, grain sorghum, rice or soybeans.
  - Do not apply later than 14 days before planting cotton, corn, soybeans or grain sorghum.
  - ~~Do not allow livestock to graze on, or feed forage, hay or straw from treated soybean fields.~~ Do not apply more than 1.0 ounce (0.0156 lbs. a.i. tribenuron-methyl/acre and 0.0313 lbs a.i. thifensulfuron-methyl/acre) of this product per acre per year.
  - **For Cotton:** Do not apply more than 0.5 ounce (0.0078 lbs. a.i. tribenuron-methyl/acre and 0.0156 lbs a.i. thifensulfuron-methyl/acre) of this product per acre per application.
  - **For Rice, Field Corn, Sorghum, or Soybeans:** Do not apply more than 0.6 ounces (0.0094 lbs. a.i. tribenuron-methyl/acre and 0.0188 lbs a.i. thifensulfuron-methyl/acre) of this product per acre per application.
- **For Preplant Burndown - Cotton**
  - Do not apply more than 0.5 ounce (0.0078 lbs. a.i. tribenuron-methyl/acre and 0.0156 lbs a.i. thifensulfuron-methyl/acre) of this product per acre per application.
  - Do not apply more than 1.0 ounce (0.0156 lbs. a.i. tribenuron-methyl/acre and 0.0313 lbs a.i. thifensulfuron-methyl/acre) of this product per acre per year.
- **For Preplant Burndown - Sugarbeets, Winter Rape and Canola:**

- Do not apply more than at 0.6 ounce (0.0094 lbs. a.i. tribenuron-methyl/acre and 0.0188 lbs a.i. thifensulfuron-methyl/acre) of this product per acre per application.
- Do not apply more than 1.0 ounce (0.0156 lbs. a.i. tribenuron-methyl/acre and 0.0313 lbs a.i. thifensulfuron-methyl/acre) of this product per acre per year.
- **For Preplant Burndown - any other crop (including corn, rice, grain sorghum or soybeans)**
  - Do not apply more than 0.6 ounce (0.0094 lbs. a.i. tribenuron-methyl/acre and 0.0188 lbs a.i. thifensulfuron-methyl/acre) of this product per acre per application.
  - Do not apply more than 1.0 ounce (0.0156 lbs. a.i. tribenuron-methyl/acre and 0.0313 lbs a.i. thifensulfuron-methyl/acre) of this product per acre per year.
- **For Preplant Application for Broadleaf Weeds and Wild Garlic Control Prior to Planting Corn, Cotton, Rice, Grain Sorghum or Soybeans**
  - Do not use this product through any type of irrigation equipment
  - Do not apply more than 0.6 ounce (0.0094 lbs. a.i. tribenuron-methyl/acre and 0.0188 lbs a.i. thifensulfuron-methyl/acre) of this product per acre per application.
  - Do not apply more than 1 ounce (0.0156 lbs. a.i. tribenuron-methyl/acre and 0.0313 lbs a.i. thifensulfuron-methyl/acre) of this product per acre per year.

### CEREALS

#### USE RATES IN CEREALS

Do not use less than 0.3 oz. of this product per acre (0.0047 lbs. a.i. tribenuron-methyl/acre and 0.0094 lbs a.i. thifensulfuron-methyl/acre).

**Wheat (including Durum), Barley and Triticale:** Apply this product at 0.3 oz. per acre (0.0047 lbs. a.i. tribenuron-methyl/acre and 0.0094 lbs a.i. thifensulfuron-methyl/acre) to 0.6 oz. per acre (0.0094 lbs. a.i. tribenuron-methyl/acre and 0.0188 lbs a.i. thifensulfuron-methyl/acre) to wheat (including durum), barley or triticale. Two applications of this product may be made provided the total amount applied does not exceed 1.0 oz. per acre (0.0156 lbs. a.i. tribenuron-methyl/acre and 0.0313 lbs a.i. thifensulfuron-methyl/acre) per year.

Use 0.3 oz. per acre (0.0047 lbs. a.i. tribenuron-methyl/acre and 0.0094 lbs a.i. thifensulfuron-methyl/acre) to 0.4 oz. of this product per acre (0.0063 lbs. a.i. tribenuron-methyl/acre and 0.0125 lbs a.i. thifensulfuron-methyl/acre) for light infestation of the weeds listed under WEEDS CONTROLLED. Conditions at application should be optimum for effective treatment of these weeds.

Use 0.5 oz. of this product per acre (0.0078 lbs. a.i. tribenuron-methyl/acre and 0.0156 lbs a.i. thifensulfuron-methyl/acre) for heavy infestation of the weeds listed under WEEDS PARTIALLY CONTROLLED.

Use 0.6 oz. of this product per acre (0.0094 lbs. a.i. tribenuron-methyl/acre and 0.0188 lbs a.i. thifensulfuron-methyl/acre) for heavy infestation of the weeds listed under WEEDS PARTIALLY CONTROLLED when application timing and environmental conditions are marginal (refer to Environmental Conditions and Biological Activity for best performance).

This product may be used as postemergence applications to triticale anytime after crop is in the 2-3 leaf stage but before the flag leaf is visible. Follow the postemergence use rate instructions listed for wheat.

Other suitable herbicides, fungicides, and insecticides registered for use on triticale may be tank mixed or used sequentially with these products providing the recommended application timing is the same. Read and follow all manufacturers' label instructions for the tank mix partner prior to use. The most restrictive provisions on either label apply.

**Oat (Spring and Winter):** Apply this product at 0.3 oz. per acre (0.0047 lbs. a.i. tribenuron-methyl/acre and 0.0094 lbs a.i. thifensulfuron-methyl/acre) to 0.4 oz. per acre (0.0063 lbs. a.i. tribenuron-methyl/acre and 0.0125 lbs a.i. thifensulfuron-methyl/acre) for control of the weeds listed in WEEDS CONTROLLED table. Do not make more than one application of this product per year on oat.

#### APPLICATION TIMING IN CEREALS

**Wheat (including Durum), Barley, Winter Oat and Triticale:** Make applications after the crop is in the 2-leaf stage, but before the flag leaf is visible.



**Spring Oat:** Make applications after the crop is in the 3-leaf stage, but before jointing. Do not use on Ogle, Porter or Premier varieties as crop injury can occur.

Since this product has very little or no soil activity, it controls only those weeds that have germinated; therefore, apply this product when all or most of the weeds have germinated. Annual broadleaf weeds should be past the cotyledon stage, actively growing, and less than 4" tall or wide. Wild garlic plants should be less than 12" tall with 2" to 4" of new growth. See Specific Weed Problems for more information.

Rainfall immediately after treatment can wash this product off of weed foliage, resulting in reduced weed control. Several hours of dry weather are needed to allow this product to be sufficiently absorbed by weed foliage.

#### **Restrictions - Wheat (including Durum), Barley and Triticale**

- Do not use less than 0.3 ounce (0.0047 lbs. a.i. tribenuron-methyl/acre and 0.0094 lbs a.i. thifensulfuron-methyl/acre) of this product per acre.
- Do not apply more than 0.6 ounce (0.0094 lbs. a.i. tribenuron-methyl/acre and 0.0188 lbs a.i. thifensulfuron-methyl/acre) of this product per acre per application.
- Do not apply more than 1 ounce (0.0156 lbs. a.i. tribenuron-methyl/acre and 0.0313 lbs a.i. thifensulfuron-methyl/acre) of this product per acre per year.
- Two applications of this product may be made provided the total amount applied does not exceed 1.0 ounce (0.0156 lbs. a.i. tribenuron-methyl/acre and 0.0313 lbs a.i. thifensulfuron-methyl/acre) per acre per year. Allow at least 7 days between applications.

#### **Restrictions – Oat (Spring and Winter)**

- Do not use less than 0.3 ounce (0.0047 lbs. a.i. tribenuron-methyl/acre and 0.0094 lbs a.i. thifensulfuron-methyl/acre) of this product per acre per application.
- Do not use more than 0.4 ounce (0.0063 lbs. a.i. tribenuron-methyl/acre and 0.0125 lbs a.i. thifensulfuron-methyl/acre) of this product per acre per application.
- Do not apply more than 0.4 ounce (0.0063 lbs. a.i. tribenuron-methyl/acre and 0.0125 lbs a.i. thifensulfuron-methyl/acre) of this product per acre per year.
- Do not make more than one application of this product per year.
- Do not use on Ogle, Porter or Premier varieties as crop injury can occur.

#### **SPECIFIC WEED PROBLEMS IN CEREALS**

**Canada thistle:** For control in wheat, barley and triticale, use 0.6 oz. per acre (0.0094 lbs. a.i. tribenuron-methyl/acre and 0.0188 lbs a.i. thifensulfuron-methyl/acre) plus surfactant when all thistles are 4" to 8" with 2" to 6" of new growth. Make the application in the spring. Control will be improved by using this product in combination with 2,4-D (refer to TANK MIXTURES).

For control in oat, use 0.4 oz. of this product per acre (0.0063 lbs. a.i. tribenuron-methyl/acre and 0.0125 lbs a.i. thifensulfuron-methyl/acre) plus 2,4-D (refer to TANK MIXTURES).

**Common cocklebur, Common ragweed, Lanceleaf sage:** In wheat, barley and triticale, apply this product at 0.4 oz. per acre (0.0063 lbs. a.i. tribenuron-methyl/acre and 0.0125 lbs a.i. thifensulfuron-methyl/acre) to 0.5 oz. per acre (0.0078 lbs. a.i. tribenuron-methyl/acre and 0.0156 lbs a.i. thifensulfuron-methyl/acre) in combination with 2,4-D at rates from 1/4 to 3/8 lb. active ingredient (ester formulations work best) when weeds are small and actively growing. When using 1/4 lb active ingredient of 2,4-D, be sure to add surfactant at the rate of 1/4 to 1/2 quart per 100 gallons of spray solution (0.06 to 0.125% v/v--use the higher rate under stress conditions).

For control in oat, use 0.4 oz. of this product per acre (0.0063 lbs. a.i. tribenuron-methyl/acre and 0.0125 lbs a.i. thifensulfuron-methyl/acre) plus 2,4-D. Refer to the TANK MIXTURES sections of this label for additional details.

**Corn gromwell, Wild buckwheat:** For control in wheat, barley and triticale, use 0.5 oz. per year (0.0078 lbs. a.i. tribenuron-methyl/acre and 0.0156 lbs a.i. thifensulfuron-methyl/acre) to 0.6 oz. of this product per acre (0.0094 lbs. a.i. tribenuron-methyl/acre and 0.0188 lbs a.i. thifensulfuron-methyl/acre) plus surfactant. For control in oat, use 0.4 oz. of this product per acre (0.0063 lbs. a.i. tribenuron-methyl/acre and 0.0125 lbs a.i. thifensulfuron-methyl/acre) plus 2,4-D,



MCPA or Buctril (refer to TANK MIXTURES).

**Kochia, Russian thistle, Prickly lettuce:** Naturally occurring resistant biotypes of these weeds are known to occur. For best results, use this product in a tank mix with dicamba and 2,4-D; or bromoxynil and 2,4-D. This product must be applied in the spring when weeds are less than 2" tall or 2" across and are actively growing. Refer to the TANK MIXTURES section of this label for additional details.

**Vetch (common and hairy):** For control in wheat, barley and triticale, use 0.5 oz. per acre (0.0078 lbs. a.i. tribenuron-methyl/acre and 0.0156 lbs a.i. thifensulfuron-methyl/acre) to 0.6 oz. of this product per acre (0.0094 lbs. a.i. tribenuron-methyl/acre and 0.0188 lbs a.i. thifensulfuron-methyl/acre) plus surfactant when vetch is less than 6" in length. For severe infestations of vetch, or when vetch is greater than 6 in length, use this product in combination with 2,4-D or MCPA (refer to the Tank Mixtures section of this label).

For control in oat, use 0.4 oz. of this product per acre (0.0063 lbs. a.i. tribenuron-methyl/acre and 0.0125 lbs a.i. thifensulfuron-methyl/acre) plus 2,4-D or MCPA (refer to TANK MIXTURES).

**Wild garlic:** For control in wheat, barley and triticale, use 0.5 oz. per acre (0.0078 lbs. a.i. tribenuron-methyl/acre and 0.0156 lbs a.i. thifensulfuron-methyl/acre) to 0.6 oz. of this product per acre (0.0094 lbs. a.i. tribenuron-methyl/acre and 0.0188 lbs a.i. thifensulfuron-methyl/acre) plus surfactant when wild garlic plants are less than 12" tall with 2" to 4" of new growth. For severe infestations, use the 0.6 oz. per acre (0.0094 lbs. a.i. tribenuron-methyl/acre and 0.0188 lbs a.i. thifensulfuron-methyl/acre) rate of this product. Plants hardened-off by cold weather and/or drought stress may be more difficult to control. Thorough spray coverage of all garlic plants is essential. Typical symptoms of dying garlic plants may not be noticeable for 2 to 5 weeks.

For control in oat, use 0.4 oz. of this product per acre (0.0063 lbs. a.i. tribenuron-methyl/acre and 0.0125 lbs a.i. thifensulfuron-methyl/acre) plus 2,4-D or MCPA (refer to TANK MIXTURES).

**Wild radish:** For best results in wheat, barley and triticale, apply 0.4 oz. per acre (0.0063 lbs. a.i. tribenuron-methyl/acre and 0.0125 lbs a.i. thifensulfuron-methyl/acre) to 0.6 oz. of this product per acre (0.0094 lbs. a.i. tribenuron-methyl/acre and 0.0188 lbs a.i. thifensulfuron-methyl/acre) plus surfactant either in the fall or spring to wild radish rosettes less than 6" in diameter. Applications made later than 30 days after weed emergence will result in partial control. For increased control of severe wild radish infestations, or wild radish emerged greater than 30 days, apply this product at 0.3 oz. per acre (0.0047 lbs. a.i. tribenuron-methyl/acre and 0.0094 lbs a.i. thifensulfuron-methyl/acre) in combination with MCPA at 1/4 lb. active ingredient per acre. Surfactant is required when tank mixing with MCPA, add 1 quart per 100 gallons of spray solution (0.25% vol/vol). Fall applications should be made prior to hardening off of plants.

For control in oat, use 0.4 oz. of this product per acre (0.0063 lbs. a.i. tribenuron-methyl/acre and 0.0125 lbs a.i. thifensulfuron-methyl/acre) plus 2,4-D or MCPA (refer to TANK MIXTURES IN CEREALS).

#### **TANK MIXTURES IN CEREALS**

This product may be tank mixed with other suitable registered herbicides to control weeds listed as suppressed, weeds resistant to this product or weeds not listed under WEEDS CONTROLLED. Read and follow all manufacturers' label instructions for the companion herbicide. If those instructions conflict with this label, do not tank mix the herbicide with this product.

This product can also be mixed with registered fungicides, insecticides, or liquid fertilizer for use on wheat, barley, triticale, oat, or fallow.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

#### **With Broadleaf Herbicides**

**With 2,4-D (amine or ester) or MCPA (amine or ester):** This product may be tank mixed with the amine and ester formulations 2,4-D and MCPA herbicides for use on wheat, barley, triticale and oat.

Refer to the of 2,4-D or MCPA herbicide ester formulations for rates specific to the Red River Valley and adjacent areas of North Dakota and Minnesota and for all other areas.

For best results in the Red River Valley and adjacent areas of North Dakota and Minnesota, add the ester formulations of 2,4-D or MCPA herbicides to the tank at 3/8 lb active ingredient (such as 3/4 pt of a 4 lb/gal product, 1/2 pt of a 6 lb/gal product). No additional surfactant is needed with this mixture.

For best results in other areas, add the ester formulations of 2,4-D or MCPA herbicides to the tank at 1/4 to 3/8 lb. active ingredient (such as 1/2 - 3/4 pt. of a 4 lbs./gal. product, 1/3 - 1/2 pt. of a 6 lbs./gal. product). Surfactant may be added to the mixture at 1/2 to 1 qt. per 100 gals. of spray solution (0.125 to 0.25% v/v); however, adding surfactant may increase the potential for crop injury, especially at the higher phenoxy rates.

Higher rates of 2,4-D or MCPA may be used, but do not exceed the highest rate allowed by those respective labels. Read and follow all label instructions on timing, precautions, and warnings for these herbicides before using these tank mixtures.

**With dicamba:** This product may be tank mixed with 1/16 to 1/8 lb. active ingredient dicamba. Use higher rates when weed infestation is heavy. Surfactant may be added to the mixture at 1/2 to 1 qt. per 100 gals. of spray solution (0.125 to 0.25% v/v); however, adding surfactant may increase the potential for crop injury. Refer to the specific dicamba label for application timing and restrictions.

Tank mixes of this product plus dicamba may result in reduced control of some broadleaf weeds.

**With 2,4-D (amine or ester) and dicamba:** This product may be applied in a 3-way tank mix with formulations of dicamba and 2,4-D. Make application of this product plus dicamba plus 2,4-D Ester or Amine. Use higher rates when weed infestation is heavy. Surfactant may be added to the mixture at 1/2 to 1 qt. per 100 gals. of spray solution (0.125 to 0.25% v/v); however, adding surfactant may increase the potential for crop injury. Consult the specific 2,4-D label, dicamba label, or local guidance for more information and restrictions.

Apply this 3-way combination to winter wheat and winter oat after the crop is tillering and prior to jointing (first node). In spring wheat (including durum) and spring oat, apply after the crop is tillering and before it exceeds the 5-leaf stage.

In spring barley, apply after the crop is tillering and before it exceeds the 4-leaf stage.

**With bromoxynil (including bromoxynil plus MCPA, bromoxynil plus 2,4-D):** This product may be tank mixed with bromoxynil containing herbicides registered for use on wheat, barley, triticale, or fallow.

Read and follow all label instructions on timing, precautions, and warnings for these herbicides before using these tank mixtures. Follow the most restrictive labeling. Tank mixes of this product plus bromoxynil may result in reduced control of Canada thistle.

**With metsulfuron:** This product may be tank mixed with Metsulfuron based on local guidance. Read and follow all label instructions on timing, precautions, and warnings for these herbicides before using this tank mixture.

**With fluroxypyr:** For improved control of Kochia (2" to 4" tall), Russian thistle, mustard species, and wild buckwheat. This product may be tank mixed with fluroxypyr. Refer to the product labels. The most restrictive provisions on either label will apply. Do not use the tank mix if any restrictions on the companion product label conflict with instructions on this herbicide label.

Other suitable registered herbicides, fungicides, and insecticides registered for use on small grains or fallow may be tank mixed or used sequentially with this mixture. Read and follow all manufacturers' label instructions for the companion herbicide. The most restrictive provisions on either label will apply.

**With Fluroxypyr plus 2,4-D:** This product can be tank mixed with fluroxypyr plus 2,4-D herbicides for improved control of broadleaf weeds in wheat, barley, fallow and oats.

For improved control of Kochia (2" to 4" tall), Russian thistle, mustard species and wild buckwheat, this product may be tank mixed with fluroxypyr plus 2,4-D herbicides. Refer to this product's label, and the companion product labels for information regarding use restrictions, labeled crops, rotational cropping intervals, sprayer cleanup, use precautions and other information. The most restrictive provisions on either label will apply. Do not use the tank mix if any restrictions on the companion product label conflict with instructions on this product's herbicide label.

**With fluroxypyr plus MCPA:** This product can be tank mixed with fluroxypyr plus MCPA herbicides for improved control of broadleaf weeds in wheat, barley, fallow and oats.

For improved control of Kochia (2" to 4" tall) Russian thistle, mustard species and wild buckwheat, this product may be tank mixed with fluroxypyr plus MCPA herbicides. Refer to this product's label, and the companion product labels for information regarding use restrictions, labeled crops, rotational cropping intervals, sprayer cleanup, use precautions and other information. The most restrictive provisions on either label will apply. Do not use the tank mix if any restrictions on the companion product label conflict with instructions on this product's label.

Other suitable registered herbicides, fungicides, and insecticides registered for use on cereal grains or fallow may be tank mixed or used sequentially with this mixture. Read and follow all manufacturers' label instructions for the companion herbicide. The most restrictive provisions on either label will apply.

**With carfentrazone-ethyl:** This product can be tank mixed with carfentrazone-ethyl herbicide for improved control of weeds in wheat, barley and triticale.

**With clopyralid, clopyralid plus 2,4-D, clopyralid plus MCPA or clopyralid plus fluroxypyr:** This product can be tank mixed with clopyralid, clopyralid plus 2,4-D, clopyralid plus MCPA or clopyralid plus fluroxypyr herbicides for improved control of weeds in wheat, barley and triticale.

**With Other Broadleaf Herbicides:** Tank mixes of this product plus metribuzin may result in reduced control of wild garlic.

#### **With Grass Herbicides**

**With diclofop-methyl:** This product may be used in combination with diclofop-methyl and bromoxynil herbicides in accordance with the diclofop-methyl label. For best results, use the three-way tank mix of this product at 0.4 oz. per acre (0.0063 lbs. a.i. tribenuron-methyl/acre and 0.0125 lbs a.i. thifensulfuron-methyl/acre) plus diclofop-methyl plus bromoxynil. Apply only to winter wheat. This tank mix should only be used under good soil conditions when wild oat is in the 1-4 leaf stage. If conditions are not ideal for the performance of diclofop-methyl, wild oat control may be reduced. Be sure to follow all warnings and cautions on the diclofop-methyl and bromoxynil labels.

**With imazamethabenz or difenzoquat methyl sulfate:** This product can be tank mixed with imazamethabenz or difenzoquat methyl sulfate. When tank mixing with imazamethabenz, always include another broadleaf weed herbicide with a different mode of action (for example: 2,4-D ester, MCPA ester). Tank-mixed applications of this product plus imazamethabenz may cause temporary crop discoloration, stunting, or injury when heavy rainfall occurs shortly after application.

**With clodinafop-propargy:** This product can be tank mixed with clodinafop-propargy for improved control of weeds in spring wheat.

**With flucarbazone-sodium:** This product can be tank mixed with flucarbazone-sodium for improved control of weeds in spring wheat.

**With sulfosulfuron:** This product can be tank mixed with sulfosulfuron for improved control of weeds in wheat.

**With fenoxaprop-p-ethyl:** This product can be tank mixed with fenoxaprop-p-ethyl for control of some annual grass weeds. This tank mix may also include MCPA ester, bromoxynil or bromoxynil/MCPA for greater spectrum of broadleaf control - see fenoxaprop-p-ethyl label for specific use directions and restrictions on tank mixes.

### **With other grass control products**

Tank mixtures of this product and grass control products may result in poor grass control. ~~Rotam~~ Albaugh LLC recommends that you first consult your state experiment station, university, or extension agent, Agricultural dealer, or ~~Rotam~~ Albaugh LLC representative as to the potential for antagonism before using the mixture. If no information is available, limit the initial use of this product and the grass product to a small area.

### **With Insecticides or Fungicides**

This product may be tank mixed or used sequentially with insecticides (or fungicides) registered for use on cereal grains. However, under certain conditions (drought stress, or if the crop is in the 2- to 4-leaf stage), tank mixes or sequential applications of this product with organophosphate insecticides (such as parathion) may produce temporary crop yellowing or, in severe cases, crop injury. Test these mixtures in a small area before treating large areas. Do not use this product plus Malathion, as crop injury will result.

### **With Liquid Nitrogen Solution Fertilizer**

Liquid nitrogen fertilizer solutions may be used as a carrier in place of water. Run a tank mix compatibility test before mixing this product in fertilizer solution. This product must first be slurried with water and then added to liquid nitrogen solutions (e.g., 28-0-0, 32- 0-0). Ensure that the agitator is running while the product is added. Use of this mixture may result in temporary crop yellowing and stunting.

If using low rates of liquid nitrogen fertilizer in the spray solution (less than 50% of the spray solution volume), the addition of surfactant is necessary. Add surfactant at 1/4 qt. -1 qt. per 100 gals. of spray solution (0.06 - 0.25% v/v) based on local guidance.

When using high rates of liquid nitrogen fertilizer solution in the spray solution, adding surfactant increases the risk of crop injury. Consult your agricultural dealer, consultant, fieldman, or ~~Rotam~~ Albaugh LLC representative for specific instructions before adding an adjuvant to these tank mixtures.

If 2,4-D or MCPA is included with this product and fertilizer mixture, ester formulations tend to be more compatible (See manufacturer's label). Additional surfactant is not needed when using this product in tank mix with 2,4-D ester or MCPA ester and liquid nitrogen fertilizer solutions.

Note: In certain areas east of the Mississippi river, unacceptable crop response may occur with use of straight or dilute nitrogen fertilizer carrier solutions where cold temperatures or widely fluctuating day/night temperatures exist. In these areas consult your agricultural dealer, consultant, field advisor, or ~~Albaugh LLC~~ Rotam representative for specific instructions before using nitrogen fertilizer carrier solutions.

**Liquid nitrogen fertilizer solutions that contain sulfur can increase crop response.**

**Do not use low rates of liquid nitrogen fertilizer solution as a substitute for a surfactant. Do not use with liquid fertilizer solutions with a pH less than 3.0.**

### **IDAHO: SPRINKLER CHEMIGATION WITH THIS PRODUCT AND MCPA PLUS BROMOXYNIL FOR POSTEMERGENCE WEED CONTROL IN WINTER & SPRING WHEAT & SPRING BARLEY IN IDAHO**

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

### **HOW TO USE THROUGH SPRINKLER CHEMIGATION**

Use 0.4 oz. per acre (0.0063 lbs. a.i. tribenuron-methyl/acre and 0.0125 lbs a.i. thifensulfuron-methyl/acre) to 0.5 oz. of this product per acre (0.0078 lbs. a.i. tribenuron-methyl/acre and 0.0156 lbs a.i. thifensulfuron-methyl/acre) in combination with labeled rates of MCPA plus bromoxynil. Apply to wheat, barley and triticale after the 3-leaf stage but before the flag leaf is visible. Make only one chemigation application of this tank mixture per crop year. For best results, apply to broadleaf weeds up to the 4-leaf stage, or 2 inch height (1 in diameter), whichever comes first. Consult all product labels for list of weeds controlled/suppressed.

### **SPRINKLER IRRIGATION APPLICATION**

Apply this tank mix through sprinkler irrigation systems including center pivot, lateral move, side (wheel) roll, solid set or hand move irrigation systems only. Do not apply these herbicides through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for this product's application to any public water system. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The sprinkler chemigation system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

#### **SPECIFIC REQUIREMENTS FOR APPLICATION THROUGH SPRINKLER IRRIGATION SYSTEMS**

1. In center pivot and continuous lateral move systems, this product plus MCPA plus bromoxynil should be applied continuously for the duration of the water application. In solid set systems, application of the tank mix should be made during the last 30 to 45 minutes of the irrigation set.
2. Set the sprinkler system to deliver approximately 0.5 inch or less of water per acre for best product performance.
3. Fill the supply tank with half of the water amount desired, add this product and agitate it well. Add the MCPA plus bromoxynil and then add the remaining water amount with agitation.
4. Agitation is recommended in the pesticide supply tank when applying this tank mix.
5. The use of a surfactant is not recommended with this tank mix application.
6. Inject this product plus MCPA plus bromoxynil solution at least 8 feet ahead of a right angle turn of irrigation pipe to insure adequate mixing. Allow sufficient time for the herbicide mixture to be flushed through the lines before turning off irrigation water.
7. Follow both this product and MCPA plus bromoxynil label instructions for spray tank cleanout both before and after application. Flush lines with clean water following application.
8. Do not apply when wind speed favors drift beyond the area intended for treatment. Avoiding spray drift is the responsibility of the applicator.

#### **MIXING INSTRUCTIONS – ALL USES**

1. Fill the tank 1/4 to 1/3 full of water.
2. While agitating, add the required amount of this product.
3. Continue agitation until the product is fully dispersed, at least 5 minutes.
4. Once the product is fully dispersed, maintain agitation and continue filling tank with water. This product should be mixed thoroughly with water before adding any other material.
5. As the tank is filling, add tank mix partners (if desired) then add the required volume of nonionic surfactant. Always add surfactant last. Do not use with spray additives that alter the pH of the spray solution below pH 5.0 or above pH 9.0, as rapid product degradation can occur. Spray solutions of pH 6.0-8.0 allow for optimum stability of this product.
6. If the mixture is not continuously agitated, settling will occur. If settling occurs, thoroughly re-agitate before using.
7. Apply product spray mixture within 24 hours of mixing to avoid product degradation.
8. If this product and a tank mix partner are to be applied in multiple loads, pre-slurry this product in clean water prior to adding to the tank. This will prevent the tank mix partner from interfering with the dissolution of this



product.

### PRODUCT MEASUREMENT – ALL USES

This product is measured using the product volumetric measuring cylinder. The degree of accuracy of this cylinder varies by  $\pm 7.5\%$ . For more precise measurement, use scales calibrated in ounces.

### CROP ROTATION - ALL USES

Labeled crops may be planted at specified time intervals following application of labeled rates of this product. Use the time intervals listed below to determine the required time interval before planting.

#### Time Interval Before Planting\* (days after treatment with Rotam Thifen + Triben 750WG herbicide)

| Crop   | Days |
|--|------|
| Barley, Rice, Triticale, and Wheat (including durum) | 0    |
| Soybeans   | 7**  |
| Cotton, Field Corn, and Grain Sorghum                | 14** |
| Sugarbeets, Winter Rape, and Canola                  | 60   |
| Any other crop                                       | 45   |

\* Refer to individual product labels to determine rotational crop restrictions when tank mixtures are used.

\*\*When this herbicide is applied on light textured soils (such as sands and loamy sands) or on high pH soils (>7.9), extend time to planting by 7 additional days.

### SURFACTANTS - ALL USES

Unless otherwise specified, add a Albaugh LLC ~~Rotam~~ recommended nonionic surfactant having at least 80% active ingredient at 1 to 2 qts. per 100 gals. of spray solution (0.25 to 0.5% v/v -refer to TANK MIXTURES for specific adjuvant instructions when this product is used in a tank mix).

For preplant burndown in Cotton, include a nonionic surfactant, petroleum based crop oil concentrate, or a vegetable-seed oil-based product (methylated seed oils are considered a vegetable seed-based oil). If another herbicide is tank mixed with this product to increase the broadleaf weed spectrum, select adjuvants based on the adjuvant limitations of the companion herbicide.

Consult your agricultural dealer, applicator, or Albaugh LLC ~~Rotam~~ representative for a listing of recommended surfactants. Antifoaming agents may be used if needed.

Do not use low rates of liquid nitrogen fertilizer solution as a substitute for surfactant.

### GROUND APPLICATION - ALL USES

- For optimum spray distribution and thorough coverage, use flat-fan or low-volume flood nozzles.
- For flat-fan nozzles, use a spray volume of at least 5 gal per acre (GPA).
- For flood nozzles on 30 spacings, use at least 10 GPA, flood nozzles no larger than TK10 (or the equivalent), and a pressure of at least 30 psi. For 40 nozzle spacings, use at least 13 GPA; for 60 spacings use at least 20 GPA. It is essential to overlap the nozzles 100% for all spacings.
- Raindrop RA nozzles are not recommended for product applications, as weed control performance may be reduced. Use screens that are 50-mesh or larger.

### AERIAL APPLICATION - ALL USES

- Use nozzle types and arrangements that provide optimum spray distribution and maximum coverage at 2 to 5 GPA. Use at least 3 GPA in Idaho, Oregon, or Utah.
- Do not apply this product by air in the state of New York. See the SPRAY DRIFT MANAGEMENT section of this label.

### GRAZING

Allow at least 7 days between application and grazing of treated forage. In addition, allow at least 7 days between application and feeding of forage from treated areas to livestock. Allow at least 30 days between application and feeding of hay from treated areas to livestock. Harvested straw may be used for bedding and/or feed. Allow at least 45 days between application and harvesting of grain. ~~Do not graze livestock in treated areas. In addition, do not feed~~



~~forage or hay from treated areas to livestock (harvested straw may be used for bedding and/or feed).~~

### **SU TOLERANT CANOLA**

(FOR USE ONLY ON SULFONYLUREA TOLERANT CANOLA THAT CONTAINS  
THE CIBUS SU CANOLA™ TRAIT)

#### **APPLICATION TIMING**

Use 0.3 oz. of this product per acre (0.0047 lbs. a.i. tribenuron-methyl/acre and 0.0094 lbs a.i. thifensulfuron-methyl/acre) when SU Canola™ is at the 2- to 5-leaf stage of development but prior to the beginning of bolting for control of weeds listed under the WEEDS CONTROLLED table.

#### **RESTRICTIONS:**

- Only for use on canola that contains the Cibus sulfonyurea herbicide tolerant trait (SU Canola™ Trait).
- Do not apply to non-sulfonylurea tolerant canola as severe crop injury or death of the plants may occur.
- Do not exceed more than 0.3 oz. of product per acre (0.0047 lbs. a.i. tribenuron-methyl/acre and 0.0094 lbs a.i. thifensulfuron-methyl/acre) applied to any one crop of SU tolerant Canola (Cibus SU Canola™) per year.
- Do not exceed more than 0.3 oz. of product per acre (0.0047 lbs. a.i. tribenuron-methyl/acre and 0.0094 lbs a.i. thifensulfuron-methyl/acre) per application.

#### **Tank Mixtures**

Other suitable herbicides, fungicides, and insecticides registered for use on canola may be tanked mixed or used sequentially with this product providing the labeled application timing is the same. Read and follow all manufacturer label instructions for the tank mix partner prior to use. The most restrictive provisions on either label must apply.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

For control of grass weeds in SU canola USE 0.3 OZ. PER ACRE (0.0047 lbs. a.i. tribenuron-methyl/acre and 0.0094 lbs a.i. thifensulfuron-methyl/acre) OF THIS PRODUCT with one of the following grass herbicides:

| Grass control product | Use Rate                 |
|-----------------------|--------------------------|
| Clethodim (26.4%)     | 4 to 6 fl. oz. per acre  |
| Clethodim (12.6%)     | 9 to 12 fl. oz. per acre |
| Sethoxydim (18%)      | 2.5 pints per acre       |

#### **SPRAY EQUIPMENT**

For specific application equipment, refer to the manufacturer's recommendations for additional information on GPA, pressure, speed, nozzle types and arrangements, nozzle heights above the target canopy, etc.

Be sure to calibrate air or ground equipment properly before application. Select a spray volume and delivery system that will ensure thorough coverage and a uniform spray pattern with minimum drift. Use higher spray volumes to obtain better coverage when crop canopy is dense. Avoid swath overlapping, and shut off spray booms while starting, turning, slowing, or stopping, to avoid injury to the crop.

Do not make applications using equipment and/or spray volumes or during weather conditions that might cause spray to drift onto nontarget sites. For additional information on spray drift refer to Spray Drift Management section of label. Continuous agitation is required to keep this product in suspension.

#### **SPRAYER CLEANUP**

The spray equipment must be cleaned before this product is sprayed. Follow the cleanup procedures specified on the labels of the previously applied products. If no directions are provided, follow the six steps outlined in AFTER SPRAYING THIS PRODUCT section.

**AT THE END OF THE DAY**

It is recommended that during periods when multiple loads of this product are applied, at the end of each day of spraying the interior of the tank be rinsed with fresh water and then partially filled, and the boom and hoses flushed. This will prevent the buildup of dried pesticide deposits which can accumulate in the application equipment.

**AFTER SPRAYING THIS PRODUCT AND BEFORE SPRAYING CROPS OTHER THAN WHEAT, BARLEY, TRITICALE, OAT and SU TOLERANT CANOLA**

To avoid subsequent injury to desirable crops, thoroughly clean all mixing and spray equipment immediately following applications of this product as follows:

1. Drain tank; thoroughly rinse spray tanks, boom, and hoses with clean water. Loosen and physically remove any visible deposits.
2. Fill the tank with clean water and 1 gal of household ammonia\* (contains 3% active) for every 100 gals. of water. Flush the hoses, boom, and nozzles with the cleaning solution. Then add more water to completely fill the tank. Circulate the cleaning solution through the tank and hoses for at least 15 min. Flush the hoses, boom, and nozzles again with the cleaning solution, and then drain the tank.
3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
4. Repeat step 2. Rinse the tank, boom, and hoses with clean water. If only Ammonia is used as a cleaner, the rinsate solution may be applied back to the crop(s) recommended on this label. Do not exceed the maximum labeled use rate. If other cleaners are used, consult the cleaner label for rinsate disposal instructions. If no instructions are given, dispose of the rinsate on site or at an approved waste disposal facility.

\*Equivalent amounts of an alternate-strength ammonia solution or a [Albaugh LLC Rotam](#)-approved cleaner can be used in the cleanout procedure. Carefully read and follow the individual cleaner instructions. Consult your Ag dealer, applicator, or [Albaugh LLC Rotam](#) representative for a listing of approved cleaners.

**Notes:**

1. IMPORTANT: Do not use chlorine bleach with ammonia as dangerous gases will form.
2. Do not clean equipment in an enclosed area.
3. Steam-cleaning aerial spray tanks is recommended prior to performing the above cleanout procedure to facilitate the removal of any caked deposits.
4. When this product is tank mixed with other pesticides, all cleanout procedures should be examined and the most rigorous procedure should be followed.
5. In addition to this cleanout procedure, all pre-cleanout guidelines on subsequently applied products should be followed as per the individual label.
6. Where routine spraying practices include shared equipment frequently being switched between applications of this product and applications of other pesticides to product-sensitive crops during the same spray season, it is recommended that a sprayer be dedicated to this product to further reduce the chance of crop injury.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

**Pesticide Disposal:** Open dumping is prohibited. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

**Container Disposal (Nonrefillable Container):** Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Contact your state regulatory agency to determine allowable practices in your state. Offer for recycling, or puncture and dispose of in a sanitary landfill, or if allowed by State and local authorities, by burning. If burned, stay out

of smoke.

**Residue Removal:** Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**SPILLS:** For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. Handle and open container in a manner as to prevent spillage. If the container is leaking, invert to prevent leakage. If container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away. You may contact the CHEMTREC Emergency Response for decontamination procedures.

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call CHEMTREC AT 1-800-424-9300.

### **CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of ~~ROTAM-AGROCHEMICAL COMPANY LTD.~~ ALBAUGH LLC, or Seller. To the extent allowed by applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold ALBAUGH LLC ~~ROTAM-AGROCHEMICAL COMPANY LTD.~~ and Seller harmless for any claims relating to such factors.

~~ALBAUGH LLC~~ ~~ROTAM-AGROCHEMICAL COMPANY LTD.~~ warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions for use under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or ~~ALBAUGH LLC~~ ~~ROTAM-AGROCHEMICAL COMPANY LTD.~~ and to the extent allowed by applicable law. Buyer and User assume the risk of any such use. **TO THE EXTENT CONSISTANT WITH APPLICABLE LAW, ALBAUGH LLC ~~ROTAM-AGROCHEMICAL COMPANY LTD.~~ MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

To the extent allowed by law, ~~ALBAUGH LLC~~ ~~ROTAM-AGROCHEMICAL COMPANY LTD.~~ or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT ALLOWED BY LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ALBAUGH LLC ~~ROTAM-AGROCHEMICAL COMPANY LTD.~~ AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ALBAUGH LLC ~~ROTAM-AGROCHEMICAL COMPANY LTD.~~ OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

~~ALBAUGH LLC~~ ~~ROTAM-AGROCHEMICAL COMPANY LTD.~~ and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of ~~ALBAUGH LLC~~ ~~ROTAM-AGROCHEMICAL COMPANY LTD.~~

LTD.

EPA Registered: 05/20/2010  
Amended (Canola): TBA

[All trademarks are the property of their respective owners.]

031825

**{LABEL HISTORY**  
[(Not included in final printed labeling)]

| File Name                   | Version Mark | Comment  |
|-----------------------------|--------------|--|
| 083100-00024.20250318.DRAFT | 031825       | Label Amendment to Add Grazing and Marketing Changes, Company Name and Address Change Throughout |
|                             |              |  |
|                             |              |  |
|                             |              |  |
|                             |              |  |
|                             |              |  |
|                             |              |  |

}