

83100-26

11-18-2010

1/33



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Chemical Safety and Pollution Prevention  
Registration Division (7505P)  
1200 Pennsylvania Ave., N.W.  
Washington, D.C. 20460

EPA Reg. Number:

83100-26

Date of Issuance:

NOV 18 2010

NOTICE OF PESTICIDE:

Registration  
 Reregistration

(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Rotam Imidacloprid 75WP  
Insecticide

Name and Address of Registrant (include ZIP Code):

Rotam Agrochemical Company Limited  
c/o IPM Resources LLC  
4032 Crockers Lake Blvd., Suite 818  
Sarasota, FL 34238

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. Make the following label changes before you release the product for shipment:
  - a. Revise the EPA Registration Number to read, "EPA Reg. No. 83100-26".
  - b. On page 10, revise "Flatheaded borers (including bronze Royal palm bugs birch borer and alder borer)" to read "Flatheaded borers (including bronze birch borer and alder borer)".

Signature of Approving Official:

Date:

NOV 18 2010

Venus Eagle, Product Manager (01)  
Insecticide-Rodenticide Branch, Registration Division (7505P)

- c. On page 10 of the label, revise “0.7 to 1.4 level teaspoons ROTAM IMIDACLOPRID 75 WP INSECTICIDE per inch of trunk diameter breast height (D.B.H.), at 4.5 feet from the ground surface, or 1 to 2 ounces per 30 cumulative inches of trunk diameter breast height (D.B.H).” to read “Apply 0.7 to 1.4 level teaspoons of ROTAM IMIDACLOPRID 75 WP INSECTICIDE per inch of trunk diameter (D.B.H.), or 1 to 2 ounces per 30 cumulative inches of trunk diameter (D.B.H.). Diameter at Breast Height (D.B.H.) = is measured at 4.5 feet from the ground.”
  - d. On page 23 of the label, revise “0.7 to 1.4 level teaspoons Rotam Imidacloprid 75 WP Insecticide per inch of trunk diameter breast height (D.B.H.), at 4.5 feet from the ground surface.” to read “Apply 0.7 to 1.4 level teaspoons of Rotam Imidacloprid 75 WP Insecticide per inch of trunk diameter (D.B.H.). Diameter at Breast Height (D.B.H.) = is measured at 4.5 feet from the ground.”
- 3. The data requirements for storage stability (830-6317) and corrosion characteristics (830-6320) have not been satisfied, and must be submitted within eighteen months of the date of this letter.
  - 4. Submit one copy of the revised final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records. If you have any questions regarding this notice, please contact Kable Bo Davis at (703) 306-0415 [davis.kable@epa.gov](mailto:davis.kable@epa.gov).

Venus Eagle  
Product Manager (01)  
Insecticide-Rodenticide Branch  
Registration Division (7505P)

Enclosure- Stamped Labeling

[Front Container Label]

# Rotam Imidacloprid 75WP Insecticide

For foliar and systemic insect control on turfgrass (including sod farms), nursery, greenhouse, commercial and residential landscape ornamentals, and, interior landscapes.

**ACTIVE INGREDIENT:**

Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine ..... 75%

**OTHER INGREDIENTS** ..... 25%

Total: ..... 100%

EPA Reg. No. 83100- TBA

EPA Est. No. 069821-CHN-005

## KEEP OUT OF REACH OF CHILDREN CAUTION

**PRECAUCION AL USUARIO:** Si usted no puede leer o entender inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente. (If you cannot read or understand English, do not use this product until the label has been fully explained to you.)

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated:**

NOV 18 2010

**Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under EPA Reg. No.**

83100-26

Manufactured by:  
Rotam Agrochemical Company Ltd  
4032 Crockers Lake Blvd.  
Suite 818  
Sarasota, FL 34238  
1-866-927-9826

NET CONTENTS: (tba) Gallons

[Page 2 inside Booklet]

**TABLE OF CONTENTS**

[TO BE ADDED]

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if swallowed, inhaled, or absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing dust or vapor. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Keep children or pets off treated area until spray is dry.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

**Applicators and Other Handlers Must Wear:**

- Long-sleeved shirt and long pants
- Chemical resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton.
- Shoes plus socks

Follow manufacturers instructions for cleaning/ maintaining personal protective equipment, PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering controls statements:**

- When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**User Safety Recommendations**

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID	
If swallowed	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
If on skin or clothing	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
If in eyes	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
In case of emergency call your local poison control center for assistance. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.	
<b>Note To Physician:</b> No specific antidote is available. Treat the patient symptomatically.	

**ENVIRONMENTAL HAZARDS**

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

**Do not formulate this product into other end-use products.**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during

application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

**Exception:** If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton.
- Shoes plus socks

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Keep children and pets off treated area until dry.

**RESISTANCE**

Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area. Consult your Cooperative Extension Service for resistance management strategies and recommended pest management practices for your area. To aid in resistance management, do not foliar apply any other chloronicotinyl insecticide following a soil application of ROTAM IMIDACLOPRID 75 WP INSECTICIDE to the same crop

## TURFGRASS APPLICATIONS

ROTAM IMIDACLOPRID 75 WP INSECTICIDE can be used for the control of listed soil inhabiting pests of turfgrass, such as Northern & Southern masked chafers, *Cyclocephala borealis*, *C. immaculata*, and/or *C. lurida*; Asiatic garden beetle, *Maladera castanea*; European chafer, *Rhizotrogus majalis*; Green June beetle, *Cotinis nitida*; May or June beetle, *Phyllophaga* spp.; Japanese beetle, *Popillia japonica*; Oriental beetle, *Anomala orientalis*; Billbugs, *Sphenophorus* spp.; Annual bluegrass weevil, *Hyperodes* spp.; Black turfgrass ataenius, *Ataenius spretulus* and *Aphodius* spp.; European Crane Fly, *Tipula paludosa*; and mole crickets, *Scapteriscus* spp.. ROTAM IMIDACLOPRID 75 WP INSECTICIDE can also be used for the suppression of cutworms and chinchbugs in turfgrass areas. ROTAM IMIDACLOPRID 75 WP INSECTICIDE can be used as directed on turfgrass and home lawns, business and office complexes, shopping complexes, multi-family residential complexes, golf courses, airports, cemeteries, parks, playgrounds; athletic fields and sod farms.

The active ingredient in ROTAM IMIDACLOPRID 75 WP INSECTICIDE has sufficient residual activity so that applications can be made preceding the egg laying activity of the target pests. High levels of control can be achieved when applications are made preceding or during the egg laying period. The need for an application can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. For control make applications prior to egg hatch of the target pests. Irrigation or rainfall must occur within 24 hours after application to move the active ingredient through the thatch. Applications must not be made when turfgrass areas are waterlogged or the soil is saturated with water. Adequate distribution of the active ingredient cannot be achieved when these conditions exist. The treated turf area must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile. Do not exceed a total of 8.6 oz (0.4 lb of active ingredient) per acre per year.

### **Application Equipment for Use on Turfgrass**

Apply ROTAM IMIDACLOPRID 75 WP INSECTICIDE in sufficient water to provide adequate distribution in the treated area. The use of accurately calibrated equipment normally used for the application of turfgrass insecticides is required. Use equipment, which will produce a uniform, coarse droplet spray, using a low pressure setting to eliminate off target drift. Check calibration periodically to ensure that equipment is working properly.



TURFGRASS APPLICATIONS			
CROP	PEST	RATE Rotam Imidacloprid 75 WP Insecticide	
Turfgrasses	<b>Larvae of:</b> Annual bluegrass weevil Asiatic garden beetle Billbug Black turfgrass ataeuius Cutworms (suppression) European chafer	European Crane Fly Green June beetle Japanese beetle Northern masked chafer Oriental beetle <i>Phyllophaga</i> spp. Southern masked chafer	6.4 to 8.6 oz per acre or 3 to 4 level teaspoons per 1000 sq ft
	<b>Application Timing:</b> For optimum control of grubs, billbugs, annual bluegrass weevil and European crane fly larvae, make application prior to egg hatch of the target pest. Be sure to read "APPLICATION EQUIPMENT" Section of this label.		
	Chinchbug (suppression) Mole Crickets		8.6 oz per acre or 4 level teaspoons per 1000 sq ft
	<b>Application Timing:</b> For suppression of chinchbugs, make application prior to hatching of the first instar nymphs. For control of mole crickets make application prior to or during the peak egg hatch period. When adults or large nymphs are present and actively tunneling, ROTAM IMIDACLOPRID 75 WP INSECTICIDE application should be accompanied by a curative insecticide. Follow label instructions for other insecticides when tank-mixing.		
Consult your local State Agricultural Experiment Station or State Extension Turf Specialists for more specific information regarding timing of application.			
<b>APPLICATION INSTRUCTIONS:</b> Irrigation or rainfall must occur within 24 hours after application to move the active ingredient through the thatch. <b>RESTRICTIONS:</b> <ul style="list-style-type: none"> <li>Do not apply more than 8.6 oz (0.4 lb of active ingredient) per acre per year.</li> <li>Do not mow turf or lawn area until after sufficient irrigation or rainfall has occurred so that uniformity of application will not be affected.</li> </ul> <b>Note:</b> 1 level teaspoon=1.4 grams ROTAM IMIDACLOPRID 75 WP Insecticide 3 level teaspoons=1 level tablespoon			

## ORNAMENTAL APPLICATIONS

### Commercial and Residential Landscape

ROTAM IMIDACLOPRID 75 WP Insecticide is for use on ornamentals in commercial and residential landscapes and interior plantscapes. ROTAM IMIDACLOPRID 75 WP INSECTICIDE is a systemic product and will be translocated upward into the plant system from root uptake. The product must be placed where the growing portion of the target plant can absorb the active ingredient. The addition of a nitrogen containing fertilizer, where applicable, into the solution may enhance the uptake of the active ingredient. Application can be made by foliar application or soil applications; including soil injection, drenches, and broadcast sprays. Foliar applications offer locally systemic activity against insect pests.

When making soil applications to plants with woody stems, systemic activity will be delayed until the active ingredient is translocated throughout the plant. In some cases, this translocation delay could take 60 days or longer. For this reason, applications should be made prior to anticipated pest infestation to achieve optimum levels of control.

For outdoor ornamentals, broadcast applications do not exceed a total of 8.6 oz (0.4 lb of active ingredient) per acre per year.

### Application Equipment for Foliar Applications

ROTAM IMIDACLOPRID 75 WP INSECTICIDE mixes readily with water and may be used in many types of application equipment. Mix product with the required amount of water and apply as desired dependent upon the selected use pattern. When making foliar applications on hard to wet foliage such as holly, pine, or ivy, the addition of a spreader/ sticker can be applied. If concentrate or mist type spray equipment is used, an equivalent amount of product must be used on the area sprayed, as would be used in a dilute application. Do not apply through any irrigation system.

**MIXING:** To prepare the spray mixture, add the appropriate amount of product into the spray tank while filling with water to the desired level. Operate the agitator while mixing.

ROTAM IMIDACLOPRID 75 WP INSECTICIDE has been found to be compatible with commonly used fungicides, miticides, liquid fertilizers, and other commonly used insecticides. Check physical compatibility using the correct proportion of products in a small jar test if local experience is unavailable.

### Ant Management Programs

Use ROTAM IMIDACLOPRID 75 WP Insecticide to control aphids, scale insects, mealybugs and other sucking pests on ornamentals to limit the honeydew available as a food source for ant populations. ROTAM IMIDACLOPRID 75 WP INSECTICIDE applications can be then be supplemented with residual sprays, bait placements or other ant control tactics to further reduce the pest population.

**Applications for use only in residential areas and indoor and outside perimeters of industrial and commercial buildings**

CROP	PEST	RATE Rotam Imidacloprid 75 WP Insecticide		
		Level measure	Water	
Trees Shrubs Evergreens Flowers Foliage plants Ground- covers Interior plantscapes	Adelgids	Mealybugs	0.25 tsp.	2.5 gal.
	Aphids	Psyllids	0.5 tsp.	5 gal.
	Japanese beetles	Sawfly larvae	1 tsp.	10 gal.
	Lace bugs	(suppression)	2.5 tsp.	25 gal.
	Leaf beetles (including elm and viburnum leaf beetles)	Thrips	5 tsp.	50 gal.
	Leafhoppers (including glassy- winged sharpshooter)	Whiteflies	3 Tbsp. + 1 tsp.	100 gal.
	White grub larvae (such as Japanese beetle larvae, Chafers, <i>Phyllophaga</i> spp. Asiatic garden beetle, Oriental beetle)		3 to 4 teaspoons per 1000 sq ft	
<p><b>APPLICATION INSTRUCTIONS:</b></p> <p><b>Foliar Applications:</b> Start treatments prior to establishment of high pest populations and reapply on an as needed basis.</p> <p><b>Broadcast Applications:</b> Mix required amount of product in sufficient water to uniformly and accurately cover the area being treated. Use 2 gallons of water or more per 1000 sq ft. Irrigate thoroughly to incorporate ROTAM IMIDACLOPRID 75 WP INSECTICIDE into the upper soil profile. Refer to use directions specific for FLOWERS and GROUNDCOVERS concerning additional use directions.</p> <p><b>RESTRICTION:</b> Do not exceed a total of 8.6 oz (0.4 lb of active ingredient) per acre per year.</p>				

**Application to Trees, Shrubs, Flowers and Groundcovers**

(Applications for use only in residential areas, indoor and outside perimeters of industrial and commercial buildings, and state, national, and private wooded and forested areas to control the insect pests listed below)

<b>Pests Controlled:</b>	
<ul style="list-style-type: none"> <li>Adelgids</li> <li>Aphids</li> <li>Armored scales (suppression)</li> <li>Black vine weevil larvae</li> <li>Eucalyptus longhorned borer</li> <li>Flatheaded borers (including bronze Royal palm bugs birch borer and alder borer)</li> <li>Japanese beetles</li> <li>Lace bugs</li> <li>Leaf beetles (including elm and viburnum leaf beetles)</li> </ul>	<ul style="list-style-type: none"> <li>Leafhoppers (including glassy winged sharpshooter)</li> <li>Leafminers</li> <li>Mealybugs</li> <li>Pine tip moth larvae</li> <li>Psyllids</li> <li>Royal palm bugs</li> <li>Sawfly larvae</li> <li>Soft scales</li> <li>Thrips (suppression)</li> <li>White grub larvae</li> <li>Whiteflies</li> </ul>
<b>Trees</b>	0.7 to 1.4 level teaspoons ROTAM IMIDACLOPRID 75 WP INSECTICIDE per inch of trunk diameter breast height (D.B.H.), at 4.5 feet from the ground surface, or 1 to 2 ounces per 30 cumulative inches of trunk diameter breast height (D.B.H).
<p><b>Soil Injection:</b> <u>Grid Application:</u> Space holes on 2.5-foot centers, in a grid pattern, extending to the drip line of the tree. <u>Circle Application:</u> Apply in holes evenly spaced in circles, (use more than one circle dependent upon the size of the tree) beneath the drip line of the tree extending in from that line. <u>Basal Application:</u> Space injection holes evenly around the base of the tree trunk no more than 6 to 12 inches out from the base.</p> <p>Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. Keep the treated area moist for 7 to 10 days. Do not use less than 4 holes per tree.</p> <p><b>No Soil Injection Applications Allowed in Nassau or Suffolk Counties of New York.</b></p> <p><b>Soil Drench:</b> Uniformly apply the dosage in no less than 10 gallons of water per 1000 square feet as a drench around the base of the tree, directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone.</p> <p><b>For Control of Specified Borers:</b> Application to trees already heavily infested may not prevent the eventual loss of the trees due to existing pest damage and tree stress.</p>	
<b>Shrubs</b>	Apply 0.7 to 1.4 level teaspoons ROTAM IMIDACLOPRID 75 WP INSECTICIDE per foot of shrub height or 1.0 to 2.0 oz per 30 cumulative feet of shrub height.

<p><b>Soil Injection:</b> Apply to individual plants using dosage indicated. Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. Keep the treated area moist for 7 to 10 days. Do not use less than 4 holes per shrub.</p> <p><b>No Soil Injection Applications Allowed in Nassau or Suffolk Counties of New York.</b></p> <p><b>Soil Drench:</b> Uniformly apply the dosage in no less than 10 gallons of water per 1000 square feet as a drench around the base of the shrub, directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone.</p> <p><b>For Control of Specified Borers:</b> Application to trees already heavily infested may not prevent the eventual loss of the trees due to existing pest damage and tree stress.</p>	
<p><b>Flowers and Groundcovers</b></p>	<p>3 to 4 level teaspoons ROTAM IMIDACLOPRID 75 WP INSECTICIDE per 1000 sq ft</p>
<p><b>APPLICATION INSTRUCTIONS:</b> Apply as a broadcast treatment and incorporate into the soil before planting or apply after plants are established. If application is made to established plants irrigate thoroughly after application.</p> <p><b>RESTRICTION:</b> Do not exceed a total of 8.6 oz (0.4 lb of active ingredient) per acre per year.</p> <p><b>Note:</b> 1 level teaspoon = 1.4 grams ROTAM IMIDACLOPRID 75 WP Insecticide 3 level teaspoons = 1 level tablespoon</p>	

**Perimeter applications only in residential areas**

CROP	PEST	RATE		
		Rotam Imidacloprid 75 WP Insecticide		
Pome Fruits Apple Crabapple Loquat Mayhaw Pear Pear (oriental). Quince	Aphids (except Woolly scale* apple aphid) Leafhoppers (including glassy winged sharpshooter) Leafminer	Mealybugs* San Jose	0.5 oz (3 Tbsp. +1 tsp.) per 100 gal	2 oz per acre'

**APPLICATION INSTRUCTIONS:**

Apply specified dosage as foliar spray as needed after petal-fall is complete. For control of rosy apple aphid, apply prior to leafrolling caused by the pest. For first generation leafminer control, make first application as soon as petal-fall is complete. Greatest leafminer control will result from the earliest possible application. For second and succeeding generations of leafminer, optimal control is obtained from applications made early in the adult flight against egg and early instar larvae. A second application may be required 10 days later if severe pressure continues or if generations are overlapping. A single application may result in suppression only. ROTAM IMIDACLOPRID 75 WP INSECTICIDE will not control late stage larvae. For San Jose Scale, time applications to the crawler stage. Treat each generation. For late season (preharvest) control of leafhopper species, apply ROTAM IMIDACLOPRID 75 WP INSECTICIDE while most leafhoppers are in the nymphal stage. For optimal control of mealybug, insure good spray coverage of the trunk and scaffolding limbs or other resting sites of the mealybug.

**RESTRICTIONS:**

- Do not apply more than 2.0 ounces per acre in a single application.
- Do not make more than 5 applications. Allow 10 or more days between applications. Allow at least 7 days between last application and harvest.
- Do not exceed a total of 8.6 oz (0.4 lb of active ingredient) per acre per year.

\*Not for use in California for control on pears.

**Perimeter applications residential areas - Continued**

Pecan*	Yellow pecan aphid Black margined aphid Pecan leaf phylloxera	Pecan spittlebug Pecan stem phylloxera	0.5 oz (3 Tbsp. +1 tsp.) per 100 gal	2.0 oz per acre <sup>1</sup>
--------	---	--	---	---------------------------------

**APPLICATION INSTRUCTIONS:**

Make foliar applications as pests begin to build before populations become extreme. Two applications at a 10 to 14 day interval may be required to achieve control. Scout and retreat if needed. Thorough uniform coverage of foliage is necessary for control. Addition of an organosilicone-based spray adjuvant at a rate not to exceed the adjuvant manufacturer's labeled use rate may improve coverage.

**RESTRICTIONS:**

- Do not apply more than a total of 6 ounces of ROTAM IMIDACLOPRID 75 WP INSECTICIDE per acre per year.
- Do not make more than 3 applications. Allow 10 or more days between applications.
- Do not exceed a total of 8.6 oz (0.4 lb of active ingredient) per acre per year.

<sup>1</sup>The amount of ROTAM IMIDACLOPRID 75 WP INSECTICIDE required per acre will depend on tree size and volume of foliage present. The rate per acre is based on a standard of 400 gallons of dilute spray solution per acre for large trees.

**Note:**

1 level teaspoon = 1.4 grams ROTAM IMIDACLOPRID 75 WP Insecticide  
3 level teaspoons = 1 level tablespoon

\*Not for use in California unless directed by supplemental labeling.

**Applications for use only in residential areas and indoor and outside perimeters of industrial and commercial buildings**

Grapes	Leafhoppers (including glassy-winged sharpshooter) Mealybugs	0.5 oz (3 Tbsp. +1 tsp.) per 100 gal	2.0 oz per acre
--------	---	---	--------------------

Apply specified dosage as a foliar spray using 200 gallons of water per acre. Do not apply more than a total of 2.0 ounces of ROTAM IMIDACLOPRID 75 WP INSECTICIDE per acre per year. Allow at least 14 days between applications. Applications may be applied up to and including day of harvest.

**RESTRICTIONS:**

- Do not graze treated areas or use clippings from treated areas for feed or forage.
- Prevent runoff or puddling of irrigation water following application.

- Keep children and pets off treated area until dry.
- Do not apply ROTAM IMIDACLOPRID 75 WP INSECTICIDE to areas which are water logged or saturated, or frozen, which will not allow penetration into the root zone of the plant.
- Do not apply more than 8.6 oz (0.4 lb of active ingredient) per acre per year.

**Food Crops:** Treated areas may be replanted with any crop specified on an imidacloprid label, or with any crop for which a tolerance exists for the active ingredient.

For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12-month plant-back interval must be observed.

### **Nurseries (field and containerized, including forest nurseries), Greenhouses, and, Interior plantscapes**

ROTAM IMIDACLOPRID 75 WP INSECTICIDE provides insect control on ornamental and vegetable plants in nurseries, greenhouses and interior plantscapes. ROTAM IMIDACLOPRID 75 WP INSECTICIDE is a systemic product and will be translocated upward within the plant. To assure optimum effectiveness, the product must be placed where the growing portion of the target plant can absorb the active ingredient. The addition of a nitrogen containing fertilizer into the solution, where applicable, may enhance uptake of the active ingredient. Application can be made by foliar application or soil applications, including soil injection, drenches, chemigation, and broadcast sprays.

When making soil applications to plants with woody stems, systemic activity will be delayed until the active ingredient is translocated throughout the plant. For this reason, applications should be made prior to anticipated pest infestation to achieve optimum levels of control.

For outdoor ornamentals, broadcast applications cannot exceed a total of 8.6 oz. (0.4 lb. of active ingredient) per acre per year.

**Bark Media:** Media with 30% or more bark content may confer a shorter period of protection when treated with ROTAM IMIDACLOPRID 75 WP INSECTICIDE.

#### **Application Equipment for Ornamentals and Vegetable Plants**

ROTAM IMIDACLOPRID 75 WP INSECTICIDE mixes readily with water and may be used in many types of application equipment. Mix product with the required amount of water and apply as desired dependent upon the selected use pattern.

When making foliar applications on hard to wet foliage such as holly, pine, or ivy, the addition of a spreader/sticker facilitates uptake of the product. If concentrate or mist type spray equipment is used, an equivalent amount of product should be used on the area sprayed, as would be used in a dilute application.

**Tank Mixes:** ROTAM IMIDACLOPRID 75 WP INSECTICIDE has been found to be compatible with commonly used liquid fertilizers, fungicides and insecticides.



Rotam recommends that physical compatibility be confirmed using the correct proportion of products in a small jar test if local experience is unavailable.

**CHEMIGATION**

Refer to DIRECTIONS FOR USE section before proceeding with chemigation application.

ROTAM IMIDACLOPRID 75 WP INSECTICIDE may be applied at labeled rates either alone or in tank mixture with other pesticides and chemicals registered for application through irrigation systems. The normal dilution ratio is 1:100 to 1:200, depending on the system. Always meter the product into the irrigation water during the first part of the irrigation cycle. The product may be mixed separately prior to injection. Agitation may be necessary if the mixture is allowed to stand more than 24 hours.

Remove scale, pesticide residue and other foreign matter from the tank and entire irrigation system.

Apply ROTAM IMIDACLOPRID 75 WP INSECTICIDE only through micro irrigation (individual spaghetti tubes), drip irrigation, overhead irrigation, ebb and flood, or hand-held or motorized calibrated irrigation equipment. Do not apply this product through any other type of irrigation system. Crop injury or lack of effectiveness can result from non-uniform distribution of treated water.

If you have any questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or a person who is under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

**SAFETY DEVICES FOR IRRIGATION SYSTEMS CONNECTED TO PUBLIC WATER SUPPLIES:**

If the source of water for your irrigation system is a public water supply, follow the instructions below.

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

**SAFETY DEVICES FOR IRRIGATION SYSTEMS NOT CONNECTED TO A PUBLIC WATER SUPPLY:**

1. The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where the pesticide distribution is adversely affected.
6. Systems must use a metering pump such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

**Application to Grassy Areas in Nurseries**

Use ROTAM IMIDACLOPRID 75 WP INSECTICIDE as directed on nursery grass in sites such as under or around field or container grown plants, on roadways or other grassy areas in or around nurseries.

**Insects controlled:** listed soil inhabiting pests of grassy areas of nurseries such as

- Northern and Southern masked chafers,  
*Cyclocephala borealis*, *C. immaculata*, and/or *C. lurida*
- Asiatic garden beetle, *Maladera castanea*
- European chafer *Rhizotroqus majalis*
- Green June beetle, *Cotinis nitida*
- May or June beetle, *Phyllophaga* spp
- Japanese beetle, *Popillia japonica*
- Oriental beetle, *Anomala orientalis*
- Billbugs, *Spherophorus* spp.
- Annual bluegrass weevil, *Hyperodes* spp.
- Black turfgrass ataenius, *Ataenius spretulus* and *Aphodius* spp.
- Mole crickets, *Scapteriscus* spp.

**Insects suppressed:** cutworms and chinchbugs.

For control make applications prior to egg hatch of the target pests. Irrigation or rainfall must occur within 24 hours after application to move the active ingredient through the thatch.

The active ingredient in ROTAM IMIDACLOPRID 75 WP INSECTICIDE has sufficient residual activity so that applications can be made preceding the egg laying activity of the target pests. Use historical monitoring data of the site, previous records or experiences, current season adult trapping or other methods to determine the need for application.

Applications must not be made when grassy areas are waterlogged or the soil is saturated with water. Applications must not be made when grassy areas are waterlogged or the soil is saturated with water. Rainfall or irrigation must be able to vertically penetrate in the soil profile. Do not apply more than a total of 8.6 oz (0.4 lb of active ingredient) per acre per year.

**Application Equipment for Use on Grassy Areas in Nurseries**

Apply ROTAM IMIDACLOPRID 75 WP INSECTICIDE in enough water to provide adequate coverage of the treated area. Use accurately calibrated equipment normally used for soil application. Use equipment that will produce a uniform, coarse droplet spray, with a low pressure setting to eliminate off-target drift. Check calibration regularly to confirm that equipment is working properly.

FIELD AND FOREST NURSERY APPLICATIONS		
CROP	PEST	RATE Rotam Imidacloprid 75 WP Insecticide
Grassy areas of Field & Forest Nurseries	<b>Larvae of:</b> Annual bluegrass weevil Asiatic garden beetle Billbugs Black turfgrass ataenius Cutworms (suppression) European chafer Green June beetle Japanese beetle Northern masked Chafer Oriental beetle <i>Phyllophaga</i> spp. Southern masked chafer	6.4 to 8.6 oz per acre or 3 to 4 level teaspoons per 1000 sq ft
	Chinchbugs (suppression) Mole crickets	8.6 oz per acre or 4 level teaspoons per 1000 sq ft

**APPLICATION INSTRUCTIONS:**  
 For control of grubs, billbugs and annual bluegrass-weevil, make application prior to egg hatch of the target pest. Be sure to read "APPLICATION EQUIPMENT" Section of this label.  
 For suppression of chinchbugs, make application prior to the hatching of the first instar nymphs. For control of mole crickets make application prior to or during the peak egg hatch period. When adults or large nymphs are present and actively tunneling, ROTAM Imidicloprid 75 WP Insecticide application should be accompanied by a curative insecticide. Follow label instructions for other insecticides when tank-mixing.

Consult your local turf, state Agricultural Experiment Station, or State Extension Service Specialist for more specific information regarding timing of application.

Apply irrigation or treat within 24 hours prior to rainfall to move the active ingredient through the thatch.

**RESTRICTIONS:**

- Do not apply more than 8.6 oz. (0.4 lb of active ingredient) per acre per year.
- Do not mow grass area until after sufficient irrigation or rainfall has occurred so that uniformity of application will not be affected.

**Note:**  
 1 level teaspoon = 1.4 grams ROTAM IMIDACLOPRID 75 WP Insecticide  
 3 level teaspoons = 1 level tablespoon

**ORNAMENTAL AND VEGETABLE\* APPLICATIONS**

For foliar and systemic insect control in and around field-grown nursery and container stock, indoor and outdoor ornamentals (including both greenhouse and interior plantscapes), ornamentals and vegetable plants grown in flats, benches or beds.

CROP	PEST	RATE Rotam Imidacloprid 75 WP Insecticide
Trees (including non-bearing fruit and nut) Shrubs Evergreens Flowers Foliage plants Ground covers Interior plantscape Vegetable plants*	Adelgids Aphids Japanese beetles (adults) Lacebugs Leaf beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassy winged sharpshooter) Leafminers Mealybugs Sawfly larvae Thrips (suppression) Whiteflies	3 tablespoons + 2 teaspoons per 100 gal of water
<p><b>Foliar Applications:</b> Start treatments prior to establishment of high pest populations and reapply on an as needed basis.</p> <p>To aid in resistance management, do not foliar apply any other chloronicotinyl insecticide following a soil application of ROTAM IMIDACLOPRID 75 WP INSECTICIDE to the same crop.</p>		
	White grub larvae (such as Japanese beetle larvae, Chafers, <i>Phyllophaga</i> spp. Asiatic garden beetle, Oriental beetle)	3 to 4 level teaspoons per 1000 sq ft
<p><b>Broadcast Applications:</b> Mix required amount of product in sufficient water to uniformly and accurately cover the area being treated. Do not use less than 2 gallons of water per 1000 sq. ft. Irrigate thoroughly to incorporate ROTAM Imidacloprid 75 WP Insecticide into the upper soil profile.</p> <p>Refer to REMARKS section for use directions specific for "Flowers and Ground Covers" concerning additional use directions.</p> <p>* Only for use on vegetable plants intended for resale including: Broccoli, Chinese Broccoli, Broccoli Raab, Brussels Sprouts, Cabbage, Chinese Cabbage, Cauliflower, Collards, Eggplant, Ground Cherry, Kale, Kohlrabi, Lettuce, Mustard Greens, Pepinos, Peppers, Potatoes, Rape, Greens, Sorghum, Sugar beets, Tomatillo, and Tomato.</p> <p><b>RESTRICTION:</b> Do not apply more than 8.6 oz. (0.4 lb of active ingredient) per acre per year.</p> <p><b>Note:</b> 1 level teaspoon = 1.4 grams ROTAM IMIDACLOPRID 75 WP Insecticide 3 level teaspoons = 1 level tablespoon</p>		

**Application to Containerized Nursery and Greenhouse Grown Plants**  
**Drench and Irrigation Applications:** For use only on ornamental and vegetable plants in greenhouses, nurseries and interior plantscapes using soil drenches, micro-irrigation, drip irrigation, overhead irrigation, ebb and flood irrigation, or hand-held or motorized calibrated irrigation equipment.

**Application Rates for Containerized Nursery and Greenhouse Grown Plants**

Pest	Use Pattern		Rate		APPLICATION INSTRUCTIONS
			Rotam Imidacloprid 75 WP Insecticide		
Adelgids Aphids Armored scale (suppression) Fungus gnats (larvae only) <sup>1</sup> Japanese Beetle (adults) Lacebugs Leaf beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassy- winged sharpshooter)	Plants in Containers	Herbaceous Species—including Vegetable plants <sup>5</sup> (one or two plants per pot)	Container Size (inches)	No. pots treated with 0.6 ounce*	Distribute 0.6 ounce of ROTAM IMIDACLOPRID 75 WP INSECTICIDE evenly in the specified number of pots, using sufficient water volume to wet potting medium without loss of liquid through leaching. Apply according to label directions. Use moderate irrigation after each application. Irrigate carefully during the next 10 days in order to avoid loss of active ingredient due to leaching.
			2	3000	
			3	2000	
			4	1500	
			5	1200	
			6	1000	
			7	850	
			8	750	
			9	675	
			10	600	
			11	550	
			12	500	

<p>Leafminers Mealybugs Psyllids Root mealybugs <sup>2</sup> Root Weevil Complex (Such as Black Vine Weevil, Apopka Weevil, Citrus Root Weevil<sup>3</sup>) Soft Scale Thrips (suppression) <sup>4</sup> White Grub larvae (such as Japanese Beetle, Masked Chafers, European Chafer, Oriental Beetle, Asiatic Garden Beetle) Whiteflies</p>		Woody Perennial Species	2 3 4 5 6 7 8 9 10 11 12	2000 1350 1000 800 650 550 500 450 400 350 300		
		Herbaceous Species – including Vegetable plants <sup>5</sup> (three or more plants per pot)	Use the above Woody Perennial Species rates			
		Ornamental and vegetable <sup>5</sup> plants grown in flats, benches, or beds	0.6 ounce* per 3000 sq ft			Mix required amount in enough water to uniformly and accurately cover the area being treated. Do not use less than 2 gallons of mixture per 1000 sq. ft. Apply as a broadcast treatment and incorporate into the medium before planting or apply after plants are established. If application is made to established plants lightly irrigated after application. Allow no leaching or runoff for 10 days after application.
		Containerized Plants	Container Size	No. Pots Treated with 0.6 Ounce*		Apply in enough water to wet the potting medium. For best control, apply before egg hatch of the target pest. Irrigate moderately after application to move the active ingredient into the root zone.
			1 gal	340 to 244		
			2 gal	280 to 210		
			3 gal	220 to 165		
			5 gal	160 to 110		

<p>White Grub larvae (such as: Japanese Beetle, Masked Chafers, European Chafer, Oriental Beetle, Asiatic Garden Beetle)</p>	<p><b>APPLICATION (Field and Forest Nurseries):</b> Mow vegetation in the area to be treated to a height of 3 inches or less before application. Mowing to the lowest possible height will provide more control. Apply May through July. For control, treatment must be followed by rainfall or irrigation. Do not use less than 2 gallons of spray volume per 1000 square feet.</p> <p>Apply as a uniform band on either side of row using a band six (6) inches wider than the actual root ball diameter to be dug. Do not allow bands in adjacent rows to overlap. Use 0.6 ounce per 1000 ft. of row or 3000 sq. ft.</p> <p>For grub control in turf areas, make a broadcast application using 0.6 ounce per 3000 sq. ft.</p>
--	--

0.6 oz. = 4 tablespoons

- 1 **Fungus gnat larvae** in the soil will be controlled by drench or incorporation. **No adult Fungus Gnat control.** Other foliar insect control is achieved by the uptake of ROTAM IMIDACLOPRID 75 WP INSECTICIDE from a healthy root system translocating the active ingredient up into the plant.
- 2 **Root Mealybug** control will require a thorough drenching of containerized media. Coverage is essential for control while minimizing the amount of leachate. Rate: 0.6 ounce in 150 gallons of water.
- 3 **Citrus Root Weevil:** For use on non-bearing citrus nursery stock.
- 4 **Thrips** suppression on foliage only. Thrips in buds and flowers will not be suppressed.
- 5 **Note:** For use on vegetable plants intended for resale only including: Broccoli, Chinese Broccoli, Broccoli Raab, Brussels Sprouts, Cabbage, Chinese Cabbage, Cauliflower, Collards, Eggplant, Ground Cherry, Kale, Kohlrabi, Lettuce, Mustard Greens, Pepinos, Peppers, Potatoes, Rape Greens, Sorghum, Sugarbeets, Tomatillo, and Tomato.

**RESTRICTIONS:**

- Do not graze treated areas or use clippings from treated areas for feed or forage.
- Do not allow runoff or puddling of irrigation water following application.
- Do not apply to soils which are water-logged or saturated, which will not allow the penetration of the insecticide into the root zone of the plants.
- Do not allow leachate runoff for the first 10 days after application, in order to retain the product and facilitate full plant uptake of the active ingredient.
- For outdoor ornamentals grown in beds or turf, applications of ROTAM IMIDACLOPRID 75 WP INSECTICIDE cannot exceed a total of 8.6 oz (0.4 lb of active ingredient) per acre per year.

**Food Crops:** Treated areas may be replanted with any crop specified on an imidacloprid label, or with any crop for which a tolerance exists for the active ingredient.

For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12 month plant-back interval must be observed.



**Application to Field Grown and Forest Nursery Plants**

PESTS	USES AND INSTRUCTIONS
<p>Adelgids Aphids Armored scales (suppression) Black vine weevil larvae Eucalyptus longhorned borers Flatheaded borers (including bronze birch and alder borer) Japanese beetles (adults) Lacebugs Leaf beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassy-winged sharpshooter) Leafminers Mealybugs Pine tip moth larvae Psyllids Royal palm bugs Sawfly larvae Soft scales Thrips (suppression) White grub larvae Whiteflies</p>	<p><b>TREES</b> 0.7 to 1.4 level teaspoons Rotam Imidacloprid 75 WP Insecticide per inch of trunk diameter breast height (D.B.H.), at 4.5 feet from the ground surface.</p> <p><b>Soil Injection: GRID SYSTEM:</b> Using a grid pattern, space holes on 2.5 foot centers, extending to the drip line of the tree. <b>CIRCLE SYSTEM:</b> Apply in holes evenly spaced in circles, (use more than one circle depending on the size of the tree) beneath the drip line of the tree extending in from that line. <b>BASAL SYSTEM:</b> Space injection holes evenly around the base of the tree trunk no more than 6 to 12 inches out from the base. Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Use low pressure and sufficient solution for distribution of the liquid into the treatment zone. Keep the treated area moist for 7 to 10 days. Use a minimum of 4 holes per tree.</p> <p><b>No Soil Injection Application Allowed in Nassau or Suffolk Counties of New York.</b></p> <p><b>Soil Drench:</b> Uniformly apply the dosage in at least 10 gallons of water per 1000 square feet as a drench around the base of the tree. Direct to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone.</p> <p><b>For Control of Specified Borers:</b> Application to trees already heavily infested may not prevent the eventual loss of the trees due to existing pest damage and tree stress.</p> <p><b>SHRUBS</b> 0.7 to 1.4 level teaspoons Rotam Imidacloprid 75 WP Insecticide per foot of shrub height</p> <p><b>Soil Injection:</b> Apply to individual plants using dosage indicated. Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. Keep the treated area moist for 7 to 10 days. Do not use less than 4 holes per shrub.</p> <p><b>No Soil Injection Application Allowed in Nassau or Suffolk Counties of New York.</b></p> <p><b>Soil Drench:</b> Uniformly apply the dosage in at least 10 gallons of water per 1000 square feet as a drench around the base of the tree, directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone.</p>

	<p><b>FLOWERS AND GROUND COVERS</b>          Apply 3 to 4 level teaspoons Rotam Imidacloprid 75 WP Insecticide per 1000 sq ft as a broadcast treatment and incorporate into the soil before planting or apply after plants are established. If application is made to established plants, optimum control will be attained if area is irrigated thoroughly after application.</p>
<p><b>RESTRICTION:</b>          Do not apply more than 8.6 oz. (0.4 lb of active ingredient) per acre per year.</p>	

**Ebb & Flood Application**

ROTAM IMIDACLOPRID 75 WP INSECTICIDE may be applied through Ebb and Flood applications. To assure accurate uptake test a minimum of 10 plants at known field capacity after allowing the plants to dry out for one or two days. Re-wet the same plants to determine how much water on average each plant will absorb to bring it back at field capacity. Use the volume absorbed per plant (keeping pot sizes uniform) multiplied by the number of pots being treated. Add to this volume a required minimum to flood your smallest treatment area. This should minimize the return back to the storage tank. Re-use the returned volume with subsequent irrigation or nutrients on the same plants.

<b>ROTAM IMIDACLOPRID 75 WP INSECTICIDE            Ebb &amp; Flood Applications</b>		
Adelgids Aphids Armored scales (suppression) Fungus Gnats (larvae only) <sup>1</sup> Japanese Beetles (adults) Lacebugs Leaf beetles (including elm and viburnum leaf beetles)	Leafhoppers (including glassy-winged sharpshooter) Leafminers Mealybugs Psyllids Root mealybugs <sup>2</sup> Root Weevil Complex: (such as Apopka Weevil, Black Vine Weevil, Citrus Root Weevil <sup>3</sup> )	Soft Scales Thrips (suppression) <sup>4</sup> Whiteflies White Grub larvae (such as Japanese Beetle, Masked Chafer, European Chafers, Oriental Beetle, Asiatic Garden Beetle)
	<b>Number of pots treated with 0.6 ounces<sup>7</sup></b>	
<b>Pot sizes (inches)</b>	<b>Herbaceous species            including vegetable plants<sup>5</sup>            (one or two plants per pot)</b>	<b>Woody perennials,            Herbaceous species            including vegetable plants<sup>5</sup>            (3 or more per pot)</b>
2	3000	2000
3	2000	1350
4	1500	1000
5	1200	800
6	1000	650
7	850	550

27  
33

8	750	500
9	675	450
10	600	400
11	550	350
12	500	300

0.6 oz. = 4 tablespoons

1 **Fungus gnat larvae** in the soil will be controlled by drench or incorporation. **No adult Fungus Gnat control.** Other foliar insect control is achieved by the uptake of ROTAM IMIDACLOPRID 75 WP INSECTICIDE from a healthy root system translocating the active ingredient up into the plant.

2 **Root Mealybug** control will require a thorough drenching of containerized media. Coverage is essential for control while minimizing the amount of leachate. Rate: 0.6 oz in 150 gallons of water.

3 **Citrus Root Weevil:** For use on non-bearing citrus nursery stock.

4 **Thrips** suppression on foliage only. Thrips in buds and flowers will not be suppressed.

5 **Note:** For use on vegetable plants intended for resale only including: Broccoli, Chinese Broccoli, Broccoli Raab, Brussels Sprouts, Cabbage, Chinese Cabbage, Cauliflower, Collards, Eggplant, Ground Cherry, Kale, Kohlrabi, Lettuce, Mustard Greens, Pepinos, Peppers, Potatoes, Rape Greens, Sorghum, Sugarbeets, Tomatillo, and Tomato.

**RESTRICTION:**

Do not apply more than 8.6 oz. (0.4 lb of active ingredient) per acre per year.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

**Pesticide Storage:** Store in a tightly closed container in a cool, dry place.

**Pesticide Disposal:** Pesticide spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Container Disposal (Nonrefillable container):** Do not reuse or refill this container. **Residue Removal:** Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or for reconditioning if appropriate, or puncture and dispose of in a sanitary landfill or by incineration. Do not burn unless allowed by state and local ordinances.

**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident,  
call CHEMTREC AT 1-800-424-9300**

**CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of ROTAM AGROCHEMICAL COMPANY LIMITED or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold ROTAM AGROCHEMICAL COMPANY LIMITED and Seller harmless for any claims relating to such factors.

ROTAM AGROCHEMICAL COMPANY LIMITED warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or ROTAM AGROCHEMICAL COMPANY LIMITED, and Buyer and User assume the risk of any such use. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW ROTAM AGROCHEMICAL COMPANY LIMITED MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

In no event shall ROTAM AGROCHEMICAL COMPANY LIMITED or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ROTAM AGROCHEMICAL COMPANY LIMITED AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ROTAM AGROCHEMICAL COMPANY LIMITED OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

ROTAM AGROCHEMICAL COMPANY LIMITED and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of ROTAM AGROCHEMICAL COMPANY LIMITED.

30  
33

[Booklet back remaining on container when Booklet is removed]

# Rotam Imidacloprid 75 WP Insecticide

For foliar and systemic insect control on turfgrass (including sod farms), nursery, greenhouse, commercial and residential landscape ornamentals, and, interior plantscapes.

**ACTIVE INGREDIENT:**

Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine ..... 75%

**OTHER INGREDIENTS** ..... 25%

Total: ..... 100%

EPA Reg. No. 83100- TBA

EPA Est. No. 069821-CHN-005

## KEEP OUT OF REACH OF CHILDREN CAUTION

**PRECAUCION AL USUARIO:** Si usted no puede leer o entender inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente. (If you cannot read or understand English, do not use this product until the label has been fully explained to you.)

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed, inhaled, or absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing dust or vapor. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Keep children or pets off treated area until spray is dry.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

##### Applicators and Other Handlers Must Wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton.
- Shoes plus socks

Follow manufacturers instructions for cleaning/ maintaining personal protective equipment, PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering controls statements:**

- When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

<b>User Safety Recommendations</b>	
User should:	
<ul style="list-style-type: none"> <li>• Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.</li> <li>• Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.</li> <li>• Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.</li> </ul>	

<b>FIRST AID</b>	
If swallowed	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
If on skin or clothing	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
If in eyes	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
In case of emergency call your local poison control center for assistance. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.	
<b>Note To Physician:</b> No specific antidote is available. Treat the patient symptomatically.	

### ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

**Do not formulate this product into other end-use products.**

[removed AGRICULTURAL USE REQUIREMENT BOX]

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

**Pesticide Storage:** Store in a tightly closed container in a cool, dry place.

**Pesticide Disposal:** Pesticide spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Container Disposal (Nonrefillable container):** Do not reuse or refill this container. **Residue Removal:** Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or for reconditioning if appropriate, or puncture and dispose of in a sanitary landfill or by incineration. Do not burn unless allowed by state and local ordinances.

**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident,  
call CHEMTREC AT 1-800-424-9300**



33  
33

Code:

Manufactured by:  
Rotam Agrochemical Company Ltd  
4032 Crockers Lake Blvd.  
Suite 818  
Sarasota, FL 34238  
1-866-927-6826

NET CONTENTS: (tba) gallons