

83222-31

8/22/2012

1/41

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460



OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

AUG 22 2012

Direct AG Source, LLC  
c/o Jane Miller, Biologic, Inc.  
115 Obtuse Hill Road  
Brookfield, CT 06804


Subject: Notification turned into amendment to convert rates from fractions to decimals, add advisory language, add "product/A" and "a.i./A" and updating storage and disposal section  
Acephate 97 % Prills  
EPA Registration No. 83222-31

Dear Ms. Miller,

The labeling referred to above submitted in connection with the Federal Insecticide, Fungicide and Rodenticide Act, as amended is acceptable.

Please submit one final printed copy of the labeling before releasing the product for shipment. A stamped copy of the labeling is enclosed for your records. If you have any questions regarding this label, please contact Autumn Metzger at (703) 305-5314.

Sincerely,

  
Venus Eagle  
Product Manager 01  
Insecticide-Rodenticide Branch  
Registration Division (7505P)

2/41

Notification to add advisory statements correct use rates and typos - 08 16 2012

GROUP	1	INSECTICIDE
-------	---	-------------

# ACEPHATE 97% PRILLS

## Insecticide

[Water Dispersible Granule]

<b>Active Ingredient</b>	By Wt
Acephate (O S Dimethyl acetylphosphoramidothioate)	97 0%
<b>Other Ingredients</b>	3 0%
<b>Total</b>	100 0%

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

SEE NEXT PAGE FOR ADDITIONAL PRECAUTIONARY STATEMENTS

<b>FIRST AID</b>	
Contains an organophosphate that inhibits cholinesterase	
If swallowed	Call poison control center or doctor immediately for treatment advice Do not induce vomiting unless told to do so by the poison control center or doctor Do not give anything by mouth to an unconscious person
If in eyes	Hold eye open and rinse slowly and gently with water for 15 20 minutes Remove contact lenses if present after the first 5 minutes then continue rinsing eye Call a poison control center or doctor for treatment advice
If on skin or clothing	Take off contaminated clothing Rinse skin immediately with plenty of water for 15 20 minutes Call a poison control center or doctor for treatment advice
If inhaled	Move person to fresh air If person is not breathing call 911 or an ambulance then give artificial respiration preferably by mouth to mouth if possible Call a poison control center or doctor for further treatment advice
Have the product container or label with you when calling a poison control center or doctor, or going for treatment FOR EMERGENCY MEDICAL ASSISTANCE CALL THE ROCKY MOUNTAIN POISON CENTER 1 866 673 6671	
NOTE TO PHYSICIAN	
Acephate is a cholinesterase inhibitor If signs of cholinesterase inhibition appear atropine is antidotal 2 PAM may also be used in conjunction with atropine but should not be used alone	

FOR CHEMICAL EMERGENCY Spill leak fire exposure or accident call  
CHEMTREC 1 800 424 9300

EPA Reg No 83222 31

EPA Est No [REDACTED]

Net Contents \_\_\_ Pounds

Batch/Lot No \_\_\_\_\_

Manufactured for  
Direct Ag Source LLC  
30473 260<sup>th</sup> Street  
Eldora IA 50627

**ACCEPTED**  
**AUG 22 2012**  
**Under the Federal Insecticide, Fungicide,  
and Rodenticide Act, as amended, for the  
pesticide registered under**



3/41

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**



**CAUTION. HARMFUL IF SWALLOWED.**

**PERSONAL PROTECTIVE EQUIPMENT (PPE):**

Some materials that are chemical-resistant to this product are butyl rubber, nitrile rubber, and neoprene rubber. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

**Applicators, mixers, loaders and other handlers must wear:**

- long-sleeved shirt and long pants
- chemical resistant gloves for all mixers and loader and for applicators using hand-held application equipment
- shoes and socks

**In addition, all mixers and loaders and all applicators using low pressure hand wand application equipment must wear:**

- A NIOSH-approved dust mist filtering respirator with MSHA/NIOSH approval number prefix TC21C or a NIOSH-approved respirator with any N, R, P, or HE filter.

See engineering controls for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Controls**

Pilots must use an enclosed cockpit in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)].

The use of human flaggers is prohibited.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to birds.

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

Exposed treated seed may be hazardous to birds and other wildlife. Dispose of all excess treated seed and seed packaging by burial away from bodies of water.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Eliminations System (NPDES) permitted and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.



4/41

**USER SAFETY RECOMMENDATIONS**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, chemical resistant gloves made of any waterproof material and shoes plus socks.

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, or greenhouses.

Do not enter or allow others to enter treated areas until sprays have dried.

**Restrictions:**

- Do not apply using low pressure handwand equipment, except for control of fire ants as a mound treatment and when used on ornamental trees, shrubs, and floral plants grown for non-agricultural or non-commercial use.
- Not for indoor residential use.
- For use on turf, use limited to sod farms and golf courses, except when applying to mound or spot treatment for fire ant and harvester ant control.
- For greenhouse/shadehouse, use is limited to commercial greenhouses/shadehouses for use on ornamental, floral and foliage plants and the tobacco floatbed application as specified in the tobacco use directions.
- Not for use on agricultural establishments in hopper-box, planter-box, slurry-box or other seed treatment applications at, or immediately before, planting.



**SPRAY DRIFT MANAGEMENT**

Observe the following precautions to minimize drift:

A variety of factors including weather conditions (e.g. wind direction, wind speed, temperature, relative humidity), and method of application (e.g. ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator and grower must evaluate all factors and make appropriate adjustments when applying this product.

- All aerial, ground and air-assisted/airblast application equipment must be properly maintained and calibrated using water as carrier. Do not apply this product as an ultralow (ULV) spray or in any carrier other than water unless specified otherwise in the directions for use.
- Use the largest droplet size consistent with good pest control. Small droplets are more prone to spray drift and can be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.
- Do not apply at wind speeds greater than 10 mph at the application site.
- Make applications when wind velocity favors on-target product deposition (approximately 3 to 10 mph).
- Apply as close to target plants as practical to obtain a good spray pattern for adequate coverage.
- For aerial applications, do not apply at heights greater than 10 feet (consistent with flight safety).
- For airblast applications, direct spray above foliage and turn off outward pointing nozzles at row ends and outer rows.
- For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used and must not exceed 75% of wing span or rotor diameter.
- For ground applications, do not apply at heights greater than 4 feet.

**CHEMIGATION**

Do not apply this product through any type of irrigation system except as specified for use on cranberries.

**USE INFORMATION**

ACEPHATE 97% PRILLS is an insecticide for control of pests on selected agricultural crops and in certain non-crop areas. The active ingredient in ACEPHATE 97% PRILLS is acephate, a water-soluble insecticide readily absorbed by plant roots and foliage to give systemic control of feeding insects. Insect pests are generally controlled more effectively by ACEPHATE 97% PRILLS through ingestion than by contact. To maintain control repeat application of ACEPHATE 97% PRILLS as directed.

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THE LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, AND/OR ILLEGAL RESIDUES.**

**NOTE:** This product is sold by weight and package is full when packed but, due to the product's nature, settling is likely to occur.

**MIXING AND COMPATIBILITY**

Fill the spray tank one-third to one-half full of clean water with agitation running and add the required amount of ACEPHATE 97% PRILLS Insecticide to the spray tank. Add remaining water to bring to the desired volume and maintain agitation during mixing, loading and application. Do not add any liquid fertilizers, micronutrients or adjuvants to the spray solution until the ACEPHATE 97% PRILLS Insecticide has completely dissolved. ACEPHATE 97% PRILLS should completely dissolve in approximately five minutes. Dissolution may be slowed by cold water, low level of agitation, or water containing high concentrations of boron or sulfur.

**FIRE ANT CONTROL**

Under prolonged hot and dry conditions, due to the location of ants deep in the nest, best results are obtained from applications made in the early morning or late afternoon when ants are most active.



6/41

Notification to add advisory statements, correct use rates and typos – 08 16 2012

**TANK MIXES:** Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator, and/or application advisor.

Read and follow the entire label of each product to be used in tank mixes with this product.

**USE DIRECTIONS FOR FOLIAR APPLICATIONS TO BEANS AND LIMA BEANS**

**—Dry and Succulent Forms—**

**Except Succulent Green Beans - Includes string beans, snap beans and wax beans**

**Ground Application:** Apply in 20 to 100 gallons of water per acre by conventional ground equipment to ensure thorough coverage of the target crop.

**Aerial Application:** Make applications of ACEPHATE 97% PRILLS Insecticide in a minimum of 2 gallons of water per acre. Use sufficient carrier volume to provide thorough, uniform coverage.

PESTS CONTROLLED	AMOUNT OF ACEPHATE 97% PRILLS PER ACRE	APPLICATION INSTRUCTIONS
Fleahoppers Grasshoppers	0.25 to 0.5 lb. (4 to 8 oz.)	Apply when eggs or insects first appear.
Aphids (excluding, Black Bean Aphid) Bean Leaf Beetle Bean Leafroller Cabbage Looper Cutworms Green Cloverworm Leafhoppers Mexican Bean Beetle Plant bugs (Lygus) Soybean Looper Thrips Whitefly (Except Sweetpotato & Silverleaf Whitefly)	0.5 to 1 lb. (8 to 16 oz.)	Repeat at 3 day spray intervals for rates up to 0.52 lb. product/A (0.5 lb. a.i./A) and at 7 day spray intervals for rates greater than 0.52 lb. product/A (0.5 lb. a.i./A) as necessary to maintain control.*  Use higher listed rates for severe insect infestations.
Armyworms (excluding Beet Armyworm) Corn Earworm European Corn Borer	0.75 to 1 lb. (12 to 16 oz.)	

**RESTRICTIONS:**

\*Do not apply more than 2.1 lbs. product/A (2 lbs. a.i./A) per season.

Snap beans— dry beans: Allow at least 14 days to elapse between final application and harvest.

Lima beans—succulent form: Allow at least one day to elapse between final application and harvest.

Do not feed treated vines to livestock.

Not for use on Succulent Green Beans - Includes string beans, snap beans and wax beans.



Notification to add advisory statements, correct use rates and typos – 08 16 2012

**USE DIRECTIONS FOR FOLIAR APPLICATIONS TO  
BRUSSELS SPROUTS & CAULIFLOWER**

**Ground Application:** Apply in 25 to 150 gallons of water per acre by conventional ground equipment to ensure thorough coverage of the target crop.

**Aerial Application:** Make applications of ACEPHATE 97% PRILLS Insecticide in a minimum of 5 gallons of water per acre. Use sufficient carrier volume to provide thorough, uniform coverage.

PESTS CONTROLLED	AMOUNT OF ACEPHATE 97% PRILLS PER ACRE	APPLICATION INSTRUCTIONS
Green Peach Aphid	0.5 to 1 lb. (8 to 16 oz.)	Apply when eggs or insects first appear.  Repeat applications at 3 day spray intervals for rates up to 0.52 lb. product/A (0.5 lb. a.i./A) and at 7 day spray intervals for rates greater than 0.52 lb. product/A (0.5 lb. a.i./A) as necessary to maintain control.*
Cabbage Looper Imported Cabbage Worm	1 lb. (16 oz.)	Use the higher listed rate when heavy infestations of aphids are present
Diamondback Moth Larvae	1 lb. (16 oz.)	This insect has demonstrated an ability to develop resistance to various classes of insecticides. Consult your local Agricultural Extension Service for current recommended control practices for this insect.
<b>RESTRICTIONS:</b> *Do not apply more than 2.1 lbs. product/A (2 lbs. a.i./A) per crop cycle. Allow at least 14 days to elapse between final application and harvest. Do not feed trimmings to livestock or allow animals to graze in treated areas.		

**USE DIRECTIONS FOR FOLIAR APPLICATIONS TO CELERY**

**Ground Application:** Apply in 50 to 100 gallons of water per acre by conventional ground equipment to ensure thorough coverage of the target crop.

**Aerial Application:** Make applications of ACEPHATE 97% PRILLS Insecticide in a minimum of 5 gallons of water per acre. Use sufficient carrier volume to provide thorough, uniform coverage.

PESTS CONTROLLED	AMOUNT OF ACEPHATE 97% PRILLS PER ACRE	APPLICATION INSTRUCTIONS
Green Peach Aphid	0.5 to 1 lb. (8 to 16 oz.)	Apply when eggs or insects first appear. Repeat at 3 day spray intervals for rates up to 0.52 lb. product/A
Cabbage Looper Fall Armyworm	1 lb. (16 oz.)	(0.52 lb. a.i./A) and at 7 day spray intervals for rates greater than 0.52 lb. product/A (0.5 lb. a.i./A) as necessary to maintain control.*
<b>RESTRICTIONS:</b> *Do not apply more than 2.1 lbs. product/A (2 lbs. a.i./A) per season. Allow at least 21 days to elapse between final application and harvest. All celery must be trimmed (tops removed) before shipment for use. Do not use trimmed tops for food or feed.		



**USE DIRECTIONS FOR COMMERCIALY TREATED COTTONSEED**

ACEPHATE 97% PRILLS contains a water soluble systemic insecticide which helps to reduce injury to cotton seedlings by several insects ACEPHATE 97% PRILLS can be mixed in the slurry tank with most of the fungicide seed treatments in common use It also can be applied as a separate treatment using enough water to give adequate coverage of the seed

PESTS CONTROLLED	AMOUNT OF ACEPHATE 97% PRILLS PER 100 LBS COTTONSEED	EFFICACY
Cotton Aphids Thrips (Including Western Flower Thrips)	0.28 - 0.4 lb (4.5 - 6.4 oz) Lower rate not approved for use in California	Gives effective reduction of thrips and cotton aphids for up to three weeks after planting
Black (Greasy) Cutworm	0.28 - 0.4 lb (4.5 - 6.4 oz) Lower rate not approved for use in California	Gives effective reduction of Black (Greasy) Cutworm from planting through the 3 <sup>rd</sup> to 4 <sup>th</sup> leaf stage of development When planting into fields where large cutworms are present (5 <sup>th</sup> instar and larger) economic damage may occur

**RESTRICTIONS**

- 1 Do not apply more than 4.0 lbs a.i./A per drop cycle This includes the uses of acephate such as commercial seed treatments in furrow spray and foliar applications
- 2 Do not use treated seed for food feed or oil production Excess treated seed may be used for ethanol production only if (1) by products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by products that are used in agronomic practice
- 3 Treated seed must be planted into the soil at the depth of 1 inch or greater
- 4 Store away from feeds and other foodstuffs
- 5 Wear long sleeved shirt long pants and chemical resistant gloves when handling treated seed
- 6 Dispose of seed packaging in accordance with local requirements
- 7 Exposed treated seed may be hazardous to birds Dispose of all excess treated seed by burial away from bodies of water Cover or incorporate spilled treated seeds

When using formulations that do not contain dye compliance with 21 CFR Chapter 1 Section 2.5 requires that all seed treated with a pesticide must be colored to distinguish from and prevent inadvertent use as food for man or feed for animals

**Seed Container Label Requirements**

Seed commercially treated with Acephate 97% PRILLS must be labeled in compliance with the Federal Seed Act In addition all bags containing treated seed must be labeled or tagged with the following information

- 1 This seed has been treated with Acephate 97% PRILLS which contains x mg\* [*to be filled in after seed treatment*] active ingredient (ai) per seed
- 2 Do not apply more than 4.0 lbs a.i./A per drop cycle This includes the uses of acephate such as commercial seed treatments in furrow spray and foliar applications
- 3 Do not use treated seed for food feed or oil production Excess treated seed may be used for ethanol production only if (1) by products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by products that are used in agronomic practice
- 4 Treated seed must be planted into the soil at the depth of 1 inch or greater
- 5 Store away from feeds and other foodstuffs
- 6 Wear long sleeved shirt long pants and chemical resistant gloves when handling treated seed
- 7 Dispose of seed packaging in accordance with local requirements
- 8 Exposed treated seed may be hazardous to birds Dispose of all excess treated seed by burial away from bodies of water Cover or incorporate spilled treated seeds



9/41

Notification to add advisory statements correct use rates and typos – 08 16 2012

Take all precautions and limitations on labeling of all products used in mixtures

**USE DIRECTIONS FOR COTTON**

**RESTRICTIONS APPLICABLE TO ALL USES ON COTTON**

Do not use treated seed for food or feed purposes or process for oil  
 Do not feed gin trash or forage to livestock  
 Do not allow animals to graze on treated areas  
 Do not apply more than 4 12 lbs product/A (4 lbs a 1/A) per season This includes the use of ACEPHATE 97% PRILLS in commercial seed treatment in furrow and foliar applications  
 When applied by air do not apply more than 1 03 lbs product/A (1 lb a 1/A) in California and Arizona Do not apply more than 0 77 lb product/A (0 75 lb a 1/A) for all other areas of the US when applied by air

**RESISTANCE MANAGEMENT**

Cotton pest control programs especially those for control of Silverleaf/Sweetpotato Whitefly populations should employ a properly designed resistance management strategy Such resistance management strategies include mixture or rotation of alternative classes of chemistry including organophosphates carbamates pyrethroids or insect growth regulators Consult your state or area agricultural extension service for local resistance management strategies and advice on alternative insecticides

**IN FURROW APPLICATION**

PESTS CONTROLLED	AMOUNT OF ACEPHATE 97% PRILLS PER ACRE	APPLICATION INSTRUCTIONS
Aphids (Excluding Cotton Aphids in AZ & CA) Black (Greasy) Cutworm (Except CA) Thrips	0 5 to 1 lb (8 to 16 oz) For the Blacklands of Texas use 1 lb product/A ACEPHATE 97% PRILLS	Apply in water a 3 to 5 gals carrier per acre as an in furrow spray Use flat fan nozzles for in furrow application Align nozzles to ensure good spray deposition into the seed furrow Cone type nozzles are not recommended since unacceptable spray deposition may occur Securely fasten spray system tubing to furrow opener and check frequently to ensure proper positioning and operation Use higher listed rates within the specified range for severe insect infestations ACEPHATE 97% PRILLS can be mixed with fungicides that are sprayed in furrow for disease control

**FOLIAR APPLICATIONS**

**Ground Application** Apply in 10 to 25 gallons of water per acre by conventional ground equipment to ensure thorough coverage of the target crop  
**Aerial Application** Make applications of ACEPHATE 97% PRILLS Insecticide in 3 to 10 gallons of water per acre (minimum of 5 gallons per acre in AZ & CA) Use sufficient carrier volume to provide thorough uniform coverage Do not apply ACEPHATE 97% PRILLS at more than 1 03 lbs product/A (1 lb a 1/A) in California and Arizona and not more than 0 77 lb product/A (0 75 lb a 1/A) for all other areas of the United States) Use sufficient carrier volume to provide thorough uniform coverage



Notification to add advisory statements correct use rates and typos – 08 16 2012

PESTS CONTROLLED	AMOUNT OF ACEPHATE 97% PRILLS PER ACRE	APPLICATION INSTRUCTIONS *
Thrips	0.16 to 0.19 lb (2.5 oz to 3.0 oz)	Apply when eggs or insects first appear Repeat applications at 3 day intervals as necessary to maintain control *
Plant bugs (Lygus)	0.25 to 1 lb (4 to 16 oz) AZ & CA 0.75 to 1 lb (12 to 16 oz)	Apply when eggs or insects first appear  Use highest listed rate for Lygus adults that have migrated into cotton  Repeat applications at 3 day spray intervals for rates up to 0.52 lb product/A (0.5 lb a.i./A) and at 7 day spray intervals for rates greater than 0.52 lb product/A (0.5 lb a.i./A) as necessary to maintain control *
Fleahoppers	0.25 lb (4 oz)	Apply when eggs or insects first appear  Repeat applications at 3 day intervals as necessary to maintain control *
Cotton Aphid (excluding AZ & CA)	0.5 to 1 lb (8 to 16 oz)	This insect may develop resistance to various classes of insecticides. Consult your local Agricultural Extension Service for current control recommendations  Repeat applications at 3 day spray intervals for rates up to 0.52 lb product/A (0.5 lb a.i./A) and at 7 day spray intervals for rates greater than 0.52 lb product/A (0.5 lb a.i./A) as necessary to maintain control *
Whitefly (excluding Sweetpotato & Silverleaf Whitefly)	0.5 to 1 lb (8 to 16 oz)	Repeat applications at 3 day spray intervals for rates up to 0.52 lb product/A (0.5 lb a.i./A) and at 7 day spray intervals for rates greater than 0.52 lb product/A (0.5 lb a.i./A) as necessary to maintain control *
Armyworms (excluding Beet Armyworm) Cabbage Looper	1 lb (16 oz)	Apply when eggs appear and repeat at 7 day spray intervals *
<b>RESTRICTIONS</b> Allow at least 21 days to elapse between final application and harvest *See restrictions applicable to all uses on cotton		



Notification to add advisory statements correct use rates and typos – 08 16 2012

**FOLIAR APPLICATIONS TO COTTON (continued)**

PESTS CONTROLLED	AMOUNT OF ACEPHATE 97% PRILLS PER ACRE	APPLICATION INSTRUCTIONS
Bollworm Tobacco Budworm Adults Larvae Eggs (DEADHATCH®)	East of Rockies 0.5 to 1 lb (8 to 16 oz) AZ & CA 1 lb (16 oz)	<p><b>Early Season</b> Use 0.5 lb product/A for light infestation  <b>Mid and Late Season</b> Use 0.75 to 1 lb product/A for moderate to severe infestations</p> <p>Apply when eggs appear and repeat at 3 day spray intervals for rates up to 0.52 lb product/A (0.5 lb a.i./A) and at 7 day spray intervals for rates greater than 0.52 lb product/A (0.5 lb a.i./A) *</p> <p>Moths of budworm larvae are controlled by direct contact with spray. Moth kill is most likely to occur when late evening applications are made.</p> <p>DEADHATCH® Control of emerging larvae by consumption of treated egg casings</p>
Stink bugs	0.75 lb (12 oz)	Apply when eggs appear and repeat at 7 day spray intervals *
Pink Bollworm (AZ & CA)	1 lb (16 oz)	Apply when insects appear and repeat at 7 day spray intervals *
Cutworms	0.75 lb (12 oz)	<p>Ground application is recommended  Aerial applications are less effective but may be used</p> <p>Control is most effective when ground application is made in the evenings and sprays are directed toward the base and lower portion of plant</p> <p>Apply when insects first appear or damage is first noted and repeat applications at 7 day spray intervals as necessary to maintain control *</p>
<p><b>RESTRICTIONS</b>  Allow at least 21 days to elapse between final application and harvest  *See restrictions applicable to all uses on cotton</p>		

**COTTON TANK MIXES FOR FOLIAR APPLICATION**

**Ground Application** Apply in 10 to 25 gallons of water per acre by conventional ground equipment to ensure thorough coverage of the target crop

**Aerial Application** Make applications of ACEPHATE 97% PRILLS Insecticide in 3 to 10 gallons of water per acre (minimum of 5 gallons per acre in AZ & CA) Use sufficient carrier volume to provide thorough uniform coverage Do not apply ACEPHATE 97% PRILLS at more than 1.03 lbs product/A (1 lb a.i./A) in California and Arizona and not more than 0.77 lb product/A (0.75 lb a.i./A) for all other areas of the United States Use sufficient carrier volume to provide thorough uniform coverage

**RESTRICTIONS APPLICABLE TO ALL USES ON COTTON**

Do not use treated seed for food or feed purposes or process for oil

Do not feed gin trash or forage to livestock

Do not allow animals to graze on treated areas

Do not apply more than 4.12 lbs product/A (4 lbs a.i./A) per season This includes the use of ACEPHATE 97% PRILLS in commercial seed treatment in furrow and foliar applications



12/41

Notification to add advisory statements, correct use rates and typos – 08 16 2012

Always read and follow all label directions when using any pesticide alone or in tank mix combinations. Observe all restrictions and precautions that appear on all product labels. The most restrictive labeling applies when using a tank mix.

PESTS CONTROLLED	AMOUNT OF ACEPHATE 97% PRILLS + TANK MIX PARTNER PER ACRE	APPLICATION INSTRUCTIONS
Sweetpotato & Silverleaf Whitefly	0.5 lb. (8 oz.) + DANITOL® 2.4 EC 8 to 16 fl. oz.	Apply when insects first appear or when damage is first noted and repeat applications at 3 day spray intervals as necessary to maintain control.*  Use a minimum rate of 10 2/3 fl. oz/acre DANITOL® 2.4 EC (0.2 lb. a.i./A) in CA and AZ.  Always read and follow all applicable directions, restrictions, and precautions on the EPA registered label for DANITOL® 2.4 EC Spray.
Armyworms (excluding Beet Armyworm) Bollworm Cabbage Looper Cotton Aphid Fleahoppers Grasshoppers Pink Bollworm Plant bugs (Lygus, Mirids) Salt Marsh Caterpillar Thrips Tobacco Budworm Whitefly (Excluding Sweetpotato & Silverleaf Whitefly)	0.25 to 1 lb. (4 to 16 oz.) + LORSBAN® 4E 1 to 2 pts.	Always read and follow all applicable directions, restrictions, and precautions on the EPA registered label for LORSBAN® 4E.  Apply when insects first appear or when damage is first noted and repeat applications at 3 day spray intervals for rates up to 0.52 lb. product/A (0.5 lb. a.i./A) and at 7 day spray intervals for rates greater than 0.52 lb. product/A (0.5 lb. a.i./A) as necessary to maintain control.*  Use higher rates within the specified range for severe insect infestations.
Cutworms	0.75 lb. (12 oz.) + LORSBAN® 4E 1 to 2 pts.	Apply when insects first appear or when damage is first noted and repeat application at 7 day spray intervals as necessary to maintain control.*  Control is most effective when ground application is made in the evenings and sprays are directed toward the base and lower portion of plant.  Directed, ground application is most effective. Aerial applications are less effective, but may be used.
<b>RESTRICTIONS:</b> Allow at least 21 days to elapse between final application and harvest. *See restrictions applicable to all uses on cotton.		



Notification to add advisory statements correct use rates and typos – 08 16 2012

**COTTON TANK MIXES WITH PYRETHROIDS FOR FOLIAR APPLICATIONS**

**Ground Application** Apply in 10 to 25 gallons of water per acre by conventional ground equipment to ensure thorough coverage of the target crop

**Aerial Application** Make applications of ACEPHATE 97% PRILLS Insecticide in 3 to 10 gallons of water per acre (minimum of 3 gallons per acre in AZ & CA) Use sufficient carrier volume to provide thorough uniform coverage Do not apply ACEPHATE 97% PRILLS at more than 1.03 lbs product/A (1 lb a i /A) in California and Arizona and not more than 0.77 lb product/A (0.75 lb a i /A) for all other areas of the United States Use sufficient carrier volume to provide thorough uniform coverage

**RESTRICTIONS APPLICABLE TO ALL USES ON COTTON**

- Do not use treated seed for food or feed purposes or process for oil
- Do not feed gin trash or forage to livestock
- Do not allow animals to graze on treated areas
- Do not apply more than 4.12 lbs product/A (4 lbs a i /A) per season This includes the use of ACEPHATE 97% PRILLS in commercial seed treatment in furrow and foliar applications

Always read and follow all label directions when using any pesticide alone or in tank mix combinations Observe all restrictions and precautions that appear on all product labels The most restrictive labeling applies when using a tank mix Synthetic Pyrethroids should be used within the guidelines of state and/or regional resistance management programs and recommendations

PESTS CONTROLLED	AMOUNT OF ACEPHATE 97% PRILLS + TANK MIX PARTNER PER ACRE	APPLICATION INSTRUCTIONS
Aphids Bollworm Cabbage Looper Cotton Leaf Perforator Fall Armyworm Fleahoppers Pink Bollworm (AZ & CA) Plant bugs Sweetpotato & Silverleaf Whitefly Thrips Tobacco Budworm Whiteflies	0.5 to 1 lb /A (8 to 16 oz) + AMMO® 2.5 EC (Follow label directions) or ASANA® XL (Follow label directions) or BAYTHROID® 2 EC (Follow label directions) or CAPTURE® 2 EC (Follow label directions) or KARATE® 1E (Follow label directions) or SCOUT® X TRA (Follow label directions)	Begin applications when eggs or insects first appear  Repeat application at 3 day spray intervals for rates up to 0.52 lb product/A (0.5 lb a i /A) and at 7 day spray intervals for rates greater than 0.52 lb product/A (0.5 lb a i /A) as necessary to maintain control *  Use higher rates within the specified range for severe infestations



Notification to add advisory statements correct use rates and typos – 08 16 2012

Cutworms	0.75 lb /A (12 oz) + Any Synthetic Pyrethroid listed above	Control is most effective when ground applications are made in the evenings and sprays are directed toward the base and lower portion of plant  Repeat applications at 7 day spray intervals as necessary to maintain control *  Directed ground applications are most effective Aerial applications are less effective but may be used
Stink Bugs	0.75 lb /A (12 oz) + Any Synthetic Pyrethroid listed above	Apply when eggs or insects first appear Repeat application at 7 day spray intervals necessary to maintain control *
<b>RESTRICTIONS</b> Allow at least 21 days to elapse between final application and harvest *See restrictions applicable to all uses on cotton		

---

**USE DIRECTIONS FOR FOLIAR APPLICATIONS TO CRANBERRY**

---

**Ground Application** Apply with conventional ground equipment in sufficient water to ensure thorough coverage of the target crop

**Aerial Application** Make aerial applications in a minimum of 2 gallons of water per acre Use sufficient carrier volume to provide thorough uniform coverage

**Chemigation Application** This product may only be applied through sprinkler irrigation systems including center pivot lateral move end tow side (wheel) roll travelers big gun solid set or hand move Do not apply this product through any other type of irrigation system Constant agitation must be maintained in the chemical supply tank during the entire period of insecticide application Inject the product with a positive displacement pump into the main line ahead of a right turn to ensure adequate mixing Application of more than label recommended quantities of irrigation water per acre may result in decreased product performance by removing the chemical from the zone of effectiveness Crop injury lack of effectiveness or illegal pesticide residues in the crop may result from non uniform distribution of treated water

**RESTRICTIONS AND PRECAUTIONS FOR APPLICATIONS THROUGH CHEMIGATION SYSTEMS**

**Do not connect an irrigation system (including greenhouse/shadehouse systems) used for pesticide application to a public water system unless the label prescribed safety devices for public water systems are in place**

- A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise
- The system must contain a functional check valve vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow
- The pesticide injection pipeline must contain a functional automatic quick closing check valve to prevent the flow of fluid back toward the injection pump
- The pesticide injection pipeline must also contain a functional normally closed solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down
- The system must contain functional interlocking controls to automatically shut off the pesticide



Notification to add advisory statements correct use rates and typos – 08 16 2012

- injection pump when the water pump motor stops
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected
- Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock
- Do not apply when wind speed favors drift beyond the area intended for treatment
- If you have questions about calibration you should contact State Extension Service specialists equipment manufacturers or other experts
- **Solid Set System** Apply specified dosage for the entire length of the irrigation period or for a 30 to 60 minute period at the end of a regular irrigation set or as a 30 to 60 minute injection as a separate application not associated with a regular irrigation Allow time for all lines to flush the pesticide through all nozzles before turning off irrigation water To ensure the lines are flushed and free of remaining pesticide a dye indicator may be injected into the lines to mark the end of the application period See Note below
- **Center Pivot Systems** Inject the specified dosage per acre continuously for one complete revolution of the system See Note below

Constant agitation must be maintained in the chemical supply tank during the entire period of insecticide application Inject the product with a positive displacement pump into the main line ahead of a right turn to ensure adequate mixing

**Application of more than label specified quantities of irrigation water per acre may result in decreased product performance by removing the chemical from the zone of effectiveness**

PESTS CONTROLLED	AMOUNT OF ACEPHATE 97% PRILLS PER ACRE	APPLICATION INSTRUCTIONS
Cranberry Blossomworm Cranberry Cutworm Gypsy Moth False Armyworm Fireworms Spanworms Sparganothis	1 lb (16 oz)	Apply when insects first appear
<b>RESTRICTIONS</b> Limit to one application per growing season Do not apply more than 1.03 lbs product/A (1 lb a 1/A) per season The Pre Harvest Interval (PHI) is 75 days in the states of MA NJ OR WA WI and 90 days in all other states Do not apply from start of bloom until all berries set		

16/41

Notification to add advisory statements correct use rates and typos – 08 16 2012

**USE DIRECTIONS FOR FOLIAR APPLICATIONS TO HEAD LETTUCE**  
**—Crisphead Type Only—**

**Ground Application** Apply in 10 to 60 gallons of water per acre by conventional ground equipment to ensure thorough coverage of the target crop

**Aerial Application** Make aerial applications in minimum of 5 gallons of water per acre Use sufficient carrier volume to provide thorough uniform coverage

PESTS CONTROLLED	AMOUNT OF ACEPHATE 97% PRILLS PER ACRE	APPLICATION INSTRUCTIONS
Aster Leafhopper Green Peach Aphid	0.5 to 1 lb (8 to 16 oz)	Repeat application at 3 day spray intervals for rates up to 0.52 lb product/A (0.5 lb a.i./A) and at 7 day spray intervals for rates greater than 0.52 lb product/A (0.5 lb a.i./A) as necessary to maintain control *
Armyworm (excluding Beet Armyworm) Cabbage Looper	1 lb (16 oz)	Use higher rates within the specified range for severe infestations

**RESTRICTIONS**

\*Do not apply more than 2.1 lbs product/A (2 lbs a.i./A) per season  
 Allow at least 21 days to elapse between final application and harvest  
 Do not feed trimmings to livestock or allow animals to graze on treated areas  
 For use on Spring Summer and early Fall crops in all areas Winter crops in Florida and Texas late Fall crops in Arizona and Winter crops in Arizona and California  
 Do not apply after first head begins to form in crops which germinate from mid September through November in desert areas of AZ & CA



Notification to add advisory statements correct use rates and typos – 08 16 2012

**USE DIRECTIONS FOR FOLIAR APPLICATIONS TO NON-BEARING CITRUS  
(Newly Planted Groves) — Agricultural Settings ONLY**

**FOLIAR APPLICATIONS**

**Ground Application** Apply in 100 to 200 gallons of water per acre by conventional ground equipment to ensure thorough coverage of the target crop

PESTS CONTROLLED	AMOUNT OF ACEPHATE 97% PRILLS PER ACRE	APPLICATION INSTRUCTIONS
Aphids Grasshoppers Katydid Mealybugs Orangedog Plant bugs Thrips Whiteflies (except Sweetpotato & Silverleaf Whitefly)	0.5 lb (8 oz)	Apply at 3 day spray intervals as necessary to maintain control  Spray individual juvenile or non bearing trees for coverage with total application not to exceed specified rate in lbs per acre Length of residual activity will depend upon spray coverage and the amount of moisture following application
Citrus Blackfly*	0.5 to 0.75 lb (8 to 12 oz)	Apply when eggs or insects first appear Use the high rate when a heavy infestation is present Repeat at 3 day spray intervals for rates up to 0.52 lb product/A (0.5 lb a 1/A) and at 7 day spray intervals for rates greater than 0.52 lb product/A (0.5 lb a 1/A)
Ants (excluding fire harvester carpenter and fire ants)	0.75 lb (12 oz)	Repeat at 7 day spray intervals as necessary to maintain control

**RESTRICTIONS**

Do not graze treated areas  
Do not harvest citrus for one year after treatment (365 days)  
\*Not registered in California for control of Citrus Blackfly

Notification to add advisory statements, correct use rates and typos - 08 16 2012

**SOIL MOUND DRENCHES**

PESTS CONTROLLED	AMOUNT OF ACEPHATE 97% PRILLS PER ACRE	APPLICATION INSTRUCTIONS
Fire Ants & Harvester Ants*	0.75 oz./ 5 gals.	<p>Apply 1 gal. of mix to each mound area by sprinkling the mound until it is wet and treat a four (4) foot diameter circle around the mound.</p> <p><b>Note:</b> For best results, apply the material in the early morning or late afternoon when the ants are most active. Applications made under prolonged dry and hot conditions may be ineffective due to the location of the ants deep within the nest.</p>
<p><b>RESTRICTIONS:</b>            Do not graze treated areas.            Do not harvest citrus for one year after treatment (365 days).            Do not treat mound more than once per season.            *Not registered in California for control of Harvester Ants.</p>		

**USE DIRECTIONS FOR NON-CROP AREAS**

**FIELD BORDERS, FENCEROWS, ROADSIDES, DITCHBANKS, AND BORROW PITS**

**Ground Application:** Apply in 10 to 20 gallons of water per acre by conventional ground equipment to ensure thorough coverage.

**Aerial Application:** Make aerial applications in minimum of 1 to 5 gallons of water per acre (minimum 5 gallons/A in CA). Use sufficient carrier volume to provide thorough, uniform coverage.

PESTS CONTROLLED	AMOUNT OF ACEPHATE 97% PRILLS PER ACRE	APPLICATION INSTRUCTIONS
Grasshoppers	0.25 lb. (4 oz.)	<p>For early to mid-season application.</p> <p>Use the higher volumes when spraying dense foliage.</p> <p>An approved drift retardant may be added to aid in controlling drift and reducing evaporation of aerial sprays.</p>
<p><b>RESTRICTIONS:</b>            Do not graze or feed vegetation cut from treated areas.</p>		



Notification to add advisory statements, correct use rates and typos – 08 16 2012

**WASTELAND**

**Ground Application:** Apply in 10 to 20 gallons of water per acre by conventional ground equipment to ensure thorough coverage.

**Aerial Application:** Make aerial applications in minimum of 1/2 gallon of water per acre (minimum 5 gallons/acre in CA). Use sufficient carrier volume to provide thorough, uniform coverage.

PESTS CONTROLLED	AMOUNT OF ACEPHATE 97% PRILLS PER ACRE	APPLICATION INSTRUCTIONS
Black Grass Bugs Grasshoppers Mormon Cricket	1.5 oz. to 2 oz.	Use the higher volumes when spraying dense foliage  An approved drift retardant may be added to aid in controlling drift and reducing evaporation of aerial sprays.  Use higher rates within the specified range for severe infestations.

**RESTRICTIONS:**  
Do not make more than one application per season.  
Do not graze or feed vegetation cut from treated areas.

**USE DIRECTIONS FOR FOLIAR APPLICATIONS TO PEANUTS**

**Ground Application:** Apply in 10 to 50 gallons of water per acre by conventional ground equipment to ensure thorough coverage.

**Aerial Application:** Make aerial applications in 5 to 10 gallons of water per acre. Use sufficient carrier volume to provide thorough, uniform coverage.

PESTS CONTROLLED	AMOUNT OF ACEPHATE 97% PRILLS PER ACRE	APPLICATION INSTRUCTIONS
Grasshoppers	0.25 to 0.5 lb. (4 to 8 oz.)	Apply when eggs or insects first appear. Repeat application at 3 day spray intervals for rates up to 0.52 lb. product/A (0.5 lb. a.i./A) and at 7 day spray intervals for rates greater than 0.52 lb. product/A (0.5 lb. a.i./A) as necessary to maintain control.*
Thrips	0.375 to 0.75 lb. (6 to 12 oz.)	
Corn Earworm Fall Armyworm Leafhoppers Loopers Velvetbean Caterpillar	0.75 to 1 lb. (12 to 16 oz.)	Use higher rates within the specified range for severe insect infestations.  ACEPHATE 97% PRILLS can be tank mixed with registered at-cracking and early post-emergence peanut herbicides, provided those products do not prohibit tank mixes, provided the most restrictive of label limitations and precautions are observed, and provided no label dosage rates are exceeded.  To determine physical compatibility, pour the specified proportions of each chemical with the same proportion of water as will be present in the chemical supply tank, into a suitable container. Mix thoroughly and allow to stand for five minutes. If the combination remains mixed, or can be remixed readily, the mixture is considered physically compatible. When mixing wettable powder or dry flowable formulations, add and disperse these first, then add liquid pesticides. Do not use the mix combination if there is any separation which cannot be readily remixed.  Combinations should be kept agitated and applied immediately. Do not allow combinations to set for prolonged periods in the chemical supply tank or irrigation lines.



Notification to add advisory statements, correct use rates and typos – 08 16 2012

**RESTRICTIONS:**

\*Do not apply more than 4.12 lbs. product/A (4 lbs. a.i./A) per season.  
Allow at least 14 days to elapse between final application and digging or lifting of peanuts.  
Do not feed treated forage or hay to livestock.  
Do not allow animals to graze on treated areas.

**USE DIRECTIONS FOR FOLIAR APPLICATIONS TO PEPPERMINT AND SPEARMINT**

**Ground Application:** Apply in 20 to 100 gallons of water per acre by conventional ground equipment to ensure thorough coverage of the target crop.

**Aerial Application:** Make aerial applications in minimum of 5 to 10 gallons of water per acre. Use sufficient carrier volume to provide thorough, uniform coverage.

PESTS CONTROLLED	AMOUNT OF ACEPHATE 97% PRILLS PER ACRE	APPLICATION INSTRUCTIONS
Alfalfa Looper Aphids Cutworms	1 lb. (16 oz.)	Apply when eggs or insects first appear. Make one repeat application at no less than 7 day spray interval, if necessary, to maintain control.*
Strawberry Root Weevil Adult Black Vine Weevil Adult	1 lb. (16 oz.)	For control, apply in water in a minimum of 40 gals. spray per acre by ground. Good spray coverage and canopy spray penetration is critical for control. Increase spray volume and pressure with tall or dense mint canopy. Apply after adult emergence is complete but prior to egg laying. Apply at dusk or during the night on a warm still evening. Two applications 10 to 14 days apart may be necessary to reduce heavy infestations.

**RESTRICTIONS:**

\*Do not apply more than 2.1 lbs. product/A (2 lbs. a.i./A) per season. Allow at least 14 days to elapse between final application and harvest.  
The grazing of animals on treated areas and the feeding of spent mint hay to animals are prohibited.



Notification to add advisory statements correct use rates and typos – 08 16 2012

**USE DIRECTIONS FOR FOLIAR APPLICATIONS TO PEPPERS**

**BELL**

**Ground Application** Apply in 25 to 150 gallons of water per acre by conventional ground equipment to ensure thorough coverage

**Aerial Application** Make aerial applications in minimum of 3 gallons of water per acre (minimum of 5 gal /A in CA) Use sufficient carrier volume to provide thorough uniform coverage

PESTS CONTROLLED	AMOUNT OF ACEPHATE 97% PRILLS PER ACRE	APPLICATION INSTRUCTIONS
Grasshoppers	0.25 to 0.5 lb (4 to 8 oz)	Apply when eggs or insects first appear
Cabbage Looper Green Peach Aphid Tobacco Hornworm	0.5 to 1 lb (8 to 16 oz)	Repeat at 3 day spray intervals for rates up to 0.52 lb product/A (0.5 lb a.i./A) and at 7 day spray intervals for rates greater than 0.52 lb product/A (0.5 lb a.i./A) as necessary to maintain insect pest populations below economically damaging numbers *
European Corn Borer	0.75 to 1 lb (12 to 16 oz)	Use higher rates within the specified range for severe infestations

**RESTRICTIONS**

\*Do not apply more than 2.1 lbs product/A (2 lbs a.i./A) per season  
Allow at least 7 days to elapse between final application and harvest

**NON BELL (For use only in Midwestern states, Eastern states, and Puerto Rico)**

**Ground Application** Apply in 40 to 150 gallons of water per acre by conventional ground equipment to ensure thorough coverage

PESTS CONTROLLED	AMOUNT OF ACEPHATE 97% PRILLS PER ACRE	APPLICATION INSTRUCTIONS
Aphids	0.5 lb (8 oz)	Repeat at 3 day spray intervals as necessary *

**RESTRICTIONS**

\*Do not apply more than 1.03 lbs product/A (1 lb a.i./A) per season  
Allow at least 7 days to elapse between final application and harvest

Notification to add advisory statements correct use rates and typos – 08 16 2012

**USE DIRECTIONS FOR SOYBEANS**

<b>PESTS CONTROLLED</b>	<b>AMOUNT OF ACEPHATE 97% PRILLS PER ACRE</b>	<b>APPLICATION INSTRUCTIONS</b>
Grasshopper Thrips	0.25 to 0.5 lb (4 to 8 oz)	Apply when eggs or insects first appear
Potato Leafhopper Stinkbugs	0.5 to 1 lb (8 to 16 oz)	Repeat applications as necessary to maintain insect pest populations below economically damaging numbers
Armyworms (except Beet) Bean Leaf Beetle Cabbage Looper Green Cloverworm Mexican Bean Beetle Soybean Aphid Threecornered Alfalfa Hopper Velvetbean Caterpillar	0.75 to 1 lb (12 to 16 oz)	Repeat application at 3 day spray intervals for rates up to 0.52 lb product/A (0.5 lb a 1/A) and at 7 day spray intervals for rates greater than 0.52 lb product/A (0.5 lb a 1/A) as necessary to maintain control  <b>Ground Application</b> Apply in 10 to 50 gallons per acre by conventional ground equipment to ensure thorough coverage  <b>Aerial Application</b> Make aerial applications in 5 to 10 gallons per acre (minimum of 5 gal /A in CA) Use sufficient carrier volume to provide thorough uniform coverage
<b>RESTRICTIONS</b> Do not apply more than 1.55 lbs product/A (1.5 lbs a 1/A) per crop cycle Allow at least 14 days to elapse between final application and harvest Do not graze or cut vines for hay or forage		



Notification to add advisory statements, correct use rates and typos - 08 16 2012

**TOBACCO PLANT BED APPLICATION (EXCEPT CA)**

PESTS CONTROLLED	AMOUNT OF ACEPHATE 97% PRILLS PER ACRE	APPLICATION INSTRUCTIONS
Cutworm Flea Beetle Green Peach Aphid Tobacco Aphid	0.75 lb. (12 oz.)	Apply to foliage at the equivalent of 3/4 tablespoon in 1 gal. of water per every 1,000 sq. ft. of bed. Apply evenly to ensure thorough coverage. Apply at 7 day intervals as needed.

**RESTRICTIONS:**  
Do not apply more than 4.12 lbs. product/A (4 lbs. a.i./A) per season. This includes the total use of ACEPHATE 97% PRILLS in transplant water, plant bed, soil, float bed (greenhouse), and foliar applications.

**TOBACCO GREENHOUSE (FLOATBED) APPLICATION (EXCEPT CA)**

PESTS CONTROLLED	AMOUNT OF ACEPHATE 97% PRILLS PER ACRE	APPLICATION INSTRUCTIONS
Cutworm Flea Beetle Green Peach Aphid Tobacco Aphid	0.75 lb. (12 oz.)	Apply to foliage at the equivalent of 3/4 tablespoon in 3 gals. water per every 1,000 sq. ft. of bed.  Apply evenly to ensure thorough coverage. Apply at 7 day intervals as needed.

**RESTRICTIONS:**  
Do not apply more than 4.12 lbs. product/A (4 lbs. a.i./A) per season. This includes the total use of ACEPHATE 97% PRILLS in transplant water, plant bed, soil, float bed (greenhouse), and foliar applications.  
Dispose of floatbed water in the transplanted field by either transplant water or foliar spray.



Notification to add advisory statements, correct use rates and typos – 08 16 2012

**USE DIRECTIONS FOR TOBACCO**

**TOBACCO TRANSPLANT WATER APPLICATION (EXCEPT CA)**

PESTS CONTROLLED	AMOUNT OF ACEPHATE 97% PRILLS PER ACRE	APPLICATION INSTRUCTIONS
Cutworms Flea Beetle Green Peach Aphid Potato Tuberworm Tobacco Aphid Tobacco Thrips	0.75 lb. (12 oz.)	Provides control of early season flea beetles, green peach aphids, tobacco aphids and cutworms for approximately 3 to 4 weeks after transplanting. For later season control of these insects, apply a foliar spray of ACEPHATE 97% PRILLS.  Apply in a minimum of 100 gals. of transplant water per acre. ACEPHATE 97% PRILLS should be pre-mixed in water to form a slurry solution prior to adding to the transplant water tank.  Do not apply more than 0.75 lb. ACEPHATE 97% PRILLS per acre as a transplant water application as some phytotoxicity may occur.

**RESTRICTIONS:**  
Do not apply more than 4.12 lbs. product/A (4 lbs. a.i./A) per season. This includes the total use of ACEPHATE 97% PRILLS in transplant water, plant bed, soil, float bed (greenhouse), and foliar applications.  
Make transplant-water applications using mechanical transplant equipment only. Using such equipment, the insecticide/water mixture is mechanically applied directly into the soil along with the transplanted plants.

**TOBACCO FOLIAR APPLICATION (EXCEPT CA)**

**Ground Application:** Apply in 10 to 50 gallons of water per acre by conventional ground equipment to ensure thorough coverage.

**Aerial Application:** Make aerial applications in minimum of 3 gallons of water per acre. Use sufficient carrier volume to provide thorough, uniform coverage.

PESTS CONTROLLED	AMOUNT OF ACEPHATE 97% PRILLS PER ACRE	APPLICATION INSTRUCTIONS
Grasshoppers	0.25 to 0.5 lb. (4 to 8 oz.)	Apply at 3 day spray intervals for rates up to 0.52 lb. product/A (0.5 lb. a.i./A) and at 7 day spray intervals for rates greater than 0.52 lb. product/A (0.5 lb. a.i./A) as needed.*
Green Peach Aphid Flea Beetle Hornworm Tobacco Thrips	0.5 lb. (8 oz.)	Use higher listed rates within the specified range for severe infestations.
Stink bugs Tobacco Aphid Vegetable Weevils	0.5 to 0.75 lb. (8 to 12 oz.)	
Budworm Cabbage Looper Cutworm Japanese Beetle	0.75 lb. (12 oz.)	

**RESTRICTIONS:**  
\*Do not apply more than 4.12 lbs. product/A (4 lbs. a.i./A) per season. This includes the total use of ACEPHATE 97% PRILLS in transplant water, plant bed, soil, float bed (greenhouse), and foliar applications.  
Allow at least 3 days to elapse between final application and harvest.



Notification to add advisory statements, correct use rates and typos - 08 16 2012

**TOBACCO SOIL MOUND DRENCH APPLICATION (EXCEPT CA)**

PESTS CONTROLLED	AMOUNT OF ACEPHATE 97% PRILLS	APPLICATION INSTRUCTIONS
Fire Ant & Harvester Ant	Mix 0.75 oz. in 5 gals. of water.	Apply 1 gal. of mix to each mound area by sprinkling the mound until it is wet and treat a four (4) foot diameter circle around the mound when insects first appear.  Note: For best results apply the material in the early morning or late afternoon when the ants are most active. Applications made under prolonged dry and hot conditions may be ineffective due to the location of the ants deep within the nest
<b>RESTRICTIONS:</b> Do not apply more than 4.12 lbs. product/A (4 lbs. a.i./A) per season. This includes the total use of ACEPHATE 97 in transplant water, plant bed, soil, float bed (greenhouse), and foliar applications. Allow at least 3 days to elapse between final application and harvest. Treat maximum of 13 mounds per acre. Do not treat more than once per season.		

**SPECIALITY USES**

**CROP TOLERANCE**

ACEPHATE 97% PRILLS Insecticide is generally not phytotoxic to most greenhouse/shadehouse or field grown plants or turf. However, it is impossible to test all plant varieties and cultural conditions. Before treating large plantings, apply to a representative group of plants and observe for two weeks to assure that a particular variety, grown under current conditions is not sensitive to ACEPHATE 97% PRILLS. Use on turfgrass is limited to sod farms and golf courses, except when applying by mound or spot treatment for fire ant and harvester ant control.

The following have been found to be sensitive to ACEPHATE 97% PRILLS:

**Trees:** Balm of Gilead, Cottonwood, Lombardy Poplar or Viburnum suspensum and Crabapple varieties, Ichonoski, Malus floribunda, Pink Perfection, Red Wine and Snow Cloud.

**Plants:** Bletchum gibbum, Cissus antarctica, Ficus triangularis, Fittonia verschaffeltii, Maranta leuconeura kerchoveana, Pachystachya lutea, Plectranthus australis, Polypodium aureus, Polystichum, Pteris ensiformis, Tolmiea menziesii. **Chrysanthemum Varieties:** Albatross, Bonnie Jean, Dixie, Garland, Gem, Iceberg, Pride, Showoff, Statesman, Tally Ho, Westward Ho and Wild Honey. Applications to Chrysanthemums and Roses with open flowers may result in flower damage.

**SPECIALITY USES**

**SMALL SPRAY EQUIPMENT**

PREPARATION OF ACEPHATE 97% PRILLS SPRAY MIX USING A 16 OUNCE MEASURING CUP		
LABEL DIRECTION PER ACRE IN 100 GALLONS OF WATER	EQUIVALENT PER ACRE IN 25 GALLONS OF WATER*	EQUIVALENT PER ACRE IN 1 GALLON WATER (TEASPOONS/GAL)
0.25 lb.	1 oz.	1/3 tsp.
0.50 lb.	2 oz.	2/3 tsp.
0.60 lb.	2.4 oz.	0.8 tsp.
0.75 lb.	3 oz.	1 tsp.
1.00 lb.	4 oz.	1 1/3 tsp.

Notification to add advisory statements correct use rates and typos - 08 16 2012

**NON BEARING CITRUS**

PLANTS	INSECTS	AMOUNT OF ACEPHATE 97% PRILLS PER ACRE	APPLICATION TIMING	APPLICATION INSTRUCTIONS
Non bearing citrus (including citrus nursery seedlings)	Aphids Grasshoppers Katydid Mealybugs Orangedogs Plant bugs (Lygus) Thrips Whiteflies	0.5 lb (8 oz)	Apply at 3 day spray intervals as needed for control of existing populations	Apply in 100 to 200 gals of water per acre. Spray individual juvenile or non bearing trees for coverage with total application not to exceed specified rate in lbs per acre. Length of residual activity will depend upon spray coverage and the amount of moisture following application.
	Citrus Blackfly	0.5 to 0.75 lb (8 to 12 oz)	Begin applications when eggs or adults first appear.  Apply at 3 day spray intervals for rates up to 0.52 lb product/A (0.5 lb a.i./A) and at 7 day spray intervals for rates greater than 0.52 lb product/A (0.5 lb a.i./A) to maintain control.  Use the higher listed rate when a heavy infestation of Citrus Blackfly is present. Begin applications when eggs or adults first appear.	
Non bearing citrus (including citrus nursery seedlings)	Ants (excluding fire, harvester carpenter and pharaoh ants)	Foliar spray 0.75 lb (12 oz)	Apply at 7 day spray intervals as needed for control of existing populations	
	Fire Ants and Harvester Ants	Mound Treatment — Drench Method	Mix 0.75 oz in 5 gals of water. Apply 1 gal of mix to each mound area by sprinkling the mound until it is wet and treat a four (4) foot diameter circle around the mound.  Applications made under prolonged dry and hot conditions may be ineffective due to the location of the ants deep within the nest.	
<b>RESTRICTIONS</b> Do not allow livestock to graze treated areas Do not harvest citrus for one year after treatment (365 days)				



Notification to add advisory statements correct use rates and typos – 08 16 2012

**NURSERY STOCK NON BEARING DECIDUOUS FRUIT TREES, NUT TREES AND VINES IN NURSERY FIELD OR NON-BEARING ORCHARDS**

**Ground Application** Apply in 50 to 100 gals of water per acre in conventional equipment to ensure thorough coverage of the target plants

**Aerial Application** Apply in a minimum of 5 gals of water per acre Use sufficient carrier volume to provide thorough coverage

PLANTS	INSECTS	AMOUNT OF ACEPHATE 97% PRILLS PER 100 GALS	APPLICATION TIMING
Nursery stock (non bearing Deciduous Fruit Trees Nut Trees Vines) Almond Apricot Apple Cherry Grape Kiwi Pear Pistachio Plum Prune Walnut	Aphids Plant Bugs (Lygus) Thrips	0.5 to 1 lb (8 to 16 oz)	Apply at 3 day spray intervals for rates up to 0.52 lb product/A (0.5 lb a.i./A) and at 7 day spray intervals for rates greater than 0.52 lb product/A (0.5 lb a.i./A) as needed for control of existing populations  Use higher rates within the specified range for severe infestations

**RESTRICTIONS**  
 For application only to non bearing trees and vines in nursery fields or in non bearing orchards Do not apply to interplants or single trees in a bearing orchard  
 Do not harvest any fruit from sprayed trees for one year after application (365 days)  
 Do not allow livestock to graze treated area  
 Do not exceed 1 lb product/A per 100 gallons of water

Notification to add advisory statements correct use rates and typos – 08 16 2012

**SPECIALTY USES — NON CROP AREAS**

CROP	INSECTS	AMOUNT OF ACEPHATE 97% PRILLS PER 100 GALS
Non crop areas (field borders fencerows roadsides ditchbanks and borrow pits)	Grasshoppers	0.25 lb (4 oz)  <b>Notes</b> For early to mid season application to grasshoppers in field borders fencerows roadsides ditch banks and borrow pits
	<b>RESTRICTIONS</b> Apply in 1 to 5 gals of water by air or in 10 to 20 gals of water with ground equipment Use the higher volumes when spraying heavier foliage Do not graze or feed vegetation cut from treated areas	
	Fire Ants and Harvester Ants	Mound Treatment —Drench Method Mix 0.75 oz in 5 gals of water Apply 1gal of mix to each mound area by sprinkling the mound until it is wet and treat a four (4) foot diameter circle around the mound  <b>Notes</b> Grass in the treated area may be injured For best results apply the material in the early morning or late afternoon when the ants are most active Applications made under prolonged dry and hot conditions may be ineffective due to the location of the ants deep within the nest
	<b>RESTRICTIONS</b> Do not treat mound more than once per season	



Notification to add advisory statements, correct use rates and typos – 08 16 2012

**CONTAINER GROWN NURSERY STOCK**

PLANTS	INSECTS	AMOUNT OF ACEPHATE 97% PRILLS PER 100 GALS.	APPLICATION TIMING	APPLICATION INSTRUCTIONS
Container Grown Nursery Stock (Arborvitae Azalea Camellia Rhododendron Roses Viburnum Yew)	Black Vine Weevil Strawberry Root Weevil	0.75 lb. (12 oz.)	Application should be made by mid-September for greenhouse/shadehouse stock and by mid- October for outdoor stock. Consult your local county extension agent for information on the identification and control of root weevils on ornamentals.	Apply the specified amount of ACEPHATE 97% PRILLS spray per 100 gallons of solution so as to thoroughly drench the root system.  Apply at 7day spray intervals to maintain control.
	<b>RESTRICTIONS:</b> Do not apply more than 0.75 lb. product/A./100 gallons of water.			
	Ants, (excluding fire, harvester, carpenter and pharaoh ants)	0.75 lb. (12 oz.)	Apply as needed to control the pest.	

**ORNAMENTAL TREES AND SHRUBS**

PLANTS	INSECTS	AMOUNT OF ACEPHATE 97% PRILLS PER 100 GALS.	APPLICATION TIMING	APPLICATION INSTRUCTIONS
Trees & Shrubs (except Flowering Crabapple and Douglas Fir and Crape Myrtle) see below	Aphids Bagworms Birch Leafminer Lace Bugs Leafrollers Tent Caterpillar*	0.25 lb. (4 oz.)	As the insects begin to appear.	Apply the specified amount of ACEPHATE 97% PRILLS in 100 gallons of water with a hydraulic sprayer as a full coverage spray.  *Mist Blower Application: Adjust rates to 1 lb. per 100 gallons of water for Gypsy Moth control and 0.75 lb. per 100 gallons of water for Tent Caterpillar control.
	Douglas Fir Tussock Moth Larvae Gypsy Moth Larvae* Webworms	0.5 lb. (8 oz.)	As the insects begin to appear.	The addition of a suitable sticker improves control of Gypsy Moth larvae. Repeat at 14-day spray intervals as necessary to maintain control.



Notification to add advisory statements, correct use rates and typos – 08 16 2012

Scales (crawlers)	0.5 lb. (8 oz.)	As crawlers begin to appear. Repeat applications at a 2-week or more interval may be necessary where there is continuous crawler production.
Ponderosa Pine Needle Miner	0.5 lb. (8 oz.)	Time of application is important. Consult your Farm Advisor or County Extension Agent.
Grasshoppers	0.5 lb. (8 oz.)	As the grasshoppers begin to appear.
California Oakworm Cankerworms (Spring & Fall)	0.25 to 0.5 lb. (4 to 8 oz.)	As the insects begin to appear. Use the higher amount when larger larvae are present.

**RESTRICTIONS:**  
 Do not apply more often than 3 times in a growing season to trees and shrubs.  
 Do not apply more than 1 lb. product/A/100 gallons of water.  
 Refer to Crop Tolerance statement.

**ORNAMENTAL TREES AND SHRUBS (continued)**

PLANTS	INSECTS	AMOUNT OF ACEPHATE 97% PRILLS PER 100 GALS.	APPLICATION TIMING	APPLICATION INSTRUCTIONS
Trees & Shrubs (except Flowering Crabapple and Douglas Fir and <u>Crape Myrtle</u> ) - see below	Nantucket Pine Tip Moth Larvae	0.75 lb. (12 oz.)	Time of application is important. Consult your Farm Advisor or County Extension Agent. Repeat applications will be required for subsequent generations.	Apply the specified amount of ACEPHATE 97% PRILLS in 100 gallons of water with a hydraulic sprayer as a full coverage spray.  Repeat at 14-day spray intervals as necessary to maintain control.
	Root Weevil Adults	0.75 lb. (12 oz.)	Apply when first feeding damage occurs, Repeat applications at four week intervals until the first heavy frost, may be necessary for complete foliage protection.	
	Box Elder Bugs Budworms Leafhoppers Sawflies	0.75 lb. (12 oz.)	As the insects begin to appear.	

Notification to add advisory statements correct use rates and typos – 08 16 2012

	Japanese Beetle	1 lb (16 oz)	As the Japanese Beetles begin to appear Repeat applications at 2 week intervals may be necessary	
	Elm Leaf Beetle (larvae)	1 lb (16 oz)	As the larvae begin to appear ACEPHATE 97% PRILLS will not prevent Elm Leaf Beetle eggs from hatching	
<b>RESTRICTIONS</b> Do not apply more often than 3 times in a growing season to trees and shrubs Do not apply more than 1 lb product/A/100 gallons of water Refer to Crop Tolerance Statement				
Douglas Fir (Christmas Trees)	Douglas Fir Needle Midge	0.5 lb (8 oz) (0.5 lb per acre)	Make applications no more than 2 weeks prior to bud burst For additional pest management information consult your County Extension Agent	Apply the specified amount of ACEPHATE 97% PRILLS in not less than 2 gallons of spray per acre by air or in 100 gallons of spray per acre by ground
<b>RESTRICTIONS</b> Do not apply more than 0.5 lb product/A/100 gallons of water Do not treat more than once per season				

**ORNAMENTAL TREES AND SHRUBS (continued)**

PLANTS	INSECTS	AMOUNT OF ACEPHATE 97% PRILLS PER 100 GALS	APPLICATION TIMING	APPLICATION INSTRUCTIONS
Flowering Crabapple	Aphids Leafrollers Tent Caterpillars	0.25 lb (4 oz)	As the insects begin to appear	Apply the specified amount of ACEPHATE 97% PRILLS in 100 gallons of water with a hydraulic sprayer  Note Phytotoxicity has occurred on the following Crabapple varieties Hops Ichonoski Malus floribunda Pink Perfection Red Wine and Snow Cloud
<b>RESTRICTIONS</b> Do not apply more often than 3 times in a growing season at 28 day intervals Do not apply more than 0.25 lb product/A/100 gallons of water				



Notification to add advisory statements correct use rates and typos – 08 16 2012

Crape Myrtle	Aphids	2 to 3 level Tsp (0.6 to 0.8 oz) per tbsp water as paint on	As aphids begin to appear	Make a paint on slurry by mixing the specified amount of ACEPHATE 97% PRILLS with 1 tablespoon of water. Remove the loose bark from the trunk areas to be treated. Completely paint a band around each trunk to a width twice its diameter. Make applications to trunks within a zone 6 to 12 inches above the ground and below the point where branching begins. For multi trunk plants be certain to treat all trunks. For either single or multi trunk plants make application as low as possible within the specified treatment zone.
--------------	--------	---	---------------------------	---

**OUTDOOR FLORAL CROPS AND GROUND COVERS**

PLANTS	INSECTS	AMOUNT OF ACEPHATE 97% PRILLS PER 100 GALS	APPLICATION TIMING	APPLICATION INSTRUCTIONS
Chrysanthemums Daisies Dahlias Easter Lily Gladioli Gypsophila Pachysandra Peony Roses Sedum Statice Strawflower Yarrow Zinnia	Aphids Lygus Thrips	0.5 lb (8 oz)	As insects begin to appear Repeat applications may be necessary	Apply the specified amount of ACEPHATE 97% PRILLS in 100 gallons of water with a hydraulic sprayer as a full coverage spray.  Repeat at 14 day spray intervals as necessary to maintain control.  Note: Multiple applications may cause slight tip burn or marginal leaf necrosis on some varieties. Test on a few plants to determine varietal susceptibility.
Boston Ivy Roses	Japanese Beetle	1 lb (16 oz)	As the Japanese Beetle begin to appear Repeat applications at 2 week intervals may be necessary	

**RESTRICTIONS**  
Do not apply more than 1 lb product/A/100 gallons of water  
Do not apply more than 0.77 lb product/A (0.75 lb a i /A) to cut flowers

Notification to add advisory statements correct use rates and typos - 08 16 2017

**COMMERCIAL GREENHOUSE/SHADEHOUSE FLORAL AND FOLIAGE PLANT CROPS**

Not for use in residential greenhouses/shadehouses

**Restrictions**

Do not apply under conditions involving possible drift to food forage or other plantings that might be damaged or the crops thereof rendered unfit for sale use or consumption

Do not apply with low pressure handwand equipment

PLANTS	INSECTS	AMOUNT OF ACEPHATE 97% PRILLS PER 100 GALS	APPLICATION TIMING	APPLICATION INSTRUCTIONS
Roses	Leafrollers	0.5 to 0.75 lb (8 to 12 oz)	As leafrollers begin to appear Use the higher amount when large larvae are present	Apply the specified amount of ACEPHATE 97% PRILLS in 100 gallons of water with a hydraulic sprayer as a full coverage spray  Use higher rates within the specified range for severe infestations  <b>Notes</b> Direct Ag Source I.C. does not recommend application to roses with open flowers

**RESTRICTIONS**

Do not apply more than 0.75 lb product/A/100 gallons of water

Do not apply at less than 14 day spray intervals

**COMMERCIAL GREENHOUSE/SHADEHOUSE FLORAL AND FOLIAGE PLANT CROPS**

Not for use in residential greenhouses/shadehouses (continued)

PLANTS	INSECTS	AMOUNT OF ACEPHATE 97% PRILLS PER 100 GALS	APPLICATION TIMING	APPLICATION INSTRUCTIONS
Anthuriums Cacti Foliage Plants Orchids Poinse	Aphids	0.25 lb (4 oz)	As aphids begin to appear	Apply the specified amount of ACEPHATE 97% PRILLS in 100 gallons of water with a hydraulic sprayer as a full coverage spray The addition of a wetting agent may be required on difficult to wet foliage
	Mealybugs Thrips Whiteflies	0.5 lb (8 oz)	As the insects begin to appear A repeat application at a 2 week interval may be necessary for control of mealybugs and whiteflies	<b>Notes</b> Phytotoxicity has occurred on the following plants Blechnum gibbum Cissus antarctica Ficus triangularis Fittonia vershaffeltii Maranta leuconeura kershoweana Plectranthus lutes Plectranthus australis Polypodium aureus Polystichum



Notification to add advisory statements correct use rates and typos – 08 16 2012

Scales (Crawlers)	0.5 lb (8 oz)	As crawlers begin to appear Repeat applications at a 2 week or more interval may be necessary where there is continuous crawler production	Pteris ensiformis Tolmiea menziesii Before treating large plantings, spray only a few plants and observe 2 weeks for varietal phytotoxicity  Application to poinsettias after bract formation may result in phytotoxicity on certain varieties
<b>RESTRICTIONS</b> Do not apply more than 0.5 lb product/A/100 gallons of water Do not apply at less than 14 day spray intervals			

**COMMERCIAL GREENHOUSE/SHADEHOUSE FLORAL AND FOLIAGE PLANT CROPS**  
 Not for use in residential greenhouses/shadehouses (continued)

PLANTS	INSECTS	AMOUNT OF ACEPHATE 97% PRILLS PER 100 GALS	APPLICATION TIMING	APPLICATION INSTRUCTIONS
Anthuriums Cacti Foliage Plants Orchids Poinsettia (continued)	Sweetpotato Silverleaf Whiteflies (except CA)	0.25 lb (4 oz) + TAME® 2.4 EC 10 2/3 fl oz (0.2 lb a.i.)	Apply when insects first appear. If a population is well established, make one application of the tank mix and follow 5 to 7 days later with TAME® 2.4 EC alone at 16 fl oz /100 gallons  See the TAME® 2.4 EC label for instructions	Apply the specified amount of ACEPHATE 97% PRILLS plus TAME® 2.4 EC. Spray as a tank mix at a volume necessary to obtain good coverage. Follow the TAME® 2.4 EC label for specific instructions on the alternation of TAME® 2.4 EC plus ACEPHATE 97% PRILLS and TAME® 2.4 EC alone and the rotation instruction to avoid potential resistance.
<b>RESTRICTIONS</b> Do not apply more than 0.25 lb product/A/100 gallons of water				

Notification to add advisory statements correct use rates and typos – 08 16 2012

**COMMERCIAL GREENHOUSE/SHADEHOUSE FLORAL AND FOLIAGE PLANT CROPS**  
**Not for use in residential greenhouses/shadehouses (continued)**

PLANTS	INSECTS	AMOUNT OF ACEPHATE 97% PRILLS PER 100 GALS	APPLICATION TIMING	APPLICATION INSTRUCTIONS
Carnations Chrysanthemums Roses	Aphids Thrips	0.5 lb (8 oz)	As aphids begin to appear As thrips begin to appear or at the tight flower bud stage Repeat applications may be necessary	<p>Apply the specified amount of ACEPHATE 97% PRILLS in 100 gallons of water with a hydraulic sprayer as a full coverage spray</p> <p><b>Notes</b> Phytotoxicity has occurred on the following Chrysanthemum varieties Albatross Bonnie Jean Dixie Garland Gent Iceberg Pride Showoff Statesman Tally Ho Westward Ho &amp; Wild Honey</p> <p>Before treating large Chrysanthemum plantings spray only a few plants and observe two weeks for varietal phytotoxicity</p> <p>Direct Ag Source LLC does not recommend application to chrysanthemums and roses with open flowers</p>
<p><b>RESTRICTIONS</b> Do not apply more often than once every 5 days for greenhouse and 14 days for shade house Do not apply more than 0.5 lb product/A/100 gallons of water</p>				



Notification to add advisory statements correct use rates and typos – 08 16 2012

**COMMERCIAL TURFGRASS GOLF COURSES AND SOD FARMS ONLY**

**Restrictions**

Do not allow livestock to graze treated areas

Do not feed treated grass to livestock

Use is limited to sod farms and golf courses except when applying by mound or spot treatment for fire ant and harvester ant control Aerial applications to turf are prohibited

PLANTS	INSECTS	AMOUNT OF ACEPHATE 97% PRILLS PER 1000 SQ FT	APPLICATION TIMING	APPLICATION INSTRUCTIONS
Turfgrass (Golf courses and Sod Farms only)	Fall Armyworm Yellow Striped Armyworm Southern Armyworm	0.4 to 0.9 oz (1 to 2.5 lbs per acre)	As the insects appear A repeat application at a 14 day interval may be necessary	Apply the specified amount of ACEPHATE 97% PRILLS per 1 000 sq ft Use a minimum of 5 gallons of water per 1 000 sq ft to obtain good coverage  Use higher rates within the specified range for severe insect infestations
	Cutworm	Golf course 0.9 to 1.5 oz (2.5 to 4.12 lbs per acre)  Sod farms 0.9 to 1.1 oz (2.5 to 3.09 lbs per acre)		
	Chinch Bugs	Golf course 0.9 to 1.5 oz (2.5 to 4.12 lbs per acre)  Sod farms 0.9 to 1.1 oz (2.5 to 3.09 lbs per acre)	Apply as needed for adult population knockdown  Apply at 10 to 14 day intervals as needed	Apply the specified amount of ACEPHATE 97% PRILLS per 1 000 sq ft Use 1 to 15 gallons of water per 1 000 sq ft to obtain good coverage  Use higher rates within the specified range for severe insect infestations
	Leafhopper	0.75 oz (2 lbs per acre)	As the leafhoppers begin to appear A repeat application at a 7 day interval may be necessary	

Notification to add advisory statements, correct use rates and typos – 08 16 2012

<p>Mole Crickets (except CA.) Spittlebug (except CA.)</p>	<p>Golf courses: 0.8 to 1.4 oz. (2.2 to 4 lbs. per acre)  Sod farms: 0.8 to 1.1 oz (2.2 to 3.09 lbs. per acre)</p>	<p>As mole crickets begin to appear. For knockdown of existing populations, more than one application may be required throughout the growing season. For heavy infestations, use the higher dosage rate. Apply at 7 to 14 day intervals as needed.</p>	<p>Apply the specified amount of ACEPHATE 97% PRILLS per 1,000 sq. ft. Use 1 to 15 gallons of water per 1,000 sq. ft. to obtain good coverage. Apply during late afternoon or early evening hours and after irrigation. Do not irrigate after application. Use higher rates within the specified range for severe insect infestations. The use of a lemon fragrance substance in the spray mix may enhance control by acting as a flushing agent and thus provide increased mole cricket contact with the acephate. The following lemon-scented products have been shown to be effective flushing agents: Lemon Joy, Lemon Palmolive, and Mighty Products manufactured-base pure lemon fragrance. The use rate for these lemon-scented products is 2 teaspoons per gallon of water for small total mix volume or 6 fl. oz. per 50 gallons of water for large mix volume.</p>
---	--	--	--

**RESTRICTIONS:**  
Do not apply more than 3.09 lbs. product/A for Sod Farms and 4.12 lbs. product/A for Golf Courses.  
For Sod Farms, allow at least 3 days to elapse between final application and harvesting sod.  
Do not apply aerially.



Notification to add advisory statements, correct use rates and typos – 08 16 2012

**COMMERCIAL TURFGRASS: GOLF COURSES AND SOD FARMS ONLY (continued)**

PLANTS	INSECTS	AMOUNT OF ACEPHATE 97% PRILLS PER 1000 SQ. FT.	APPLICATION TIMING	APPLICATION INSTRUCTIONS
Turfgrass (Golf courses and Sod Farms only)	Sod Webworm (Crambus spp.)	0.4 to 0.8 oz. (1 to 2.2 lbs. per acre)	As sod webworms begin to appear use the higher amount when quick knockdown is needed or with heavy infestations.  Repeat application may be needed.  Apply at 7 to 14 day intervals as needed.	Apply the specified amount of ACEPHATE 97% PRILLS per 1,000 sq. ft. Use 1 to 15 gallons of water per 1,000 sq. ft. to obtain good coverage.  Use higher rates within the specified range for severe insect infestations.
	Greenbug (Shizaphis graminum) Grasshoppers	0.4 oz. (1 lb. per acre)	Apply when insects or their damage first appear.  Apply at 7 to 14 day intervals as needed.	Apply the specified amount of ACEPHATE 97% PRILLS. Use 4 gallons of water per 1,000 sq. ft. to obtain good coverage. Do not mow turfgrass for at least 24 hours after application.
	Black Turfgrass Ataenius (except CA.)	Golf courses: 0.9 to 1.4 oz. (2.5 to 4 lbs. per acre)  Sod Farms: 0.9 to 1.1 oz. (2.5 to 3.09 lbs. per acre)	Apply when insects or their damage first appear.  Apply at 7 to 14 day intervals as needed.	Apply the specified amount of ACEPHATE 97% PRILLS per 1,000 sq. ft. Use a minimum of 5 gallons of water per 1,000 sq. ft. Irrigate lightly after application (not more than 1/2 inch). Use higher rates within the specified range for severe insect infestation
Dichondra (Golf Courses and Sod Farms only)	Cutworm Flea Beetle Southern Armyworm Yellow Striped Armyworm	Golf courses: 0.8 to 1.4 oz. (2.2 to 4 lbs. per acre)  Sod Farms: 0.8 to 1.1 oz. (2.2 to 3.09 lbs. per acre)	Apply when the insects appear.  Repeat at 14 day intervals as necessary.	Apply the specified amount of ACEPHATE 97% PRILLS per 1,000 sq. ft. Use a minimum of 15 gallons of water per 1,000 sq. ft. to obtain good coverage. Use higher rates within the specified range for severe insect infestations.

**RESTRICTIONS:**

Do not apply more than 3.09 lbs. product/A for Sod Farms and 4.12 lbs. product/A for Golf Courses.  
For Sod Farms, allow at least 3 days to elapse between final application and harvesting sod.  
Do not apply aerially.

Notification to add advisory statements correct use rates and typos – 08 16 2012

**MOUND TREATMENT OF FIRE ANTS AND HARVESTER ANTS IN TURFGRASS**

Do not allow livestock to graze treated areas Do not feed treated grass to livestock

PLANTS	INSECTS	AMOUNT ACEPHATE 97% PRILLS	APPLICATION TIMING	APPLICATION INSTRUCTIONS
Turfgrass and Non Crop Areas	Imported Fire Ants and Harvester Ants	Mound Treatment Drench Method Mix 0.75 oz in 5 gallons of water Apply 1 gallon of mix to each mound area by sprinkling the mound until it is wet and treat a four (4) foot diameter circle around the mound	Apply when insects first appear	Apply the specified amount of ACEPHATE 97% PRILLS as directed Grass in treated areas may be injured  For best results apply the material in the early morning or late afternoon when the ants are most active Applications made under prolonged dry and hot conditions may be ineffective due to the location of the ants deep within the nest

**RESTRICTIONS**  
Do not treat mound more than once per season  
For Sod Farms Allow at least 3 days to elapse between last application and harvesting

**OUTDOOR AND PERIMETER SPRAY EXCLUDING RESIDENTIAL TURF**

LOCATION	INSECTS	AMOUNT ACEPHATE 97% PRILLS PER GALLON	APPLICATION TIMING	APPLICATION INSTRUCTIONS
Outdoor and Perimeter area (excluding residential turf)	Wasps	1.2 oz	Treat early or late in the day as wasps are generally less active during these times	Apply the specified amount of ACEPHATE 97% PRILLS per each gallon of water used  Apply as a spot treatment to the nest nest entrance and surrounding areas where the wasps alight
	Ants (excluding fire harvester carpenter and pharaoh ants) Crickets Earwigs Pillbugs	1.2 oz	As the insects appear	Apply the specified amount of ACEPHATE 97% Prills per each gallon of water used Apply to a band of soil 6 to 10 feet adjacent to the structure and to a height of 2 to 3 feet on the foundation where pests may be active or may find entrance Also apply as a residual spray or with a paint brush to surfaces of buildings window frames shutters entry ways screens eaves patios garages carports around garbage areas where these pests congregate

**RESTRICTIONS**  
The use of low pressure handwand equipment for perimeter or wasp treatments is prohibited



40/41

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Keep pesticide in original container. Do not put concentrate or dilute product into food or drink containers. Store in cool, dry place. Protect from excessive heat. Do not contaminate food or foodstuffs. Do not store near feed or food.

**PESTICIDE DISPOSAL:** To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

**CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**IMPORTANT INFORMATION  
READ BEFORE USING PRODUCT**

**CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product reflect the opinion of experts based on field use and tests, and must be followed carefully. It is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Direct Ag Source, LLC. or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of Direct Ag Source, LLC and Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Direct Ag Source, LLC and Seller harmless for any claims relating to such factors.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, DIRECT AG SOURCE, LLC AND SELLER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ON THIS LABEL.**

To the extent consistent with applicable law, Direct Ag Source, LLC or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product and **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF DIRECT AG SOURCE, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF DIRECT AG SOURCE, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Direct Ag Source, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of Direct Ag Source, LLC



41/41

*Notification to add advisory statements, correct use rates and typos – 08 16 2012*

- AMMO® -- Reg. TM of FMC Corporation for cypermethrin insecticide.
- ASANA® -- Reg. TM of E.I. duPont de Nemours & Co., Inc. for esfenvalerate insecticide.
- BAYTHROID® -- Reg. TM of Bayer Crop Protection for cyfluthrin synthetic pyrethroid.
- CAPTURE® -- Reg. TM of FMC Agri Chemical Group Corporation for bifenthrin insecticide-miticide.
- DANITOL® -- Reg. TM of Valent USA Corporation for fenpropathrin insecticide-miticide.
- KARATE® -- Reg. TM of Syngenta Crop Protection, Inc. for lambda-cyhalothrin insecticide.
- LORSBAN® -- Reg. TM of Dow Agro Sciences for chlorpyrifos insecticide.
- SCOUT® -- Reg. TM of Aventis Crop Science for tralomethrin pyrethroid insecticide.
- TAME® -- Reg. TM of Valent USA Corporation for fenpropathrin insecticide.