
 <p style="text-align: center;"><b>U.S. ENVIRONMENTAL PROTECTION AGENCY</b> Office of Pesticide Programs Registration Division (7505P) Ariel Rios Building 1200 Pennsylvania Ave., NW Washington, D.C. 20460</p> <p style="text-align: center;">NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration (under FIFRA, as amended)</p>	EPA Reg. Number: 83278-5	Date of Issuance: NOV 30 2006
	Term of Issuance: conditional	
	Name of Pesticide Product: PRE-AMINE 0.58% Plus	
Name and Address of Registrant (include ZIP Code): Envanage, LLC 115 Obtuse Hill Road Brookfield, CT 06804		
<p><b>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</b></p> <p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:</p> <ol style="list-style-type: none"> <li>1. Add the phrase "EPA Registration No. 83278-5" to the label before you release the product for shipment.</li> </ol> <p>COMMENTS CONTINUED ON PAGES 2 and 3 OF THIS NOTICE OF REGISTRATION.</p> <p>If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product under the enclosed stamped copy of the label constitutes acceptance of these conditions.</p> <p>Enclosures</p>		
Signature of Approving Official:  Joanne I. Miller Product Manager 23 Herbicide Branch Registration Division (7505P)	 Date: NOV 30 2006	

page 2

EPA Reg. No. 83278-5

Comments Continued:

2. Delete the descriptor "Selective", or name the plants that prodiamine controls and the plants that are not controlled (grass species that make-up the turf). "Selective" occurs on page 1 and 3.
3. Replace the chemical expression of prodiamine with the Chemical Abstracts Index name in compliance with PR Notice 97-5.
4. Add an EPA establishment number and the "Net Content" of package to the label.
5. Revise the "Precautionary Statements" by adding as the first statement: "Harmful if absorbed through skin.", followed by "Harmful if inhaled", *followed by* "Causes moderate eye irritation (wear protective eyewear)"; *followed by*:  
Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals"
6. Add a "Note to Physician" to the First Aid Statements. This note should give -technical information on symptomatology, -use of supportive treatments to maintain life functions, -medicine that will counteract the specific physiological effects of the pesticide; -company telephone number to specific medical personnel who can provide specialized medical advice; and the statement: "Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-xxx-xxxx for emergency medical treatment information.
7. On page 3, revise the sentence that reads: "It controls susceptible weeds by inhibiting weed seeds, germination and not root development." to read: It controls susceptible weeds by inhibiting germination, not root development".
8. On page 3, add the map indicating: "This map shows approximate crabgrass germination dates."
9. On page 4, correct the "maximum annual rate" chart by either deleting all reference to Lbs/acre per year or adding Worker Protection Language with reentry information. If you wish add the Worker Protection Language to this label, you must Correct the two rates so they are the same based on a 43, 560 square foot acre.
10. On page 5, Under "Landscape Ornamentals" add "listed on this label. " to the statement, i.e. "This product may.....ornamentals **listed on this label**".

Comments Continued

11. On page 6, under "Tolerant Ornamental Species" revise the 1st sentence to read: "PRE-AMINE 0.58% Plus will not harm trees, shrubs, vines and flowers listed on this label".
  12. On page 1, add the word "many" after the claim "Stops **Many** Grasses....."
  13. Revise the "Conditions of Sale and Warranty" by adding the statement: "To the extent consistent with applicable law" at the beginning of the disclaimer of liability section.
  - 14.. Submit one (1) copy of the final printed labeling before you release this product for shipment.
  - 15.. Submit and/or cite all data required for the registration of this product when the Agency requires all registrants of similar products to submit data; and submit acceptable responses required for reregistration of this product under FIFRA, section 4.
-

4/13

# PRE-AMINE 0.58% PLUS

For Selective Pre-Emergence Control of Grass and Broadleaf Weeds in Lawns and Ornamental Turf (excluding golf course putting greens), and Landscape Ornamentals (including Established Perennials and Wildflower Plantings)

*Stops Weed Grasses Before They Grow  
Prevents Crabgrass, Annual Bluegrass, Foxtail & More*

**ACTIVE INGREDIENT:**

Prodiamine: [N<sup>3</sup>,N<sup>3</sup>-Di-n-propyl-2,4-dinitro-6-(trifluoromethyl)-m-phenylenediamine] .....0.58%

**INERT INGREDIENTS:** ..... 99.42%

TOTAL: ..... 100.00%

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See back panel for additional precautionary statements

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated:**

NOV 30 2006  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

EPA REG NO. 83278-XXX  
EPA EST. NO.  
NET CONTENT: \_\_\_\_\_

83278-5

Manufactured by:  
Envanage, LLC  
Brookfield, CT 06804

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or repeated skin contact with product may cause allergic reactions in some individuals.

<b>FIRST AID</b>	
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>● Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>● Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>● Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>● Take off contaminated clothing.</li> <li>● Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>● Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>● Call a poison control center or physician immediately for treatment advice.</li> <li>● Have person sip a glass of water if able to swallow.</li> <li>● Do not induce vomiting unless told to do so by the poison control center or physician.</li> <li>● Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"> <li>● Move person to fresh air.</li> <li>● If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>● Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>HOT LINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	

**ENVIRONMENTAL HAZARDS**

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

If material is spilled: Contain and sweep up material of spill for disposal or use per label instructions. Wash cleanup equipment to prevent contamination elsewhere. Excess quantities above normal application rates could result in damage to plants subject to spill. Water-in remainder, disc under, cover with layer of soil, or leave exposed to sunlight as deemed appropriate to the situation.

### DIRECTION FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

PRE-AMINE 0.58% Plus is a selective pre-emergence herbicide that provides residual control of many grass and broadleaf weeds in established lawns and ornamental turf (excluding golf course putting greens), landscape ornamentals and established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seeds, germination and root development.

### WHEN TO APPLY TO ESTABLISHED TURF

This product must be applied prior to germination of weed seeds, as it will not control established weeds.

### APPLICATION DIRECTIONS

Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days application. See map below for approximate crabgrass germination dates.

### [MAP OF USA]

### THIS MAP SHOWS APPROXIMATE CRABGRASS GERMINATION DATES

### USE PRECAUTIONS

The following precautions apply to the use of this product in turf grasses and lawns.

- Application of this product may thin emerged annual bluegrass and newly overseeded grasses.
- Do not apply newly set sod until the following year.
- Application to turf stressed by drought, low fertility, or pest damage may result in turf injury.
- Disturbing the herbicide barrier with cultural practices (such as core aeration) may result in reduced weed control.
- Do not apply to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.
- Do not apply through any type of irrigation system.
- Do not apply aerially.
- Do not graze or feed livestock forage cut from treated areas.

### RATES OF APPLICATION

PRE-AMINE 0.58% Plus may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rates listed for longer periods of weed control.

PRE-AMINE 0.58% Plus is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not apply more than 259 lbs./A of this product per calendar year.

MAXIMUM APPLICATION RATES	
Turf Species	Maximum Annual Rate <sup>1</sup>
Bahiagrass Bermudagrass <sup>2</sup> Centipedegrass Seashore Paspalum St. Augustinegrass Tall Fescue (including turf-type) Zoysiagrass	6 lbs/1,000 sq. ft. per year (259 lbs/acre per year)
Buffalograss Kentucky Bluegrass Perennial Ryegrass	4 lbs/1,000 sq. ft. per year (173 lbs/acre per year)
Creeping Red Fescue	2.9 lbs/1,000 sq. ft. per year (130 lbs/acre per year)
Creeping Bentgrass	2.5 lbs/1,000 sq. ft. per year (112 lbs/acre per year)

<sup>1</sup>These are the maximum rates per calendar year.

<sup>2</sup>May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 2 pounds/1,000 square feet (87.5 pounds/acre). Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced rates used in sprigging situations.

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rates listed for longer periods of weed control.

**WEEDS CONTROLLED**

When used according to label instructions, PRE-AMINE 0.58% Plus will control the following weeds:

Barnyardgrass	Kochia
Bluegrass, Annual (Poa annua)	Lambsquarter, Common
Carpetweed	Lovegrass
Chickweed, Common	Panicum (Texas, Fall, Browntop)
Chickweed, Mouseear (from seed)	Pigweed
Crabgrass (Large, Smooth)	Puslane, Common
Crowfootgrass	Pusley, Florida
Cupgrass, Woolly	Rescuegrass <sup>3</sup>
Foxtails, Annual	Shepard's Purse <sup>2</sup>
Goosegrass <sup>1</sup>	Signalgrass, Broadleaf
Henbit	Speedwell, Persian
Itchgrass	Sprangletop
Johnsongrass (from seed)	Spurge, Prostrate
Junglerice	Witchgrass
Knotweed	Woodsorrel, Yellow (from seed)

<sup>1</sup>In many areas a single application of 2.5 to 6 pounds/1,000 square feet (112 to 259 pounds/acre) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 2.5 to 3.9 pounds/1,000 square feet (112 to 173 pounds/acre), followed after 60 to 90 days by a second application at doses that would not exceed the maximum annual rate (see "Maximum Application Rate" table in the Rates of Application section of this label).

<sup>2</sup>For this weed, apply in late summer, fall or winter prior to germination.

<sup>3</sup>Suppression only.

**WHEN TO APPLY AFTER OVERSEEDING TURF**

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil (not thatch plus soil).

**WHEN TO OVERSEED AFTER APPLICATION**

PRE-AMINE 0.58% Plus will inhibit germination of turf seed if overseeded too soon after application. Follow rates and intervals in table for best overseeding/reseeding results.

USE RATE (lbs./A)	MONTHS BEFORE OVERSEEDING		
	North	Transition	South
1.9 pounds/1,000 sq. ft. (86 pounds/acre)	4	4	4
2.5 pounds/1,000 sq. ft. (112 pounds/acre)	5	4	4
1.9 pounds/1,000 sq. ft. (86 pounds/acre)	6	5	5
3.16 pounds/1,000 sq. ft. (138 pounds/acre)	-	6	6
4 pounds/1,000 sq. ft. (173 pounds/acre)	-	7	7
4.5 pounds/1,000 sq. ft. (197 pounds/acre)	-	-	9
5.14 pounds/1,000 sq. ft. (224 pounds/acre)	-	-	10
6.0 pounds/1,000 sq. ft. (259 pounds/acre)	-	-	12

RECOMMENDED SPREADER SETTINGS						
These suggested spreader settings are not intended to replace calibration.						
Please calibrate your spreader before applying product.						
Light Rate (2.5 lbs./1,000 sq. ft.)			X lbs. treats X sq. ft. at the 112 lbs. product/A			
Low Rate (2.9 lbs./1,000 sq. ft.)			X lbs. treats X sq. ft. at the 130 lbs. product/A			
Medium Rate (4.0 lbs./1,000 sq. ft.)			X lbs. treats X sq. ft. at the 173 lbs. product/A			
High Rate (6.0 lbs./1,000 sq. ft.)			X lbs. treats X sq. ft. at the 259 lbs. product/A			
Spreader	Ground or PTO Speed	Width of Coverage	Spreader Settings			
			Light Rate	Low Rate	Medium Rate	High Rate
aa	X mph	XX ft.	X	X	X	X
bb	X mph	XX ft.	X	X	X	X
cc	X mph	XX ft.	X	X	X	X

**LANDSCAPE ORNAMENTALS (including established perennials and wildflower plantings):** This product may be applied for residual pre-emergence weed control in ornamentals.

**RATES OF APPLICATION**

Apply at the rate of 2.5 to 6.0 pounds/1,000 square feet (112 to 259 pounds/acre) in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 56.0 pounds/1,000 square feet (259 pounds/acre) per year.

**APPLICATION TIMING AND INFORMATION**

PRE-AMINE 0.58% Plus is a pre-emergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application.



Best weed control is obtained when applied to soil free of clods, weeds and debris such as leaves. Prior to application, control existing vegetation by hand weeding, cultivation or the use of an appropriate post-emergence herbicide.

#### **ESTABLISHED LANDSCAPE ORNAMENTALS**

PRE-AMINE 0.58% Plus may be applied at any time to established ornamentals as a broadcast, over the top or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate the product in the soil.

#### **NEWLY TRANSPLANTED LANDSCAPE ORNAMENTALS**

PRE-AMINE 0.58% Plus may be applied to newly transplanted ornamentals as a broadcast, over the top or directed application. Delay application to allow soil to settle and water thoroughly before applying. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union.

#### **TOLERANT ORNAMENTAL SPECIES**

PRE-AMINE 0.58% Plus will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.

#### **SCIENTIFIC NAME**

Abies spp.  
Abelia grandiflora  
Acer palmatum  
A. Platanoides  
Achillea spp.  
Actindia chinensis  
Agapanthus africanus  
Agapanthus orientalis  
Akebiaq quintata  
Allium cemuum  
Anemone hybrida  
Aquilegia spp.  
Arctostaphylos densiflora  
Arctotheca calendula  
Artemesia spp.  
Aster spp.  
Aster X Frikartii  
Athrium Filiz-femina  
Aucuba japonica  
Begonia spp.  
Berberis gladwynensis  
B. Julianae  
B. mentorensis  
B. Thunbergil  
B. Verruculosa  
Bergenia cordifolia  
Boltonia asteroides  
Buddleia davidii  
Buxus microphylla  
Callistemon viminalis  
Calluna vulgaris

#### **COMMON NAME**

Fir species\*\* (Balsam, Frasier, Noble, etc.)  
Abelia: Sherwood  
Japanese Maple  
Norway Maple  
Yarrow: King Edward  
Kiwi\*  
Lily-of-the-Nile (African Lily)  
  
Five-leaf or Chocolate Vine  
Lady's Leek, Nodding Onion  
Japanese Anemone  
Aquilegia: Red and Gold  
Vine Hill Manzanita  
Cape Weed  
Wormwood; Silver Mound, Castle  
Aster: Bonnie Blue, Purple Dome  
  
Lady Fern; Fern Lady  
Japanese Aucuba  
Fibrous Begonia: Hardy Grandis  
Barberry  
Wintergreen Barberry  
Mentor Barberry  
Japanese Barberry  
Warty Barberry  
  
Snowbank  
Butterfly-Bush (Dwarf Blue); Royal Red  
Japanese Boxwood  
Weeping Bottlebrush  
Scotch Heather

Campanula carpatica  
 Campis X Tagliabuana  
 Carpobrotus edulis  
 Cassia artemisoides  
 Ceanothus rigidus  
 Ceratostigma plumbagonoides  
 Chamaecyparis pisifera  
 Chrysanthemum nipponicum  
 Cleyera japonica  
 Citrus spp.  
 Coreopsis spp.  
 Cornus florida  
 C. Stolonifera  
 Cortaderia seloana  
 Cotoneaster apiculatus  
 C. buxifolius  
 C. dammeri  
 C. Microphyllus  
 Cretaegus spp.  
 Crocosmia spp.  
 Cupressus sempervirens  
 Delosperma alba  
 Delosperma spp.  
 Delphinium spp.  
 Dianthus deltoides  
 D. gratianopolitanus  
 Dodonea viscosa  
 Echinacea purpurea  
 Elaeagnus pungens  
 Euonymus fortunei  
 E. japonica  
 E. kiautschovica  
 Fatsia japonica  
 Forsythia intermedia  
 Forsythia suspensa  
 F. viridissima  
 Gaillardia spp.  
 Gardenia jasminoides  
 Gaura spp.  
 Gentiana dahurica  
 Geranium cinereum  
 Gladiolus spp.  
 Gypsophilia repens  
 Hedera helix  
 Hellanthemum spp.  
 Hemerocallis spp.  
 Hibiscus  
 Hibiscus spp.  
 Hibiscus rosa-sinensis  
 Hosta plantaginea  
 H. Sieboldiana  
 Houttuynia cordata var. variegata  
 Hydrangea macrophylla  
 Ilex Bernyi  
 Ilex cornuta  
 I. crenata  
 Tussock Bellflower, (White Clips)  
 Trumpet Flower, Madame Galen  
 Hottentot Fig (Ice Plant)  
 Feathery Cassia  
 Wild Lilac  
  
 False Cypress  
  
 Cleyera  
 Citrus species\*  
 Coreopsis (Calliopsis) Early Sunrise, Moonbeam  
 Flowering Dogwood  
 American Dogwood  
 Pampas Grass  
 Cranberry Cotoneaster  
 Cotoneaster  
 Bearberry Cotoneaster  
 Rockspray Cotoneaster  
 Hawthorne  
 Lucifer  
 Italian Cypress  
 White Trailing Ice Plant  
 Cooperi Pink  
 Larkspur: Blue Elf  
 Dianthus: Maiden Pinks "Zing"  
 Cheddar Pink  
 Hop Bush  
 Coneflower, Purple; Magnus  
 Silverberry  
 Wintercreeper  
 Evergreen Euonymus  
 Spreading Euonymus  
 Japanese Aralia  
 Border Forsythia  
 Weeping Forsythia  
 Greenstem Forsythia  
 Gaillardia, Blanket Flower: "Goblin"  
 Gardenia, Cape-Jasmine  
  
 Gentian  
 Cranesbill  
 Gladiolus species\*\*  
 Baby's Breath  
 English Ivy  
 Sunrose  
 Daylily: Aztec Gold, Stella de Oro, Tender Love  
 Rose of Sharon\*\*  
 Mallow: Disco Belle White  
 Chinese Hibiscus  
 Hosta, Plantain Lily (Fragrant)  
 Hosta, "Seersucker"  
  
 Bigleaf Hydrangea  
 Holly  
 Chinese Holly  
 Japanese Holly

<i>I. opaca</i>	American Holly
<i>I. vomitoria</i>	Yaupon Holly, Schillings
<i>Inula ensifolia</i>	
<i>Iris ensata</i>	Sword-Leaved Iris; Jodlesong
<i>Iris siberica</i>	Siberian Iris; Cabernet
<i>Iris spp.</i>	Iris species**
<i>Jasminium nudiflorum</i>	Winter Jasmine
<i>Juniperus chinensis</i>	Chinese Juniper; Nick's Compact, Parsonnii
<i>J. conferta</i>	Shore Juniper; Blue Pacific
<i>J. davurica</i>	Parsoni
<i>J. horizontalis</i>	Creeping Juniper
<i>Juglans spp.</i>	Walnut*
<i>Justicia brandegeana</i>	Shrimp Plant
<i>Lagerstromia indica</i>	Crape Myrtle
<i>Lagerstromia indica X Fauriei</i>	Crape Myrtle; Tuscarora
<i>Lantana montevidensis</i>	Weeping Lantana
Lavender spp.	Lavender; Munstead
<i>Leontopodium alpinum</i>	Edelweiss
<i>Ligustrum amurense</i>	Amur Privet
<i>Ligustrum sinense</i>	Chinese Privet; Variegata
<i>L. japonicum</i>	Japanese Privet
<i>L. lucidum</i>	Glossy Privet (Waxleaf)
<i>Lilium spp.</i>	Lily; Jazz
<i>Liriope muscari</i>	Big Blue Lily, Silver Mound, Evergreen Giant
<i>Liriope muscari var. variegata</i>	Liriope, Variegated
<i>Liriope spicata</i>	Liriope, Creeping
<i>Lobelia Cardinalis</i>	Cardinal Flower, Indian Pink
<i>Lonicera japonica</i>	Japanese Honeysuckle
<i>L. tatarica</i>	Tatarian Honeysuckle
<i>Lorapetalum chinense</i>	Burgundy
<i>Lythrum spp.</i>	Loosestrife; Modern Pink
<i>Magnolia spp.</i>	Magnolia species**
<i>Maleophora luteola</i>	Ice Plant
<i>Malus spp.</i>	Crabapple*
<i>Miscanthus sinensis</i>	Yaku Jima, Silberfeder
<i>Nandina domestica</i>	Heavenly Bamboo
<i>Narcissus spp.</i>	Narcissus species**
<i>Nerium oleander</i>	Oleander
<i>Olea europaea</i>	Olive*
<i>Ophiopogon japonicus</i>	Mondo Grass
<i>Osmanthus heterophyllus</i>	Osmanthus (False Holly); Gulf Tide
<i>Osteospermum fruticosum</i>	Trailing African Daisy
<i>Oxydendrum arboreum</i>	Sourwood
<i>Paeonia suffruticosa</i>	Tree Peony
<i>Pennisetum setaceum</i>	Fountain Grass (Dwarf)
<i>Perovskia atriplicifolia</i>	
<i>Persea americana</i>	Avocado*
<i>Photinia Fraseri</i>	Fraser's Photinia (Redtip)
<i>Physotegia Virginiana</i>	Dragonhead, False; Vivid
<i>Picea spp.</i>	Spruce species** (Colorado Blue, Norway, etc.)
<i>Pieris japonica</i>	Lily-of-the-Valley Bush
<i>Pinus brutia</i>	Calabrian Pine
<i>P. canariensis</i>	Canary Island Pine
<i>P. Elliottii</i>	Slash Pine
<i>P. halepensis</i>	Aleppo Pine
<i>P. nigra</i>	Austrian Pine

<i>P. palustris</i>	Longleaf Pine
<i>P. radiata</i>	Monterey Pine
<i>P. strobus</i>	Eastern White Pine
<i>P. sylvestris</i>	Scotch Pine
<i>P. taeda</i>	Loblolly Pine
<i>P. Thunbergiana</i>	Japanese Black Pine
<i>P. Virginiana</i>	Virginia Pine
Pistachio spp.	Pistachio*
<i>Pittosporum rhombifolium</i>	Queensland Pittosporum
<i>P. tobira</i>	Japanese Pittosporum
<i>Podocarpus macrophyllus</i>	Japanese Yew
<i>Prunus laurocerasus</i>	English Laurel
<i>Prunus</i> spp.	Almond, Apricot, Nectarine, Peach, Plum and Prune*
<i>Pseudotsuga menziesii</i>	Douglas Fir**
<i>Pyracantha coccinea</i>	Firethorn (Scarlet)
<i>P. fortuneana</i>	Firethorn
<i>P. koidzumii</i>	Firethorn
<i>Pyrus</i> spp.	Bradford Pear spp.
<i>Quercus Shumardii</i>	Oak, Shumard's Red
<i>Quercus</i> spp.	Oak species
<i>Raphiolepis indica</i>	Indian Hawthorne
<i>Raphiolepis umbellata</i>	Yedda Hawthorne
<i>Rhododendron</i> (including Azalea)	'Coral Bells,' 'Formosa,' 'Hino-crimson,' 'PJM' 'Delaware Valley White,' 'Roseum Elegans,' 'Flame Creeper,' 'Girard Crimson,' 'Wakeiebisu' 'White Gumbo,' 'George L. Tabor'
<i>Rosa banksiae</i>	Lady Bank's Rose
<i>Rosmarinus officinalis</i>	Rosemary*
<i>Rudbeckia</i> spp.	Black-Eyed Susan; Goldsturm
<i>Santolina virens</i>	Santolina
<i>Saxifraga</i> spp.	Saxifrage; Purple Dome
<i>Scabiosa</i> spp.	Pincushion Flower
<i>Sedum album</i>	Stonecrop
<i>Sedum cauticola</i>	Stonecrop; Lidakense
<i>Sedum dasyphyllum</i>	Stonecrop
<i>Sedum spurium</i>	Stonecrop; Dragon's Blood
<i>Spiraea bumalda</i>	Spirea; Anthony Waterer
<i>Syzygium paniculatum</i>	Australian Brushcherry
<i>Taxus cuspidata</i>	Japanese Yew
<i>T. media</i>	Yew
<i>Teucrium</i> spp.	Germander
<i>Thalictrum diptercarpum</i>	Meadow Rue
<i>Thuja occidentalis</i>	American Arborvitae
<i>Trachelospermum asiaticum</i>	Star Jasmine
<i>Tsuga canadensis</i>	Canada Hemlock
<i>Tulipa</i> spp.	Tulip species
<i>Veronica</i> spp.	Veronica, Speedwell; Sunny Border
<i>Viburnum japonicum</i>	Japanese Viburnum
<i>V. odoratissimum</i>	Sweet Viburnum
<i>V. plicatum</i>	Japanese Snowball
<i>V. rigidum</i>	Canary Island Viburnum
<i>V. suspensum</i>	Arrowood Viburnum
<i>V. tinus</i>	Laurustinus
<i>V. trilobium</i>	Cranberry Bush
<i>V. wrightii</i>	Leatherleaf Viburnum

Vinca major  
 Vinca Minor  
 Vitis spp.  
 Weigela florida  
 Yucca aloifolia  
 Y. filametosa

Greater Periwinkle, Vinca  
 Common Periwinkle, Vinca  
 Grape\*  
 Old Fashioned Weigela  
 Spanish Bayonet, Yucca  
 Adam's Needle, Yucca

\*Do not use on food producing trees, vines or plants.

\*\*Not for use on container grown plants.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in original container in a locked storage area out of reach of children and domestic animals.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, of, if allowed by State and local authorities, by burning. If burned, stay out of smoke

#### CONDITIONS OF SALE AND WARRANTY

The DIRECTIONS FOR USE of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop, injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of Envamage, LLC or the Seller. All such risks shall be assumed by the Buyer.

Envamage, LLC warrants that this product conforms to the chemical description on the label, and is reasonably fit for the purposes set forth in the DIRECTIONS FOR USE, when it is used in accordance with such directions, subject to the inherent risks mentioned above.

ENVAMAGE, LLC NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESSED OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. THIS WARRANTY EXTENDS TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNING, OR CAUTIONS. BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ENVAMAGE, LLC and the seller offer this product, and the Buyer and User accept it, subject to the foregoing CONDITIONS OF SALE AND WARRANTY.