

8 3529-11

7/10/2012

1/30



**UNITED STATES ENVIRONMENTAL PROTECTION AGENCY**  
WASHINGTON, DC 20460

OFFICE OF  
CHEMICAL SAFETY AND  
POLLUTION PREVENTION

**JUL 10 2012**

Sharda USA LLC  
% Lori Kohler  
Wagner Regulatory Associates Inc  
P O Box 640  
Hockessin DE 19707

Product Name	Shar Teb 3 6FL Fungicide
EPA Reg No	83529 11
Subject	Your notification dated April 10 2012 - revised label
OPP Decision Number	464246

Dear Ms Kohler

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98 10. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98 10 and finds that the action(s) requested fall within the scope of PRN 98 10.

The label submitted with the application has been stamped Notification and will be placed in our records.

If you have any questions please contact Robert Westin by phone at (703) 305 5721 or via email at [westin.robert@epa.gov](mailto:westin.robert@epa.gov)

Sincerely

A handwritten signature in cursive script, reading "Mary L Waller", is positioned above the typed name.

Mary L Waller  
Product Manager (21)  
Fungicide Branch  
Registration Division (7504P)



United States  
**Environmental Protection Agency**  
 Washington DC 20460

Registration

Amendment

Other

OPP Identifier Number

464246

**Application for Pesticide Section I**

1 Company/Product Number 83529 11	2 EPA Product Manager Mary Waller	3 Proposed Classification <input checked="" type="checkbox"/> None Restricted
4 Company/Product (Name) Sharda USA LLC/ Shar Teb 3 6FL Fungicide	PM# 21	
5 Name and Address of Applicant (Include Zip Code) Sharda USA LLC c/o Wagner Regulatory Associates Inc P O Box 640 Hockessin DE 19707 <input type="checkbox"/> Check if this is a new address	6 Expedited Review In accordance with FIFRA Section 3(c)(3) (b)(I) my product is similar or identical in composition and labeling to EPA Reg No Product Name	

**Section II**

<input type="checkbox"/> Amendment Explain below	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> Me Too Application
<input checked="" type="checkbox"/> Notification Explain below	<input type="checkbox"/> Other Explain below

**Explanation** Use additional page(s) if necessary (For Section I and Section II)  
 Notification to add State specific language per PR Notice 98 10 This notification is consistent with the provisions of PR Notice 98 10 and EPA regulations at 40 CFR 152 46 and no other changes have been made to the labeling or the confidential statement of formula of this product I understand that it is a violation of 18 U S C Sec 1001 to willfully make any false statement to EPA I further understand that if this notification is not consistent with the terms of PR Notice 98 10 and 40 CFR 152 46 this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA


**Section - III**

<b>1 Material This Product Will Be Packaged In</b>			
Child Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No If Yes Unit Packaging wgt No per container	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No If Yes Package wgt No per container	2 Type of Container <input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) HDPE lined bags
<b>Certification must be submitted</b>			
3 Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container	4 Size(s) Retail Container 2 5 gallons	5 Location of Label Direction <input type="checkbox"/> On Label <input checked="" type="checkbox"/> On Labeling accompanying product	
6 Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other _____	

**Section IV**

<b>1 Contact Point (Complete items directly below for identification of individual to be contacted if necessary to process this application)</b>			
Name Lori Kohler	Title Agent for Sharda USA LLC	Telephone No (Include Area Code) (302) 635 7281	
<b>Certification</b> I certify that the statements I have made on this form and all attachments thereto are true accurate and complete I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law			6 Date Application Received  (Stamped)
2 Signature 	3 Title Agent for Sharda USA LLC		
4 Typed Name Lori Kohler	5 Date April 10 2012		

3/30  
4/23



April 10 2012

Document Processing Desk (NOTIF)  
Attn Mary Waller PM 21  
Registration Division (7504P)  
U S Environmental Protection Agency  
Office of Pesticide Programs  
Room S 4900 One Potomac Yard  
2777 South Crystal Drive  
Arlington VA 22202 4501

Dear Ms Waller

Subject Shar Teb 3 6FL Fungicide  
[ABN Tebu Crop 3 6F Tebusha 3 6FL Fungicide Tebu Turf 3 6F]  
EPA Registration Number 83529 11  
**Notification to add State specific language per PR Notice 98 10**


Wagner Regulatory Associates Inc on behalf of Sharda USA LLC submits the enclosed Notification to add State specific language per PR Notice 98 10 The following has been added to the front page of sub label A and B Not registered for sale in the State of New York No other changes were made to this label

In support of this notification the following documents are attached

- Letter from Sharda USA LLC appointing Wagner Regulatory Associates Inc as its agent
- Application for Pesticide Notification (8570 1)
- Draft label 2 copies (1 clean and 1 with changes highlighted) and CD
- Certification with Respect to Label Integrity *u s s c a x*

Please feel free to contact me via email at [lori@wagnerreg.com](mailto:lori@wagnerreg.com) or telephone at (302) 635 7281 if you have any questions

Respectfully submitted

  
Lori Kohler  
Agent for Sharda USA LLC

4/30

**SHAR-TEB 3 6FL FUNGICIDE****[ABN TEBUSHA 3 6FL FUNGICIDE, Tebu-Turf 3 6F, Tebu-Crop 3 6F]****Active Ingredient**

Tebuconazole alpha [2 (4 chlorophenyl)ethyl] alpha(1 1 dimethylethyl) 1 H 1 2 4 triazole 1 ethanol

Other Ingredients

**Total**

Contains 3 6 pounds tebuconazole per gallon

38 7%

61 3%

100 0%

**NOTIFICATION****JUL 10 2012****KEEP OUT OF REACH OF CHILDREN****CAUTION**

Si usted no entiende la etiqueta busque a alguien para que se la explique a usted en detalle (If you do not understand the label find someone to explain it to you in detail )

See additional precautionary statements and directions for use in booklet

**STOP Read the label before use****FIRST AID**

<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice</li> <li>• Have person sip a glass of water if able to swallow</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor</li> <li>• Do not give anything by mouth to an unconscious person</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing</li> <li>• Rinse skin immediately with plenty of water for 15 20 minutes</li> <li>• Call a poison control center or doctor for treatment advice</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 20 minutes</li> <li>• Remove contact lenses if present after the first 5 minutes then continue rinsing eye</li> <li>• Call a poison control center or doctor for treatment advice</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air</li> <li>• If person is not breathing call 911 or an ambulance then give artificial respiration preferably by mouth to mouth if possible</li> <li>• Call a poison control center or doctor for further treatment advice</li> </ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment	
<b>NOTE TO PHYSICIAN</b> No specific antidote Treat symptomatically	
<b>Symptoms of Poisoning</b> The compound does not cause any definite symptoms that would be diagnostic	
Contact with the eyes may cause irritation	
<b>EMERGENCY NUMBERS</b>	
For 24 hour medical emergency assistance (human or animal) call 1 800 222 1222 For chemical emergency assistance (spill leak fire or accident) call CHEMTREC at 1 800 424 9300	

**Not registered for sale in the state of New York****EPA Reg No** 83529 11**EPA Est No****Net Contents** 2 5 Gallons**Manufactured for**

Sharda USA LLC

PO Box 640

Hockessin DE 19707

5/30

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Causes moderate eye irritation Harmful if swallowed inhaled or absorbed through skin Avoid contact with skin eyes and clothing Avoid breathing vapor or spray mist

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical resistant to this product are listed below If you want more options follow the instructions for category C on an EPA chemical resistance category selection chart

**Applicators and other handlers must wear**

- Long sleeved shirt and long pants
- Chemical resistant gloves such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton and
- Shoes plus socks

**User Safety Requirements**

Follow manufacturer s instructions for cleaning/maintaining PPE If no such instructions for washables exist use detergent and hot water Keep and wash PPE separately from other laundry

**ENGINEERING CONTROLS STATEMENTS**

When handlers use closed systems enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170 240(d)(4 6)] the handler PPE requirements may be reduced or modified as specified in the WPS

**USER SAFETY RECOMMENDATIONS**

**Users should**

- Wash hands before eating drinking chewing gum using tobacco or using the toilet
- Remove clothing immediately if pesticide gets inside Then wash thoroughly and put on clean clothing
- Remove PPE immediately after handling this product Wash the outside of gloves before removing As soon as possible wash thoroughly and change into clean clothing

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to mammals fish and aquatic invertebrates Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark Runoff may be hazardous to aquatic organisms in neighboring areas Do not contaminate water when disposing of equipment washwater or rinsate

**Groundwater Advisory** Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use Use of this chemical in areas where soils are permeable particularly where the water table is shallow may result in groundwater contamination

**Surface Water Advisory** This product may contaminate water through drift of spray in wind This product has a high potential for runoff for several months or more after application Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product A level well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds streams, and springs will reduce the potential for contamination of water from rainfall runoff Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling Do not apply this product in a way that will contact workers or other persons either directly or through drift Only protected handlers may be in the area during application For any requirements specific to your State or Tribe consult the agency responsible for pesticide regulation

6/30

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) specified in the use directions for each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

**INFORMATION FOR AGRICULTURAL USE****SHAKE WELL BEFORE USING**

**SPRAY DRIFT MANAGEMENT** Do not allow this product to drift.

**Foliar Spray Drift Management** Avoiding spray drift from foliar applications is the responsibility of the applicator. Similar to aerial spray drift, the interaction of many equipment- and weather-related factors determine the potential for spray drift from foliar applications. To protect water resources, the applicator and the grower are responsible for considering all these factors when making decisions.

**Aerial Spray Drift Management** Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed  $\frac{3}{4}$  the length of the wingspan or rotor.
  2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.
- Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the [Aerial Drift\\* Reduction Advisory Information](#).

**AERIAL DRIFT REDUCTION ADVISORY**

This section is advisory in nature and does not supersede the mandatory label requirements.

**INFORMATION ON DROPLET SIZE** The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature, and Humidity, and Temperature Inversions).

**CONTROLLING DROPLET SIZE**

- **Volume** Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of nozzles** Use minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.

**BOOM LENGTH** For some use patterns, reducing the effective boom length to less than  $\frac{3}{4}$  of the wingspan or rotor length may further reduce drift without reducing swath width.

**APPLICATION HEIGHT** Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

**SWATH ADJUSTMENT** When applications are made with a crosswind, the swath will be displaced downward. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.).

**WIND** Drift potential is lowest between wind speeds of 2–10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential.

**NOTE** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

**TEMPERATURE AND HUMIDITY** When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

**TEMPERATURE INVERSIONS** Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

**SENSITIVE AREAS** The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

**Chemigation** Do not apply this product through any type of irrigation system.

**Spray Volume** Apply Tebu Crop 3 6F with ground or aerial equipment using sufficient volume of spray to provide thorough coverage. Apply Tebu Crop 3 6F in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment. Check equipment calibration frequently. Continuous agitation is required to keep the material in suspension.

Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied. If necessary, increase the spray volume per acre for complete crop coverage. Use the higher rate under conditions of severe disease pressure. Also, see local State Extension Service recommendations for application schedules.

**Mixing** Add specified amount of Tebu Crop 3 6F into the spray tank while filling with water to the desired level. Operate the agitator while mixing. If other materials are added to the spray tank, the Tebu Crop 3 6F should be thoroughly dispersed prior to the addition of other materials. Do not tank mix with products containing a prohibition against tank mixing. Follow the most restrictive labeling requirements of any tank mix product.

**Compatibility** To determine the compatibility of Tebu Crop 3 6F with other products, the following procedure should be followed. Pour the recommended proportions of the products into a suitable container of water, mix thoroughly and allow to stand at least five minutes. If the combination remains mixed or can be re-mixed readily, the mixture is considered physically compatible.

#### **OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS AND ESTUARIES**

Apply only during alternate years in fields adjacent to aquatic areas listed above. Do not apply by ground or air within 100 feet of aquatic areas listed above. Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip. **Spray Drift Management** For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used and must not exceed 75% of the wingspan or rotor diameter. Use the largest droplet size consistent with pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure. Apply in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment. Spray should be released at the lowest possible height consistent with good pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided. Make aerial or ground applications when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

**Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.** Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature. Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

**ROTATIONAL CROPS** Treated areas may be replanted with any crop specified on this label as soon as practical after last application. Any crop not specified on this label may be planted into treated areas 20 days after last application.

**Comments** For optimum disease control, the lowest specified rate of a spray surfactant should be tank mixed with Tebu Crop 3 6F. Tebu Crop 3 6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tebu Crop 3 6F will be resistant to weathering. Tebu Crop 3 6F is a demethylation inhibitor (DMI) fungicide (Group 3).



9/30

CROP	DISEASE	Tebu-Crop 3.6F APPLICATION RATE (fl. oz. per acre)
Asparagus	Rust ( <i>Puccinia</i> spp )	4 – 6

**Notes** Apply Tebu Crop 3 6F as a foliar spray to the developing ferns after harvest of spears is completed. Apply at the earliest sign of rust pustules or when weather conditions are conducive for rust development. Apply 4 to 6 fl oz of Tebu Crop 3 6F (0.11 to 0.17 lb of active ingredient per acre) in alternation with another effective fungicide. Under conditions of severe rust pressure use the higher rate. Repeat applications on a 14 day interval as necessary to maintain control of rust. Do not apply to harvestable spears. Do not make more than three foliar applications per season (18 fl oz/acre or 0.51 lb of active ingredient per acre).

**Comments** Applications may be made using ground or aerial application equipment. A 50 foot spray drift buffer zone is required for all aerial applications. For optimum disease control the lowest labeled rate of a spray surfactant should be tank mixed with Tebu Crop 3 6F. Tebu Crop 3 6F is a sterol demethylation inhibitor (DMI) fungicide (Group 3). Alternating Tebu Crop 3 6F with other DMI fungicides may lead to resistance.

**Restricted entry interval (REI) = 12 hours**

**Pre harvest interval (PHI) = 100 days (California) 180 days (all other states)**

CROP	DISEASE	Tebu-Crop 3.6F APPLICATION RATE (fl. oz. per acre)
Barley	Rusts ( <i>Puccinia</i> spp ) Head blight ( <i>Fusarium</i> spp ) Suppression	4

#### NOTES for Barley

Apply Tebu Crop 3 6F in a minimum of 10 gallons of spray solution per acre by ground or in a minimum of 5 gallons of spray solution per acre by air. A maximum of 4 fl oz of Tebu Crop 3 6F may be applied per acre per crop per season. Do not apply within 30 days of harvest. Straw cut after harvest may be fed or used for bedding. Grazing livestock or feeding of green forage is permitted 6 or more days after the last application of Tebu Crop 3 6F. Barley fields should be observed closely for early disease symptoms particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development.

#### Application timing Directions

**Rusts** Apply Tebu Crop 3 6F at the earliest sign of rust pustules on foliage.

**Fusarium head blight** Optimal timing of Tebu Crop 3 6F for Fusarium head blight suppression is when main stem heads have fully emerged (Feekes 10.5) on 50% of the plants.

#### Comments

For optimum disease control the lowest specified rate of a spray surfactant should be tank mixed with Tebu Crop 3 6F. Tebu Crop 3 6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time Tebu Crop 3 6F will be resistant to weathering. Tebu Crop 3 6F is a demethylation inhibitor (DMI) fungicide (Group 3).

**Restricted entry interval (REI) = 12 hours**

10/30

CROP	DISEASE	Tebu-Crop 3 6F APPLICATION RATE (fl. oz. per acre)
------	---------	--

**Beans**  
(fresh & dry except succulent shelled)

**Rust**  
(*Uromyces appendiculatus*)

4 6

**Notes** Apply Tebu Crop 3 6F in a protective spray schedule or when weather conditions are favorable for rust development Repeat applications at 14 day intervals or as necessary to maintain control Fresh beans Do not apply more than 24 fl oz of Tebu Crop 3 6F per acre per crop season Dry beans Do not apply more than 12 fl oz of Tebu Crop 3 6F per acre per crop season

**Comments** For optimum disease control the lowest labeled rate of a spray surfactant should be tank mixed with Tebu Crop 3 6F Tebu Crop 3 6F must have two to four hours of drying time on bean foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs After this period of time Tebu Crop 3 6F will be resistant to weathering Tebu Crop 3 6F is a demethylation inhibitor (DMI) fungicide (Group 3)

**Restricted entry interval (REI) = 12 hours**

**Pre harvest interval (PHI) = 7 days (fresh beans) 14 days (dry beans)**

CROP	DISEASE	Tebu-Crop 3 6F APPLICATION RATE (fl. oz. per acre)
------	---------	--

**Corn**  
(sweet corn field corn field corn grown for seed and popcorn)

**Rust (*Puccinia spp*)**  
**Northern Leaf Blight**  
(*Helminthosporium turcicum*)  
**Southern Leaf Blight**  
(*Helminthosporium maydis*)  
**Northern Leaf Spot**  
(*Helminthosporium carbonum*)  
**Gray Leaf Spot**  
(*Cercospora zeae maydis*)

4 – 6

**Notes** Apply Tebu Crop 3 6F in a protective spray schedule or when weather conditions are favorable for disease development Repeat applications at 7 to 14 day intervals or as necessary to maintain control A maximum of 24 fl oz (1 5 pint) of Tebu Crop 3 6F may be applied per acre per crop season

**Comments** For optimum disease control the lowest labeled rate of a spray surfactant should be tank mixed with Tebu Crop 3 6F Tebu Crop 3 6F must have two to four hours of drying time on corn foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs After this period of time Tebu Crop 3 6F will be resistant to weathering Tebu Crop 3 6F is a demethylation inhibitor (DMI) fungicide (Group 3)

**Restricted entry interval (REI) for sweet corn = 19 days**

**Pre harvest interval (PHI) for sweet corn = 7 days (ears or forage) 49 days (fodder)**

**Restricted entry interval (REI) for all corn except sweet corn = 12 hours**

**Pre harvest interval (PHI) for field seed or popcorn = 21 days (forage) 36 days (grain or fodder)**

11/30

CROP	DISEASE	Tebu-Crop 3.6F APPLICATION RATE (fl. oz. per acre)
Cotton	Southwestern Cotton Rust ( <i>Puccinia cacabata</i> )	6 – 8

**Notes** Apply Tebu Crop 3.6F in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 7 to 14 day intervals or as necessary to maintain control. Do not apply more than 24 fl. oz. of Tebu Crop 3.6F per acre per crop season.

**Comments** For optimum disease control the lowest labeled rate of a spray surfactant should be tank mixed with Tebu Crop 3.6F. Tebu Crop 3.6F must have two to four hours of drying time on cotton foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time Tebu Crop 3.6F will be resistant to weathering. Tebu Crop 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3).

**Restricted entry interval (REI) = 12 hours**

**Pre harvest interval (PHI) = 30 days**

CROP	DISEASE	Tebu-Crop 3.6F APPLICATION RATE (fl. oz. per acre)
<b>Cucurbit Vegetables Group</b>	<b>Powdery Mildew</b>	
Chayote	( <i>Sphaerotheca fuliginea</i> /	4 6
Chinese Waxgourd	<i>Podosphaera xanthi</i> )	
Citron Melon	( <i>Erysiphe cichoracearum</i> )	
Cucumber		
Gherkin	<b>Gummy Stem Blight – suppression</b>	
Edible Gourd	( <i>Didymella bryonae</i> )	8
(includes Hyotan Cucuzza	(watermelon squash pumpkin and	
Hechima and Chinese Okra)	melons only)	
<i>Momordica spp</i>		
(includes Balsam Apple Balsam	<b>Notes</b> apply specified dosage in a protective spray schedule to	
Pear Bitter Melon and Chinese	foliage and fruit. Repeat applications at 10 to 14 day intervals. Do	
Cucumber)	not apply more than 24 fl. oz. of Tebu Crop 3.6F per acre per crop	
Muskmelon	season	
(includes Cantaloupe Casaba	<b>Comments</b> for optimum disease control the lowest labeled rate of a	
Crenshaw Melon Golden Pershaw	spray surfactant should be tank mixed with Tebu Crop 3.6F. Tebu	
Melon Honeydew Melon Honey	Crop 3.6F must have two to four hours of drying time for the active	
Balls Mango Melon Persian Melon	ingredient to move systematically into plant tissue before rain or	
Pineapple Melon Santa Clause	irrigation occurs. After this period of time Tebu Crop 3.6F will be	
Melon and Snake Melon)	resistant to weathering. Tebu Crop 3.6F is a demethylation inhibitor	
Pumpkin	(DMI) fungicide (Group 3)	
Summer Squash		
(includes Crookneck Squash	<b>Restricted entry interval (REI) = 12 hours</b>	
Scallop Squash Straightneck	<b>Pre harvest interval (PHI) = 7 days</b>	
Squash Vegetable Marrow and		
Zucchini)		
Winter Squash		
(includes Butternut Squash		
Calabaza Hubbard Squash Acorn		
Squash and Spaghetti Squash)		
Watermelon		

CROP	DISEASE	Tebu-Crop 3.6F APPLICATION RATE (fl. oz. per acre)
Dry Bulb Onion	White Rot	White rot 20.5 fl oz per acre
Garlic Great headed (Elephant)	( <i>Sclerotium cepivorum</i> )	applied in a 4 to 6 inch band over/into each furrow
Garlic	Rust ( <i>Puccinia allii</i> <i>Puccinia porri</i> )	4 – 6
Shallot	Purple Blotch ( <i>Alternaria porri</i> )	

**White Rot** For the control of white rot make one application in the furrow at the time of planting. Make the in furrow application at the rate of 20.5 fl oz Tebu Crop 3.6F per acre. Apply the entire per acre rate in a 4 to 6 inch band over/into each furrow. Additional control may be obtained by including two foliar applications at 4 – 6 fl oz/acre.

**Rust** For the control of rust make foliar applications at the rate of 4 – 6 fl oz of Tebu Crop 3.6F per acre per application. Repeat at an interval of 10 – 14 days. Apply Tebu Crop 3.6F in a protective spray schedule or when weather conditions are favorable for rust development.

**Notes** Do not apply more than 32.5 fl oz Tebu Crop 3.6F per acre per season if an in furrow treatment is made. If Tebu Crop 3.6F is not applied as an in furrow treatment then do not apply more than 12 fl oz/acre per season as a foliar spray.

**Comments** For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank mixed with Tebu Crop 3.6F. Tebu Crop 3.6F must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time Tebu Crop 3.6F will be resistant to weathering. Tebu Crop 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3).

**Restricted entry interval (REI) = 12 hours**

**Pre harvest interval (PHI) = 7 days**

CROP	DISEASE	Tebu-Crop 3.6F APPLICATION RATE (fl. oz. per acre)
Garden Beet	Cercospora Leaf Spot	3 – 7.2
roots and tops (leaves)	( <i>Cercospora beticola</i> )	

**Notes** Make applications on 14 day intervals. Do not apply more than 28.8 fl oz of Tebu Crop 3.6F per acre per season.

**Comments** For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest labeled rate of a spray surfactant may be tank mixed with Tebu Crop 3.6F. Tebu Crop 3.6F must have two to four hours of drying time on beet foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time Tebu Crop 3.6F will be resistant to weathering. Tebu Crop 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3).

**Restricted entry interval (REI) = 12 hours**

**Pre harvest interval (PHI) = 7 days**

CROP	DISEASE	Tebu-Crop 3.6F APPLICATION RATE (fl. oz. per acre)
------	---------	---

Grasses Grown for Seed	Rust ( <i>Puccinia spp</i> ) Powdery Mildew	4 – 8
------------------------	--	-------

**Information** Apply the specified rate of Tebu Crop 3 6F in a minimum of 20 gallons of water per acre with ground sprayers or in a minimum of 10 gallons of water per acre with aircraft. Thorough coverage is important for optimum disease control. For optimum benefit the lowest specified rate of a spray surfactant should be tank mixed with Tebu Crop 3 6F. Do not apply more than 16 fl. oz. Tebu Crop 3 6F (45 lb a.i.) per acre per crop season. Chaff screenings and straw from treated areas may be used for feed purposes. Do not forage cut green crop or use seed for feed purposes. Regrowth may be grazed starting 17 days after last application.

**Rust** Apply the specified rate of Tebu Crop 3 6F as soon as weather conditions are favorable for rust development or when first rust pustules are present. Repeat applications at 14 to 16 day intervals. Under heavy disease pressure use 6 to 8 fl. oz. /A and apply at shorter spray intervals.

**Powdery Mildew** Apply specified rate of Tebu Crop 3 6F when powdery mildew first appears on the leaves. Repeat applications at 14 to 16 day intervals. Under heavy disease pressure use 6 to 8 fl. oz. /A and apply at shorter spray intervals.

**Pre harvest interval (PHI) 4 days**

**Restricted entry interval (REI) =12 hours**

CROP	DISEASE	Tebu-Crop 3.6F APPLICATION RATE (fl. oz. per acre)
------	---------	---

Green Onion	White Rot	4 – 6
Leek	( <i>Sclerotium cepivorum</i> ) (suppression only)	
	Rust	
Spring Onion	( <i>Puccinia allii</i> <i>Puccinia porri</i> )	
Welsh Onion	Purple Blotch	
	( <i>Alternaria porii</i> )	

**Notes** For the control of diseases make foliar applications using an interval of 10 – 14 days. Apply Tebu Crop 3 6F in a protective spray schedule or when weather conditions are favorable for rust development. Do not apply more than 24 fl. oz. Tebu Crop 3 6F per acre per season.

**Comments** For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank mixed with Tebu Crop 3 6F. Tebu Crop 3 6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time Tebu Crop 3 6F will be resistant to weathering. Tebu Crop 3 6F is a demethylation inhibitor (DMI) fungicide (Group 3).

**Restricted entry interval (REI) =12 hours**

**Pre harvest interval (PHI) = 7 days**

14/30

CROP	DISEASE	Tebu-Crop 3.6F APPLICATION RATE (fl. oz. per acre)
Hops	Powdery Mildew ( <i>Sphaerotheca humuli</i> / <i>Sphaerotheca macularis</i> )	4 – 8

**Notes** Apply the specified dosage in a protective spray schedule to foliage Repeat applications at 10 to 14 day intervals Do not apply more than 32 fl oz of Tebu Crop 3 6F per acre per crop season Increase the spray volume and the application rate as vine growth increases during the season

**Comments** For optimum disease control the lowest labeled rate of a spray surfactant should be tank mixed with Tebu Crop 3 6F Tebu Crop 3 6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs After this period of time Tebu Crop 3 6F will be resistant to weathering Tebu Crop 3 6F is a demethylation inhibitor (DMI) fungicide (Group 3)

**Restricted entry interval (REI) = 12 hours**

**Pre harvest interval (PHI) = 14 days**

CROP	DISEASE	Tebu-Crop 3.6F APPLICATION RATE (fl. oz. per acre)
<b>Leafy Brassica Greens</b>	Cercospora Leaf Spot	
Broccoli Raab	( <i>Cercospora brassicicola</i> )	
Chinese Cabbage (Bok Choy)	Powdery Mildew	
Collards	( <i>Erysiphe cruciferarum</i> )	3 4
Kale	Alternaria Leaf Spot	
Mizuma	( <i>Alternaria brassicicola</i> )	
Mustard Greens		
Mustard Spinach		
Rape Greens		
Turnip Greens		

**Notes** Do not apply more than 16 fl oz Tebu Crop 3 6F per acre per season

**Comments** For optimum results use as a preventative treatment Begin applications as soon as crop and/or environmental conditions become favorable for disease development The lowest recommended rate of a spray surfactant may be tank mixed with Tebu Crop 3 6F Tebu Crop 3 6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs After this period of time Tebu Crop 3 6F will be resistant to weathering Tebu Crop 3 6F is a demethylation inhibitor (DMI) fungicide (Group 3)

**Restriction** Application to turnip greens is limited to east of the Rockies

**Reapplication interval** Do not apply more than once every 10 days

**Restricted entry interval (REI) = 12 hours**

**Pre harvest interval (PHI) = 7 days**

CROP	DISEASE	Tebu-Crop 3.6F APPLICATION RATE (fl. oz. per acre)
------	---------	--

Lychee	Anthrachnose <i>gloeosporioides</i>	4 – 6 ( <i>Colletotrichum</i> )
--------	--	---------------------------------

**Notes** Begin first application of Tebu Crop 3 6F as panicle emerges Spray up to 6 fl oz Tebu Crop 3 6F per acre every 10 days thereafter for a total of 8 sprayings Apply specified dosage in a minimum of 50 gallons of spray solution per acre by ground only Do not apply more than 48 fl oz Tebu Crop 3 6F per acre per season

**Comments** For optimum disease control the lowest labeled rate of a non ionic spray surfactant should be tank mixed with Tebu Crop 3 6F Tebu Crop 3 6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs After this period of time Tebu Crop 3 6F will be resistant to weathering Tebu Crop 3 6F is a demethylation inhibitor (DMI) fungicide (Group 3)

**Restricted entry interval (REI) = 2 days**

**Pre harvest interval (PHI) = 0 (zero) days**

CROP	DISEASE	Tebu-Crop 3.6F APPLICATION RATE (fl. oz. per acre)
------	---------	--

Okra	Cercospora Leaf Spot <i>spp</i>	4 – 6 ( <i>Cercospora</i> )
------	------------------------------------	-----------------------------

**Notes** Apply specific dosage of Tebu Crop 3 6F in a preventative spray program Use the highest rate when disease conditions are favorable and in areas where high disease pressure is expected Applications may be repeated at 14 day intervals in order to maintain control of the disease Apply specified dosage as a foliar spray in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air Do not apply more than 24 fl oz Tebu Crop 3 6F per acre per season

**Comments** For optimum disease control the lowest labeled rate of spray surfactant should be tank mixed with Tebu Crop 3 6F Tebu Crop 3 6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs After this period of time Tebu Crop 3 6F will be resistant to weathering Tebu Crop 3 6F is a demethylation inhibitor (DMI) fungicide (Group 3)

**Restricted entry interval (REI) = 12 hours**

**Pre harvest interval (PHI) = 3 days**

CROP	DISEASE	Tebu-Crop 3.6F APPLICATION RATE (fl. oz. per acre)
------	---------	--

Peanut	<b>SOILBORNE</b> Cylindrodadium Black Rot (suppression) Rhizoctonia Limb Rot Rhizoctonia Pod Rot (Virginia and North Carolina only) Sclerotium Stem and Pod Rot (White Mold Southern Blight Southern Stem Rot) <b>FOLIAR</b> Early Leaf Spot Late Leaf Spot Leaf Rust Pepper Spot ( <i>Leptosphaerulina</i> ) Web Blotch ( <i>Phoma</i> )	7.2
--------	--	-----



**Information** For optimum control of the specified soilborne diseases four consecutive applications of Tebu Crop 3 6F must be made at 14 day intervals. A maximum of 28.8 fl. oz. (8.1 lbs a.i.) of Tebu Crop 3 6F may be applied per crop season. Do not feed hay or threshings or allow livestock to graze in treated areas.

Tebu Crop 3 6F is a sterol demethylation inhibitor (DMI) fungicide. Chlorothalonil may be tank mixed at the rate of 12 ounces of active ingredient with Tebu Crop 3 6F as a leaf spot resistance management strategy. A spray surfactant is not necessary when Tebu Crop 3 6F is tank mixed with chlorothalonil. Mixing or alternating Tebu Crop 3 6F with other DMI fungicides may lead to resistance.

Tebu Crop 3 6F must be carried by rainfall or irrigation into the root and pod zone for control of root and pod rots caused by *Sclerotium rolfsii* and *Rhizoctonia solani*. Drought conditions will decrease the effectiveness of Tebu Crop 3 6F against the root and pod rots. Use Tebu Crop 3 6F in conjunction with cultural practices that are known to reduce the severity of soilborne diseases such as proper crop rotation practices.

**FOUR APPLICATION SPRAY PROGRAM** Apply the specified rate in a preventive spray schedule. See table below for proper timing of applications. Applications of chlorothalonil should be made prior to and following applications of Tebu Crop 3 6F to discourage development of resistant strains of fungi. For optimum control of foliar diseases such as leaf rust, web blotch, and pepper spot, the lowest label specified rate of a spray surfactant should be tank mixed with Tebu Crop 3 6F.

**LEAF SPOT ADVISORY SCHEDULE** For control of soil borne diseases in an advisory schedule, apply Tebu Crop 3 6F in the first advisory spray in July and continue Tebu Crop 3 6F applications at 14 day intervals. When applying Tebu Crop 3 6F after August 15, tank mix with chlorothalonil for resistance management purposes.

**Application Timing of Tebu Crop 3 6F for Optimum Control of White Mold and Rhizoctonia Limb and Pod Rot**

Spray Program	Tebu Crop 3 6F Application No	Chlorothalonil Application No
7 applications	3, 4, 5 and 6	1, 2 and 7

Restricted entry interval (REI) = 12 hours

Pre harvest interval (PHI) = 14 days

CROP	DISEASE	Tebu-Crop 3 6F APPLICATION RATE (fl. oz. per acre)
Pecan	Brown Leaf Spot ( <i>Sirosporum diffusum</i> )	4 – 8
	Downy Spot ( <i>Mycosphaerella caryigena</i> )	
	Liver Spot ( <i>Gnomonia caryae</i> )	
	Scab ( <i>Cladosporium caryigenum</i> )	
	Vein Spot ( <i>Gnomonia nerviseda</i> )	
	Zonate Leaf Spot ( <i>Grovesinia pyramidalis</i> )	



**Notes** Apply Tebu Crop 3 6F in a preventative spray schedule beginning at early bud break (young leaves unfolding) and continue applications at 10 to 14 day intervals through the pollination period. Tebu Crop 3 6F should be applied at 4 fl oz /acre in a tank mix with the labeled rate of Super Tin® in cover sprays. Follow label directions for the use of Super Tin®. Do not add a surfactant to the spray solution when tank mixing Tebu Crop 3 6F with Super Tin®. Apply Tebu Crop 3 6F in a spray volume of 15 or more gallons per acre by air or 50 or more gallons per acre by ground. Apply 7 – 8 fl oz /acre of Tebu Crop 3 6F to full size mature trees and 4 – 6 fl oz Tebu Crop 3 6F per acre to smaller trees. Apply the high rate to varieties that are highly susceptible to the indicated diseases or when severe disease conditions exist. The lowest labeled rate of a surfactant may be added to the spray solution for optimum control of the indicated diseases. Do not apply after shucks begin to split. Do not apply more than 32 fl oz Tebu Crop 3 6F per acre per crop season. Do not cut cover crops in treated areas for feed or allow livestock to graze treated areas.

**Comments** For optimum disease control the lowest labeled rate of a spray surfactant should be tank mixed with Tebu Crop 3 6F. Tebu Crop 3 6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time Tebu Crop 3 6F will be resistant to weathering. Tebu Crop 3 6F is a demethylation inhibitor (DMI) fungicide (Group 3). It may be applied in a tank mix or alternated (every other spray application) with a non DMI fungicide as a resistance management strategy.

**Restricted entry interval (REI) = 12 hours**

**Pre harvest interval (PHI) = Do not apply after shucks begin to split**

CROP	DISEASE	Tebu-Crop 3.6F APPLICATION RATE (fl. oz. per acre)
Soybean	Rust ( <i>Phakopsora pachyrhizi</i> )	3 – 4
	Powdery Mildew ( <i>Microsphaera diffusa</i> )	

**Notes** Apply Tebu Crop 3 6F as a broadcast foliar spray as a preventative spray or at first visible symptoms of disease. Repeat applications on a 10 to 14 day spray interval if environmental conditions are favorable for continued disease development. Use of the higher rates and shorter spray intervals are recommended when disease pressure is severe. The lowest label recommended rate of a spray surfactant may be tank mixed with Tebu Crop 3 6F. Apply Tebu Crop 3 6F in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons per acre by aircraft spray equipment. Do not apply more than 12 fl oz Tebu Crop 3 6F per acre per use season. Do not make more than three applications per season.

**Restricted entry interval (REI) = 12 hours**

**Pre harvest interval (PHI) = 21 days**

CROP	DISEASE	Tebu-Crop 3.6F APPLICATION RATE (fl. oz. per acre)
Sunflower	Rust ( <i>Puccinia helianthi</i> )	4 – 6

**Notes** Apply specific dosage of Tebu Crop 3 6F at the earliest sign of infection (rust pustules developing) or when weather conditions are favorable for rust development. Apply higher rate to highly susceptible varieties and/or under severe disease conditions. Application may be repeated at 14 days if necessary to maintain control of the disease. Apply specified dosage in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. Do not apply more than 16 fl oz Tebu Crop 3 6F per acre per season.

**Comments** For optimum disease control the lowest labeled rate of a spray surfactant should be tank mixed with Tebu Crop 3 6F. Contact your state extension service for a list of approved surfactants. Tebu Crop 3 6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time Tebu Crop 3 6F will be resistant to weathering. Tebu Crop 3 6F is a demethylation inhibitor (DMI) fungicide (Group 3).

**Restricted entry interval (REI) = 12 hours**

**Pre harvest interval (PHI) = 50 days**

18/30

CROP	DISEASE	Tebu-Crop 3.6F APPLICATION RATE (fl. oz. per acre)
------	---------	--

Turnip	Cercospora Leaf Spot	4 – 7.2
--------	----------------------	---------

(Application is limited to east of the Rockies) (*Cercospora brassicicola*)

**Notes** Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 12 to 14 day intervals. Do not apply more than 28.8 fl. oz. Tebu Crop 3.6F per acre per crop season.

**Comments** For optimum disease control the lowest labeled rate of a spray surfactant should be tank mixed with Tebu Crop 3.6F. Tebu Crop 3.6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time Tebu Crop 3.6F will be resistant to weathering. Tebu Crop 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3).

**Restricted entry interval (REI) = 12 hours**

**Pre harvest interval (PHI) = 7 days**

CROP	DISEASE	Tebu-Crop 3.6F APPLICATION RATE (fl. oz. per acre)
------	---------	--

Wheat	Rusts leaf, stem, and stripe ( <i>Puccinia</i> spp.)	4
	Head blight or scab ( <i>Fusarium</i> spp.) Suppression	

**NOTES for Wheat** Wheat fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. A maximum of 4 fl. oz. of Tebu Crop 3.6F may be applied per acre per crop per season. Straw cut after harvest may be fed or used for bedding. Do not allow livestock to graze or feed green forage to livestock prior to 6 days after treatment with Tebu Crop 3.6F. Apply Tebu Crop 3.6F in a minimum of 10 gallons of spray solution per acre by ground or in a minimum of 5 gallons of spray solution per acre by air.

**Application timing Directions**

**Rusts** Apply Tebu Crop 3.6F at the earliest sign of rust pustules on foliage.

**Fusarium head blight** Optimal timing of Tebu Crop 3.6F for Fusarium head blight suppression is the beginning of flowering on main stem heads (Feekes 10.51).

**Restricted entry interval (REI) = 12 hours**

**Pre harvest interval (PHI) = 30 days**

**SEED TREATMENT – Corn** (Sweet Corn, Field Corn Grown for Seed, and Popcorn)  
For control of soilborne and seedborne Fusarium and soilborne and seedborne head smut

**SEED LABELING** To meet U.S. Federal Seed Act requirements, all seed treated with Tebu Crop 3.6F must be labeled

**TREATED SEED DO NOT USE FOR FOOD, FEED, OR OIL PURPOSES**

**Treated with Tebuconazole**

**USE PRECAUTION** When using formulations that do not contain dye to comply with 40 CFR 153.155, all seed treated with an economic poison must be colored to distinguish and prevent subsequent inadvertent use as a food for man or feed for animals.

DISEASE	RATE Fl. Oz./CWT	DIRECTIONS FOR USE
<b>Soilborne and Seedborne</b> Fusarium	0.071	Apply as a seed treatment using standard slurry or mist type seed treatment equipment. Uniform application of seed is necessary to ensure seed safety and best disease protection. Seed should be sound and well cured prior to treatment. Product should be diluted with sufficient water to ensure complete seed coverage. Consult a seed treatment specialist regarding slurry rates recommended for the crop to be treated with Tebu Crop 3.6F. The length of control will vary depending on the rate used.
<b>Soilborne and Seedborne Head</b> Smut ( <i>Sphacelotheca reiliana</i> )	0.27 – 0.54	

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage** Store in the original container in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store out of the reach of children, preferably in a locked storage area. Open and handle container in a manner as to prevent spillage. If container is leaking, invert to prevent leakage. If the container is leaking or material is spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away.

**Pesticide Disposal** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal** Nonrefillable container. Do not refill or reuse container. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate in application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials, resistant strains, or other influencing factors in the use of the product. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Sharda USA LLC, Manufacturer and Seller, harmless for any claims relating to such factors.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SHARDA USA LLC AND MANUFACTURER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ON THIS LABEL. To the extent consistent with applicable law, Sharda USA LLC, Manufacturer or Seller, shall not be liable for any incidental, consequential, or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE

20/30

083529 00011 20120410 V2

Sharda USA Sub Label A\_Notif

EXCLUSIVE LIABILITY OF SHARDA USA LLC MANUFACTURER AND SELLER FOR ANY AND ALL CLAIMS LOSSES INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY CONTRACT NEGLIGENCE TORT STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR AT THE ELECTION OF SHARDA USA LLC MANUFACTURER OR SELLER THE REPLACEMENT OF THE PRODUCT

Sharda USA LLC Manufacturer and Seller offer this product subject to the foregoing conditions of sale and limitations of warranty and of liability which may not be modified except by written agreement signed by a duly authorized representative of Sharda USA LLC

c

e

**SHAR TEB 3 6FL FUNGICIDE****[ABN TEBUSHA 3 6FL FUNGICIDE, Tebu Turf 3 6F, Tebu-Crop 3 6F]****Active Ingredient**

Tebuconazole alpha [2 (4 chlorophenyl)ethyl] alpha(1 1 dimethylethyl) 1 H 1 2 4 triazole 1 ethanol

38 7%

**Other Ingredients**

61 3%

**Total****100 0%**

Contains 3 6 pounds tebuconazole per gallon

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

Si usted no entiende la etiqueta busque a alguien para que se la explique a usted en detalle (If you do not understand the label find someone to explain it to you in detail )

See additional precautionary statements and directions for use in booklet

**STOP Read the label before use****FIRST AID**

<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice</li> <li>• Have person sip a glass of water if able to swallow</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor</li> <li>• Do not give anything by mouth to an unconscious person</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing</li> <li>• Rinse skin immediately with plenty of water for 15 20 minutes</li> <li>• Call a poison control center or doctor for treatment advice</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 20 minutes</li> <li>• Remove contact lenses if present after the first 5 minutes then continue rinsing eye</li> <li>• Call a poison control center or doctor for treatment advice</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air</li> <li>• If person is not breathing call 911 or an ambulance then give artificial respiration preferably by mouth to mouth if possible</li> <li>• Call a poison control center or doctor for further treatment advice</li> </ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment	
<b>NOTE TO PHYSICIAN</b> No specific antidote Treat symptomatically	
<b>Symptoms of Poisoning</b> The compound does not cause any definite symptoms that would be diagnostic Contact with the eyes may cause irritation	
<b>EMERGENCY NUMBERS</b>	
For 24 hour medical emergency assistance (human or animal) call 1 800 222 1222 For chemical emergency assistance (spill leak fire or accident) call CHEMTREC at 1 800 424 9300	

**Not registered for sale in the state of New York****EPA Reg No 83529 11****EPA Est No****Net Contents 2 5 Gallons****Manufactured for**

Sharda USA LLC

PO Box 640

Hockessin DE 19707

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Causes moderate eye irritation Harmful if swallowed inhaled or absorbed through skin Avoid contact with skin eyes and clothing Avoid breathing vapor or spray mist

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical resistant to this product are listed below If you want more options follow the instructions for category C on an EPA chemical resistance category selection chart

**Applicators and other handlers must wear**

- Long sleeved shirt and long pants
- Chemical resistant gloves such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton and
- Shoes plus socks

Follow manufacturer s instructions for cleaning/maintaining PPE If no such instructions for washables exist use detergent and hot water Keep and wash PPE separately from other laundry

**ENGINEERING CONTROLS STATEMENTS**

When handlers use closed systems enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170 240(d)(4 6)] the handler PPE requirements may be reduced or modified as specified in the WPS

**USER SAFETY RECOMMENDATIONS**

**Users should**

- Wash hands before eating drinking chewing gum using tobacco or using the toilet
- Remove clothing immediately if pesticide gets inside Then wash thoroughly and put on clean clothing
- Remove PPE immediately after handling this product Wash the outside of gloves before removing As soon as possible wash thoroughly and change into clean clothing

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to mammals fish and aquatic invertebrates Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark Runoff may be hazardous to aquatic organisms in neighboring areas Do not contaminate water when disposing of equipment washwater or rinsate

**Groundwater Advisory** Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use Use of this chemical in areas where soils are permeable particularly where the water table is shallow may result in groundwater contamination

**Surface Water Advisory** This product may contaminate water through drift of spray in wind This product has a high potential for runoff for several months or more after application Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product A level well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds streams and springs will reduce the potential for contamination of water from rainfall runoff Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling Do not apply this product in a way that will contact workers or other persons either directly or through drift Only protected handlers may be in the area during application For any requirements specific to your State or Tribe consult the agency responsible for pesticide regulation

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) specified in the use directions for each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

**NON AGRICULTURAL USE REQUIREMENTS**

The requirements in this box only apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Golf Course Turf and Landscape Uses: Keep children and pets out of treated areas until sprays have dried.

**INFORMATION FOR GOLF COURSE TURF AND ORNAMENTAL USE**

**Chemigation:** Do not apply this product through any type of irrigation system.

**OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES, OR NATURAL PONDS, AND ESTUARIES**

- Do not apply within 100 feet of aquatic areas listed above.
- Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetation filter strip.
- See Spray Drift Management section for further information.

**Spray Drift Management**

Make ground application when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperatures.

Do not make ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

**Spray Volume** For best results Tebu Turf 3 6F may be applied in 66 132 gallons of water per acre for turf using ground based equipment For ornamentals 50 300 gallons of finished spray per acre are recommended depending upon equipment plant species and plant growth stage at time of application For the most effective results equipment calibration should be checked regularly When using lower spray volumes be sure to maintain uniform application and full crop coverage so as to ensure effective control Increase spray volume to ensure proper application if required

### Compatibility Test for Mix Components

Before mixing components always perform a compatibility jar test For 66 gallons per acre spray volume use 5 cups of water in a clear clean mixing jar For other spray volumes adjust accordingly Only use water from the intended source at the source temperature Add components in the sequence indicated below in Mixing Order using 3 teaspoons for each pound of dry product or 1½ teaspoon for each pint of liquid product of recommended label rate per acre Always cap the jar and invert 10 cycles between component additions When the components have all been added to the jar and fully mixed let the solution stand for 15 minutes Evaluate the solution for uniformity and stability The spray solution should not have free oil on the surface nor fine particles that precipitate to the bottom nor thick (clabbered) texture If the spray solution is not compatible repeat the compatibility test with the addition of a suitable compatibility agent and use the compatibility agent as directed on its label

**Mixing** Continuous agitation is required during mixing When mixing this product and water use the specified application rates as listed for each crop on this label Before combining any other substances with the mixture ensure that the Tebu Turf 3 6F is completely dispersed in the mixture

### Recommended Mixing Procedure

- 1 Water Add three quarters of the required volume to a thoroughly clean sprayer tank
- 2 Agitation Start agitation and maintain constant agitation throughout mixing and application
- 3 Inductor If an inductor is used rinse it thoroughly after each component has been added
- 4 Products in PVA Bags Place any product contained in water soluble PVA bags into the mixing tank Wait until all water soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing
- 5 Water Dispersible Products Including dry flowables (DF) wettable powders (WP) suspension concentrates (SC) or suspo emulsions (SE)
- 6 Water soluble products
- 7 Emulsifiable concentrates (such as oil concentrates when applicable)
- 8 Water soluble additives (such as AMS or UAN when applicable)
- 9 Remaining quantity of water

### Resistance Management Information

The active ingredient in Tebu Turf 3 6F is a member of the DMI (Demethylation Inhibitor) fungicide group (FRAC grouping 3) and exhibits no known cross resistance to products with the same mode of action when used repeatedly in the same location or in successive years as the primary method of control for targeted diseases Because the speed and scope of resistant population development cannot be predicted the use of this product should conform to resistance management strategies established for the crop and use area Such strategies may include the rotation and/or tank mixing with products utilizing different modes of action or limiting the number of applications per season Contact your local university or extension specialist and/or manufacturer for fungicide resistance management recommendations

### DISEASE CONTROL IN GOLF COURSE TURF

For use on all Golf turf applications of cool season and warm season grasses (such as Bentgrasses Bluegrasses Fescues Ryegrasses St Augustine grasses and Zoysia) or their mixtures Tebu Turf 3 6F is not phytotoxic to any of the above mentioned grasses when used in accordance with the label

**Note** Bermudagrass can be sensitive to Tebu Turf 3 6F under certain conditions Do not apply consecutive applications during or just after dormancy break Avoid applications when temperatures are expected to exceed 85 degrees F



Tebu Turf 3 6F can be used for the prevention and control of the diseases mentioned in table below. Begin applications when conditions favor disease development and repeat applications as long as these conditions persist. Preventative treatments can be applied using 28 day intervals as indicated. When treating golf greens, always treat aprons and approaches. Spray uniformly over the area to be treated with properly calibrated equipment.

Apply the specified amount of Tebu Turf 3 6F in sufficient water for thorough coverage. A volume of 66 132 gallons per acre (1 5 3 0 gallons per 1 000 sq ft) is recommended. Apply using properly calibrated low volume hand held mechanical or motorized ground broadcast equipment. Application to small areas may be made with low pressure handwand or backpack equipment. Maintain constant agitation during application.

Depending on the disease, Tebu Turf 3 6F should be watered into the crown and active root zone for best results. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. For best results, use spray mixture the same day it is prepared.

#### **TURF USE RESTRICTIONS AND PRECAUTIONS**

- For use on golf course turf only
- Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (i.e., elementary, middle and high school), campgrounds, churches, and theme parks.
- Not for homeowner use
- Not for use on turf being grown for sale or commercial use as sod
- Do not use clippings for animal feed
- Do not exceed 3 6 fl. oz of Tebu Turf 3 6F per 1 000 sq ft per year
- Do not apply more than 6 applications per year

## Golf Course Turf Disease Control

DISEASE	RATE of Tebu Turf 3 6F (Fl oz/1000 Sq Ft)	NOTES
Dollar Spot ( <i>Sclerotinia homoeocarpa</i> )  Copper Spot ( <i>Gloeocercospora sorghi</i> )  Powdery Mildew ( <i>Erysiphe graminis</i> )  Corticium Red Thread ( <i>Laetisaria fuciformis</i> )  Rusts ( <i>Puccinia</i> spp) Brown Patch/Rhizoctonia Blight Large Patch ( <i>Rhizoctonia solani</i> )  Brown Ring Patch ( <i>R. circinata</i> ) Anthracnose Basal and Foliar ( <i>Colletotrichum cereal</i> )  Red Thread ( <i>Laetisaria fuciformis</i> )  Pink Patch ( <i>Limonomyces rosipellis</i> )	0.6	For prevention begin applications when conditions are favorable for disease development. Do not make two consecutive applications of Tebu Turf 3 6F. Alternate with another fungicide with a different mode of action. A second application may be made after 28 days.
Bermuda Grass decline ( <i>Gaeumannomyces graminis</i> var <i>graminis</i> )	0.6	Immediately after fungicide is applied irrigate the area with sufficient water to move the active ingredient down into the crown and root zone of the turf. The amount of water is dependent on the depth of the root zone. For prevention begin applications two or four weeks prior to the historical appearance of disease symptoms. Initiate cultural control practices at the same time the fungicide is applied. Refer to your local County Extension Service for this information. Apply subsequent applications at 28 day intervals.
Take All Patch ( <i>Gaeumannomyces graminis</i> )	0.6	For prevention apply in the fall when soil temperature reaches 55-65 F and again in the spring under similar soil temperature conditions. Applications in both fall and spring may be necessary. Immediately after fungicide is applied irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.

DISEASE	RATE of Tebu Turf 3 6F (Fl oz/1000 Sq Ft)	NOTES
Gray Leaf Spot ( <i>Pyricularia grisea</i> )	0.6	Apply when conditions are favorable for disease development at 28 day intervals. If using under conditions favoring moderate to heavy disease pressure, Tebu Turf 3 6F can be tank mixed with a registered contact fungicide at the label rate.
Stripe Smut ( <i>Ustilago striiformis</i> )	0.6	Make a single application to historical disease areas in spring as grass growth begins.
Spring Dead Spot ( <i>Leptosphaeria korrea</i> , <i>L. narmari</i> , <i>Ophiosphaerella herpotricha</i> , <i>Gaeumannomyces graminis</i> )	0.6	For prevention, apply in fall when soil temperature reaches 65 F and again in spring under similar soil temperature conditions or after dormancy break. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
Necrotic Ring Spot ( <i>Leptosphaeria korrea</i> )		
Fusarium Patch ( <i>Fusarium roseum</i> )	0.6	Apply first application in mid June or 28 days prior to time this blight normally becomes evident. Make applications at no less than 28 day intervals.
Summer Patch ( <i>Magnaporthe poae</i> )	0.6	Apply beginning in the spring. Do not make two consecutive applications of Tebu Turf 3 6F. Alternate with another fungicide with a different mode of action. Make second and third applications at 28 day intervals. See local university recommendations for suggested timing. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
Zoysia Patch Large Patch of zoysia ( <i>Rhizoctonia solani</i> )	0.6	Make first application in early fall (mid September to mid October) prior to development of disease symptoms. A second application in early spring may be necessary in areas where disease pressure is known to be heavy.
Gray Snow Mold/ Typhula Blight ( <i>Typhula incarnate</i> )	0.6	Apply in the fall before anticipated turf dormancy and before first snow cover. If turf breaks dormancy during winter months, a second application may be made. Do not apply over snow cover or when turf is dormant. It is recommended that Tebu Turf 3 6F be tank mixed with other registered snow mold products for best season long results.
Pink Snow Mold/Microdochium Patch ( <i>Microdochium nivalis</i> )		

NOTE: Apply the specified amount of Tebu Turf 3 6F in 1.5 to 3.0 gallons of water per 1000 sq. ft. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. Do not use clippings for animal feed. Do not exceed 3.6 fl. oz. of Tebu Turf 3 6F per 1000 sq. ft. per year. Do not exceed 6 applications per year.

**DISEASE CONTROL IN FIELD, NURSERY AND CONTAINER ORNAMENTALS AND COMMERCIAL and RESIDENTIAL LANDSCAPES**

Tebu Turf 3 6F can be used in a preventative and curative disease control program for the listed plant types and disease in the table below. Optimum disease management is obtained when Tebu Turf 3 6F is used in conjunction with sound disease management practices.

Apply material with properly calibrated hand held mechanical or motorized spray equipment. Begin applications when disease first appears and repeat at 14-21 day intervals during the growing season. Use the shortest interval when conditions are unusually favorable for the development of disease. For hand held mechanical or motorized applications, mix as directed below and apply as a foliage coverage spray to drip for the prevention and control of the diseases listed below. Choose a finished spray volume appropriate for the size of the plants and amount of foliage, which will provide thorough coverage throughout the canopy. Allow sprays to dry before overhead irrigation is applied.

Apply Tebu Turf 3 6F at rates of 4-10 fl oz per acre in 100 gallons of water. Spray volume may range from 50 up to 300 gallons of finished spray per acre depending upon equipment, plant species and plant growth stage at time of application.

**Note:** The Directions for Use of this product reflect the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before a foliar treatment is applied to any plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results. This product is not recommended for use on African Violets, Begonias, Boston Fern, and Geraniums.

**ORNAMENTAL USE RESTRICTIONS AND PRECAUTIONS**

- For use on ornamental plants only, not for woodlands or forest management
- Not for homeowner use
- Do not apply more than 10 fl oz per acre in a single application
- Do not apply more than 0.31 gallons (40 fl oz) of Tebu Turf 3 6F (equal to 1.13 lbs of tebuconazole) per acre per year
- Do not make more than 4 applications per year at highest rate
- Do not apply to bearing fruit trees

**Ornamentals Disease Control**

PLANTS	DISEASE	APPLICATION	
		To Prevent Diseases	To Treat Existing Disease
Roses	Black Spot Powdery Mildew Rust	Apply every 14 21 days during the growing season starting when leaves first appear	Apply every 14 days for a total of 3 applications beginning at the first sign of disease
Flowers	Leaf Spot Powdery Mildew Rust Southern Blight	Apply at least 3 times per year 14 21 days apart beginning with Spring bud break Rotation or Tank mixing with barrier protectant fungicides is recommended for resistance management	
Crabapples (Ornamental) Dogwoods and Other Landscape (Ornamental) Trees	Anthrachnose Leaf Spot Powdery Mildew Rust Scab		
Azaleas Camellias Rhododendrons and Other Landscape (Ornamental) Shrubs	Anthrachnose Black Spot Leaf Spot Petal Blight Powdery Mildew	Petal Blight Apply 2 3 times per week into the flowers as the open and develop color	
Ground Covers and Vines	Rust Southern Blight		
HOW MUCH TO USE FOR SMALL PLANTINGS ADD 1 TEASPOON TO 2 5 GALLONS OF WATER			

**Pump Style Sprayers**

- 1 Add the appropriate amounts of concentrate and water to the sprayer tank
- 2 Close the sprayer shake well and pressurize
- 3 Adjust nozzle to a coarse spray pattern and apply
- 4 Occasionally re pressurize the sprayer if needed to maintain a good spray pattern

**STORAGE AND DISPOSAL**

Do not contaminate water food or feed by storage or disposal

**Pesticide Storage** Store in the original container in a cool dry place and in such a manner as to prevent cross contamination with other pesticides fertilizers food and feed Store out of the reach of children preferably in a locked storage area Open and handle container in a manner as to prevent spillage If container is leaking invert to prevent leakage If the container is leaking or material is spilled for any reason or cause carefully dam up spilled material to prevent runoff Refer to Precautionary Statements on label for hazards associated with the handling of this material Do not walk through spilled material Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below In spill or leak incidents keep unauthorized people away

**Pesticide Disposal** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility

**Container Disposal** Nonrefillable container Do not refill or reuse container Triple rinse as follows Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip Fill the container ¼ full with water and recap Shake for 10 seconds Pour rinsate in application equipment or a mix tank or store rinsate for later use or disposal Drain for 10 seconds after the flow begins to drip Repeat this procedure two more times Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or incineration or if allowed by state and local authorities by burning If burned stay out of smoke

**CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

The Directions for Use of this product must be followed carefully It is impossible to eliminate all risks inherently associated with the use of this product Crop injury ineffectiveness or other unintended consequences may result because of such factors as manner of use or application weather or crop conditions presence of other materials resistant strains or other influencing factors in the use of the product To the extent consistent with applicable law all such risks shall be assumed by Buyer and User and Buyer and User agree to hold Sharda USA LLC Manufacturer and Seller harmless for any claims relating to such factors TO THE EXTENT CONSISTENT WITH APPLICABLE LAW SHARDA USA LLC AND MANUFACTURER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ON THIS LABEL To the extent consistent with applicable law Sharda USA LLC Manufacturer or Seller shall not be liable for any incidental consequential or special damages resulting from the use or handling of this product TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF SHARDA USA LLC MANUFACTURER AND SELLER FOR ANY AND ALL CLAIMS LOSSES INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY CONTRACT NEGLIGENCE TORT STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR AT THE ELECTION OF SHARDA USA LLC MANUFACTURER OR SELLER THE REPLACEMENT OF THE PRODUCT

Sharda USA LLC Manufacturer and Seller offer this product subject to the foregoing conditions of sale and limitations of warranty and of liability which may not be modified except by written agreement signed by a duly authorized representative of Sharda USA LLC