



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs  
Registration Division (7505C)  
1200 Pennsylvania Ave., N.W.  
Washington, D.C. 20460

EPA Reg. Number:

83529-14

Date of Issuance:

JAN 28 2008

NOTICE OF PESTICIDE:

Registration  
 Reregistration

(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Sharda Imidacloprid 5SC

Name and Address of Registrant (include ZIP Code):

Sharda USA LLC  
c/o SynTech Global  
P.O. Box 640  
Hockessin, DE 19707

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit such data.
2. The data requirements for storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) are not satisfied. A one year study is required to satisfy these data requirements. You have 18 months from the date of registration to provide these data.

Continued on page 2

Signature of Approving Official:

Venus Eagle, Product Manager  
Insecticide-Rodenticide Branch, Registration Division (7505P)

Date:

JAN 28 2008

3. Make the following label changes before you release the product for shipment:

- a. Revise the EPA Registration Number to read, "EPA Reg. No. **83529-14**."
- b. NOTE: It is recommended that you add the complete company mailing address to the label.
- c. On page 1, under the First Aid section, add the following text:
  - **NOTE TO PHYSICIAN: No specific antidote is available. Treat the patient symptomatically.**
- d. On page 2, under the User Safety Recommendations, revise the second statement so it reads, "Remove clothing/**PPE** immediately if pesticide gets inside. Then wash thoroughly..."
- e. On page 10 in the Rotational Plant-Back Intervals section, modify the second sentence so it reads, "The following plant-back intervals **must** be observed for listed crops."
- f. On page 10, in the Rotational Plant-Back Intervals table, correct the spelling of "**triticale**."
- g. On page 10, in the Rotational Plant-Back Intervals table, modify the text of the second asterisked note so it reads, "For all other crops..., a 12-month plantback [*sic*] interval **must** be observed."
- h. NOTE: The spelling of "**plant-back**" is inconsistent. In some cases, it appears without the hyphen.
- i. On page 10, revise the first sentence of the third paragraph within the "Note" section so it reads, "Labels for commercially treated seeds **must** include the following addition to the Environmental Hazards statements:..."
- j. On page 10, revise the second sentence under the Use Restriction section so it reads, "Rapeseed grown and harvested from treated seed is only for industrial uses and **must** not be used for edible oil or any other human/feed consumption."
- k. Add the following, bolded text to the third sentence of the Disclaimer of Warranties section so it reads, "**To the extent consistent with applicable law, Sharda USA LLC disclaims any liability whatsoever...**"

4. Submit one copy of the revised final printed label for the record before you release the product for shipment.

5. NOTE: For future submissions, make the following changes to your Confidential Statement of Formulation (Item 4 of EPA Form 8570-4):

- a. Revise the EPA Reg. No. so it reads, "EPA Reg. No. **83529-14**"

3 7 14

Page 3  
EPA Reg. No. 83529-14

**If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.**

A stamped copy of the label is enclosed for your records. If you have any questions, please contact Julie Chao at 703-308-8735 or [chao.julie@epa.gov](mailto:chao.julie@epa.gov).

Venus Eagle  
Product Manager  
Insecticide-Rodenticide Branch  
Registration Division (7505P)

Enclosure

4 8 14

Sharda Imidacloprid 5SC

**Active Ingredient:**

Imidacloprid 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine ..... 48.7%

**Other Ingredients:** ..... 51.3%

**Total** ..... 100.0%

Contains 5 lbs. active ingredient per gallon (600 grams per liter).

EPA Reg. No.

EPA Est. No.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

| <b>FIRST AID</b>   |  |
|--|--|
| <b>If swallowed</b>  | <ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul> |
| <b>If on skin or clothing</b>  | <ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>  |
| <b>If inhaled</b>  | <ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>  |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. |  |
| In case of an emergency involving this product, call CHEMTREC at (800) 424-9300                                      |  |
| SEE INSIDE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS   |  |

**Manufactured for:**

Sharda USA LLC  
Hockessin, DE 19707

**ACCEPTED**  
**with COMMENTS**  
**In EPA Letter Dated:**  
**JAN 28 2008**

**Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under EPA Reg. No.  
83529-14**

5 7 14

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

Caution. Harmful if swallowed, absorbed through skin or inhaled. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for Category C on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

1. Long-sleeved shirt and long pants,
2. Socks and shoes, and
3. Chemical-resistant gloves (such as nitrile, butyl, neoprene, barrier laminate, polyvinyl chloride or Viton).

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately, from other laundry.

**USER SAFETY RECOMMENDATIONS**

Users should:

1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
2. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change clothing.

**ENVIRONMENTAL HAZARDS**

This product is highly toxic to birds and aquatic invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Cover or incorporate spilled treated seeds.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For use in commercial seed treaters only, with the exception of application to canola, cotton (delinted seed), field corn, sorghum, wheat and barley, which may be made either by commercial seed treatment or as an end-use seed treatment on agricultural establishments at, or immediately before, planting. This product is to be used in liquid or slurry treaters.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Mix thoroughly before use or use entire container at one time. All tank mixes should be pre-tested to determine physical compatibility between formulations. Observe all cautions and limitations on labeling of all products used in mixtures.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

**Exception.** If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

1. Coveralls
2. Chemical-resistant gloves (such as nitrile, butyl, neoprene, barrier laminate, polyvinyl chloride, or Viton)
3. Shoes plus socks
4. Protective eyewear.

### STORAGE AND DISPOSAL

**Prohibitions:** Do not contaminate water, food or feed by storage, disposal, or by cleaning of equipment. Do not reuse empty container. Open dumping is prohibited.

**Pesticide Storage:** Store in a cool place. Do not store in direct sunlight. Protect from freezing temperatures.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Contact your chemical supplier for container return instructions. If container is not returned in accordance with instructions, triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**STORED SEED PROTECTION**

To protect stored seed of crops listed on this label apply a minimum of 0.8 fluid ounces of Sharda Imidacloprid 5SC per 100 pounds of seed. Pests controlled by this application include Indian Meal Moth (*Plodia interpunctella*), Rice Weevil (*Sitophilus oryzae*), Red Flour Beetle (*Tribolium castaneum*), and Lesser Grain Borer (*Rhizopertha dominica*). If seed being treated is known to have existing pest infestation fumigation may be necessary prior to bagging.

**EARLY SEASON PROTECTION AGAINST SUCKING INSECTS**

| Crop                              | Pest   | Rate (fl. oz./hundred weight seed) | Comment  |
|-----------------------------------|--|------------------------------------|--|
| Canola, Rapeseed and Mustard Seed | Aphids, Flea beetles, Wireworms  | 10.24 – 25.6                       | Use higher rate when pest populations are above historical levels.                   |
|                                   | Lygus Bugs (including Lygus spp.)<br>Seedpod Weevil Larvae (including Ceutorhynchus assimilis) | 15.36 – 25.6                       | Suppression only. Use higher rate when pest populations are above historical levels. |

**Canola, Rapeseed and Mustard Seed (For end use application at agricultural establishments):**

Shake product well before making slurry and apply using a HCBT or Batch Treater. Apply 5.1 to 12.8 fl. oz. of Sharda Imidacloprid 5SC per 50 pound bag. May be diluted with an approved fungicide for disease control. Treat 1/2 of seed with 1/2 of slurry then add remainder of seed and slurry and continue mixing until all seed are covered.

| Crop       | Pest   | Rate (Fl. oz. per 80,000 units of seed)                 | Comment   |
|------------|--|---|---|
| Field Corn | Corn root worm (including Northern, Western, Southern, and Mexican) <sup>(1)</sup> ,<br>Flea beetle,<br>Chinch bug,<br>Southern green stinkbug,<br>White grub,<br>Seed corn maggot,<br>Thrips,<br>Wireworm,<br>Corn leaf aphid,<br>Imported fire ant,<br>Southern corn leaf beetle,<br>Billbug <sup>(2)</sup> ,<br>Grape colaspis <sup>(2)</sup> ,<br>Black cutworm <sup>(3)</sup> | 6.0<br>(Equivalent to 1.34 milligrams a.i. per kernel)  | <sup>(1)</sup> In areas of heavy to severe corn rootworm populations, protection will not be adequate. Use only in areas of light to moderate corn root-worm populations. Consult your State Agricultural Extension Service on levels of corn rootworm populations.<br><br><sup>(2)</sup> Reduces early season feeding damage.<br><br><sup>(3)</sup> Will reduce feeding damage caused by leaf feeding black cutworms that are ½ inch or less in length.<br><br><sup>(4)</sup> Reduces feeding damage during emergence and seedling stages. |
|            | Flea beetle,<br>Chinch bug,<br>Seed corn maggot,<br>Thrips,<br>Wireworm,<br>Corn leaf aphid<br>Imported fire ant,<br>Grape colaspis <sup>(2)</sup> ,<br>White grub   | 2.70<br>(Equivalent to 0.60 milligrams a.i. per kernel) |   |
|            | Seed corn maggot (seed protection only),<br>Wireworm (seed protection only),<br>Flea beetle (through 1-leaf stage),<br>Imported fire ant,<br>White grub <sup>(4)</sup>   | 0.72<br>(Equivalent to 0.16 milligrams a.i. per kernel) |   |

**Field Corn (For end use application at agricultural establishments):**

Shake product well before making slurry and apply using a HCBT or Eight-Bag Batch Treater. May be diluted with an approved fungicide for disease control. Mix slurry so that rate is 8-10 fluid ounces per 50 pound bag of seed. Treat ½ of seed with ½ of slurry then add remainder of seed and slurry and continue mixing until all seed are covered. After slurry treatment, the addition of dry talc at the rate of ¾ -1 ounce per 50 pounds of seed is recommended to minimize risk moisture buildup in planter.



| Crop                                | Pest  | Rate (fl. oz./hundred weight seed) | Comment   |
|-------------------------------------|---|------------------------------------|---|
| Wheat, Barley, Oats, Rye, Triticale | Aphids (including Bird cherry-oat, English grain, Greenbug, and Russian wheat aphid), Hessian fly | 0.8 - 2.4                          | <p>Reduces possible yellow dwarf virus disease from aphid vectoring.</p> <p>Use the higher rate to increase time of protection and/or when insect pressure is heavy.</p>  |
|                                     | Wireworm  | 0.13 – 0.26                        | <p>Suppression only.</p> <p>Use the higher rate to increase time of protection and/or when insect pressure is heavy.</p>  |
|                                     | Grasshopper   | 1.2 – 2.4                          | <p>Treated seed may be planted as a 50- to 60-foot border around the perimeter of the field to help minimize early season damage from grasshoppers.</p> <p>Consult your local agricultural extension agent for grasshopper control information specific to your area.</p> <p>Use the higher rate to increase time of protection and/or when insect pressure is heavy.</p> |

**Wheat, Barley, Oats, Rye, Triticale (For End-Use Application at Agricultural Establishments):**

Make application using a Total Slurry Treater (TST), Farmer Applied Seed Treater (F.A.S.T.) or other on-farm seed treating equipment capable of accurately applying low rates of Sharda Imidacloprid 5SC. Apply 0.8 – 2.4 fl. oz. per hundredweight of seed. Sharda Imidacloprid 5SC may be mixed with a fungicide product for protection against diseases. Sharda Imidacloprid 5SC may be applied to seed previously treated with a fungicide. In this case, dilution is necessary.

Do not graze or feed livestock on treated areas for 45 days after planting.

| Crop    | Pest   | Rate (fl. oz. per hundred weight seed) | Comments  |
|---------|--|--|---|
| Sorghum | Aphids (including Corn leaf, English grain, Greenbug, and Yellow sugar cane aphid), Chinch bugs, Imported Fire ants, Wireworms | 6.4                                    | Thorough coverage is necessary for optimal control. |

**Sorghum (For End-Use Application at Agricultural Establishments):**

Shake product well before making slurry and apply using a HCBT or Eight-Bag Batch Treater. May be diluted with an approved fungicide for disease control. Mix slurry so that rate is 8-10 fl. oz. per 50 pound bag of seed. Treat 1/2 of seed with 1/2 of slurry then add remainder of seed and slurry and continue mixing until all seed are covered. After slurry treatment, the addition of dry talc at the rate of 3/4 - 1 ounce per 50 pounds of seed is recommended to minimize risk moisture buildup in planter.

Do not graze or feed livestock on treated areas for 45 days after planting.

| Crop                        | Pest            | Rate (fl. oz. per hundred weight seed)         | Comments   |
|-----------------------------|-----------------|--|--|
| Cotton (Delinted Seed Only) | Thrips & Aphids | 12.8 (Equivalent to 0.375 milligram a.i./seed) | Regardless of the type of application (seed treatment, soil, or foliar) do not apply more than a total of 0.5 lb. of imidacloprid/acre/cropping cycle. |

**Cotton (Delinted Seed Only) (For End-Use Application at Agricultural Establishments):**

Shake product well before making slurry and apply using a HCBT or Eight-Bag Batch Treater. May be diluted with an approved fungicide for disease control. Mix slurry so that rate is 8 - 10 fluid ounces per 50 pound bag of seed. Treat 1/2 of seed with 1/2 of slurry then add remainder of seed and slurry and continue mixing until all seed are covered.

| Crop                                       | Pest   | Rate (fl. oz. per hundred weight seed) <sup>(1)</sup> | Comments  |
|--|--|---|---|
| Sugar Beets<br>(commercial seed treatment) | Whitefly, Aphids, Leafhoppers <sup>(2)</sup> , Root aphid, Thrips, Wireworms | 2.4 – 5.0   | <p><sup>(1)</sup>On a unit of pelleted sugar beet seed with a weight ratio of 2:1 pelleting mixture to raw seed (seed count 100,000 seed – approximately 1 kilogram by weight).</p> <p><sup>(2)</sup>Including to reduce risk of spread of Curly top and Yellow mosaic virus.</p> <p>If applying to non-pelleted raw seed, do not exceed a rate of 2.4 fluid ounces per 100,000 seed (approximately 1Kg by weight).</p> |

| Crop                                   | Pest   | Rate (fl. oz. per hundred weight seed) | Comments   |
|--|--|--|--|
| Sweet Corn (commercial seed treatment) | Flea beetle, Early season corn leaf aphid, Seed corn maggot, Wireworm                  | 6.4                                    | Mix slurry so that it is applied at a rate of 16 – 20 fl. oz. of diluted solution per hundredweight of seed. |
|  | Imported fire ant, Early season corn leaf aphid, Seed corn maggot, Wireworm            | 3.2 – 6.4                              |  |
|  | Imported fire ant<br>Seed corn maggot (seed protection),<br>Wireworm (seed protection) | 1.6 – 3.2                              |  |
|  | Wireworm (seed protection)   | 0.8 – 1.6                              |  |
| Popcorn (commercial seed treatment)    | Flea beetles   | 6.4                                    |  |

| Crop    | Pest   | Rate (fl. oz. per hundred weight seed) | Comments   |
|---------|--|--|--|
| Soybean | Seed corn maggot,<br>Soybean aphids,<br>Over-wintering bean leaf beetles | 1.6 – 3.2                              | Use higher rates to provide increased length of protection and for heavy insect pressure.<br><br>Can be used as an over-treatment.<br><br>Do not graze or feed livestock on soybean forage or hay. |

| Crop  | Pest  | Rate (fl. oz. per hundred weight seed) | Comments   |
|---|---|--|--|
| Adzuki Bean,<br>Asparagus Bean,<br>Broad Bean (Succulent or Dry),<br>Catjang Bean, Chinese<br>Longbean, Field Bean, Guar<br>Bean, Jackbean, Kidney Bean,<br>Lablab Bean, Lima Bean<br>(Succulent or Dry),<br>Moth Bean (Succulent or Dry),<br>Mung Bean, Navy Bean, Pinto<br>Bean, Rice Bean, Runner Bean,<br>Snap Bean, Sword Bean,<br>Tepary Bean, Urd Bean, Wax<br>Bean, Yardlong Bean, Blackeye<br>Pea (Succulent or Dry),<br>Chickpea, Cowpea (Succulent<br>or Dry), Crowder Pea, Dwarf<br>Pea, Edible-Pod Pea, English<br>Pea, Field Pea, Garden Pea,<br>Green Pea, Pigeon Pea<br>(Succulent or Dry), Snow Pea,<br>Southern Pea (Succulent or<br>Dry), Sugar Snap Pea, Grain<br>Lupin, Sweet Lupin, White<br>Lupin, White Sweet Lupin,<br>Lentil* | Wireworm,<br>Bean leaf beetle,<br>Imported fire ant,<br>Aphid | 1.6 – 3.2                              | *Seed-and-pod vegetable seed treated in California must be destined for planting in states other than California and is not to be planted in California. |
| Carrot  | Seed corn maggot,<br>Wireworm                                 | 6.4                                    |  |

13 8 14

### ROTATIONAL CROPS

Treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient, as soon as practical following the last application. The following plant-back intervals should be observed for listed crops:

| Rotational Plant-Back Intervals*   |   |              |                    |   |  |
|--|---|--------------|--------------------|---|--|
| Immediate Plantback  |   |              |                    | 30-Day Plantback  |  |
| Artichoke  | Cranberry                               | Mustard Seed | Soybean            | Cereals, including:<br>buckwheat<br>Millet<br>Rice<br>Safflower |  |
| Barley   | Cucurbits                               | Oats         | Strawberry         |   |  |
| Brassica (cole)  | Eggplant                                | Okra         | Sugarbeet          |   |  |
| Leafy vegetable  | Ground cherry                           | Pepinos      | Tomatillo          |   |  |
| Canola   | Leaf petiole                            | Pepper       | Tomato             |   |  |
| Cilantro   | vegetable                               | Popcorn      | Triticvale         |   |  |
| Corn, Field  | Leafy vegetables                        | Potato       | Vegetable Root and |   |  |
| Corn, Sweet  | Legume vegetable                        | Rapeseed     | Tuber              |   |  |
| Cotton   | (succulent or dried,<br>except soybean) | Rye          | Watercress         |   |  |
|  |   | Sorghum      | Wheat              |   |  |
| *Cover crops for soil building or erosion control may be planted any time, but do not graze or harvest for food or feed.   |   |              |                    |   |  |
| *For all other crops not listed on an Imidacloprid label, or for crops for which no tolerances for the active ingredient has been established, a 12-month plantback interval should be observed. |   |              |                    |   |  |

Notification of the crop rotational restriction must be conveyed to the grower by appropriate seed tag labeling or bag printing on all seed units.

**NOTE:** The purchaser of this product is responsible for ensuring that all seed treated with this product are adequately dyed with a suitable color to prevent its accidental use as food for man or feed for animals. Refer to 21 CFR, Part 2.25. Any dye or colorant added to treated seed must be cleared for use under 40 CFR, Part 180.1001. Federal regulations have established official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates.

Treated seed must not be used for or mixed with food or animal feed or processed for oil. Seed commercially treated with Sharda Imidacloprid 5SC must be labeled in accordance with all applicable requirements of the Federal Seed Act.

Labels for commercially treated seed should include the following addition to the Environmental Hazards statements:

Exposed treated seed may be hazardous to birds. Dispose of all excess treated seed and seed packaging by burial away from bodies of water. Cover or incorporate spilled treated seeds.

#### Use Restriction

Rape greens grown and harvested from Sharda Imidacloprid 5SC treated seed must not be used for human and feed consumption. Rapeseed grown and harvested from treated seed is only for industrial uses and can not be used for edible oil or any other human/feed consumption.

**Wheat, Barley, Oats, Rye, Triticale, Sorghum:** Do not graze or feed livestock on treated areas for 45 days after planting.

**Cotton (delinted seed only):** Regardless of the type of application (seed treatment, soil, or foliar) do not apply more than a total of 0.5 lb. imidacloprid per acre per cropping cycle.

#### **Conditions of Sale and Limited Warranty Statement**

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Sharda USA LLC, no claims are made to guarantee germination of carry-over seed.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Sharda USA LLC. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW SHARDA USA LLC MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Sharda USA LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. SHARDA USA LLC DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT SHARDA USA LLC'S ELECTION, THE REPLACEMENT OF PRODUCT.