3529-14

0 10/05/200



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

OCT 0 5 2009

Ms. Cheryl Wagner. Warner Regulatory Associates, Inc. P.O. Box 640 7460 Lancaster Pike, Suite 9 Hockessin, Delaware 19707

Subject: Revised Storage and Disposal Label Language

Dear Ms. Wagner:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 dated September 8, 2009 for:

EPA Registration No. 83529-14

Sharda Imidacloprid 5SC

The Registration Division (RD) has conducted a review of this request for applicability under PRN 2007-4 and finds that the label change(s) requested falls within the scope of PRN-2007-4. The label has been date-stamped "Notification" and will be placed in our records.

If you have any questions, call me at 703 305-5409 or electronically at daniel.dani@epa.gov.

Sincerely,

Dani Daniel Registration Division (7505P) Insecticide/Rodenticide Branch

Please read instructions on reverse	United Vironmental P Washington	States OC rotection	FICATION T 0 5 2009 Agency	X ·	<u>red. OMB No. 20</u> Registration Amendme Other	on OPP Ide	al expires 05-31-5 ntifier Number
	Appli	cation for	Pesticide -	Sectio	n l	_	
1. Company/Product Number 83529-14			2. EPA Produc Venus Eagle			3. Proposed (Classification
4. Company/Product (Name) Sharda Imidacloprid 5SC			PM# 1	<u>-</u>	-	🗙 None	Restricte
 Name and Address of Applic Sharda USA LLC c/o Wagner Regulatory Ass P.O. Box 640 Hockessin, DE 19707 Check if a check if a check set of the check of				uct is simi		with FIFRA Sec in composition a	
		Sec	tion - II				
Armendment - Explain belo Resubmission in response Notification - Explain below Explanation: Use additional pa Notification of label change per PR 156.10, 154.140, 156.144, 156.146 I understand that it is a violation of consistent with the terms of PR Not	to Agency letter dated v. age(s) if necessary. Notice 2007-4. This no , and 156.156. No othe 18 U.S.C. Sec. 1001 to	(For Section I tification is cons r changes have willfully make a	Agency "Me Too IX Other - and Section II.) istent with the prov been made to the ny false statement	v letter date o" Applicati Explain be visions of F labeling or t to EPA. I t	ion. low. PR Notice 2007-4 the confidential s further understan	statement of formu	la of this product ation is not
penalties under Sections 12 and 14			tion - III				
Yes* X No	Be Packaged In: Unit Packaging Yes No If "Yes" Unit Packaging wgt.	No. per If container P	ackage wgt	ckaging No. per container		ontainer Metal Plastic Glass Paper Other (Specify) H	1DPE lined baç
Contain of Net Contents Info X Label Conta		4. Size(s) R 1 gallon	etail Container	5	Location of Location of Location of Location		ying product
6. Manner in Which Label is Af	fixed to Product	X Pap	ograph		Otheradhes	sive backed labe	91
	din all talents		tion - IV				
1. Contact Point (Complete iter Name	ns directly below for Title		oi individual to be	e contacte		v, to process this e No. (Include A	
CHERYL WAGNER I certify that the statements I have I acknowledge that any knowingly both under applicable law.	Certifi made on this form and	Agent for SI cation I all attachments tement may be			(302) 234	8551 د ب 6. Date Applicat د Received د ز ک (Stan	ion ເ ກຸped)
2. Signature	Drema	3. Title Agent for S	harda USA LLC			((ι) ι(ι) () ι (ι) (ι) (ι) (ι) (ι)	ι ι
- mup (S S	ίτι τζί	(

(10000m) MUUUU || Nr

Wagner Regulatory Associates, Inc. P.O. Box 640 7460 Lancaster Pike, Suite 9 Hockessin, Delaware 19707

September 8, 2009

Document Processing Desk (NOTIF) ATTN: Ms. Venus Eagle, PM 1 Registration Division (7504P) U.S. Environmental Protection Agency Room S-4900, One Potomac Yard 2777 South Crystal Drive Arlington, Virginia 22202-4501

(i

Dear Ms. Eagle:

Re: Sharda Imidacloprid 5SC EPA Registration Number 83529-14 Notice of Revised Storage & Disposal Label Language

Wagner Regulatory Associates, Inc., on behalf of Sharda USA LLC, hereby notifies the Agency that the storage and disposal section of the subject label as been revised in accordance with PR Notice 2007-4. Enclosed for the Agency's file is:

- Letter from Sharda USA LLC authorizing Wagner Regulatory to serve as Agent
- EPA Notification form (EPA Form 8570-1)
- One copy of revised labeling with the revised text bolded for convenience.

Please feel free to contact me at (302) 234-8551 if you have any questions or require additional information.

Respectfully submitted,

of wagne

Cheryl Wagner Agent for Sharda USA LLC

.....

Sharda Imidacloprid 5SC

OCT 0 5 2009

YIIU

Active Ingredient:	
Imidacloprid 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine	48.7%
Other Ingredients:	51.3%
5	100.0%

Contains 5 lbs. active ingredient per gallon (600 grams per liter).

Ć

Net Contents: 1 gallon EPA Reg. No. 83529-14 EPA Est. No. _____

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detaile. (If you do not understand the label, find someone to explain it to you in detail).

· · · · · · · · · · · · · · · · · · ·	FIRST AID				
If swallowed	• Call a poison control center or doctor immediately for				
	treatment advice.				
	• Have person sip a glass of water if able to swallow.				
	• Do not induce vomiting unless told to do so by a poison				
	control center or doctor.				
	• Do not give anything by mouth to an unconscious person.				
If on skin or clothin	ng • Take off contaminated clothing.				
	• Rinse skin immediately with plenty of water for 15-20 minutes.				
	• Call a poison control center or doctor for treatment advice.				
If inhaled	• Move person to fresh air.				
	• If person is not breathing, call 911 or an ambulance, then give				
	artificial respiration, preferably mouth-to-mouth if possible.				
	• Call a poison control center or doctor for further treatment				
······································	advice.				
NOTE TO PHYSIC symptomatically.	CIAN: No specific antidote is available. Treat the patient				
Have the product con	ntainer or label with you when calling a poison control center or				
doctor, or going for	treatment.				
For 24-hour medical	emergency assistance (human or animal) call 1-800-222-1222. For				
• •	v assistance (spill, leak, fire, or accident) call ChemTrec at 1-800-424-				
9300.	<u>c</u>				
SEE INSIDE BOO	DKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS				
	د در د در در د در در در در در				
	Manufactured for:				
	Sharda USA LLC				
	7460 Lancaster Pike, Suite 9				

Hockessin, DE 19707

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution. Harmful if swallowed, absorbed through skin or inhaled. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and was before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for Category C on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- 1. Long-sleeved shirt and long pants,
- 2. Socks and shoes, and
- 3. Chemical-resistant gloves (such as nitrile, butyl, neoprene, barrier laminate, polyvinyl chloride or Viton).

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately, from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- 1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- 2. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- 3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change clothing.

ENVIRONMENTAL HAZARDS

This product is highly toxic to birds and aquatic invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Cover or incorporate spilled treated seeds.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For use in commercial seed treaters only, with the exception of application to canola, cotton (delinted seed), field corn, sorghum, wheat and barley, which may be made either by commercial seed treatment or as an enduse seed treatment on agricultural establishments at, or immediately before, planting. This product is to be used in liquid or slurry treaters.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Mix thoroughly before use or use entire container at one time. All tank mixes should be pre-tested to determine physical compatibility between formulations. Observe all cautions and limitations on labeling of all products used in mixtures.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Exception. If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- 1. Coveralls
- 2. Chemical-resistant gloves (such as nitrile, butyl, neoprene, barrier laminate, polyvinyl chloride, or Viton)
- 3. Shoes plus socks
- 4. Protective eyewear.

STORAGE AND DISPOSAL

Prohibitions: Do not contaminate water, food or feed by storage, disposal, or by cleaning of equipment. Do not reuse empty container. Open dumping is prohibited.

Pesticide Storage: Store in a cool place. Do not store in direct sunlight. Protect from freezing temperatures.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Nonrefillable container. Do not refill or reuse container. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incideration; Gr, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Control of the contro

STORED SEED PROTECTION

,

To protect stored seed of crops listed on this label apply a minimum of 0.8 fluid ounces of Sharda Imidacloprid 5SC per 100 pounds of seed. Pests controlled by this application include Indian Meal Moth (*Plodia interpunctella*), Rice Weevil (*Sitophilus oryzea*), Red Flour Beetle (*Tribiolium castaneum*), and Lesser Grain Borer (*Rhizopertha dominica*). If seed being treated is known to have existing pest infestation fumigation may be necessary prior to bagging.

Сгор	Pest	Rate (fl. oz./hundred weight seed)	Comment
Canola, Rapeseed and Mustard Seed	Aphids, Flea beetles, Wireworms	10.24 – 25.6	Use higher rate when pest populations are above historical levels.
	Lygus Bugs (including Lygus spp.) Seedpod Weevil Larvae (including Ceutorhynchus assimilis)	15.36 – 25.6	Suppression only. Use higher rate when pest populations are above historical levels.

EARLY SEASON PROTECTION AGAINST SUCKING INSECTS

Canola, Rapeseed and Mustard Seed (For end use application at agricultural establishments):

Shake product well before making slurry and apply using a HCBT or Batch Treater. Apply 5.1 to 12.8 fl. oz. of Sharda Imidacloprid 5SC per 50 pound bag. May be diluted with an approved fungicide for disease control. Treat ½ of seed with ½ of slurry then add remainder of seed and slurry and continue mixing until all seed are covered.

..... Draft Sharda İmidacloprid 5SC September 4, 2009 (CC) Page 4 of 11

Сгор	Pest	Rate (Fl. oz. per 80,000 units of seed)	Comment
Field Corn	Corn root worm (including Northern, Western, Southern, and Mexican) ⁽¹⁾ , Flea beetle, Chinch bug, Southern green stinkbug, White grub, Seed corn maggot, Thrips, Wireworm, Corn leaf aphid, Imported fire ant, Southern corn leaf beetle, Billbug ⁽²⁾ , Grape colaspis ⁽²⁾ , Black cutworm ⁽³⁾	6.0 (Equivalent to 1.34 milligrams a.i. per kernel)	 ⁽¹⁾In areas of heavy to severe corn rootworm populations, protection will not be adequate. Use only in areas of light to moderate corn root-worm populations. Consult your State Agricultural Extension Service on levels of corn rootworm populations. ⁽²⁾Reduces early season feeding damage. ⁽³⁾Will reduce feeding damage caused by leaf feeding black cutworms that are ½ inch or less in length. ⁽⁴⁾Reduces feeding damage during emergence and seedling stages.
	Flea beetle, Chinch bug, Seed corn maggot, Thrips, Wireworm, Corn leaf aphid Imported fire ant, Grape colaspis ⁽²⁾ , White grub Seed corn maggot (seed protection only), Wireworm (seed protection only), Flea beetle (through 1-leaf stage), Imported fire ant, White grub ⁽⁴⁾	2.70 (Equivalent to 0.60 milligrams a.i. per kernel) 0.72 (Equivalent to 0.16 milligrams a.i. per kernel)	

Field Corn (For end use application at agricultural establishments):

.

Shake product well before making slurry and apply using a HCBT or Eight-Bag Batch Treater. May be diluted with an approved fungicide for disease control. Mix slurry so that rate is 8-10 fluid ounces per 50 pound bag of seed. Treat ½ of seed with ½ of slurry then add remainder of seed and slurry and continue mixing until all seed are covered. After slurry treatment, the addition of dry talc at the rate of ¾ -1 ounce per 50 pounds of seed is recommended to minimize risk moisture buildup in planter.

8/14

Сгор	Pest	Rate (fl. oz./hundred weight seed)	Comment
Wheat, Barley, Oats, Rye, Triticale	Aphids (including Bird cherry-oat, English grain, Greenbug, and Russian wheat aphid), Hessian fly	0.8 - 2.4	Reduces possible yellow dwarf virus disease from aphid vectoring. Use the higher rate to increase time of protection and/or when insect pressure is heavy.
	Wireworm	0.13 - 0.26	Suppression only. Use the higher rate to increase time of protection and/or when insect pressure is heavy.
	Grasshopper	1.2 - 2.4	Treated seed may be planted as a 50- to 60-foot border around the perimeter of the field to help minimize early season damage from grasshoppers.
			Consult your local agricultural extension agent for grasshopper control information specific to your area.
			Use the higher rate to increase time of protection and/or when insect pressure is heavy.

(

Wheat, Barley, Oats, Rye, Triticale (For End-Use Application at Agricultural Establishments):

Make application using a Total Slurry Treater (TST), Farmer Applied Seed Treater (F.A.S.T.) or other on-farm seed treating equipment capable of accurately applying low rates of Sharda Imidacloprid 5SC. Apply 0.8 - 2.4 fl. oz. per hundredweight of seed. Sharda Imidacloprid 5SC may be mixed with a fungicide product for protection against diseases. Sharda Imidacloprid 5SC may be applied to seed previously treated with a fungicide. In this case, dilution is necessary.

Do not graze or feed livestock on treated areas for 45 days after planting.

Crop	Pest	Rate (fl. oz. per hundred weight seed)	Comments
Sorghum	Aphids (including Corn leaf, English grain, Greenbug, and Yellow sugar cane aphid), Chinch bugs, Imported Fire ants, Wireworms	6.4	Thorough coverage is necessary for optimal control.

Sorghum (For End-Use Application at Agricultural Establishments):

Shake product well before making slurry and apply using a HCBT or Eight-Bag Batch Treater. May be diluted with an approved fungicide for disease control. Mix slurry so that rate is 8-10 fl. oz. per 50 pound bag of seed. Treat $\frac{1}{2}$ of seed with $\frac{1}{2}$ of slurry then add remainder of seed and slurry and continue mixing until all seed are covered. After slurry treatment, the addition of dry talc at the rate of $\frac{3}{4}$ - 1 ounce per 50 pounds of seed is recommended to minimize risk moisture buildup in planter.

Do not graze or feed livestock on treated areas for 45 days after planting.

Сгор	Pest	Rate (fl. oz. per hundred weight seed)	Comments
Cotton (Delinted Seed Only)	Thrips & Aphids	12.8 (Equivalent to 0.375 milligram a.i./seed	Regardless of the type of application (seed treatment, soil, or foliar) do not apply more than a total of 0.5 lb. of imidacloprid/acre/cropping cycle.

Cotton (Delinted Seed Only) (For End-Use Application at Agricultural Establishments):

Shake product well before making slurry and apply using a HCBT or Eight-Bag Batch Treater. May be diluted with an approved fungicide for disease control. Mix slurry so that rate is 8 - 10 fluid ounces per 50 pound bag of seed. Treat $\frac{1}{2}$ of seed with $\frac{1}{2}$ of slurry then add remainder of seed and slurry and continue mixing until all seed are covered.

Сгор	Pest	Rate (fl. oz. per hundred weight seed) ⁽¹⁾	Comments
Sugar Beets (commercial seed treatment)	Whitefly, Aphids, Leafhoppers ⁽²⁾ , Root aphid, Thrips, Wireworms	2.4 – 5.0	⁽¹⁾ On a unit of pelleted sugar beet seed with a weight ratio of 2:1 pelleting mixture to raw seed (seed count 100,000
			seed – approximately 1 kilogram by weight).
			⁽²⁾ Including to reduce risk of spread of Curly top and Yellow mosaic virus.
			If applying to non- pelleted raw seed, do not exceed a rate of 2.4 fluid
			ounces per 100,000 seed (approximately 1Kg by weight).

(

•

.

(

.

Сгор	Pest	Rate (fl. oz. per hundred weight seed)	Comments
Sweet Corn (commercial seed treatment)	Flea beetle, Early season corn leaf aphid, Seed corn maggot, Wireworm	6.4	Mix slurry so that it is applied at a rate of 16 – 20 fl. oz. of diluted solution per
	Imported fire ant, Early season corn leaf aphid, Seed corn maggot, Wireworm	3.2 - 6.4	hundredweight of seed.
	Imported fire ant Seed corn maggot (seed protection), Wireworm (seed protection)	1.6 - 3.2	
	Wireworm (seed protection)	0.8 - 1.6	
Popcorn (commercial seed treatment)	Flea beetles	6.4	<u> </u>

N/M

Сгор	Pest	Rate (fl. oz. per hundred weight seed)	Comments
Soybean	Seed corn maggot, Soybean aphids, Over-wintering bean leaf beetles	1.6 - 3.2	Use higher rates to provide increased length of protection and for heavy insect pressure.
			Can be used as an over- treatment.
			Do not graze or feed livestock on soybean forage or hay.

(

(

· ·

		Rate (fl. oz. per hundred	
Сгор	Pest	weight seed)	Comments
Adzuki Bean,	Wireworm,	1.6 – 3.2	*Seed-and-pod vegetable
Asparagus Bean,	Bean leaf beetle,		seed treated in California
Broad Bean (Succulant or Dry),	Imported fire ant,		must be destined for
Catjang Bean, Chinese	Aphid		planting in states other
Longbean, Field Bean, Guar			than California and is not
Bean, Jackbean, Kidney Bean,			to be planted in
Lablab Bean, Lima Bean			California.
(Succulent or Dry),			
Moth Bean (Succulent or Dry),			
Mung Bean, Navy Bean, Pinto			
Bean, Rice Bean, Runner Bean,			
Snap Bean, Sword Bean,			
Tepary Bean, Urd Bean, Wax			
Bean, Yardlong Bean, Blackeye			
Pea (Succulent or Dry),			
Chickpea, Cowpea (Succulent			
or Dry), Crowder Pea, Dwarf			
Pea, Edible-Pod Pea, English			
Pea, Field Pea, Garden Pea,			
Green Pea, Pigeon Pea			
(Succulent or Dry), Snow Pea,			
Southern Pea (Succulent or			
Dry), Sugar Snap Pea, Grain			
Lupin, Sweet Lupin, White			
Lupin, White Sweet Lupin,			
Lentil*			
Carrot	Seed corn maggot,	6.4	
	Wireworm		

12/14

ROTATIONAL CROPS

Treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient, as soon as practical following the last application. The following plant-back intervals must be observed for listed crops:

Rotational Plant-Back Intervals*				
Immediate Plant-back				30-Day Plant-back
Artichoke	Cranberry	Mustard Seed	Soybean	Cereals, including:
Barley	Cucurbits	Oats	Strawberry	Buckwheat
Brassica (cole)	Eggplant	Okra	Sugarbeet	Millet
Leafy vegetable	Ground cherry	Pepinos	Tomatillo	Rice
Canola	Leaf petiole	Pepper	Tomato	Safflower
Cilantro	vegetable	Popcorn	Triticale	
Corn, Field	Leafy vegetables	Potato	Vegetable Root and	
Corn, Sweet	Legume vegetable	Rapeseed	Tuber	
Cotton	(succulent or dried,	Rye	Watercress	
	except soybean)	Sorghum	Wheat	

*Cover crops for soil building or erosion control may be planted any time, but do not graze or harvest for food or feed.

*For all other crops not listed on an Imidacloprid label, or for crops for which no tolerances for the active ingredient has been established, a 12-month plant-back interval must be observed.

Notification of the crop rotational restriction must be conveyed to the grower by appropriate seed tag labeling or bag printing on all seed units.

NOTE: The purchaser of this product is responsible for ensuring that all seed treated with this product are adequately dyed with a suitable color to prevent its accidental use as food for man or feed for animals. Refer to 21 CFR, Part 2.25. Any dye or colorant added to treated seed must be cleared for use under 40 CFR, Part 180.1001. Federal regulations have established official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates.

Treated seed must not be used for or mixed with food or animal feed or processed for oil. Seed commercially treated with Sharda Imidacloprid 5SC must be labeled in accordance with all applicable requirements of the Federal Seed Act.

Labels for commercially treated seed must include the following addition to the Environmental Hazards statements:

Exposed treated seed may be hazardous to birds. Dispose of all excess treated seed and seed packaging by burial away from bodies of water. Cover or incorporate spilled treated seeds.

Use Restriction

Rape greens grown and harvested from Sharda Imidacloprid 5SC treated seed must not be used for human and feed consumption. Rapeseed grown and harvested from treated seed is only for industrial uses and must not be used for edible oil or any other human/feed consumption.

Wheat, Barley, Oats, Rye, Triticale, Sorghum: Do not graze or feed livestock on treated areas for 45 days after planting.

Cotton (delinted seed only): Regardless of the type of application (seed treatment, soil, or foliar) do not apply more than a total of 0.5 lb. imidacloprid per acre per cropping cycle.

Conditions of Sale and Limited Warranty Statement

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Sharda USA LLC, no claims are made to guarantee germination of carry-over seed.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Sharda USA LLC. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW SHARDA USA LLC MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Sharda USA LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SHARDA USA LLC DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT SHARDA USA LLC'S ELECTION, THE REPLACEMENT OF PRODUCT.

Draft Sharda Imidacloprid 5SC September 4, 2009 Page 11 of 11