



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVETION

APR 29 2010

Ms. Catherine M. Byrd Sharda USA LLC c/o Wagner Regulatory Associates, Inc. P.O. Box 640 Hockessin, DE 19707

Subject: Revised Storage and Disposal Language

Dear Ms. Byrd:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 dated April 9, 2010 for:

EPA Registration 83529-14

Sharda Imidacloprid 5SC

The Registration Division (RD) has conducted a review of this request for applicability under PRN 2007-4 and finds that the label change(s) requested falls within the scope of PRN-2007-4. The label has been date-stamped "Notification" and will be placed in our records.

If you have any questions, call me at 703 305-5409 or electronically at daniel.dani@epa.gov.

Sincerely,

Dani Daniel

Registration Division (7504P)

Insecticide/Rodenticide Branch

Please read instructions on revers	e before comple/form	7.	IOTIFICATT	Approv	Registration		Approval expires 05-31-98 PP Identifier Number
	United				Negistiati		
SEPA E	nvironmental Pı		APR 2 9 2010 Agency		Amendme	nt	
	Washington			Χ	Other		
	Appli	cation for	Pesticide - Se	ction	n I		
1. Company/Product Number			2. EPA Product M	lanage	er	3. Propo	osed Classification
83529-14			Venus Eagle				
4. Company/Product (Name)		<u>.</u>	PM#			 ⊠ No	ne Restricted
Sharda Imidacloprid 5SC			1				
5. Name and Address of Appli	icant (Include Zip Cod	le)	6. Expedited Rev				
Sharda USA LLC			to:	13 31111	iai oi identicai	iii compos	stion and labeling
c/o Wagner Regulatory As P.O. Box 640	sociates, Inc.		EPA Reg. No.				
Hockessin, DE 19707			Product Name				
· —	f this is a new address	3		•			
			ction - II				
Amendment - Explain be	low.		Final printe		s in response to	<u> </u>	
Resubmission in respons	se to Agency letter dated	٠,	Agency lett _				
Notification - Explain belo			Other - Exp				
Explanation: Use additional p	nage(s) if necessary	(For Section	I and Section II \				
Notification of label change per PF	R Notice 2007-4. This no	tification is cons	sistent with the provision	ns of P	R Notice 2007-4	and EPA R	Regulations at 40 CFR
156.10, 154.140, 156.144, 156.146 I understand that it is a violation of							
consistent with the terms of PR No							
and penalties under Sections 12 a	nd 14 of FIFRA.						
		Sec	ction - III				
1. Material This Product Will					10 = 10		
Child-Resistant Packaging Yes*	Unit Packaging Yes	l v	Water Soluble Packa Yes	iging	2. Type of C	ontainer Metal	
	 	:	\vdash		! .	vietai Plastic	
X No	X No	<u> </u>	X No				
* 0466464	If "Yes"	'. l		per	l ——	Glass	i
* Certification must	Unit Packaging wgt.	container F	Package wgt con	tainer	l	⊃aper Other (Spe	osify UDDE lined have
be submitted					\	omer (Spe	ecify) HDPE lined bags
3. Location of Net Contents In	formation	4. Size(s) R	Retail Container	5.	Location of L	abel Direc	tions
X Label Con	tainer	1 gallon, bul	lk	- 1	X On Lab	el	
					On Lab	eling acco	ompanying product
6. Manner in Which Label is A	Affixed to Product	Lith	nograph	7 (Other adhe	sive backe	ed label
			per glued		 ,		0 0 0
		Ste	enciled				0000 0000
		Sec	ction - IV				0000
1. Contact Point (Complete items directly below for identification of individual to be contacted, if neces&ary of corrects this application.)							
Name	Title				Telephon	e No. (Incl	lude Area Code)
Catherine M. Byrd Agent for Sharda USA LLC (302) 238 2780							
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. ိုင္ငံ Received							
I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or						Č	
both under applicable law. (Stamped)					(Stamped)		
2. Signature	Jan Dund	3. Title					00CL
Cathww	whyd	Agent for S	Sharda USA LLC				
4. Typed Name	•	5. Date					
Catherine M. Byrd		April 9, 201	10		1		

JOHN WRA

April 9, 2010

Document Processing Desk (NOTIF) ATTN: Ms. Venus Eagle, PM 1 Registration Division (7505P) Office of Pesticide Programs U.S. Environmental Protection Agency Room S-4900, One Potomac Yard 2777 South Crystal Drive Arlington, Virginia 22202-4501 Wagner Regulatory Associates, Inc. P.O. Box 640 7460 Lancaster Pike, Suite 9 Hockessin, Delaware 19707

Subject:

Sharda Imidacloprid 5SC

EPA Reg. No. 83529-14

Revised Storage and Disposal Language Notification per PR Notice 2007-4

Dear Ms. Eagle,

On behalf of Sharda USA LLC, I am submitting this notification of revised storage and disposal language per PR Notice 2007-4 for the above referenced product. No additional changes were made to the label.

In support of this notification I have enclosed the following documents:

- Letter from Sharda USA LLC authorizing Wagner Regulatory to serve as Agent;
- Application for Pesticide Registration (EPA Form 8570-1); and
- One copy of the revised label with changes clearly marked.

If you have any questions concerning this submission please contact the undersigned.

Sincerely,

Catherine M. Byrd

Agent for Sharda USA LLC

Cotherine M. Byrd

enclosures

83529-15-V3-9APR10

Sharda Imidacloprid 5SC

Revised Storage and Disposal (Large and Small Containers)

= - LVIQ	
Active Ingredient:	
Imidacloprid 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine	48.7%
Other Ingredients:	51.3%
Total	100.0%
Contains 5 lbs. active ingredient per gallon (600 grams per liter).	

Net Contents: 1 gallon, Bulk EPA Reg. No. 83529-14 EPA Est. No.

KEEP OUT OF REACH OF CHILDREN **CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detaile. (If you do not understand the label, find someone to explain it to you in detail).

	FIRST AID			
If swallowed	Call a poison control center or doctor immediately for treatment advice.			
	Have person sip a glass of water if able to swallow.			
	• Do not induce vomiting unless told to do so by a poison control center or doctor.			
	• Do not give anything by mouth to an unconscious person.			
If on skin or clothing	Take off contaminated clothing.			
	• Rinse skin immediately with plenty of water for 15-20 minutes.			
	• Call a poison control center or doctor for treatment advice.			
If inhaled	Move person to fresh air.			
	• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.			
	Call a poison control center or doctor for further treatment advice.			
NOTE TO PHYSICIA symptomatically.	N: No specific antidote is available. Treat the patient			
Have the product contain	ner or label with you when calling a poison control center or			
doctor, or going for treat	tment.			
	ergency assistance (human or animal) call 1-800-222-1222. For co			
	istance (spill, leak, fire, or accident) call ChemTrec at 1-800-424-			
9300.	T)			
SEE INSIDE BOOKL	ET FOR ADDITIONAL PRECAUTIONARY STATEMEN FS			

Manufactured for:

Sharda USA LLC 7460 Lancaster Pike, Suite 9 Hockessin, DE 19707

> Sharda Imidacloprid 5SC April 9, 2010 Page 1 of 11

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution. Harmful if swallowed, absorbed through skin or inhaled. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and was before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for Category C on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- 1. Long-sleeved shirt and long pants,
- 2. Socks and shoes, and
- 3. Chemical-resistant gloves (such as nitrile, butyl, neoprene, barrier laminate, polyvinyl chloride or Viton).

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately, from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- 1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- 2. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- 3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change clothing.

ENVIRONMENTAL HAZARDS

This product is highly toxic to birds and aquatic invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Cover or incorporate spilled treated seeds.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For use in commercial seed treaters only, with the exception of application to canola, cotton (delinted seed), field corn, sorghum, wheat and barley, which may be made either by commercial seed treatment or as an enduse seed treatment on agricultural establishments at, or immediately before, planting. This product is to be used in liquid or slurry treaters.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Mix thoroughly before use or use entire container at one time. All tank mixes should be pre-tested to determine physical compatibility between formulations. Observe all cautions and limitations on labeling of all products used in mixtures.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Exception. If the seed is treated with the product and the treated seed is soil-injected or soilincorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- 1. Coveralls
- 2. Chemical-resistant gloves (such as nitrile, butyl, neoprene, barrier laminate, polyvinyl chloride, or Viton)
- 3. Shoes plus socks
- 4. Protective eyewear.

STORAGE AND DISPOSAL

Prohibitions: Do not contaminate water, food or feed by storage, disposal, or by cleaning of equipment. Do not reuse empty container. Open dumping is prohibited.

Pesticide Storage: Store in a cool place. Do not store in direct sunlight. Protect from freezing temperatures.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal:

NONREFULABLE CONTAINER (EQUAL TO OR LESS THAN 5 GALLONS): Donot reuse container. Intiple tinse container (or equivalent) promptly after emptying. Triple tinse as follows: Empty the remaining contents into application equipment or a mixtank and dram for 10 seconds after the flow begins to drip. Fill the container of a mixtank and dram for 10 seconds after the flow begins to drip. Fill the container of a mixtank or store full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mixtank or store tinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure; two more times. Ofter for recycling, if available.

NONREFILE ABLE CONTAINER (GREATER THAN 5 GALLONS). Donot reuse container. Briple tinse

container (for equivalent) promptly after emptying. Triple tinse as follows: Empty the renativing contents into application equipment of atmix tank. Fill the container & full with water. Replace and tighten elosures; till container on its side and foll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Furn the container over onto its other one and tip it back and forth several times. Furn the container over onto its other one and tip it back and forth several times. Empty the disate into application equipment or a mix tank or store tinsate for later use or disposal.

Repeat this procedure two more times. Offer for necycling, if available.

STORED SEED PROTECTION


To protect stored seed of crops listed on this label apply a minimum of 0.8 fluid ounces of Sharda Imidacloprid 5SC per 100 pounds of seed. Pests controlled by this application include Indian Meal Moth (*Plodia interpunctella*), Rice Weevil (*Sitophilus oryzea*), Red Flour Beetle (*Tribiolium castaneum*), and Lesser Grain Borer (*Rhizopertha dominica*). If seed being treated is known to have existing pest infestation fumigation may be necessary prior to bagging.

EARLY SEASON PROTECTION AGAINST SUCKING INSECTS

Crop	Pest	Rate (fl.	Comment
		oz./hundred weight	
		seed)	
Canola, Rapeseed	Aphids, Flea	10.24 – 25.6	Use higher rate when pest
and Mustard Seed	beetles, Wireworms		populations are above historical
			levels.
	Lygus Bugs (including Lygus spp.) Seedpod Weevil Larvae (including Ceutorhynchus	15.36 – 25.6	Suppression only. Use higher rate when pest populations are above historical levels.
	assimilis)		

Canola, Rapeseed and Mustard Seed (For end use application at agricultural establishments):

Shake product well before making slurry and apply using a HCBT or Batch Treater. Apply 5.1 to 12.8 fl. oz. of Sharda Imidacloprid 5SC per 50 pound bag. May be diluted with an approved fungicide for disease control. Treat ½ of seed with ½ of slurry then add remainder of seed and slurry and continue mixing until all seed are covered.



Crop	Pest	Rate (Fl. oz. per 80,000 units of seed)	Comment
Field Corn	Corn root worm (including Northern, Western, Southern, and Mexican) ⁽¹⁾ , Flea beetle, Chinch bug, Southern green stinkbug, White grub, Seed corn maggot, Thrips, Wireworm, Corn leaf aphid, Imported fire ant, Southern corn leaf beetle, Billbug ⁽²⁾ , Grape colaspis ⁽²⁾ , Black cutworm ⁽³⁾	6.0 (Equivalent to 1.34 milligrams a.i. per kernel)	(1) In areas of heavy to severe corn rootworm populations, protection will not be adequate. Use only in areas of light to moderate corn root-worm populations. Consult your State Agricultural Extension Service on levels of corn rootworm populations. (2) Reduces early season feeding damage. (3) Will reduce feeding damage caused by leaf feeding black cutworms that are ½ inch or less in length. (4) Reduces feeding damage during emergence and seedling stages.
	Flea beetle, Chinch bug, Seed corn maggot, Thrips, Wireworm, Corn leaf aphid Imported fire ant, Grape colaspis ⁽²⁾ , White grub	2.70 (Equivalent to 0.60 milligrams a.i. per kernel)	
	Seed corn maggot (seed protection only), Wireworm (seed protection only), Flea beetle (through 1-leaf stage), Imported fire ant, White grub ⁽⁴⁾	0.72 (Equivalent to 0.16 milligrams a.i. per kernel)	, , , ,

Field Corn (For end use application at agricultural establishments):

Shake product well before making slurry and apply using a HCBT or Eight-Bag Batch Treater. May be diluted with an approved fungicide for disease control. Mix slurry so that rate is 8-10 fluid ounce of 50 pound bag of seed. Treat ½ of seed with ½ of slurry then add remainder of seed and slurry and continue mixing until all seed are covered. After slurry treatment, the addition of dry talc at the rate of ¾ -1 ounce per 50 pounds of seed is recommended to minimize risk moisture buildup in planter.

		6600
	Rate (fl	,,,,,
·	Nate (II.	

0000

0220


83529-15-V3-9APR10

Revised Storage and Disposal (Large and Small Containers)

Crop	Pest	oz./hundred weight	Comment
Wheat, Barley, Oats, Rye, Triticale	Aphids (including Bird	seed) 0.8 - 2.4	Reduces possible yellow dwarf virus disease from aphid vectoring.
	cherry-oat, English grain, Greenbug, and Russian wheat aphid), Hessian fly		Use the higher rate to increase time of protection and/or when insect pressure is heavy.
	Wireworm	0.13 - 0.26	Suppression only.
			Use the higher rate to increase time of protection and/or when insect pressure is heavy.
	Grasshopper	1.2 – 2.4	Treated seed may be planted as a 50-to 60-foot border around the perimeter of the field to help minimize early season damage from grasshoppers.
			Consult your local agricultural extension agent for grasshopper control information specific to your area.
			Use the higher rate to increase time of protection and/or when insect pressure is heavy.

Wheat, Barley, Oats, Rye, Triticale (For End-Use Application at Agricultural Establishments):

Make application using a Total Slurry Treater (TST), Farmer Applied Seed Treater (F.A.S.T.) or other on-farm seed treating equipment capable of accurately applying low rates of Sharda Imidacloprid 5SC. Apply 0.8-2.4 fl. oz. per hundredweight of seed. Sharda Imidacloprid 5SC may be mixed with a fungicide product for protection against diseases. Sharda Imidacloprid 5SC may be applied to seed previously treated with a fungicide. In this case, dilution is necessary.



Sharda Imidacloprid 5SC April 9, 2010 Page 6 of 11 Crop Pest Rate (fl. oz. per hundred weight seed) Comments

Sorghum Aphids (including Corn leaf, English grain, Greenbug, and Yellow sugar cane aphid), Chinch Greenbug, and Yellow sugar cane aphid), Chinch

Do not graze or feed livestock on treated areas for 45 days after planting.

Sorghum (For End-Use Application at Agricultural Establishments):

Wireworms

bugs, Imported Fire ants.


Shake product well before making slurry and apply using a HCBT or Eight-Bag Batch Treater. May be diluted with an approved fungicide for disease control. Mix slurry so that rate is 8-10 fl. oz. per 50 pound bag of seed. Treat ½ of seed with ½ of slurry then add remainder of seed and slurry and continue mixing until all seed are covered. After slurry treatment, the addition of dry talc at the rate of ¾ - 1 ounce per 50 pounds of seed is recommended to minimize risk moisture buildup in planter.

Do not graze or feed livestock on treated areas for 45 days after planting.

Crop	Pest	Rate (fl. oz. per hundred weight seed)	Comments
Cotton (Delinted Seed Only)	Thrips & Aphids	12.8 (Equivalent to 0.375 milligram a.i./seed	Regardless of the type of application (seed treatment, soil, or foliar) do not apply more than a total of 0.5 lb. of imidacloprid/acre/cropping cycle.

Cotton (Delinted Seed Only) (For End-Use Application at Agricultural Establishments):

Shake product well before making slurry and apply using a HCBT or Eight-Bag Batch Treater. May be diluted with an approved fungicide for disease control. Mix slurry so that rate is 8 – 10 fluid ounces per 50 pound bag of seed. Treat ½ of seed with ½ of slurry then add remainder of seed and slurry and continue mixing until all seed are covered.



		Rate (fl. oz. per hundred	
Crop	Pest	weight seed) ⁽¹⁾	Comments
Sugar Beets	Whitefly,	2.4 - 5.0	(1)On a unit of pelleted
(commercial seed	Aphids,		sugar beet seed with a
treatment)	Leafhoppers ⁽²⁾ , Root		weight ratio of 2:1
	aphid, Thrips, Wireworms		pelleting mixture to raw
			seed (seed count 100,000
,			seed – approximately 1
			kilogram by weight).
			(2)Including to reduce risk of spread of Curly top and Yellow mosaic virus.
			If applying to non-
			pelleted raw seed, do not
			exceed a rate of 2.4 fluid
			ounces per 100,000 seed
			(approximately 1Kg by
			weight).

Crop	Pest	Rate (fl. oz. per hundred weight seed)	Comments
Sweet Corn (commercial seed treatment)	Flea beetle, Early season corn leaf aphid, Seed corn maggot, Wireworm	6.4	Mix slurry so that it is applied at a rate of 16 – 20 fl. oz. of diluted solution per
	Imported fire ant, Early season corn leaf aphid, Seed corn maggot, Wireworm	3.2 – 6.4	hundredweight of seed.
	Imported fire ant Seed corn maggot (seed protection), Wireworm (seed protection)	1.6 – 3.2	0 (C O O O O O O O O O O O O O O O O O O
	Wireworm (seed protection)	0.8 – 1.6	000000 0 0 0 0 0 0 0 0 0 0 0 0 0
Popcorn (commercial seed treatment)	Flea beetles	6.4	00000 0000

Pest	Rate (fl. oz. per hundred weight seed)	Comments
Seed corn maggot, Soybean aphids, Over-wintering bean leaf beetles	1.6 – 3.2	Use higher rates to provide increased length of protection and for heavy insect pressure. Can be used as an overtreatment. Do not graze or feed livestock on soybean forage or hay.
	Seed corn maggot, Soybean aphids, Over-wintering bean leaf	Pest weight seed) Seed corn maggot, 1.6 – 3.2 Soybean aphids, Over-wintering bean leaf

		Rate (fl. oz. per hundred	
Crop	Pest	weight seed)	Comments
Adzuki Bean,	Wireworm,	1.6 - 3.2	*Seed-and-pod vegetable
Asparagus Bean,	Bean leaf beetle,		seed treated in California
Broad Bean (Succulant or Dry),	Imported fire ant,		must be destined for
Catjang Bean, Chinese	Aphid		planting in states other
Longbean, Field Bean, Guar			than California and is not
Bean, Jackbean, Kidney Bean,			to be planted in
Lablab Bean, Lima Bean			California.
(Succulent or Dry),			
Moth Bean (Succulent or Dry),			
Mung Bean, Navy Bean, Pinto			
Bean, Rice Bean, Runner Bean,			
Snap Bean, Sword Bean,			
Tepary Bean, Urd Bean, Wax			
Bean, Yardlong Bean, Blackeye			
Pea (Succulent or Dry),			
Chickpea, Cowpea (Succulent			
or Dry), Crowder Pea, Dwarf			
Pea, Edible-Pod Pea, English	,		
Pea, Field Pea, Garden Pea,			
Green Pea, Pigeon Pea			
(Succulent or Dry), Snow Pea,			0000
Southern Pea (Succulent or			0000
Dry), Sugar Snap Pea, Grain			0000
Lupin, Sweet Lupin, White			000000
Lupin, White Sweet Lupin,			0 0 0 0
Lentil*			00000
Carrot	Seed corn maggot,	6.4	0 0 0
	Wireworm		00000
			(0000

ROTATIONAL CROPS

Treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient, as soon as practical following the last application. The following plant-back intervals must be observed for listed crops:

Rotational Plant-Back Intervals*				
Immediate Plant-back				30-Day Plant-back
Artichoke	Cranberry	Mustard Seed	Soybean	Cereals, including:
Barley	Cucurbits	Oats	Strawberry	Buckwheat
Brassica (cole)	Eggplant	Okra	Sugarbeet	Millet
Leafy vegetable	Ground cherry	Pepinos	Tomatillo	Rice
Canola	Leaf petiole	Pepper	Tomato	Safflower
Cilantro	vegetable	Popcorn	Triticale	
Corn, Field	Leafy vegetables	Potato	Vegetable Root and	
Corn, Sweet	Legume vegetable	Rapeseed	Tuber	
Cotton	(succulent or dried,	Rye	Watercress	
	except soybean)	Sorghum	Wheat	

^{*}Cover crops for soil building or erosion control may be planted any time, but do not graze or harvest for food or feed.

Notification of the crop rotational restriction must be conveyed to the grower by appropriate seed tag labeling or bag printing on all seed units.

NOTE: The purchaser of this product is responsible for ensuring that all seed treated with this product are adequately dyed with a suitable color to prevent its accidental use as food for man or feed for animals. Refer to 21 CFR, Part 2.25. Any dye or colorant added to treated seed must be cleared for use under 40 CFR, Part 180.1001. Federal regulations have established official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates.

Treated seed must not be used for or mixed with food or animal feed or processed for oil. Seed commercially treated with Sharda Imidacloprid 5SC must be labeled in accordance with all applicable requirements of the Federal Seed Act.

Labels for commercially treated seed must include the following addition to the Environmental Hazards statements:

Exposed treated seed may be hazardous to birds. Dispose of all excess treated seed and seed packaging by burial away from bodies of water. Cover or incorporate spilled treated seeds.

Use Restriction

Rape greens grown and harvested from Sharda Imidacloprid 5SC treated seed must not be used for chuman and feed consumption. Rapeseed grown and harvested from treated seed is only for industrial uses and must not be used for edible oil or any other human/feed consumption.

Wheat, Barley, Oats, Rye, Triticale, Sorghum: Do not graze or feed livestock on treated areas for 45 days after planting.

^{*}For all other crops not listed on an Imidacloprid label, or for crops for which no tolerances for the active ingredient has been established, a 12-month plant-back interval must be observed.

Cotton (delinted seed only): Regardless of the type of application (seed treatment, soil, or foliar) do not apply more than a total of 0.5 lb. imidacloprid per acre per cropping cycle.

Conditions of Sale and Limited Warranty Statement

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.


By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Sharda USA LLC, no claims are made to guarantee germination of carry-over seed.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Sharda USA LLC. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW SHARDA USA LLC MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Sharda USA LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SHARDA USA LLC DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT SHARDA USA LLC'S ELECTION, THE REPLACEMENT OF PRODUCT.



Sharda Imidacloprid 5SC April 9, 2010 Page 11 of 11