

83529-20

11/18/2009

10f67



**U.S. ENVIRONMENTAL PROTECTION AGENCY**  
**Office of Pesticide Programs**  
**Registration Division (7505P)**  
**Ariel Rios Building**  
**1200 Pennsylvania Ave., NW**  
**Washington, D.C. 20460**

EPA Reg. Number:

83529-20

Date of Issuance:

NOV 18 2009

## NOTICE OF PESTICIDE:

Registration  
 Reregistration  
 (under FIFRA, as amended)

Term of Issuance: Unconditional

Name of Pesticide Product:

Sharda Glyphosate 41 SL

## Name and Address of Registrant (include ZIP Code):

Sharda USA, LLC c/o Wagner Regulatory Associates, Inc.  
 7460 Lancaster Pike, Suite 9 P.O. Box 640  
 Hockessin, DE 19707 Hockessin, DE 19707

**Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.**

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA provided that you:

1. Submit and/or cite all data required for registration review/reregistration of your product when the Agency requires all registrants of similar products to submit data.
2. Submit one-year Storage Stability (Guideline 830.6317) and Corrosion Characteristics (Guideline 830.6320) studies within one-year from the date of this notice.
3. Make the following label changes:
  - a. Change the EPA Reg. No. to "83529-20".
  - b. Include the EPA Establishment Number.
  - c. Revise the Net Contents from "1,000 liters" to "264 gallons (1,000 liters)", since the net contents must be declared in conventional U.S. units.
  - d. Remove pages 1 & 2 as the information is incomplete and redundant, or provide justification to support the need for the separated pages.

Continued on Page 2

Signature of Approving Official:

Jim Tompkins  
 Product Manager 25  
 Herbicide Branch  
 Registration Division (7505P)

Date:

NOV 18 2009

- e. Revise the Storage and Disposal statement to read “Do not contaminate water, food, or feed by storage and disposal.”
- f. Add the heading “CONTAINER HANDLING” to the subheadings “NONREFILLABLE CONTAINERS” and “RETURNABLE / REFILLABLE CONTAINERS”.
- g. Revise the Container Handling directions for non-refillable containers to language suitable for containers larger than 5 gallons as stated in PR Notice 2007-4, since the net contents for this product is greater than 5 gallons.
- h. On page 9, correct the typo from “directations” to “directions” in all instances.
- i. On pages 10 & 11 under For Aerial Applications in Mississippi, move the block of text from “The distance of the outermost nozzles on the boom...” to “3. Winds blowing from 5 to 10 miles per hour...in excess of 500 feet.” to the section under For Aerial Application in Arkansas Only.

The basic confidential statement of formula (CSF) dated July 30, 2009 is acceptable.

A stamped copy of the label is enclosed for your records. Submit one (1) copy of the revised final printed label before you release the product for shipment. Products shipped after eighteen (18) months from the date of this notice or the next printing of the label, whichever occurs first, must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA §6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Enclosure

30967

**Sharda Glyphosate 41 SL**

**Herbicide For Use in Certain Cropping Systems, Including Roundup Ready® Corn, Cotton, and Soybeans  
For Reduced Tillage and Fallow Systems and Non-Crop Uses**

<b>Active Ingredient:</b>	
Glyphosate, N-(phosphonomethyl)-glycine, as isopropylamine salt*	41.0%
<b>Other Ingredients:</b>	59.0%
<b>Total:</b>	100.0%

\*Contains 480 grams per liter or 4 pounds per U.S. gallon of the active ingredient glyphosate, in the form of isopropylamine salt. Equivalent to 358 grams per liter or 3 pounds per U.S. gallon of the acid, glyphosate.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See Attached Booklet for Additional Precautionary Statements and Directions for Use

FIRST AID	
<b>If In Eyes</b>	<ul style="list-style-type: none"> <li>- Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>- Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If Swallowed</b>	<ul style="list-style-type: none"> <li>- Call a poison control center or doctor immediately for treatment advice.</li> <li>- Have person sip a glass of water if able to swallow.</li> <li>- Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>- Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If Inhaled</b>	<ul style="list-style-type: none"> <li>- Move person to fresh air.</li> <li>- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>- Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If on Skin or Clothing</b>	<ul style="list-style-type: none"> <li>- Take off contaminated clothing.</li> <li>- Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>- Call a poison control center or doctor for treatment advice.</li> </ul>
<p><b>Emergency Numbers</b> For 24-hour medical emergency assistance (human or animal) call 1-800-222-1222. For chemical emergency assistance (spill, leak, fire, accident) call ChemTrec at 1-800-424-9300.</p>	
<p>Have the product label with you when calling a poison control center, or doctor, or going for treatment.</p>	

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION.** Causes moderate eye irritation. Avoid contact with eyes or clothing.

**AGRICULTURAL USE REQUIREMENTS**


Use Sharda Glyphosate 41 SL only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of, agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of Sharda Glyphosate 41 SL that are covered by the Worker Protection Standard.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

NOV 18 2009

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

83529-20



**STORAGE AND DISPOSAL**

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

**PESTICIDE STORAGE:** STORE ABOVE 10°F (-12°C) TO KEEP PRODUCT FROM CRYSTALLIZING. Crystals will settle to the bottom. If allowed to crystallize, place in a warm room 68°F (20°C) for several days to redissolve and shake or roll to mix well before using.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of Sharda Glyphosate 41 SL that cannot be used or chemically reprocessed must be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures. Emptied container retains vapor and product residue. Observe all label safeguards until container is destroyed.

**NONREFILLABLE CONTAINERS:** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling, if available.


**RETURNABLE / REFILLABLE CONTAINERS:** Refill this container with Sharda Glyphosate 41% SL only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect thoroughly for damage such as cracks, punctures, abrasions and damaged or worn out threads on closure devices. Do not refill or transport damaged or leaking containers. Check for leaks after refilling and before transportation. If the container is not being refilled, return it to the point of purchase. Prior to disposal or refilling, triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

EPA Reg. No. 83529-

EPA Est. No.

Net Contents: 1,000 liters

Manufactured by: Sharda USA LLC  
7460 Lancaster Pike Ste. 9, Hockessin DE 19707



50967

# Sharda Glyphosate 41 SL

## Herbicide For Use in Certain Cropping Systems, Including Roundup Ready® Corn, Cotton, and Soybeans

### For Reduced Tillage and Fallow Systems and Non-Crop Uses

Active Ingredient:	
Glyphosate, N-(phosphonomethyl)-glycine, as isopropylamine salt*	41.0%
Other Ingredients:	59.0%
<b>Total:</b>	<b>100.0%</b>

\*Contains 480 grams per liter or 4 pounds per U.S. gallon of the active ingredient glyphosate, in the form of isopropylamine salt. Equivalent to 358 grams per liter or 3 pounds per U.S. gallon of the acid, glyphosate.

EPA Reg. No. 83529-

EPA Est. No.

Net Contents: 1,000 liters

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

FIRST AID	
<b>If In Eyes</b>	<ul style="list-style-type: none"> <li>- Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>- Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If Swallowed</b>	<ul style="list-style-type: none"> <li>- Call a poison control center or doctor immediately for treatment advice.</li> <li>- Have person sip a glass of water if able to swallow.</li> <li>- Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>- Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If Inhaled</b>	<ul style="list-style-type: none"> <li>- Move person to fresh air.</li> <li>- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>- Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If on Skin or Clothing</b>	<ul style="list-style-type: none"> <li>- Take off contaminated clothing.</li> <li>- Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>- Call a poison control center or doctor for treatment advice.</li> </ul>
<b>Emergency Numbers</b> For 24-hour medical emergency assistance (human or animal) call 1-800-222-1222. For chemical emergency assistance (spill, leak, fire, accident) call ChemTrec at 1-800-424-9300.	
Have the product label with you when calling a poison control center, or doctor, or going for treatment.	

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**DOMESTIC ANIMALS:** This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

**CAUTION.** Causes moderate eye irritation. Avoid contact with eyes or clothing.

### Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning of equipment or disposing of equipment washwaters or rinsate.

60967

**PERSONAL PROTECTIVE EQUIPMENT (PPE):**

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Protective eyewear
- Shoes plus socks

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with Sharda Glyphosate 41 SL's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**ENGINEERING CONTROL STATEMENTS**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**USER SAFETY DIRECTIONS**

Users must:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.


**PHYSICAL OR CHEMICAL HAZARDS**

Spray solutions of Sharda Glyphosate 41 SL must be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers. DO NOT MIX, STORE OR APPLY SHARDA GLYPHOSATE 41 SL OR SPRAY SOLUTIONS OF SHARDA GLYPHOSATE 41 SL IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. Sharda Glyphosate 41 SL or spray solutions of Sharda Glyphosate 41 SL react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use Sharda Glyphosate 41 SL in any manner inconsistent with its labeling.

Do not apply Sharda Glyphosate 41 SL in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.



70967

**AGRICULTURAL USE REQUIREMENTS**

Use Sharda Glyphosate 41 SL only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of Sharda Glyphosate 41 SL that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical Resistant Gloves Category A such as butyl rubber, neoprene rubber, natural rubber, or nitril rubber  $\geq$  14 mils
- Protective eyewear
- Shoes plus socks

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of Sharda Glyphosate 41 SL that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when Sharda Glyphosate 41 SL is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Keep people and pets off treated areas until spray solution has dried.

**PRODUCT INFORMATION**

**DO NOT APPLY SHARDA GLYPHOSATE 41 SL USING AERIAL SPRAY EQUIPMENT EXCEPT UNDER CONDITIONS AS SPECIFIED WITHIN THIS LABEL OR CURRENT SUPPLEMENTAL LABELING ISSUED BY MANUFACTURER.**

Sharda Glyphosate 41 SL, a water soluble liquid, mixes readily with water to be applied as a foliar spray for the control or destruction of most herbaceous plants. It may be applied through most standard industrial or field-type sprayers after dilution and thorough mixing with water in accordance with label instructions. Hand-held sprayers may also be used.

Sharda Glyphosate 41 SL moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days, but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of Sharda Glyphosate 41 SL and delay visual effects of control. Visible effects are a gradual wilting and yellowing of the plant which advances to complete browning of above-ground growth and deterioration of underground plant parts. Unless otherwise specified on this label, delay application until vegetation has emerged and reached the stages described for control of such vegetation under the "WEEDS CONTROLLED" section of this label. Unemerged plants arising from unattached underground rhizomes or root stocks of perennials will not be affected by the herbicide and will continue to grow. For this reason, best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity.

Always use the higher rate of Sharda Glyphosate 41 SL per acre within the specified range when (1) weed growth is heavy or dense, or (2) weeds are growing in an undisturbed (noncultivated) area.

Reduced or unacceptable control may result if weeds or brush are treated under poor growing conditions such as drought stress, disease or insect damage. Reduced results may also occur when treating weeds or brush heavily covered with dust. Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed, or cut, and have not been allowed to regrow to the specified stage for treatment.

80967

Rainfall or irrigation occurring within 6 hours after application may reduce effectiveness. Heavy rainfall or irrigation within 2 hours after application may wash the chemical off the foliage and a repeat treatment may be required.

Sharda Glyphosate 41 SL does not provide residual weed control. For subsequent residual weed control, it is instructed that a residual herbicide program specified on this label be used. Read and carefully observe the precautionary statements and all other information appearing on the labels of all herbicides used.

Mixing Sharda Glyphosate 41 SL with herbicides or other materials not specified in this label may result in reduced performance. However, unless otherwise prohibited on this label or the label of an intended tank mix product, Sharda Glyphosate 41 SL may be applied in combination with any herbicide registered for the same site, timing, and method of application. Observe the most restrictive label statements of various tank mix products used. LIABILITY FOR CROP INJURY, HERBICIDE NONPERFORMANCE OR OTHER LOSS OR DAMAGE RESULTING FROM A TANK MIXTURE NOT SPECIFIED ON THIS LABEL, OR SUPPLEMENTAL LABELING DISTRIBUTED FOR SHARDA GLYPHOSATE 41 SL, IS SPECIFICALLY DISCLAIMED BY MANUFACTURER. BUYER AND ALL USERS ARE RESPONSIBLE FOR ALL LOSS OR DAMAGE IN CONNECTION WITH THE USE OR HANDLING OF MIXTURES OF SHARDA GLYPHOSATE 41 SL OR OTHER MATERIALS THAT ARE NOT EXPRESSLY SPECIFIED IN THIS LABEL.

For best results, spray coverage must be uniform and complete. Do not spray weed foliage to the point of runoff.

DOMESTIC ANIMALS: Sharda Glyphosate 41 SL is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of Sharda Glyphosate 41 SL or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

#### **ATTENTION**

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING SHARDA GLYPHOSATE 41 SL TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of Sharda Glyphosate 41 SL can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of injury occurring from the use of Sharda Glyphosate 41 SL is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions, including lesser wind velocities, will allow spray drift to occur. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) which are likely to drift. AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.

NOTE: Use of Sharda Glyphosate 41 SL in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences. When not in use, keep container closed to prevent spills and contamination.

#### **MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS**

APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES. DO NOT APPLY WHEN WIND OR OTHER CONDITIONS FAVOR DRIFT. HAND-GUN APPLICATIONS MUST BE PROPERLY DIRECTED TO AVOID SPRAYING DESIRABLE PLANTS.

NOTE: REDUCED RESULTS MAY OCCUR IF WATER CONTAINING SOIL IS USED, SUCH AS WATER FROM PONDS AND UNLINED DITCHES.

#### **MIXING**

Sharda Glyphosate 41 SL mixes readily with water. Mix spray solutions of Sharda Glyphosate 41 SL as follows:

1. Fill the mixing or spray tank with the required amount of water.
2. Add the specified amount of Sharda Glyphosate 41 SL (see the "DIRECTIONS FOR USE" and "WEEDS CONTROLLED" sections of this label) near the end of the filling process and mix well. Use caution to avoid siphoning back into the carrier source. Use approved anti-back-siphoning devices where required by state or local regulations. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, avoid the use of mechanical



90967

agitators, terminate by-pass and return lines at the bottom of the tank and, if needed, use an approved anti-foam or defoaming agent.

### TANK MIXTURES

Always predetermine the compatibility of labeled tank mixtures of Sharda Glyphosate 41 SL with water carrier by mixing small proportional quantities in advance. Mix labeled tank mixtures of Sharda Glyphosate 41 SL with water as follows:

1. Place a 20 to 35 mesh screen or wetting basket over filling port.
2. Through the screen, fill the spray tank one-half full with water and start agitation.
3. If a wettable powder is used, make a slurry with the water carrier, and add it SLOWLY through the screen into the tank. Continue agitation.
4. If a flowable formulation is used, premix one part flowable with one part water. Add diluted mixture SLOWLY through the screen into the tank. Continue agitation.
5. If an emulsifiable concentrate formulation is used, premix one part emulsifiable concentrate with two parts water. Add diluted mixture slowly through the screen into the tank. Continue agitation.
6. Continue filling the spray tank with water and add the required amount of Sharda Glyphosate 41 SL near the end of the filling process.
7. Add individual formulations to the spray tank as follows: wettable powder, flowable, emulsifiable concentrate, drift control additive, water soluble liquid.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed. Keep by-pass line on or near bottom of tank to minimize foaming. Screen size in nozzle or line strainers must be no finer than 50 mesh. Carefully select proper nozzle to avoid spraying a fine mist. For best results with conventional ground application equipment, use flat fan nozzles. Clean sprayer and parts immediately after using Sharda Glyphosate 41 SL by thoroughly flushing with water.

### ADDITIVES

**AMMONIUM SULFATE:** The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of Sharda Glyphosate 41 SL, and Sharda Glyphosate 41 SL plus 2,4-D, Banvel®, dicamba or residual herbicide tank mixtures on annual and perennial weeds. The improvement in performance may be apparent where environmental stress is a concern. Low-quality ammonium sulfate may contain material that will not readily dissolve, which could result in nozzle tip plugging. To determine quality, perform a jar test by adding 1/3 cup of ammonium sulfate to 1 gallon of water and agitate for 1 minute. If undissolved sediment is observed, predissolve the ammonium sulfate in water and filter prior to addition to the spray tank. If ammonium sulfate is added directly to the spray tank, add slowly with agitation. Adding too quickly may clog outlet line. Ensure that ammonium sulfate is completely dissolved in the spray tank before adding herbicides. Thoroughly rinse the spray system with clean water after use to reduce corrosion. Agriculturally-approved adjuvants containing equivalent amounts of ammonium sulfate may be used instead of dry ammonium sulfate.

### MAXIMUM USE RATE

The maximum use rates stated throughout this labeling apply to Sharda Glyphosate 41 SL combined with the use of all other herbicides containing glyphosate or sulfosate as the active ingredient, whether applied as mixtures or separately. Calculate the application rates and ensure that the total use of this and other glyphosate or sulfosate containing products does not exceed stated maximum use rate.

### COLORANTS OR DYES

Agriculturally-approved colorants or marking dyes may be added to Sharda Glyphosate 41 SL. Colorants or dyes used in spray solutions of Sharda Glyphosate 41 SL may reduce performance, especially at lower rates or dilutions. Use colorants or dyes according to the manufacturer's directions.

### APPLICATION EQUIPMENT AND TECHNIQUES

Do not apply Sharda Glyphosate 41 SL through any type of irrigation system.

Sharda Glyphosate 41 SL may be applied with the following application equipment:

- Aerial - Fixed Wing and Helicopter
- Broadcast Spray
- Controlled Droplet Applicator (CDA) - Hand-held or boom-mounted applicators which produce a

100 of 67

spray consisting of a narrow range of droplet sizes.

- Hand-Held and High-Volume Spray Equipment - Knapsack and backpack sprayers, pump-up pressure sprayers, handguns, handwands, mistblowers\*, lances and other hand-held and motorized spray equipment used to direct the spray onto weed foliage.

\*Sharda Glyphosate 41 SL is not registered in California or Arizona for use in mistblowers.

Selective equipment - Recirculating sprayers, shielded sprayers and wiper applicators. See the appropriate part of this section for specific instructions and rates of application.

### AERIAL EQUIPMENT

Use the specified rates of Sharda Glyphosate 41 SL in 3 to 15 gallons of water per acre unless otherwise specified on this label. See the "WEEDS CONTROLLED" section of this label for specific rates. Unless otherwise specified, do not exceed 1 quart per acre. Aerial applications of Sharda Glyphosate 41 SL may be made in annual cropping conventional tillage systems, fallow and reduced tillage systems, Roundup Ready® crops, noncrop areas and preharvest applications. Refer to the individual use area sections of this label for specified volumes and application rates. For specific state information, consult following section and state pesticide regulatory agency.

### STATE INFORMATION ON AERIAL APPLICATIONS

**ARKANSAS:** AVOID DRIFT. DO NOT APPLY INTO STILL AIR WHERE THERE IS A TEMPERATURE INVERSION LAYER LOW ENOUGH FOR FINE SPRAY PARTICLES TO BECOME SUSPENDED AND MOVE OUTSIDE THE TARGET AREA WHEN THE INVERSION LAYER MOVES. DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION THAT FAVORS DRIFT. DRIFT IS LIKELY TO CAUSE DAMAGE TO ANY VEGETATION CONTACTED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Use the specified rate of Sharda Glyphosate 41 SL in 3 to 15 gallons of water per acre.

Use sufficient carrier volume and appropriate equipment set-up to form droplets large enough to avoid drift potential. Coarse droplets in the 300 to 500 (VMD) micron range are directed.

Applications must typically be made with the nozzle release point at 8 to 15 feet above the top of the target plants unless a greater height is required for aircraft safety.

The distance of the outermost nozzles on the boom must not exceed 75% of the length of the wingspan or rotor, in many cases, reducing this distance to 65% of the length of the wingspan or rotor will improve drift control without affecting the swath width. Nozzles must always discharge backward parallel with the air stream and never discharge downwards more than 45 degrees on fixed wing aircraft or forward of the prevailing air flow on rotary winged aircraft. Avoid the use of nozzles with wide angle discharge. Do not apply Sharda Glyphosate 41 SL when winds are in excess of 10 miles per hour.

Do not apply when there is a low-level inversion where fine spray particles could be suspended in still air and move outside the target area when the inversion layer moves. These conditions may occur when wind speeds are less than 2 mph. Use the following guidelines when applications are made near crops or other desirable vegetation:

1. Do not apply within 100 feet of any desirable vegetation or crops.
2. If wind up to 5 miles per hour is blowing toward desirable vegetation or crops, do not apply within 500 feet of the desirable vegetation or crops.
3. Winds blowing from 5 to 10 miles per hour toward desirable vegetation or crops will likely require buffer zones in excess of 500 feet.

**CALIFORNIA- Statewide:** Aerial applications of Sharda Glyphosate 41 SL are allowed in the following situations:

1. In fallow and reduced tillage systems prior to the emergence or transplanting of labeled crops.
2. In cotton, prior to harvest. Refer to the specific preharvest application instructions.

Do not plant subsequent crops other than those listed in the label booklet for Sharda Glyphosate 41 SL for 30 days following application.

When applied as specified, under the conditions described, Sharda Glyphosate 41 SL controls annual and perennial weeds listed in the label.

110907

**DO NOT EXCEED A MAXIMUM RATE OF 2 QUARTS PER ACRE OF SHARDA GLYPHOSATE 41 SL WHEN MAKING APPLICATIONS BY AIR IN FALLOW AND REDUCED TILLAGE SYSTEMS.**

**DO NOT EXCEED A MAXIMUM RATE OF 1 QUART PER ACRE OF SHARDA GLYPHOSATE 41 SL WHEN MAKING APPLICATIONS BY AIR IN COTTON, PRIOR TO HARVEST.**

Use the specified rates of Sharda Glyphosate 41 SL in 3 to 15 gallons of water per acre.

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

**AVOID DRIFT - DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION WHICH WILL ALLOW DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.**

Use the following guidelines when aerial applications are made near crops or desirable perennial vegetation after bud break and before total leaf drop, and/or near other desirable vegetation or annual crops.

1. Do not apply within 100 feet of all desirable vegetation or crop(s).
2. If wind up to 5 miles per hour is blowing toward desirable vegetation or crop(s), do not apply within 500 hundred feet of the desirable vegetation or crop(s).
3. Winds blowing from 5 to 10 miles per hour toward desirable vegetation or crop(s) may require buffer zones in excess of 500 feet.
4. Do not apply when winds are in excess of 10 miles per hour or when inversion conditions exist. Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine spray droplets. Do not angle nozzles forward into the airstream and do not increase nozzle pressure.

Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information and all other information appearing on the additive label. Ensure uniform application - to avoid streaked, uneven or overlapped application, use appropriate marking devices.

**FOR AERIAL APPLICATION IN CALIFORNIA ONLY (including Fresno County, California)**

**CROP USES**

Aerial applications of Sharda Glyphosate 41 SL are allowed in the following situations:

1. Over-the-top applications in RoundUp Ready® corn and cotton. Refer to the Sharda Glyphosate 41 SL booklet and/or supplemental labels for specific application instructions for over-the-top applications in these crops.

Do not plant subsequent crops other than those listed in the label booklet for 30 days following application.

When applied as specified under the conditions described, Sharda Glyphosate 41 SL controls annual and perennial weeds listed in the label booklet.

Do not exceed a maximum rate of 2 quarts per acre when making applications by air to Roundup Ready® cotton.

**CALIFORNIA - Fresno County:** This section applies to aerial applications in Fresno County from February 15 through March 31 only. For aerial application outside of these dates, refer to the "CALIFORNIA ~ Statewide" section.

**APPLICABLE AREA:** This supplement only applies to the area contained inside the following boundaries within Fresno County, California only. **North:** Fresno County line, **South:** Fresno County line, **East:** State Highway 99, **West:** Fresno County line.

**PRODUCT INFORMATION:** Always read and follow the label directions and precautionary statements for all products used in the aerial application. Observe the following directions to minimize off-site movement during aerial application of this herbicide. Minimization of off-site movement is the responsibility of the grower, Pest Control Advisor, and aerial applicator.

**WRITTEN DIRECTATIONS:** A written directation **MUST** be submitted by or on behalf of the applicator to the Fresno County Agricultural Commissioner 24 hours prior to the application. This written directation **MUST**

120667

state the proximity of surrounding crops, and that conditions of each manufacturer's applicable product label(s) and this label have been satisfied.

**AERIAL APPLICATOR TRAINING AND EQUIPMENT:** Aerial application of Sharda Glyphosate 41 SL is limited to pilots who have successfully completed a Fresno County Agricultural Commissioner and California Department of Pesticide Regulation approved training program for aerial application of herbicides. All aircraft must be inspected, critiqued in flight, and certified at a Fresno County Agricultural Commissioner approved fly-in. Test and calibrate spray equipment at intervals sufficient to insure that proper rates of herbicides and adjuvants are being applied during commercial use. Applicator must document such calibrations and testing. Demonstration of performance at Fresno County Agricultural Commissioner approved "fly-ins" constitutes such documentation, or other written records showing calculations and measurements of flight and spray parameters acceptable to the Fresno County Agricultural Commissioner.

**APPLICATION AT NIGHT:** Do not apply Sharda Glyphosate 41 SL by air earlier than 30 minutes prior to sunrise and/or later than 30 minutes after sunset without prior permission from the Fresno County Agricultural Commissioner.

To report known or suspected misuse of Sharda Glyphosate 41 SL, or for additional information on the proper aerial application of Sharda Glyphosate 41 SL, call 1-800-852-5234.

Avoid direct application to any body of water.

**AVOID DRIFT - DO NOT APPLY DURING LOW-LEVEL INVERSION CONDITIONS, WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION WHICH FAVORS DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.**

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine spray droplets.

Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure above the manufacturer's direction.

Drift control additives may be used. When a drift control additive is used, read and carefully observe the precautionary statements and all other information appearing on the additive label.

#### **SPRAY DRIFT MANAGEMENT**

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they must be observed.

The applicator must be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information.

**Importance of Droplet Size:** The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversion section of this label).

#### **Controlling Droplet Size**

**Volume:** Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

130 of 107

**Pressure:** Use the lower spray pressures specified for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

**Number of nozzles:** Use the minimum number of nozzles that provide uniform coverage.

**Nozzle Orientation:** Orienting nozzles so that the spray is released backwards, parallel to the airstream will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.

**Nozzle Type:** Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.

**Boom Length:** For some use patterns, reducing the effective boom length to less than  $\frac{3}{4}$  of the wingspan or rotor length may further reduce drift without reducing swath width.

**Application:** Applications must not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

### Swath Adjustment

When applications are made with a cross-wind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance must increase, with increasing drift potential (higher wind, smaller drops, etc.).

### Wind

Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application must be avoided below 2 mph due to variable wind direction and high inversion potential. Note: Local terrain can influence wind patterns. Every applicator must be familiar with local wind patterns and how they affect drift.

### Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

### Temperature Inversions

Applications must not occur during a temperature inversion, because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a connected cloud (under low wind conditions) indicates an inversion, while smoke that moves upwards and rapidly dissipates indicates good vertical air mixing.

### Sensitive Areas

Sharda Glyphosate 41 SL must only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

Ensure uniform application - To avoid streaked, uneven or overlapped application, use appropriate marking devices. Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of Sharda Glyphosate 41 SL accumulated during spraying or from spills. PROLONGED EXPOSURE OF SHARDA GLYPHOSATE 41 SL\* TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR ARE MOST SUSCEPTIBLE. The maintenance of an organic coating (paint), which meets aerospace specification MIL-C-38413, may prevent corrosion.

\*Sharda Glyphosate 41 SL plus Oust®, Banvel®, dicamba or 2,4-D tank mixtures may not be applied by air in California.

14067

## BROADCAST EQUIPMENT

For control of annual or perennial weeds listed on this label using broadcast equipment - Use the specified rates of Sharda Glyphosate 41 SL in 3 to 40 gallons of water per acre as a broadcast spray unless otherwise specified on this label. See the "WEEDS CONTROLLED" section of this label for specific rates. As density of weeds increases, spray volume must be increased within the specified range to ensure complete coverage. Carefully select proper nozzle to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

## CONTROLLED DROPLET APPLICATION (CDA)

The rate of Sharda Glyphosate 41 SL applied per acre by vehicle-mounted CDA equipment must not be less than the amount specified in this label when applied by conventional broadcast equipment. For vehicle-mounted CDA equipment, apply 3 to 15 gallons of water per acre.

For the control of labeled annual weeds with hand-held CDA units, apply a 20 percent solution of Sharda Glyphosate 41 SL at a flow rate of 2 fluid ounces per minute and a walking speed of 1.5 MPH (1 quart per acre). For the control of labeled perennial weeds, apply a 20 to 40 percent solution of Sharda Glyphosate 41 SL at a flow rate of 2 fluid ounces per minute and a walking speed of 0.75 mph (2 to 4 quarts per acre). Controlled droplet application equipment produces a spray pattern which is not easily visible. Extreme care must be exercised to avoid spray or drift contacting the foliage or any other green tissue of desirable vegetation, as damage or destruction may result.

## HAND-HELD AND HIGH-VOLUME EQUIPMENT

Use Coarse Sprays Only.

Mix Sharda Glyphosate 41 SL in clean water and apply to foliage of vegetation to be controlled. For applications made on a spray-to-wet basis, spray coverage must be uniform and complete. Do not spray to the point of runoff.

For control of annual weeds listed on this label, apply a 0.5 percent solution of Sharda Glyphosate 41 SL to weeds less than 6 inches in height or runner length. Apply prior to seedhead formation in grass or bud formation in broadleaf weeds. Allow three or more days before tillage or mowing. For annual weeds over 6 inches tall, or unless otherwise specified, use a 1 percent solution. For best results, use a 2 percent solution on harder-to-control perennials, such as bermudagrass, dock, field bindweed, hemp dogbane, milkweed and Canada thistle. When using application methods which result in less than complete coverage, use a 5 percent solution for annual and perennial weeds and a 5 to 10 percent solution for woody brush and trees.

Prepare the desired volume of spray solution by mixing the amount of Sharda Glyphosate 41 SL in water as shown in the following table:

### Spray Solution

AMOUNT OF SHARDA GLYPHOSATE 41 SL						
Desired Volume	1/2%	1%	1-1/2%	2%	5%	10%
1 Gal.	2/3 oz.	1-1/3 oz.	2oz.	2-2/3 oz.	6-1/2 oz.	13 oz.
25 Gal.	1 qt.	1qt.	1-1/2 qt.	2qt.	5qt.	10 qt.
100 Gal.	2qt.	1 gal.	1-1/2 gal.	2 gal.	5 gal.	10 gal.

2 tablespoons = 1 fluid ounce

For use in knapsack sprayers, it is suggested that the specified amount of Sharda Glyphosate 41 SL be mixed with water in a larger container. Fill sprayer with the mixed solution.

## SELECTIVE EQUIPMENT

Sharda Glyphosate 41 SL may be applied through a recirculating spray system, a shielded applicator, or a wiper applicator after dilution and thorough mixing with water to listed weeds growing in any noncrop site specified on this label and only when specifically directed in cropping systems.

- A recirculating spray system directs the spray solution onto weeds growing above desirable vegetation, while spray solution not intercepted by weeds is collected and returned to the spray tank for reuse.
- A shielded applicator directs the herbicide solution onto weeds, while shielding desirable vegetation from the herbicide.
- A wiper applicator applies the herbicide solution onto weeds by rubbing the weed with an absorbent material containing the herbicide solution.

### AVOID CONTACT WITH DESIRABLE VEGETATION

Contact of the herbicide solution with the desirable vegetation may result in damage or destruction. Applicators used above desired vegetation must be adjusted so that the lowest spray stream or wiper contact point is at least 2 inches above the desirable vegetation. Droplets, mist, foam, or splatter of the herbicide solution settling on desirable vegetation may result in discoloration, stunting or destruction.

Applications made above the crops (such as wiper applications) must be made when the weeds are a minimum of 6 inches above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations or when the height of the weeds varies so that not all weeds are contacted. In these instances, repeat treatment may be necessary.

### SHIELDED APPLICATORS

When applied as directed under conditions described for shielded applicators, Sharda Glyphosate 41 SL will control those weeds listed in the "WEEDS CONTROLLED" section of this label. Use the following equation to convert from a broadcast rate per acre to a band rate per acre.

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Herbicide Broadcast RATE per acre} = \text{Herbicide Band RATE per acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast VOLUME of solution per acre} = \text{Band VOLUME of solution per acre}$$

Use nozzles that provide uniform coverage within the treated area. Keep shields on shielded sprayers adjusted to protect desirable vegetation. EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT WITH DESIRABLE VEGETATION.

For specific rates of application and instructions for control of various annual and perennial weeds, see the "WEEDS CONTROLLED" section of this label.

### WIPER APPLICATORS

Wiper applicators are devices that physically wipe appropriate amounts of Sharda Glyphosate 41 SL directly onto the weed. Equipment must be designed, maintained and operated to prevent the herbicide solution from contacting desirable vegetation. Operate this equipment at ground speeds no greater than 5 mph. Performance may be improved by reducing speed in areas of heavy weed infestations to ensure adequate wiper saturation. Better results may be obtained if 2 applications are made in opposite directions.

Avoid leakage or dripping onto desirable vegetation. Adjust height of applicator to ensure adequate contact with weeds. Keep wiping surfaces clean. Be aware that, on sloping ground, the herbicide solution may migrate, causing dripping on the lower end and drying of the wicks on the upper end of a wiper applicator. Do not use wiper equipment when weeds are wet.

Mix only the amount of solution to be used during a 1-day period, as reduced activity may result from use of leftover solutions. Clean wiper parts immediately after using Sharda Glyphosate 41 SL by thoroughly flushing with water.

Do not add surfactant to the herbicide solution.

**For Rope or Sponge Wick Applicators** - Mix 1 gallon of Sharda Glyphosate 41 SL in 2 gallons of water to prepare a 33 percent solution. Apply this solution to weeds listed in this "WIPER APPLICATORS" section.

**For Porous-Plastic Applicators** - Solutions ranging from 33 to 100 percent of Sharda Glyphosate 41 SL in water may be used in porous-plastic wiper applicators.

When applied as specified under the conditions described for "WIPER APPLICATORS", Sharda Glyphosate 41 SL controls the following weeds:

Annual Grasses	Annual Broadleaves
Corn ( <i>Zea mays</i> ) Panicum, Texas ( <i>Panicum texanum</i> ) Rye, common ( <i>Secale cereale</i> ) Shattercane ( <i>Sorghum bicolor</i> )	Sicklepod ( <i>Cassia obtusifolia</i> ) Spanishneedles ( <i>Bidens bipinnata</i> ) Starbur, bristly ( <i>Acanthospermum hispidum</i> )

160267

When applied as specified under the conditions described for "WIPER APPLICATORS", Sharda Glyphosate 41 SL SUPPRESSES the following weeds:

Annual Broadleaves	Perennial Grasses	Perennial Broadleaves
Beggarwood, Florida ( <i>Desmodium tortuosum</i> )	Bermudagrass ( <i>Cynodon dactylon</i> )	Dogbane, hemp ( <i>Apocynum cannabinum</i> )
Pigweed, redroot ( <i>Amaranthus retroflexus</i> )	Johnsongrass ( <i>Sorghum halepense</i> )	Milkweed ( <i>Asclepias syriaca</i> )
Ragweed, giant ( <i>Ambrosia trifida</i> )	Smutgrass ( <i>Sporobolus poiretii</i> )	Nightshade, silverleaf ( <i>Solanum elaeagnifolium</i> )
Thistle, musk ( <i>Carduus nutans</i> )	Vaseygrass ( <i>Paspalum urvillei</i> )	Thistle, Canada ( <i>Cirsium arvense</i> )
Dogfennel ( <i>Eupatorium capilliflorum</i> )	Guineagrass ( <i>Panicum maximum</i> )	
Ragweed, common ( <i>Ambrosia artemisiifolia</i> )		
Sunflower ( <i>Helianthus annuus</i> )		
Velvetleaf ( <i>Abutilon theophrasti</i> )		

**WEEDS CONTROLLED**

This herbicide controls many annual and perennial grasses and broadleaf weeds.

**ANNUAL WEEDS.**

- Apply to actively growing grass and broadleaf weeds.
- Allow at least 3 days after treatment before tillage.
- For maximum agronomic benefit, apply when weeds are 6 inches or less in height.
- To prevent seed production, applications must be made prior to seedhead formation.
- Sharda Glyphosate 41 SL does not provide residual control; therefore, delay application until maximum weed emergence. Repeat treatments may be necessary to control later germinating weeds.

**LOW-VOLUME BROADCAST APPLICATION (LOW-RATE TECHNOLOGY):** When applied as directed under the conditions described, Sharda Glyphosate 41 SL will control the weeds listed below when water carrier volumes of 3 to 10 gallons per acre for ground applications and 3 to 5 gallons per acre for aerial applications are directed. (See the "AERIAL APPLICATION" section of this label for approved sites).

**NOTE:** The addition of 2 percent dry ammonium sulfate by weight or 17 pounds per 100 gallons of water may increase the performance of Sharda Glyphosate 41 SL on annual weeds. The improvement in performance may be apparent where environmental stress is a concern. Refer to the "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" section of this label.

- Do not tank-mix with soil residual herbicides when using these rates unless otherwise specified.
- For weeds that have been mowed, grazed or cut, allow regrowth to occur prior to treatment.
- Refer to the 'TANK MIXTURES' portion of this section for control of additional broadleaf weeds.

Weed Species	Maximum Height/Length	Rate per Acre* (fl. oz.)
Foxtail ( <i>Setaria</i> spp.)	12"	8 oz.
Barnyard grass ( <i>Echinochloa crus-galli</i> )	6"	12 oz.
Bluegrass, annual ( <i>Poa annua</i> )	(0 to 4")	16 oz. <sup>1</sup>
Brome, downy** ( <i>Bromus tectorum</i> )	(4 to 6")	24 oz. <sup>1</sup>
Mustard, blue ( <i>Chorispora tenella</i> )		
Mustard, tansy ( <i>Descurainia pinnata</i> )		
Mustard, tumble ( <i>Sisymbrium altissimum</i> )	6"	12 oz.
Mustard, wild ( <i>Sinapis arvensis</i> )	(0 to 4")	16 oz. <sup>1</sup>
Spurry, umbrella ( <i>Holosteum umbellatum</i> )	(4 to 6")	24 oz. <sup>1</sup>



Weed Species	Maximum Height/Length	Rate per Acre* (fl. oz.)
Barley ( <i>Hordeum vulgare</i> ) Rye ( <i>Secale cereale</i> ) Shattercane ( <i>Sorghum bicolor</i> ) Stinkgrass ( <i>Eragrostis ciliaris</i> )	12"	
Wheat ( <i>Triticum aestivum</i> )	18"	
Morningglory ( <i>Ipomoea spp.</i> ) Sicklepod ( <i>Cassia obtusifolia</i> )	2"	16 oz.
Bluegrass, bulbous ( <i>Poa bulbosa</i> ) Cheat ( <i>Bromus secalinus</i> ) Chickweed, mouseear ( <i>Cerastium vulgatum</i> ) Corn ( <i>Zea mays</i> ) Goatgrass, jointed ( <i>Aegilops cylindrical</i> ) Groundsel, common ( <i>Senecio vulgaris</i> ) Henbit ( <i>Lamium amplexicaule</i> ) Horseweed/Marestail ( <i>Conyza Canadensis</i> ) Lambsquarters, common ( <i>Chenopodium album</i> ) Pennycress, field ( <i>Fanweed Thlaspi arvense</i> ) Rocket, London ( <i>Sisymbrium irio</i> ) Ryegrass, Italian ( <i>Lolium multiflorum</i> ) Shepherdspurse ( <i>Capsella burs-pastoris</i> ) Spurge, annual ( <i>Euphorbia spp.</i> )	6"	16 oz.
Buttercup ( <i>Ranunculus spp.</i> ) Cocklebur ( <i>Xanthium strumarium</i> ) Crabgrass ( <i>Digitaria spp.</i> ) Dwarf dandelion ( <i>Krigia cespitosa</i> ) Falseflax, smallseed ( <i>Camelina microcarpa</i> ) Foxtail, Carolina ( <i>Alopecurus carolinianus</i> ) Johnsongrass, seedling ( <i>Sorghum halepense</i> ) Oats, wild ( <i>Avena fatua</i> ) Panicum, Fall ( <i>Panicum dichotomiflorum</i> ) Panicum, Texas ( <i>Panicum texanum</i> ) Pigweed, redroot ( <i>Amaranthus retroflexus</i> ) Pigweed, smooth ( <i>Amaranthus hybridus</i> ) Witchgrass ( <i>Panicum capillare</i> )	12"	16 oz.
Sicklepod ( <i>Cassia obtusifolia</i> ) Signalgrass, broadleaf ( <i>Brachiaria platyphylla</i> )	3 to 4" 4"	24 oz.

180f67

Weed Species	Maximum Height/Length	Rate per Acre* (fl. oz.)
Horseweed/Marestail ( <i>Conyza Canadensis</i> ) Lambsquarters, common ( <i>Chenopodium album</i> ) Spurge, annual ( <i>Euphorbia spp.</i> )	7 to 12"	
Rice, Red ( <i>Oryza sativa</i> ) Teawood ( <i>Sida spinosa</i> )	4"	32 oz.
Sprangletop ( <i>Leptochloa spp.</i> )	6"	
Geranium, Carolina ( <i>Geranium carolinianum</i> ) Goosegrass ( <i>Eleusine indica</i> ) Primrose, cutleaf evening ( <i>Oenothera lacinate</i> ) Pusley, Florida ( <i>Richardia scabra</i> )	12"	
Sicklepod ( <i>Cassia obtusifolia</i> ) Spanishneedles ( <i>Bidens bipinnata</i> )	5 to 12"	32 oz.
Filaree ( <i>Erodium spp.</i> ) Sprangletop ( <i>Laptochloa spp.</i> )	12 "	48 oz.

<sup>1</sup>Use these rates to control banyardgrass in Alabama, Arkansas, Mississippi, Missouri, Louisiana, and Texas for preplant treatments.

\*For those rates less than 32 fluid ounces per acre, Sharda Glyphosate 41 SL at rates up to 32 fluid ounces per acre may be used where heavy weed densities exist.

\*\*For control in no-till systems, use 16 fluid ounces per acre.

**TANK MIXTURES**

**Sharda Glyphosate 41 SL plus BANVEL® or dicamba**

**Sharda Glyphosate 41 SL plus 2,4-D**

**DO NOT APPLY BANVEL®, DICAMBA OR 2,4-D TANK MIXTURES BY AIR IN CALIFORNIA.**

Sharda Glyphosate 41 SL may be tank-mixed with the listed herbicides provided the specific product is registered for use on these sites. These tank mixtures are directed for use in fallow and reduced tillage areas only. Follow use directions as given in the "LOW-VOLUME BROADCAST APPLICATION" section.

Sharda Glyphosate 41 SL plus Banvel®, dicamba or 2,4-D will control the annual grasses and broadleaf weeds listed for Sharda Glyphosate 41 SL alone at the indicated heights (except 8 fluid ounces per acre applications), plus the following broadleaf weeds. For those weeds previously listed at 8 fluid ounces of Sharda Glyphosate 41 SL alone per acre, use 12 fluid ounces in these tank mixtures.

**NOTE:** Refer to the specific product labels for crop rotation restrictions and precautionary statements of all products used in tank mixtures. Some crop injury may occur if Banvel® or dicamba is applied within 45 days of planting. The addition of Banvel® or dicamba in a mixture with Sharda Glyphosate 41 SL may provide short-term residual control of selected weed species.

19067

Weed Species	Maximum weed height to obtain maximum control	Application rate w/0.25 lb. a.i. of Banvel® or Dicamba	Application rate w/0.5 lb. a.i. of 2,4-D
Cocklebur ( <i>Xanthium strumarium</i> )	12"	12 to 16 fl. oz.	12 to 16 fl. oz.
Lettuce, prickly ( <i>Lactuca serriola</i> )	6"		
Morningglory ( <i>Ipomoea spp.</i> )	6"		
Pigweed, smooth ( <i>Amaranthus hybridus</i> )	12"		
Kochia* ( <i>Kochia scoparia</i> )	6"		
Marestail/Horseweed ( <i>Conyza Canadensis</i> )	6"		
Pigweed, redroot ( <i>Amaranthus retroflexus</i> )	12"		
Thistle, Russian ( <i>Salsola kali</i> )	12"		
Lambsquarters ( <i>Chenopodium album</i> )	12"		

\*Controlled with Banvel® or dicamba tank mixture only.

Weed Species	Maximum weed height to obtain maximum control	Application rate w/0.5 lb. a.i. of 2,4-D
Ragweed, Common ( <i>Ambrosia artemisiifolia</i> )	6"	16 fl. oz.
Ragweed, Giant ( <i>Ambrosia trifida</i> )		
Smartweed, Pennsylvania ( <i>Polygonum pensylvanicum</i> )		
Velvetleaf ( <i>Abutilon theophrasti</i> )		

### HIGH-VOLUME BROADCAST APPLICATIONS

If weeds have been mowed, grazed or cut, allow adequate time for new growth to reach specified stages prior to treatment. These rates will also provide control of weeds listed in the "LOW-VOLUME BROADCAST APPLICATION" section.

Weed Species	Water Carrier Volume for ground application	Application rate for weeds less than 6" tall	Application rate for weeds greater than 6" tall
Balsamapple* ( <i>Momordica charantia</i> )	10 to 40 gallons	1 quart per acre	1.5 quarts per acre
Brome ( <i>Bromus spp.</i> )			
Fleabane, hairy ( <i>Conyza bonariensis</i> )			
Kochia ( <i>Kochia scoparia</i> )			
Bassia, fivehook ( <i>Bassia hyssopifolia</i> )			
Fiddleneck ( <i>Amsinckia spp.</i> )			
Fleabane ( <i>Erigeron spp.</i> )			

Weed Species	Water Carrier Volume for ground application	Application rate for weeds less than 6" tall	Application rate for weeds greater than 6" tall
Lettuce, prickly ( <i>Lactuca serriola</i> )	10 to 40 gallons	1 quart per acre	1.5 quarts per acre
Panicum ( <i>Panicum</i> )			
Ragweed, Common ( <i>Ambrosia artemisiifolia</i> )			
Smartweed, Pennsylvania ( <i>Polygonum pennsylvanicum</i> )			
Sowthistle, annual ( <i>Sonchus oleraceus</i> )			
Sunflower ( <i>Helianthus annuus</i> )			
Thistle, Russian ( <i>Salsola kali</i> )			
Velvetleaf ( <i>Abutilon theophrasti</i> )			

**PERENNIAL WEEDS**

Apply Sharda Glyphosate 41 SL as follows to control or destroy most perennial weeds:

**NOTE:** If weeds have been mowed or tilled, do not treat until plants have resumed active growth and have reached the specified stages. Repeat treatments may be necessary to control weeds regenerating from underground parts or seed. Repeat treatments must be made prior to crop emergence. The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of Sharda Glyphosate 41 SL on perennial weeds. The improvement in performance may be apparent where environmental stress is a concern. Refer to the "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" section of this label.

When applied as specified under the conditions described, Sharda Glyphosate 41 SL WILL CONTROL the following PERENNIAL WEEDS:

Alfalfa ( <i>Medicago sativa</i> )	Cattail ( <i>Typha spp.</i> )	Kikuyugrass ( <i>Pennisetum clandestinum</i> )	Quackgrass ( <i>Agropyron repens</i> )
Alligatorweed* ( <i>Alternanthera philoxeroides</i> )	Clover, Red ( <i>Trifolium pretense</i> )	Knapweed ( <i>Centaurea repens</i> )	Redvine* ( <i>Brunnichia ovata</i> )
Anise (fennel) ( <i>Foeniculum vulgare</i> )	Clover, White ( <i>Trifolium repens</i> )	Lantana ( <i>Lantana camara</i> )	Reed, giant ( <i>Arundo donax</i> )
Artichoke, Jerusalem ( <i>Helianthus tuberosus</i> )	Cogongrass ( <i>Imperata cylindrical</i> )	Lespedeza ( <i>Lespedeza spp.</i> )	Ryegrass, perennial ( <i>Lolium perenne</i> )
Bahiagrass ( <i>Paspalum notatum</i> )	Dallisgrass ( <i>Paspalum dilatatum</i> )	Milkweed ( <i>Asclepias spp.</i> )	Smartweed, swamp ( <i>Polygonum coccineum</i> )
Bentgrass ( <i>Agrostis spp.</i> )	Dandelion ( <i>Taraxacum officinate</i> )	Muhly, wirestem ( <i>Muhlenbergia frondosa</i> )	Spurge, leafy* ( <i>Euphorbia esula</i> )
Bermudagrass ( <i>Cynodon dactylon</i> )	Dock, curly ( <i>Rumex crispus</i> )	Mullein, common ( <i>Verbascum Thapsus</i> )	Starthistle, yellow ( <i>Centaurea solstitialis</i> )
Bermudagrass, water (knotgrass) ( <i>Paspalum distichum</i> )	Dogbane, hemp ( <i>apocynum cannabinum</i> )	Napierrgrass ( <i>Pennisetum purpureum</i> )	Sweet potato, wild* ( <i>Ipomoea pandurata</i> )

Fescues ( <i>Festuca spp.</i> )	Nightshade, silverleaf ( <i>Solanum elaeagnifolium</i> )	Thistle, Canada ( <i>Cirsium arvense</i> )	Bindweed, field ( <i>Convolvulus arvensis</i> )
Fescue, tall ( <i>Festuca arundinacea</i> )	Nutsedge; purple, yellow ( <i>Cyperus rotundus</i> ; <i>Cyperus esculentus</i> )	Thistle, artichoke ( <i>Cynara cardunculus</i> )	Blueweed, Texas ( <i>Helianthus ciliaris</i> )
Guineagrass ( <i>Panicum maximum</i> )	Orchardgrass ( <i>Dactylis glomerata</i> )	Timothy ( <i>Phleum pratense</i> )	Horsenettle ( <i>Solanum carolinense</i> )
Pampasgrass ( <i>Cortaderia spp.</i> )	Torpedograss* ( <i>Panicum repens</i> )	Brackenfern ( <i>Pteridium aquilinum</i> )	Horseradish ( <i>Armoracia rusticana</i> )
Paragrass ( <i>Brachiaria mutica</i> )	Trumpetcreeper* ( <i>Campsis radicans</i> )	Bromegrass, smooth ( <i>Bromus inermis</i> )	Ice plant ( <i>Mesembryanthemum, Crystallinum</i> )
Phragmites* ( <i>Phragmites spp.</i> )	Vaseygrass ( <i>Paspalum urvillei</i> )	Velvetgrass ( <i>Holcus spp.</i> )	Canarygrass, reed ( <i>Phalaris arundinacea</i> )
Johnsongrass, ( <i>Sorghum halepense</i> )	Poison hemlock ( <i>Conium maculatum</i> )	Wheatgrass, western ( <i>Agropyron smithii</i> )	

\*Partial control

See "DIRECTIONS FOR USE" and "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" sections of this label for labeled uses and specific application instructions.

Weed	Directions for Use
Alfalfa	Apply 1 quart of Sharda Glyphosate 41 SL per acre in 3 to 10 gallons of water per acre. Make application after the last hay cutting in the fall. Allow alfalfa to regrow to a height of 6 to 8 inches or more prior to treatment. Applications must be followed with deep tillage at least 7 days after treatment, but before soil freeze-up.
Alligatorweed	Apply 4 quarts of Sharda Glyphosate 41 SL per acre or apply a 1.5 percent solution with hand-held equipment to provide partial control. Apply when most of the plants are in bloom. Repeat applications will be required to maintain such control.
Anise (fennel)/Poison hemlock	Apply a 1 to 2 percent solution of Sharda Glyphosate 41 SL as a spray-to-wet treatment. Optimum results are obtained when plants are treated at the bud to full-bloom stage of growth. Repeat applications may be needed in succeeding years to control plants arising from seeds.
Bentgrass	For suppression in grass seed production areas. For ground applications only, apply 1.5 quarts of Sharda Glyphosate 41 SL in 10 to 20 gallons of water per acre. Ensure entire crown area has resumed growth prior to a fall application. Bentgrass must be actively growing and have at least 3 inches of growth. Tillage prior to treatment must be avoided. Tillage 7 to 10 days after application is directed for best results. Failure to use tillage after treatment may result in unacceptable control.
Bermudagrass	For control, apply 5 quarts of Sharda Glyphosate 41 SL per acre. For partial control, apply 3 quarts per acre. Treat when bermudagrass is actively growing and seedheads are present. Retreatment may be necessary to maintain control Allow 7 or more days after application before tillage.

Weed	Directions for Use
Bermudagrass, water (knotgrass)	<p>Apply 1.5 quarts of Sharda Glyphosate 41 SL in 5 to 10 gallons of water per acre. Apply when water bermudagrass is actively growing and 12 to 18 inches in length. Allow 7 or more days before tilling, flushing or flooding the field. Fall applications only - Apply 1 quart of Sharda Glyphosate 41 SL in 5 to 10 gallons of water per acre. Fallow fields must be tilled prior to application. Apply prior to frost on water bermudagrass that is actively growing and 12 to 18 inches in length. Allow 7 or more days before tillage.</p>
Bindweed, field	<p>For control, apply 4 to 5 quarts of Sharda Glyphosate 41 SL per acre west of the Mississippi River and 3 to 4 quarts east of the Mississippi River. Apply when the weeds are actively growing and are at or beyond full bloom. Do not treat when weed is under drought stress as good soil moisture is necessary for active growth. For best results, apply in late summer or fall. Fall treatments must be applied before a killing frost. Allow 7 or more days after application before tillage.</p> <p>Also for control, apply 2 quarts of Sharda Glyphosate 41 SL plus 0.5 pound a.i. of Banvel® or dicamba in 10 to 20 gallons of water per acre. At these rates, apply using ground application only.</p> <p>The following tank mixtures with 2,4-D may be applied using aerial application equipment (except in California) in fallow and reduced tillage systems only.</p> <p>For suppression on irrigated agricultural land, apply 1 to 2 quarts of Sharda Glyphosate 41 SL plus 1 pound a.i. of 2,4-D in 10 to 20 gallons of water per acre with ground equipment only. Applications must be made following harvest or in fall fallow ground when the bindweed is actively growing and the majority of runners are 12 inches or more in length. The use of at least one irrigation will promote active bindweed growth.</p> <p>For suppression, apply 16 fluid ounces of Sharda Glyphosate 41 SL plus 0.5 pound a.i. of 2,4-D in 3 to 10 gallons of water per acre for ground applications and 3 to 5 gallons of water per acre for aerial applications. Applications must be delayed until maximum emergence has occurred and when vines are between 6 to 18 inches in length.</p> <p>In California only, apply 1 to 5 quarts of Sharda Glyphosate 41 SL per acre. Actual rate needed for suppression or control will vary within this range depending on local conditions.</p> <p>For suppression on irrigated land where annual tillage is performed, apply 1 quart of Sharda Glyphosate 41 SL in 3 to 10 gallons of water per acre. Apply to actively growing bindweed that has reached a length of 12 inches or greater. Allow maximum weed emergence and runner growth. Do not treat when weeds are under drought stress as good soil moisture is necessary for active growth. Allow 3 or more days after application before tillage.</p>
Bluegrass, Kentucky / Bromegrass, smooth / Orchard-grass	<p>Apply 2 quarts of Sharda Glyphosate 41 SL in 10 to 40 gallons of water per acre when the grasses are actively growing and most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 1 to 1.5 quarts of Sharda Glyphosate 41 SL in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height. Allow 7 or more days after application before tillage.</p>

23 of 67

Weed	Directions for Use
Orchardgrass (sods going to no-till corn)	Apply 1 to 1.5 quarts of Sharda Glyphosate 41 SL per acre in 3 to 10 gallons of water per acre. Apply to orchardgrass that is a minimum of 12 inches tall for spring applications and 6 inches tall for fall applications. Allow at least 3 days following application before planting. A sequential application of atrazine will be necessary for optimum results.
Blueweed, Texas	Apply 4 to 5 quarts of Sharda Glyphosate 41 SL per acre west of the Mississippi River and 3 to 4 quarts per acre east of the Mississippi River. Apply when weed is actively growing and is at or beyond full bloom. Do not treat when weed is under drought stress as good soil moisture is necessary for active growth. New leaf development indicates active growth. For best results, apply in late summer or fall. Fall treatments must be applied before a killing frost. Allow 7 or more days after application before tillage.
Brackenfern	Apply 3 to 4 quarts of Sharda Glyphosate 41 SL per acre as a broadcast spray or as a 1 to 1.5 percent solution with hand-held equipment. Apply to fully expanded fronds which are at least 18 inches long.
Bursage, woollyleaf	For control, apply 2 quarts of Sharda Glyphosate 41 SL plus 0.5 lb. a.i. of Banvel® or dicamba per acre. For partial control, apply 1 quart of Sharda Glyphosate 41 SL plus 0.5 lb. a.i. of Banvel® or dicamba per acre. Apply when plants are producing new active growth which has been initiated by moisture for at least 2 weeks and when plants are at or beyond flowering.
Canarygrass, reed/Timothy / Wheatgrass, western	Apply 2 to 3 quarts of Sharda Glyphosate 41 SL per acre. For best results, apply to actively growing plants when most have reached the boot-to-head stage of growth. Allow 7 or more days after application before tillage.
Cogongrass	Apply 3 to 5 quarts of Sharda Glyphosate 41 SL in 10 to 40 gallons of water per acre. Apply when Cogongrass is at least 18 inches tall and actively growing in late summer or fall. Allow 7 or more days after application before tillage or mowing. Due to uneven stages of growth and the dense nature of vegetation preventing good spray coverage, repeat treatments may be necessary to maintain control.
Dandelion/Dock, curly	Apply 3 to 5 quarts of Sharda Glyphosate 41 SL per acre when plants are actively growing and most have reached the early bud stage of growth. Allow 7 or more days after application before tillage. Also for control, apply 16 fluid ounces of Sharda Glyphosate 41 SL plus 0.5 pound a.i. 2,4-D in 3 to 10 gallons of water per acre.
Dogbane, hemp	<p>Apply 4 quarts of Sharda Glyphosate 41 SL per acre. Apply when actively growing and when most weeds have reached the late bud to flower stage of growth. Following crop harvest or mowing, allow weeds to regrow to a mature stage prior to treatment. For best results, apply in late summer or fall. Allow 7 or more days after application before tillage.</p> <p>For suppression, apply 16 fluid ounces of Sharda Glyphosate 41 SL plus 0.5 pound a.i. of 2,4-D in 3 to 10 gallons of water per acre for ground applications and 3 to 5 gallons of water per acre for aerial applications. Delay applications until maximum emergence of dogbane has occurred.</p>

240967

Weed	Directions for Use
Fescue, tall	<p>Apply 3 quarts of Sharda Glyphosate 41 SL in 10 to 40 gallons of water per acre to actively growing plants when most have reached boot-to-early seedhead stage of development.</p> <p>Fall applications only-Apply 1 quart of Sharda Glyphosate 41 SL in 3 to 10 gallons of water per acre. Apply to fescue in the fall when actively growing and plants have 6 to 12 inches of new growth. Allow 7 or more days after application before tillage. A sequential application of 1 pint per acre of Sharda Glyphosate 41 SL will improve long-term control and control seedlings germinating after fall treatments or the following spring.</p>
Guineagrass	<p>Apply 3 quarts of Sharda Glyphosate 41 SL per acre or use a 1 percent solution with hand-held equipment. Apply to actively growing guineagrass when most has reached at least the 7-leaf stage of growth. Ensure thorough coverage when using hand-held equipment. Allow 7 or more days after application before tillage.</p>
Johnsongrass/Ryegrass, perennial	<p>Apply 1 to 3 quarts of Sharda Glyphosate 41 SL per acre. In annual cropping systems apply 1 to 2 quarts of Sharda Glyphosate 41 SL per acre. Apply 1 quart of Sharda Glyphosate 41 SL in 3 to 10 gallons of water per acre. Use 2 quarts of Sharda Glyphosate 41 SL when applying 10 to 40 gallons of water per acre. In noncrop, or areas where annual tillage (no-till) is not performed, apply 2 to 3 quarts of Sharda Glyphosate 41 SL in 10 to 40 gallons of water per acre. For best results, apply to actively growing plants when most have reached the boot-to-head stage of growth or in the fall prior to frost. Allow 7 or more days after application before tillage. Do not tank-mix with residual herbicides when using the 1 quart per acre rate.</p> <p>For burndown of Johnsongrass, apply 1 pint per acre in 3 to 10 gallons of water per acre before the plants reach a height of 12 inches. For this use, allow at least 3 days after treatment before tillage.</p> <p>Spot treatment (partial control or suppression) - Apply a 1 percent solution of Sharda Glyphosate 41 SL when Johnsongrass is 12 to 18 inches in height. Coverage must be uniform and complete.</p>
Kikuyugrass	<p>Apply 2 to 3 quarts of Sharda Glyphosate 41 SL per acre. Spray when most kikuyugrass is at least 8 inches in height (3 or 4-leaf stage of growth) and actively growing. Allow 3 or more days after application before tillage.</p>
Knapweed/Horseradish	<p>Apply 4 quarts of Sharda Glyphosate 41 SL per acre. Apply when actively growing and when most weeds have reached the late bud to flower stage of growth. Following crop harvest or mowing, allow weeds to regrow to a mature stage prior to treatment. For best results, apply in late summer or fall. Allow 7 or more days after application before tillage.</p>
Lantana	<p>Apply Sharda Glyphosate 41 SL as a 1 to 1.25 percent solution using hand-held equipment only. Apply to actively growing lantana at or beyond the bloom stage of growth. Use the higher application rate for plants that have reached the woody stage of growth. Allow 7 or more days after application before tillage.</p>
Milkweed, common	<p>Apply 3 quarts of Sharda Glyphosate 41 SL per acre. Apply when actively growing and most of the milk-weed has reached the late bud to flower stage of growth. Following small grain harvest or mowing, allow milkweed to regrow to a mature stage prior to treatment. Allow 7 or more days after application before tillage.</p>



Weed	Directions for Use
Muhly, wirestem	Apply 1 to 2 quarts of Sharda Glyphosate 41 SL per acre. Use 1 quart of Sharda Glyphosate 41 SL in 3 to 10 gallons of water per acre. Use 2 quarts of Sharda Glyphosate 41 SL when applying 10 to 40 gallons of water per acre or in pasture, sod, or noncrop areas. Spray when the wirestem muhly is 8 inches or more in height and actively growing. Do not till between harvest and fall applications or in the fall or spring prior to spring applications. Allow 3 or more days after application before tillage. Sharda Glyphosate 41 SL will not provide residual control of wirestem muhly from seeds which germinate after application of Sharda Glyphosate 41 SL. Do not tank mix with residual herbicides when using the 1 quart per acre rate.
Nightshade, silverleaf	For control, apply 2 quarts of Sharda Glyphosate 41 SL in 3 to 10 gallons of water per acre. Applications must be made when at least 60 percent of the plants have berries. Fall treatments must be applied before a killing frost. Allow 7 or more days after application before tillage. Do not treat when weed is under drought stress as good soil moisture is necessary for active growth.
Nutsedge; purple, yellow	<p>Apply 3 quarts of Sharda Glyphosate 41 SL per acre as a broadcast spray, or apply a 1 to 2 percent solution from handheld equipment to control existing nutsedge plants and immature nutlets attached to treated plants. Treat when plants are in flower or when new nutlets can be found at rhizome tips. Nutlets which have not germinated will not be controlled and may germinate following treatment. Repeat treatments will be required for long-term control of ungerminated tubers.</p> <p>Sequential applications of 1 to 2 quarts of Sharda Glyphosate 41 SL in 3 to 10 gallons of water per acre will provide control. Make applications when a majority of the plants are in the 3 to 5-leaf stage (less than 6 inches tall). Repeat this application, as necessary, when newly emerging plants reach the 3 to 5-leaf stage. Subsequent applications will be necessary for long-term control.</p> <p>For suppression to partial control of existing plants, apply 1 pint to 2 quarts of Sharda Glyphosate 41 SL per acre in 3 to 40 gallons of water per acre. Treat when plants have 3 to 5 leaves and most are less than 6 inches tall. Repeat treatments will be required to control subsequent emerging plants or regrowth of existing plants. Wait 7 days after treatment before tillage or mowing.</p>
Pampasgrass/ice plant	Apply Sharda Glyphosate 41 SL as a 1.5 to 2 percent solution using hand-held equipment. Apply to plants that are actively growing. Pampasgrass must be at or beyond the boot stage of growth. Thorough coverage is necessary for best control.
Phragmites	For partial control of phragmites in Florida and the counties of other states bordering the Gulf of Mexico, apply 5 quarts per acre as a broadcast spray or apply a 2 percent solution from hand-held equipment, in other areas of the U.S., apply 3 quarts per acre as a broadcast spray or apply a 1 percent solution from hand-held equipment for partial control. For best results, treat during late summer or fall months or when plants are actively growing and in full bloom. Treatment before or after this stage may lead to reduced control. Due to the dense nature of the vegetation, which may prevent good spray coverage or uneven stages of growth, repeat treatments may be necessary to maintain control. Visual control symptoms will be slow to develop.

260967

Weed	Directions for Use
Quackgrass, In Annual Cropping Systems, or in Pastures and Sods Followed by Deep Tillage	Apply 1 to 2 quarts of Sharda Glyphosate 41 SL per acre. Do not tank mix with residual herbicides when using the 1 quart rate. Spray when quackgrass is 6 to 8 inches in height and actively growing. Do not till between harvest and fall applications or in fall or spring prior to spring application. Allow 3 or more days after application before tillage. In pastures or sods, for best results use a moldboard plow.
Quackgrass, Pasture or Sod or Other Noncrop Areas Where Deep Tillage is Not Planned Following Application	Apply 2 to 3 quarts in 10 to 40 gallons of water per acre. Spray when the quackgrass is greater than 8 inches tall and actively growing. Do not till between harvest and fall application or in fall or spring prior to spring application. Allow 3 or more days after application before tillage.
Redvine	For suppression, apply 24 fluid ounces of Sharda Glyphosate 41 SL per acre at each of two applications 7 to 14 days apart or a single application of 2 quarts per acre. Apply specified rates in 5 to 10 gallons of water per acre. Apply in late September or early October to actively growing plants, which are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least 1 week before a killing frost.
Reed, giant	For control of giant reed, apply a 2 percent solution of Sharda Glyphosate 41 SL when plants are actively growing. Best results are obtained when applications are made in late summer to fall.
Smartweed, swamp	Apply 3 to 5 quarts of Sharda Glyphosate 41 SL per acre when plants are actively growing and most have reached the early bud stage of growth. Allow 7 or more days after application before tillage.  Also for control, apply 16 fluid ounces of Sharda Glyphosate 41 SL plus 0.5 pound active ingredient of 2,4-D in 3 to 10 gallons of water per acre in the late summer or fall. Apply when plants are actively growing and most have reached the early bud stage of growth. Allow 7 or more days after application before tillage.
Spurge, leafy	For suppression, apply 16 fluid ounces of Sharda Glyphosate 41 SL plus 0.5 pound active ingredient 2,4-D in 3 to 10 gallons of water per acre in the late summer or fall. Apply when plants are actively growing. If mowing has occurred prior to treatment, apply when most of the plants are 12 inches tall. Allow 7 or more days after application before tillage.
Starthistle, yellow	Best results are obtained when applications are made during periods of active growth, including the rosette, bolting and early flower stages. For spray-to-wet applications, apply Sharda Glyphosate 41 SL as a 2 percent solution. For broadcast applications, apply 2 quarts per acre in 10 to 40 gallons per acre of water carrier.
Sweet Potato, wild/Thistle, artichoke	Apply Sharda Glyphosate 41 SL as a 2 percent solution using hand-held equipment. Apply to actively growing weeds that are at or beyond the bloom stage of growth. Repeat applications may be required. Allow the plant to reach the specified stage of growth before retreatment. Allow 7 or more days before tillage.
Thistle, Canada	Apply 2 to 3 quarts of Sharda Glyphosate 41 SL per acre. Apply to actively growing thistles when most are at or beyond the bud stage of growth. After harvest, mowing or tillage in the late summer or fall, allow at least 4 weeks for initiation of active growth and rosette development prior to the application of Sharda Glyphosate 41 SL. Fall treatments must be applied before a killing frost. Allow 3 or more days after application before tillage.

Weed	Directions for Use
Thistle, Canada (cont'd.)	For suppression of Canada thistle, apply 1 quart per acre of Sharda Glyphosate 41 SL, or 1 pint of Sharda Glyphosate 41 SL plus 0.5 pound a.i. 2,4-D per acre in 3 to 10 gallons of water per acre in the late summer or fall after harvest, mowing or tillage. Allow rosette regrowth to a minimum of 6 inches in diameter before treating. Applications can be made as long as leaves are still green and plants are actively growing at the time of application. Allow 3 or more days after application before tillage.
Torpedograss	Apply 4 to 5 quarts of Sharda Glyphosate 41 SL per acre to provide partial control of torpedograss. Apply to actively growing torpedograss when most plants are at or beyond the seedhead stage of growth. Repeat applications will be required to maintain control. Fall treatments must be applied before frost. Allow 7 or more days after application before tillage.
Trumpetcreeper	For control, apply 2 quarts of Sharda Glyphosate 41 SL per acre in 5 to 10 gallons of water per acre. Apply to actively growing plants in late September or October, which are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least 1 week before a killing frost.
Other perennials listed on this label	Apply 3 to 5 quarts of Sharda Glyphosate 41 SL per acre. Apply when actively growing and most have reached early head or early bud stage of growth. Allow 7 or more days after application before tillage.

**WOODY BRUSH AND TREES**

When applied as specified under the conditions described, Sharda Glyphosate 41 SL CONTROLS or PARTIALLY CONTROLS the following woody brush, plants and trees:

Alder ( <i>Alnus spp.</i> )	Coyote brush ( <i>Baccharis consanguinea</i> )	Sugar ( <i>Acer saccharum</i> )	Salmonberry ( <i>Rubus spectabilis</i> )
Ash* ( <i>Fraxinus spp.</i> )	Creeper, Virginia* ( <i>Parthenocissus quinquefolia</i> )	Vine* ( <i>Acer circinatum</i> )	Salt cedar ( <i>Tamarix spp.</i> )
Aspen, quaking ( <i>Populus tremuloides</i> )	Dewberry ( <i>Rubus, trivialis</i> )	Monkey Flower* ( <i>Mimulus guttatus</i> )	Sassafras ( <i>Sassafras aibidum</i> )
Bearmat (Bearclover) ( <i>Chamaebatia foliolosa</i> )	Dogwood* ( <i>Comus spp.</i> )	Oak, Black* ( <i>Quercus velutina</i> )	Sourwood ( <i>Oxydendrum arboretum</i> )
Beech ( <i>Fagus grandifolia</i> )	Elderberry ( <i>Sambucus spp.</i> )	Oak, Northern Pin ( <i>Quercus palustris</i> )	Sumac, Poison* ( <i>Rhus vernix</i> )
Birch ( <i>Betula spp.</i> )	Elm* ( <i>Ulmus spp.</i> )	Oak, Post ( <i>Quercus stellata</i> )	Sumac, Smooth* ( <i>Rhus glabra</i> )
Blackberry ( <i>Rubus spp.</i> )	Eucalyptus ( <i>Eucalyptus spp.</i> )	Oak, Red ( <i>Quercus rubra</i> )	Sumac, winged* ( <i>Rhus copallina</i> )
Blackgum ( <i>Nyssa spp.</i> )	Gorse ( <i>Ulex europaeus</i> )	Oak, Southern Red ( <i>Quercus falcata</i> )	Sweetgum ( <i>Liquidambar styraciflua</i> )
Bracken ( <i>peridium spp.</i> )	Hasardia* ( <i>Haplopappus squamosus</i> )	Oak, White* ( <i>Quercus alba</i> )	Swordfern* ( <i>polystichum munitum</i> )
Broom, French ( <i>Cytisus monspessulanus</i> )	Hawthorn ( <i>Crataegus spp.</i> )	Persimmon* ( <i>Diospyros spp.</i> )	Tallowtree, Chinese ( <i>Saplum sebiferum</i> )
Broom, Scotch ( <i>Cytisus scoparius</i> )	Hazel ( <i>Corylus spp.</i> )	Pine ( <i>Pinus spp.</i> )	Tan Oak ( <i>Lithocarpus densiflorus</i> )
Buckwheat, California* ( <i>Eriogonum fasciculatum</i> )	Hickory* ( <i>Carya spp.</i> )	Poison Ivy ( <i>Rhus radicans</i> )	Thimbleberry ( <i>Rubus parviflorus</i> )

Cascara* ( <i>Rhamnus purshiana</i> )	Holly, Florida/Brazilian Peppertree* ( <i>Schinus terebinthifolius</i> )	Poison Oak ( <i>Rhus toxicodendron</i> )	Tobacco, tree* ( <i>Nicotiana glauca</i> )
Catsclaw* ( <i>Acacia greggi</i> )	Honeysuckle ( <i>Lonicera spp.</i> )	Poplar, yellow* ( <i>Liriodendron tulipifera</i> )	Trumpetcreeper ( <i>Campsis radicans</i> )
Ceanothus* ( <i>Acacia greggi</i> )	Hornbeam, American* ( <i>Carpinus caroliniana</i> )	Raspberry ( <i>Rubus, spp.</i> )	Waxmyrtle, southern* ( <i>Myrica cerifera</i> )
Ceanothus* ( <i>Ceanothus spp.</i> )	Kudzu ( <i>Pueraria lobata</i> )	Redbud, eastern ( <i>Cercis Canadensis</i> )	Willow ( <i>Salix spp.</i> )
Chamise ( <i>Adenostoma fasciculatum</i> )	Locust, black* ( <i>Robinia pseudoacacia</i> )	Rose, multiflora ( <i>Rosa multiflora</i> )	
Cherry, Bitter ( <i>Prunus emarginata</i> )	Madrone ( <i>Arutus menziesli</i> )	Russian-Olive ( <i>Elaeagnus angustifolia</i> )	
Cherry, Black ( <i>Prunus serotina</i> )	Manzanita ( <i>Arctostaphylos spp.</i> )	Sage: black, white ( <i>Salvia spp.</i> )	
Cherry, Pin ( <i>Prunus pensylvanica</i> )	Maple, Red ( <i>Acer rubrum</i> )	Sagebrush, California ( <i>Artemisia californica</i> )	

\*Partial Control

\*\*See below for control or partial control instructions.

**NOTE:** If brush has been mowed or tilled or trees have been cut, treatment will not be effective until regrowth has reached the specified stages of growth.

Apply Sharda Glyphosate 41 SL when plants are actively growing and, unless otherwise directed, after full leaf expansion. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation.

In arid areas, best results are obtained when application is made in the spring to early summer when brush species are at high moisture content and are flowering.

Ensure thorough coverage when using hand-held equipment. Symptoms may not appear prior to frost or senescence with fall treatments.

Allow 7 or more days after application before tillage, moving or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if treatments are made following a frost.

See "DIRECTIONS FOR USE" AND "MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS" sections of this label for labeled uses and specific application instructions.

Apply Sharda Glyphosate 41 SL as follows to control or partially control the following woody brush and trees:

Weed	Directions for Use
Alder/Dewberry/Honeysuckle/Post Oak/Raspberry	Apply 3 to 4 quarts per acre of Sharda Glyphosate 41 SL as a broadcast spray or as a .1 to 1.5 percent solution with hand-held equipment.
Aspen, quaking/Cherry: bitter, black, pin/Hawthorn/Oak, southern red/Sweetgum/Trumpetcreeper	Apply 2 to 3 quarts of Sharda Glyphosate 41 SL per acre as a broadcast spray or as a 1 to 1.5 percent solution with hand-held equipment.
Birch/ Elderberry/ Hazel/ Salmonberry/ Thimbleberry	Apply 2 quarts per acre of Sharda Glyphosate 41 SL as a broadcast spray or as a 1 percent solution with hand-held equipment.

Weed	Directions for Use
Blackberry	Apply 3 to 4 quarts per acre of Sharda Glyphosate 41 SL as a broadcast spray, or 1 to 1.5 percent solution with hand-held equipment. Make application after plants have reached full leaf maturity. Best results are obtained when applications are made in late summer or fall. After berries have set or dropped in late fall, blackberry can be controlled by applying a 3/4 percent solution of Sharda Glyphosate 41 SL with hand-held equipment. For control of blackberries after leaf drop and until killing frost or as long as stems are green, apply 3 to 4 quarts of Sharda Glyphosate 41 SL in 10 to 40 gallons of water per acre.
Broom: French, Scotch	Apply a 1.5 to 2 percent solution with hand-held equipment.
Buckwheat, California/Hasardia/Monkey Flower/ Tobacco, tree	For partial control of these species, apply a 1 to 2 percent solution of Sharda Glyphosate 41 SL as a foliar spray with hand-held equipment. Thorough coverage of foliage is necessary for best results.
Catsclaw	For partial control, apply as a 1 to 1.5 percent solution with hand-held equipment.
Coyote Brush	For control, apply a 1.5 to 2 percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed.
Eucalyptus	For control of eucalyptus resprouts, apply a 2 percent solution of Sharda Glyphosate 41 SL with hand-held equipment when resprouts are 6 to 12 feet tall. Ensure complete coverage. Apply when plants are growing actively. Avoid application to drought-stressed plants.
Kudzu	For control, apply 4 quarts of Sharda Glyphosate 41 SL per acre as a broadcast spray or as a 2 percent solution with hand-held equipment. Repeat applications will be required to maintain control.
Madrone resprouts	For suppression or partial control, apply a 2 percent solution of Sharda Glyphosate 41 SL to resprouts less than 3 to 6 feet tall. Best results are obtained with spring/early summer treatments.
Maple, red	For control, apply as a 1 to 1.5 percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed. For partial control, apply 2 to 4 quarts of Sharda Glyphosate 41 SL per acre as a broadcast spray.
Maple, sugar/Oak, northern pin/Oak, red	For control, apply as a 1 to 1.5 percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed.
Poison Ivy/Poison Oak	For control, apply 4 to 5 quarts of Sharda Glyphosate 41 SL per acre as a broadcast spray or as a 2 percent solution with hand-held equipment. Repeat applications may be required to maintain control. Fall treatments must be applied before leaves lose green color.
Rose, multiflora	For control, apply 2 quarts of Sharda Glyphosate 41 SL per acre as a broadcast spray or as a 1 percent solution with hand-held equipment. Treatments must be made prior to leaf deterioration by leaf-feeding insects.
Sage, black/Sagebrush, California/Chamfse/Tallowtree, Chinese	For control of these species, apply a 1 percent solution of Sharda Glyphosate 41 SL as a foliar spray with hand-held equipment. Thorough coverage of foliage is necessary for best results.
Tan oak resprouts	For suppression or partial control, apply a 2 percent solution of Sharda Glyphosate 41 SL to resprouts less than 3 to 6 feet tall. Best results are obtained with fall applications.
Weed	Directions for Use
Willow	For control, apply 3 quarts of Sharda Glyphosate 41 SL per acre as a broadcast spray or as a 1 percent solution with hand-held equipment.

Other Woody Brush and Trees listed on this label	For partial control, apply 2 to 5 quarts of Sharda Glyphosate 41 SL per acre as a broadcast spray or as a 1 to 2 percent solution with hand-held equipment.
--	---

**NONCROP USES**

See "PRODUCT INFORMATION" and "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" sections of this label for essential product performance information and the following "NONCROP" sections for specific uses.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF SPRAY WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE TURFGRASSES, TREES, SHRUBS OR OTHER DESIRABLE VEGETATION SINCE SEVERE DAMAGE OR DESTRUCTION MAY RESULT.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seeds. Where repeat applications are necessary, do not exceed 10.6 quarts of Sharda Glyphosate 41 SL per acre per year.

Sharda Glyphosate 41 SL does not provide residual weed control. For subsequent residual weed control, it is directed that a residual herbicide program specified on this label be used. Read and carefully observe the precautionary statements and all other information appearing on the labels of all herbicides used.

**INDUSTRIAL, RECREATIONAL AND PUBLIC AREAS**

When applied as directed for "NONCROP USES", under conditions described, Sharda Glyphosate 41 SL controls annual and perennial weeds listed on this label growing in areas such as:

Airports	Parks
Around Farm, Ranch, Commercial or Industrial Structures	Petroleum & Other Tank Farms
Around Ornamental Gardens	Pumping Installations
Around Ornamental Trees & Shrubs	Pipeline, Power, Telephone & Utility Rights-of-Way
Ditch Banks	Preplant to Turf & Ornamental Plants
Driveways & Ramps	Railroads
Dry Ditches & Canals Fences & Fencerows	Schools
Golf Courses	Sidewalks
Gravel or Ground Bark Mulches	Similar Sites
Habitat Restoration & Management Areas	Storage Areas
Highways & Roadsides (including aprons, medians & guardrails)	Uncropped Farmstead Areas
Industrial Plant Sites	Utility Substations
Lanes, Trails & Access Roads	Vacant Lots & Wastelands
Lumberyards	
Parking Areas	

For specific rates of application and instructions for control of particular annual weeds, perennial weeds, woody brush and trees, see the "WEEDS CONTROLLED" section of this label. These applications may be made to large affected areas or as spot treatments. For use in small areas, see alternative instructions below under "Small Area Treatment With Hand-held Sprayers". Sharda Glyphosate 41 SL is a nonselective herbicide that is diluted and applied to the foliage of actively growing weeds as a spot or broadcast application. It is absorbed by the leaves and moves throughout the stem and roots to control the entire plant. Visible symptoms may require a week or more to appear, with burndown usually occurring in 2 to 4 weeks.

310967

Symptoms are a gradual wilting and yellowing of the sprayed plant followed by deterioration of both shoots and roots. Sharda Glyphosate 41 SL has no herbicidal activity in the soil and will not wash or leach to affect nearby vegetation. Any ornamental species may be planted in treated areas 7 days or more after application. For most effective results, delay mowing, clipping, tilling, planting or sodding of treated areas for at least 7 days after application. This allows time for Sharda Glyphosate 41 SL to move within the plant.

Unless the "Agricultural Use Requirements" on this label are observed, the following restrictions apply: Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

SHARDA GLYPHOSATE 41 SL CAN INJURE OR DESTROY ALL VEGETATION CONTACTED. WHEN USED AS A SPOT TREATMENT IN LAWNS, ALL VEGETATION CONTACTED WILL BE DAMAGED. AVOID SPRAY DRIFT CONTACT WITH DESIRABLE LAWN GRASSES, FLOWERS, VEGETABLES, SHRUBS OR TREES. DO NOT CONTACT GREEN BARK OF TREES OR SHRUBS. IF DESIRABLE VEGETATION IS CONTACTED, WASH IMMEDIATELY WITH WATER.

Depending on the type of noncrop application, Sharda Glyphosate 41 SL may be applied with boom equipment, high-volume spray equipment and hand-held sprayers as described in the respective portions of the "APPLICATION EQUIPMENT and TECHNIQUES" section of the label. Additionally, the product may be applied with recirculating sprayers, shielded applicators, or wiper applicators in any noncrop site specified on this label. See the "Selective Equipment" part of "APPLICATION EQUIPMENT and TECHNIQUES" section of this label for information on proper use and calibration of this equipment.

Small Area Treatment With Hand-held Sprayers - Add 3 to 6 fluid ounces of Sharda Glyphosate 41 SL to 1 gallon of clean water. Use the lower rate for many grasses and annual weeds. Use the higher specified rate for control of perennials and brush. Use pump-up sprayer, backpack sprayer or other sprayer suitable for small areas. Adjust equipment to deliver a coarse spray pattern. USE OF HOSE-END SPRAYERS OR SPRINKLER-TYPE DEVICES MAY RESULT IN POOR AND/OR ERRATIC RESULTS.

**TANK MIXTURES FOR INDUSTRIAL SITES AND FORESTRY SITE PREPARATIONS**

Use on industrial sites including airports, industrial plants, lumberyards, petroleum tank farms, pumping stations, railroads, roadsides, storage areas or other similar sites where bare ground is desired.

This tank mixture may also be used as a site preparation treatment for sites to be planted to jack pine, loblolly pine, red pine, slash pine and Virginia pine. When applied as directed for "NONCROP USES" under the conditions described, Sharda Glyphosate 41 SL plus Oust® provides control of annual weeds listed in the "WEEDS CONTROLLED" section of the label for Sharda Glyphosate 41 SL and Oust®, and control or partial control of the perennial weeds listed below.

Apply 1 to 2 quarts of Sharda Glyphosate 41 SL with 2 to 4 ounces of Oust® in 10 to 40 gallons of spray solution per acre as a broadcast spray to actively growing weeds. This mixture may be applied by aerial equipment in site prep operations. When applied by air, use the specified rates in 5 to 15 gallons of spray solution per acre. Sharda Glyphosate 41 SL plus Oust® tank mixtures may not be applied by air in California.

For control of annual weeds, use the lower rates of these products.

For control of the listed perennial weeds, use the higher rates of both products. For partial control, use the lower rates.

Bahiagrass ( <i>Paspalum notatum</i> )	Dock, Curly ( <i>Rumex crispus</i> )	Johnsongrass** ( <i>Sorghum halepense</i> )	Trumpetcreeper* ( <i>Campsis radicans</i> )
Bermudagrass* ( <i>Cynodon dactylon</i> )	Dogfennel ( <i>Eupatorium capilliflorum</i> )	Poorjoe** ( <i>Diodia teres</i> )	Vaseygrass ( <i>Paspalum urvillei</i> )
Broomsedge ( <i>Andropogon virginicus</i> )	Fescue, tall ( <i>Festuca arundinaces</i> )	Quackgrass ( <i>Agropyron repens</i> )	Vervain, blue ( <i>Verbena hastate</i> )

\*Suppression at the higher rates only.

\*\*Control at the lower rates.

Read and carefully observe the precautionary statements and all other information appearing on the labels of all herbicides used.

**TANK MIXTURES NONCROP SITES**

When applied as a tank mixture, Sharda Glyphosate 41 SL provides control of the emerged annual weeds and partial control of the emerged perennial weeds listed in this label. When applied as a tank mixture, the following residual herbicides will provide preemergence control of the weeds listed in the individual product labels.

- Sharda Glyphosate 41 SL plus DIURON®
- Sharda Glyphosate 41 SL plus KROVAR® I®
- Sharda Glyphosate 41 SL plus KROVAR® II®
- Sharda Glyphosate 41 SL plus RONSTAR® 50WP®
- Sharda Glyphosate 41 SL plus PRINCEP®\* CALIBER®\* 90
- Sharda Glyphosate 41 SL plus SIMAZINE 4L, 80W or 90DF
- Sharda Glyphosate 41 SL plus SURFLAN®\* 75W or AS

See the "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" section of this label before preparing these tank mixtures.

Read and carefully observe the label claims, precautionary statements, specified rates and all other information on the labels of all products used in these tank mixtures. Use according to the most restrictive label directions for each product in the mixture.

**CONTROL OF EMERGED WEEDS**

**Note:** For backpack sprayer and handgun applications, see the "HAND-HELD AND HIGH VOLUME EQUIPMENT" section for specified rates.

**Annual Weeds** - Apply 1 quart per acre of Sharda Glyphosate 41 SL in these tank mixtures when weeds are less than 6 inches tall and 1.5 quarts per acre when weeds are more than 6 inches tall.

**Perennial Weeds** - For partial control of perennial weeds using these tank mixtures, apply 2 to 5 quarts per acre of Sharda Glyphosate 41 SL. Follow the directions in the "WEEDS CONTROLLED" section of this label for stage of growth and rate of application for specific perennial weeds.

**PREEMERGENCE WEED CONTROL**

For preemergence weed control, refer to the individual product labels for specific noncrop sites, rates, carrier volumes and precautionary statements. Mix only the quantity of spray solution which can be used during the same day. Do not allow these tank mixtures to stand overnight as this may result in reduced weed control.

**FARMSTEAD WEED CONTROL**

When applied as directed for "NONCROP USES", under conditions described, Sharda Glyphosate 41 SL controls undesirable vegetation listed on this label around areas such as farmstead building foundations, along and in fences, shelterbelts and for nonselective farmstead weed control. For specific rates of application and instructions for control of various annual and perennial weeds, see the "WEEDS CONTROLLED" section of this label.

**FARM DITCHES**

Sharda Glyphosate 41 SL will suppress perennial grasses along farm ditches. Apply Sharda Glyphosate 41 SL at a rate of 6 to 8 fluid ounces per acre. Use 8 fluid ounces per acre when treating tall (coarse) fescue, fine fescue, orchardgrass or quackgrass covers. For best suppression of these species, add ammonium sulfate at a rate of 1.7 pounds per 10 gallons of spray solution. Use 6 fluid ounces per acre without ammonium sulfate when treating Kentucky bluegrass. Apply treatments in 10 to 20 gallons of spray solution per acre to actively growing perennial grass covers. For best spray distribution and coverage, use flat fan nozzles. Where broadleaf weed control or suppression is desired, tank mix Sharda Glyphosate 41 SL with an appropriate, labeled broadleaf weed herbicide.

**CONSERVATION RESERVE PROGRAM (CRP ACRES)**

Sharda Glyphosate 41 SL can be used to control undesirable vegetation when rotating out of CRP acres or to suppress competitive growth and seed production of undesirable vegetation in CRP acres. For specific rates of application for various annual and perennial weeds, see the "WEEDS CONTROLLED" section of this label. CRP applications may be made with wiper applicators or conventional spray equipment. For selective applications with broadcast spray equipment, apply 12 to 16 ounces per acre of Sharda Glyphosate 41 SL in early spring before desirable CRP grasses, such as crested and tall wheatgrass, break



33 of 67

dormancy and initiate green growth. Late fall applications can be made after desirable perennial grasses have reached dormancy. Some stunting of CRP perennial grasses will occur if applications are made when plants are not dormant.

**DORMANT RANGELAND**

Sharda Glyphosate 41 SL will control or suppress many weeds, including downy brome, cheat grass, cereal rye, medusahead rye and jointed goatgrass in dormant rangeland. Apply 8 to 16 ounces per acre of Sharda Glyphosate 41 SL in the early spring when the weeds have greened up, but desirable grasses, such as crested and tall wheatgrass are still truly dormant. Slight discoloration of the desirable grasses may occur, but they will regreen and regrow under moist soil conditions as effects of Sharda Glyphosate 41 SL wear off. Do not use additional surfactant or ammonium sulfate when spraying dormant rangeland grasses with Sharda Glyphosate 41 SL.

**HABITAT MANAGEMENT**

Sharda Glyphosate 41 SL is directed for the restoration and/or maintenance of native habitats and in wildlife management areas. Apply as specified in the "NONCROP USES" section of this label.

Use	Directions for Use
Habitat Restoration and Maintenance	When applied as directed, exotic and other undesirable vegetation may be controlled in habitat management areas. Applications can be made to allow recovery of native plant species, prior to planting desirable native species, and for similar broad spectrum vegetation control requirements in habitat management areas. Spot treatments can be made to selectively remove unwanted plants for habitat maintenance and enhancement. For spot treatments, care must be exercised to keep spray off of desirable plants.
Wildlife Food Plots	Sharda Glyphosate 41 SL may be used as a site preparation treatment prior to planting wildlife food plots. Any wildlife food species may be planted after applying Sharda Glyphosate 41 SL, or native species may be allowed to repopulate the area. If tillage is needed to prepare a seedbed, wait 7 days after applying Sharda Glyphosate 41 SL before tilling.

**ORNAMENTALS, TREE NURSERIES AND CHRISTMAS TREES**

**SHARDA GLYPHOSATE 41 SL IS NOT DIRECTED FOR USE AS AN OVER-THE-TOP BROADCAST SPRAY ON ORNAMENTALS AND CHRISTMAS TREES.**

**Note:** Desirable plants may be protected from the spray solution by using shields or coverings made of cardboard or other impermeable material.

When applied as instructed for the conditions described for "NONCROP USES", Sharda Glyphosate 41 SL controls undesirable vegetation listed on this label prior to planting, within and around greenhouses and shadehouses, and as a postdirected spray around established ornamentals and Christmas trees.

For specific rates of application and instructions for control of various annual and perennial weeds, see the "WEEDS CONTROLLED" section of this label. Where repeat applications are necessary, do not exceed 10.6 quarts of Sharda Glyphosate 41 SL per acre per year.

Use	Directions for Use
Site Preparation	Following preplant applications of Sharda Glyphosate 41 SL, any ornamental, nursery species or Christmas tree species may be planted. Precautions must be taken to protect nontarget plants during site preparation applications.

34 of 67

Greenhouse/Shadehouse Use	Sharda Glyphosate 41 SL may be used to control weeds listed on this label which are growing in greenhouses. Desirable vegetation must not be present during application and air circulation fans must be turned off.
Postdirected Spray	Use as a postdirected spray around established woody ornamental species, nursery species or Christmas trees such as those listed below. Care must be exercised to avoid contact of spray, drift or mist with foliage of or green bark of established ornamental species.

Arborvitae ( <i>Thuja spp.</i> )	Fir ( <i>Abies spp.</i> <i>Pseudotsuga spp.</i> )	Magnolia ( <i>Magnolia spp.</i> )	Pine ( <i>Pinus spp.</i> )
Azalea ( <i>Rhododendron spp.</i> )	Johanna ( <i>Simmondsia chinensis</i> )	Maple ( <i>Acer spp.</i> )	Spruce ( <i>Picea spp.</i> )
Crabapple ( <i>Malus spp.</i> )	Hollies ( <i>Ilex spp.</i> )	Oak ( <i>Quercus spp.</i> )	Yew ( <i>Taxus spp.</i> )
Euonymus ( <i>Euonymus spp.</i> )	Lilac ( <i>Syringa spp.</i> )	Privet ( <i>Ligustrum spp.</i> )	

**BROADCAST APPLICATIONS FOR WEED CONTROL IN CHRISTMAS TREE PLANTATIONS**

**NOTE:** IF SHARDA GLYPHOSATE 41 SL IS IMPROPERLY APPLIED, IT HAS THE POTENTIAL TO CAUSE SEVERE INJURY TO CHRISTMAS TREES. FOLLOW ALL LABELED DIRECTIONS.

Sharda Glyphosate 41 SL may be applied as a broad-cast spray over established Christmas trees. To prevent drift onto nearby desirable crops or vegetation, ensure that adequate buffers are maintained.

The following Christmas tree species are approved for this application:

- Douglas Fir (*Pseudotsuga menziesii*)
- Fir species (*Abies spp.*)
- Spruce species (*Picea spp.*)

Do not apply Sharda Glyphosate 41 SL until trees have completed at least a full growing season since planting or transplanting. Do not apply within 1 full year prior to tree harvest. In the fall, applications may only be made after the formation of final conifer resting buds. Final resting buds must be in the dormant stage and fully hardened, if applications are made at any other time, unacceptable Christmas tree injury may occur.

Avoid spray pattern overlap, as injury may result.

Apply 1 quart of Sharda Glyphosate 41 SL per acre in 5 to 30 gallons of water per acre,

**NOTE:** ADDING SURFACTANTS, ADDITIVES CONTAINING SURFACTANTS, OR ANY OTHER ADDITIVES TO SHARDA GLYPHOSATE 41 SL MAY RESULT IN SEVERE CHRISTMAS TREE INJURY.

In some areas, Sharda Glyphosate 41 SL may be used at rates from 1 to 2 quarts per acre. Contact your local Sharda USA LLC representative for specific directions if you require rates that exceed 1 quart per acre.

Drift control additives are not directed as they may increase Christmas tree injury. Using other herbicides tank mixed with Sharda Glyphosate 41 SL is not directed as Christmas trees could be severely injured.

**SILVICULTURAL SITES AND RIGHTS-OF-WAY**

**NOTE:** NOT DIRECTED FOR USE AS AN OVER-THE-TOP BROADCAST SPRAY IN SILVICULTURAL NURSERIES

When applied as directed for "NONCROP USES" under conditions described, Sharda Glyphosate 41 SL controls undesirable vegetation listed on this label. Sharda Glyphosate 41 SL also suppresses or controls undesirable vegetation listed on this label when applied at specified rates for release of established coniferous species listed on this label. For specific rates of application and instructions for control of various brush, annual and perennial weeds, see the "WEEDS CONTROLLED" section of this label. For specific rates of application for release of listed coniferous species, see the "CONIFER RELEASE" part of this

35067

section of the label. Where repeat applications are necessary, do not exceed 10.6 quarts of Sharda Glyphosate 41 SL per acre per year.

**Aerial Application**

Sharda Glyphosate 41 SL may be applied using aerial spray equipment for silvicultural site preparation, conifer release and rights-of-way treatments. See the "APPLICATION EQUIPMENT and TECHNIQUES" part of the "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" section of this label for information on how to apply Sharda Glyphosate 41 SL by air.

DO NOT APPLY SHARDA GLYPHOSATE 41 SL BY AIR TO RIGHTS-OF-WAY SITES IN THE STATE OF CALIFORNIA.

**SITE PREPARATION**

Following preplant applications of Sharda Glyphosate 41 SL, any silvicultural species may be planted.

**POSTDIRECTED SPRAY**

In established silvicultural sites, use as a spray on the foliage of undesirable vegetation. Care must be exercised to avoid contact of spray, drift or mist with foliage or green bark of desirable species.

**CONIFER RELEASE**

For release, apply only where conifers have been established for more than one year. Vegetation must not be disturbed prior to treatment or until visual symptoms appear after treatment. Symptoms of treatment are slow to appear, especially in woody species treated in late fall. Injury may occur to conifers treated for release, especially where spray patterns overlap or the higher rates are applied or when applications are made during periods of active conifer growth. **Do not use additional surfactant with conifer release applications.** Applications must be made after formation of final conifer resting buds in the fall or prior to initial bud swelling in spring. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Use the following rates for conifer release to control or partially control the weeds listed in the "WEEDS CONTROLLED" section of this label.

Crop	Directions for Use
Douglas Fir ( <i>Pseudotsuga menziesii</i> ) Hemlock ( <i>Tsuga spp.</i> ) Pines ( <i>Pinus spp.</i> )— all species except eastern white, loblolly shortleaf, longleaf or slash pines Spruce ( <i>Picea spp.</i> ) Fir ( <i>Abies spp.</i> )	Apply 1.5 to 2 quarts of Sharda Glyphosate 41 SL per acre except in Washington and Oregon, west of the crest of the Cascade Mountains. For spring treatments west of the crest of the Cascade Mountains, apply 1 quart of Sharda Glyphosate 41 SL per acre before conifer bud swell for control of annual weeds. For fall treatments in Washington and Oregon, west of the crest of the Cascade Mountains, apply 1 to 1.5 quarts of Sharda Glyphosate 41 SL per acre before any major leaf drop of deciduous species.  For release of western hemlock, apply 1 quart of Sharda Glyphosate 41 SL per acre.

For Release of the Following Conifer Species:	Directions for Use
Loblolly Pine ( <i>Pinus taeda</i> ) Eastern White Pine ( <i>Pinus strobus</i> ) Slash pine ( <i>Pinus elliotii</i> )	<u>Late Season Application</u> Apply 1.5 to 2 quarts of Sharda Glyphosate 41 SL in a minimum of 5 gallons of spray solution per acre during early autumn. Applications made prior to September 1 or when conditions are conducive to rapid growth of conifers will create the potential for increased injury in the form of tip and/or needle burn, injury may decrease with later applications. Some autumn colors are acceptable at time of application. Apply prior to frost or leaf drop of undesirable plants.

360967

Applications made according to label directions will release loblolly pine, eastern white pine and slash pine by reducing competition from the following species:

Ash ( <i>Fraxinus spp.</i> )	Maple, red ( <i>Acer rubrum</i> )	Persimmon ( <i>Diospyros spp.</i> )	Poison Sumac ( <i>Rhus vemix</i> )
Black Cherry ( <i>Prunus serotina</i> )	Black Oak ( <i>Quercus velutina</i> )	Yellow Poplar ( <i>Liriodendron tulipifera</i> )	Smooth Sumac ( <i>Rhus glabra</i> )
Pin ( <i>Prunus Pensylvanica</i> )	Post ( <i>Quercus stellata</i> )	Sassafras ( <i>Sassafras albidum</i> )	Winged Sumac ( <i>Rhus copallina</i> )
Elm ( <i>Ulmus spp.</i> )	Southern Red ( <i>Quercus falcata</i> )	Sourwood ( <i>Oxydendrum arboreum</i> )	Sweetgum ( <i>Liquidambar styraciflua</i> )
Black Locust ( <i>Robinia pseudoacacia</i> )			

Apply only to those sites where woody brush and trees listed in this label constitute the majority of the undesirable species.

**Sharda Glyphosate 41 SL plus Oust® Tank Mixtures for Conifer Release from Herbaceous Weeds**

To release loblolly pines from herbaceous weeds, tank mixtures of Sharda Glyphosate 41 SL with Oust® will provide control of annual weeds listed in the "WEEDS CONTROLLED" section of this and the Oust® label, and partial control of the perennial weeds listed below. Apply 16 to 24 fluid ounces of Sharda Glyphosate 41 SL with 2 to 4 ounces of Oust® in 10 to 30 gallons of spray solution per acre. Make application to actively growing weeds as a broadcast spray over the top of the young loblolly pines. **Sharda Glyphosate 41 SL plus Oust® tank mixtures may not be applied by air in California.** This tank mixture may be applied using aerial equipment. When applying by air, use the specified rate in 5 to 15 gallons of spray solution per acre.

For control of annual weeds below 12 inches in height (or runner length on annual vines), use the lower rates of both products. Use the higher rates of both products when annual weeds are in more advanced stages of growth and approaching flower or seed formation. Use the higher rates of both products for partial control of the following perennial weeds. Use the lower rates for suppression of growth.

Bahiagrass ( <i>Paspalum notatum</i> )	Dogfennel ( <i>Eupatorium capilliflorum</i> )	Poorjoe* ( <i>Diodia teres</i> )	Vaseygrass ( <i>Paspalum urvillei</i> )
Broomsedge ( <i>Andropogon virginicus</i> )	Fescue, tall ( <i>Festuca arundinacea</i> )	Trumpet creeper** ( <i>Campsis radicans</i> )	Vervain, blue ( <i>Verbena hastata</i> )
Curly dock ( <i>Rumex crispus</i> )	Johnsongrass* ( <i>Sorghum halepense</i> )		

\*Control at the higher rates.

\*\*Suppression at the higher rates only.

Pine damage may occur or can be accentuated if treatment takes place when young trees are under stress from drought, flood water insects or disease. Read and observe the cautionary statements and all other information appearing on the labels of all herbicides used.

**NOTE:** Sharda Glyphosate 41 SL must not be used in areas where adverse impact on federally designated endangered/threatened plant or aquatic species is likely. Prior to making applications, the user of Sharda Glyphosate 41 SL must determine that no such species are located in or immediately adjacent to the area to be treated.

**CUT STUMP TREATMENTS**

Woody vegetation may be controlled by treating freshly cut stumps of trees and resprouts with Sharda Glyphosate 41 SL. Apply Sharda Glyphosate 41 SL using suitable equipment to ensure coverage of the entire cambium. Cut vegetation close to the soil surface. Apply a 50 to 100 percent solution of Sharda Glyphosate 41 SL to the freshly-cut surface immediately after cutting. Delays in application may result in reduced performance. For best results, applications must be made during periods of active growth and full leaf expansion.

3/10/07

When used according to directions for cut stump application, Sharda Glyphosate 41 SL will CONTROL, PARTIALLY CONTROL or SUPPRESS many types of woody brush and tree species, some of which are listed below:

Alder ( <i>Alnus spp.</i> )	Oak ( <i>Quercus spp.</i> )	Saltcedar ( <i>Tamarisk spp.</i> )	Tan Oak ( <i>Lithocarpus densiflorus</i> )
Eucalyptus ( <i>Eucalyptus spp.</i> )	Giant Reed ( <i>Arundo donax</i> )	Sweetgum ( <i>Liquidambar styraciflua</i> )	Willow ( <i>Salix spp.</i> )
Madrone ( <i>Arbutus menziesii</i> )			

**INJECTION AND FRILL APPLICATIONS**

Woody vegetation may be controlled by injection or frill application of Sharda Glyphosate 41 SL. Apply Sharda Glyphosate 41 SL using suitable equipment which must penetrate into the living tissue. Apply the equivalent of 1 ml of Sharda Glyphosate 41 SL per each 2 to 3 inches of trunk diameter (DBH). This is best achieved by applying a 50 to 100 percent concentration of this material either to a continuous frill around the tree or as cuts evenly spaced around the tree below all branches. As tree diameter increases in size, better results are achieved by applying diluted material to a continuous frill or more closely spaced cuttings. Avoid application techniques that allow runoff to occur from frill or cut areas in species that exude sap freely after frills or cutting. In species such as this, make frill or cut at an oblique angle so as to produce a cupping effect and use undiluted material. For best results, application must be made during periods of active growth and after full leaf expansion.

This treatment WILL CONTROL the following woody species:

- Oak (*Quercus spp.*)
- Poplar (*Populus spp.*)
- Sweetgum (*Liquidambar styraciflua*)
- Sycamore (*Platanus occidentalis*)

This treatment WILL SUPPRESS the following woody species:

- Dogwood (*Cornus spp.*)
- Maple, red (*Acer rubrum*)
- Black gum (*Nyssa sylvatica*)
- Hickory (*Carya spp.*)

**TURFGRASSES AND GRASSES FOR SEED PRODUCTION PREPLANT AND RENOVATION**

When applied as directed for "NONCROP USES", under conditions described, Sharda Glyphosate 41 SL controls most existing vegetation prior to the planting or renovation of either turfgrasses or grass seed production areas.

For specific rates of application and instructions for control of various annual and perennial weeds, and woody brush and trees, see the "WEEDS CONTROLLED" section of this label. For maximum control of existing vegetation, delay planting to determine if any regrowth from escaped underground plant parts occurs. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm-season grasses, such as bermudagrass, summer or fall applications provide best control. DO NOT DISTURB SOIL OR UNDERGROUND PLANT PARTS BEFORE TREATMENT. Tillage or renovation techniques such as vertical mowing, coring or slicing must be delayed for 7 days after application to allow proper translocation into underground plant parts.

**TURFGRASSES**

Where existing vegetation is growing in a field or unmowed situation, apply Sharda Glyphosate 41 SL to actively growing weeds at the stages of growth listed in the "WEEDS CONTROLLED" section of this label. Where existing vegetation is growing under mowed turfgrass management, apply Sharda Glyphosate 41 SL after omitting at least one regular mowing to allow sufficient growth for good interception of the spray. Desirable turfgrasses may be planted following the above procedures.

**GRASSES FOR SEED PRODUCTION**

Apply Sharda Glyphosate 41 SL to actively growing weeds at the stages of growth specified in the "WEEDS CONTROLLED" section of this label prior to planting or renovation of turf or forage grass areas grown for seed production. DO NOT feed or graze treated areas within 8 weeks after application.

**ANNUAL WEED CONTROL IN DORMANT BERMUDAGRASS AND BAHIAGRASS TURF**

When applied as directed for "NONCROP USES" under the conditions described, Sharda Glyphosate 41 SL will provide control or suppression of many winter annual weeds and tall fescue for effective release of dormant bermudagrass and bahiagrass turf. Refer to the rate table for Sharda Glyphosate 41 SL alone under the "RELEASE OF BERMUDAGRASS and BAHIAGRASS" section of this label for specified rates and volumes on the species to be suppressed or controlled. Treat only when turf is dormant and prior to spring greenup. Spot treatments or broadcast applications of Sharda Glyphosate 41 SL in excess of 16 fluid ounces per acre may result in injury or delayed greenup in highly maintained turfgrass areas; i.e., golf courses, lawns, etc. DO NOT APPLY TANK MIXTURES of Sharda Glyphosate 41 SL plus Oust® in highly maintained turfgrass areas.

**RELEASE OF BERMUDAGRASS OR BAHIAGRASS**

**NOTE:** Use only in areas where bermudagrass or bahiagrass are desirable ground covers and where some temporary injury or discoloration can be tolerated. Use tank mixtures of Sharda Glyphosate 41 SL plus Oust® only on railroads, highways, utility plant sites, or other right-of-way areas. When applied as directed for "NONCROP USES" under the conditions described, Sharda Glyphosate 41 SL will provide control or suppression of many winter annual weeds and tall fescue for effective release of dormant bermudagrass or bahiagrass. Sharda Glyphosate 41 SL may be tank-mixed with Oust® as specified for residual control. Make applications to dormant bermudagrass or bahiagrass. Tank mixtures of Sharda Glyphosate 41 SL plus Oust® may delay greenup. To avoid delays in greenup and minimize injury, do not add more than 1 ounce per acre of Oust® on bermudagrass or more than 0.5 ounce per acre on bahiagrass, or treat when these grasses are in a semi-dormant condition. For best results on winter annuals, treat when plants are in an early growth stage (below 6 inches in height) after most have germinated. For best results on tall fescue, treat when fescue is in or beyond the 4 to 6-leaf stage.

**WEEDS CONTROLLED**

Rate directions for control or suppression of winter annuals and tall fescue are listed below:

Apply the specified rates of Sharda Glyphosate 41 SL alone or as a tank mixture in 10 to 25 gallons of water per acre.

**WEEDS CONTROLLED OR SUPPRESSED WITH SHARDA GLYPHOSATE 41 SL ALONE\***

**NOTE:** C = Control S = Suppression

SHARDA GLYPHOSATE 41 SL Fluid Ounces Per Acre:						
Weed	8	12	16	24	32	64
Barley, little ( <i>Hordeum pusillum</i> )	S	C	C	C	C	C
Bedstraw, catchweed ( <i>Galium aparine</i> )	S	C	C	C	C	C
Bluegrass, annual ( <i>Poa annua</i> )	S	C	C	C	C	C
Chervil ( <i>Chaerophyllum tainturieri</i> )	S	C	C	C	C	C
Chickweed, common ( <i>Stellaria media</i> )	S	C	C	C	C	C
Clover, crimson ( <i>Trifolium incarnatum</i> )	--	S	S	C	C	C

SHARDA GLYPHOSATE 41 SL Fluid Ounces Per Acre:						
Weed	8	12	16	24	32	64
Clover, large hop ( <i>Trifolium campestre</i> )	--	S	S	C	C	C
Fescue, tall ( <i>Festuca arundinaceae</i> )	--	--	--	--	S	S
Geranium, Carolina ( <i>Geranium carolinianum</i> )	--	--	S	S	C	C
Henbit ( <i>Lamium amplexicaule</i> )	--	S	C	C	C	C
Ryegrass, Italian ( <i>Lolium multiflorum</i> )	--	--	S	C	C	C
Speedwell, corn ( <i>Veronica arvensis</i> )	S	C	C	C	C	C
Vicia ( <i>sativa</i> )	--	--	S	C	C	C

\*These rates apply only to sites where an established competitive turf is present.

39 of 67

**WEEDS CONTROLLED OR SUPPRESSED WITH SHARDA GLYPHOSATE 41 SL PLUS OUST®**

NOTE: C = Control S = Suppression

SHARDA GLYPHOSATE 41 SL Fluid Ounces Per Acre Plus Oust®:							
Sharda Glyphosate 41 SL (Fl. oz/A)	8	12	12	16	16	12	16
+	+	+	+	+	+	+	+
OUST® (oz./A)	1/4	1/4	1/2	1/4	1/2	1	1
<b>Weed</b>							
Barley, little ( <i>Hordeum pusillum</i> )	C	C	C	C	C	C	C
Bedstraw, catchweed ( <i>Galium aparine</i> )	C	C	C	C	C	C	C
Bluegrass, annual ( <i>Poa annua</i> )	S	C	C	C	C	C	C
Chervil ( <i>Chaerophyllum tainturieri</i> )	C	C	C	C	C	C	C
Chickweed, common ( <i>Stellaria media</i> )	S	C	C	C	C	C	C
Clover, crimson ( <i>Trifolium incarnatum</i> )	S	S	S	S	C	C	C
Clover, large hop ( <i>Trifolium campestre</i> )	--	--	S	S	S	C	C
Fescue, tall ( <i>Festuca arundinaceae</i> )	--	--	--	--	--	S	S
Geranium, Carolina ( <i>Geranium carolinianum</i> )	--	S	S	C	C	C	C
Henbit ( <i>Lamium amplexicaule</i> )	--	S	C	C	C	C	C
Ryegrass, Italian ( <i>Lolium multiflorum</i> )	--	S	S	C	C	C	C
Speedwell, corn ( <i>Veronica arvensis</i> )	S	C	C	C	C	C	C
Vicia ( <i>sativa</i> )	C	C	C	C	C	C	C

\*These rates apply only to sites where an established competitive turf is present.

**RELEASE OF ACTIVELY GROWING BERMUDAGRASS**

When applied as directed, Sharda Glyphosate 41 SL will aid in the release of bermudagrass by providing control of annual species listed in the "WEEDS CONTROLLED" section of this and the Oust® label, and suppression or partial control of certain perennial weeds. For control or suppression of those annual species listed on this label, use 1 to 3 pints of Sharda Glyphosate 41 SL as a broadcast spray in 10 to 25 gallons of spray solution per acre. Use the lower rate when treating annual weeds below 6 inches in height (or length of runner in annual vines). Use the higher rate as weeds increase in size or as they approach flower or seedhead formation.

Use the higher rate of Sharda Glyphosate 41 SL for partial control of the following perennial species. Use the lower rates for suppression of growth. For best results, see the "WEEDS CONTROLLED" section of this label for proper stage of growth.

Bahiagrass ( <i>Paspalum notatum</i> )	Bluestem, silver ( <i>Andropogon saccharoides</i> )	Fescue, tall ( <i>Festuca arundinaceae</i> )
Johnsongrass* ( <i>Sorghum halepense</i> )	Trumpet creeper** ( <i>Campsis radicans</i> )	Vaseygrass ( <i>Paspalum urvillei</i> )

\* Control at the higher rates.  
 \*\* Suppression at higher rates only.

Sharda Glyphosate 41 SL may be tank-mixed with Oust®. If tank-mixed, use no more than 1 to 2 pints per acre of Sharda Glyphosate 41 SL with 1 to 2 ounces of Oust® per acre. Use the lower rates of both mixtures to control annual weeds below 6 inches in height (or runner length in annual vines) that are listed in the "WEEDS CONTROLLED" section of this booklet and the Oust® label. Use the higher rates as annual weeds increase in size and approach the flower or seedhead stages. Use the higher rates of Sharda Glyphosate 41 SL to provide partial control of the following perennial weeds. Use the lower rates for suppression of growth.

40 of 67

Bahiagrass ( <i>Paspalum notatum</i> )	Bluestem, silver ( <i>Andropogon saccharoides</i> )	Broomsedge ( <i>Andropogon virginicus</i> )
Dock, curly ( <i>Rumex crispus</i> )	Dogfennel ( <i>Eupatorium capilliflorum</i> )	Fescue, tall ( <i>Festuca arundinacea</i> )
Johnsongrass* ( <i>Sorghum halepense</i> )	Poorjoe** ( <i>Diodia teres</i> )	Trumpetcreeper** ( <i>Campsis radicans</i> )
Vaseygrass ( <i>Paspalum urvillei</i> )	Vervain, blue ( <i>Verbena hastata</i> )	

\*Suppression at higher rates only.  
 \*\*Control at the higher rates.

Use only on well-established bermudagrass. Bermudagrass injury may result from the treatment but regrowth will occur under moist conditions. Repeat applications in the same season are not directed, since severe injury may result. Read and carefully observe the precautionary statements and all other information appearing on the labels of all herbicides used.

**COOL SEASON TURF GROWTH REGULATION**

When applied as directed, Sharda Glyphosate 41 SL will suppress growth and seedhead development of listed turf species in industrial sites. Sharda Glyphosate 41 SL is directed for management of coarse turf on roadside rights-of-way or other industrial areas. Do not use on high-quality turf or other areas where some turf color changes cannot be tolerated. Slight turf discoloration may occur but turf will regreen and regrow under moist conditions as effects of Sharda Glyphosate 41 SL wear off. Apply 4 to 6 fluid ounces of Sharda Glyphosate 41 SL per acre alone or in a specified tank mixture. Spray volumes of 10 to 40 gallons per acre are directed.

**Sharda Glyphosate 41 SL can be used for growth and seedhead suppression of Tall Fescue and Smooth Brome**

For best results, apply Sharda Glyphosate 41 SL in a specified tank mixture to actively growing turfgrasses after greenup in the spring of the year. For suppression of seedheads, applications must be made before boot-to-seedhead stage of development. Applications made from seedhead emergence until maturity may result in turf discoloration or injury. After mowing or removal of seedheads, Sharda Glyphosate 41 SL in a specified tank mixture may also be used to suppress the growth of certain turfgrasses. Allow turf to recover from stress caused by heat, drought or mowing before making applications. Applications made to turf under stress may increase the potential for discoloration or injury.

**ANNUAL GRASSES**

For growth suppression of some annual grasses such as annual ryegrass, wild barley and wild oats, apply 3 to 4 fluid ounces of Sharda Glyphosate 41 SL in 10 to 40 gallons of spray solution per acre. Applications must be made when annual grasses are actively growing and before the seedheads are in the boot stage of development. Treatments made after seedhead emergence may cause injury to the desired grasses.

**TANK MIXTURES**

For the following tank mixtures, consult each product label for weeds controlled and the correct stage of application. Do not treat turf under stress.

**Tank mixtures plus 2,4-D Amine:** For additional weed control benefits, up to 1 pound a.i. per acre of 2,4-D amine may be added to the following tank mixtures. Consult the label for 2,4-D amine for weeds controlled.

Sharda Glyphosate 41 SL may be tank-mixed with the listed herbicides provided the specific product is registered for use on these sites.

**TALL FESCUE**

**Sharda Glyphosate 41 SL plus Telar®:** For suppression of tall fescue growth and seedheads, and control or partial control of some annual weeds, apply this tank mixture after greenup and prior to boot-to-seedhead stage of development. Use up to 0.5 ounce of Telar® per acre. This tank mixture can also be applied after mowing or removal of tall fescue seedheads for turf growth suppression. Make only one of the above applications per growing season.

**Sharda Glyphosate 41 SL plus Oust®:** For suppression of tall fescue growth and seedheads, and control or partial control of some annual weeds, apply this tank mixture after greenup and prior to boot-to-seedhead stage of development. Use up to 0.25 ounce of Oust® per acre.



4/10/07

**Sharda Glyphosate 41 SL plus Escort®:** This tank mixture can be applied after mowing or removal of tall fescue seedheads for turf growth suppression and control or partial control of some annual weeds. Use up to 1/3 ounce of Escort® per acre.

**NOTE:** Sharda Glyphosate 41 SL is not registered for use with Escort® in California.

**SMOOTH BROME**

**Sharda Glyphosate 41 SL plus Oust®:** For suppression of smooth brome growth and seedheads and control or partial control of some annual weeds, apply this tank mixture after greenup and prior to boot-to-seedhead stage of development. Use up to 0.25 ounce of Oust® per acre.

**BAHIAGRASS SEEDHEAD AND VEGETATIVE SUPPRESSION**

When applied as directed in the indicated noncrop areas (roadsides, airports, golf course roughs, plant sites and similar areas that are not high quality turfgrasses), Sharda Glyphosate 41 SL will provide significant inhibition of seedhead emergence and will suppress vegetative growth for a period of approximately 45 days with single applications and approximately 120 days with sequential applications. Apply Sharda Glyphosate 41 SL 1 to 2 weeks after full greenup of bahiagrass or after the bahiagrass has been mowed to a uniform height of 3 to 4 inches. Applications must be made prior to seedhead emergence. Apply 6 fluid ounces per acre of Sharda Glyphosate 41 SL in 10 to 25 gallons of water per acre.

Sequential applications of Sharda Glyphosate 41 SL may be made at approximately 45 day intervals to extend the period of seedhead and vegetative growth suppression. For continued seedhead suppression, sequential applications must be made prior to seedhead emergence. Apply no more than 2 sequential applications per year. As a first sequential application, apply 4 fluid ounces of Sharda Glyphosate 41 SL per acre. A second sequential application of 2 to 4 fluid ounces per acre may be made approximately 45 days after the last application. A tank mixture of Sharda Glyphosate 41 SL plus Oust® may be applied only on roadsides for seedhead inhibition and vegetative suppression. Apply 6 fluid ounces per acre of Sharda Glyphosate 41 SL plus 0.25 ounce per acre of Oust® 1 to 2 weeks following an initial spring mowing. When using Sharda Glyphosate 41 SL plus Oust® for suppression of bahiagrass, make only 1 application per year.

**CROPPING SYSTEMS**

When applied as directed for "CROPPING SYSTEMS", under the conditions described, Sharda Glyphosate 41 SL controls annual and perennial weeds listed on this label, prior to the emergence of direct seeded crops or prior to transplanting of crops listed on this label. In-crop application to Roundup Ready® soybeans and cotton may be made according to the directions given in those respective sections below. See "PRODUCT INFORMATION" and "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" sections of this label for essential product performance information. See the following "CROPPING SYSTEMS" sections for specific directed uses.

**EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF SPRAY WITH FOLIAGE, GREEN STEMS OR FRUIT OF DESIRABLE CROPS, PLANTS, TREES OR OTHER DESIRABLE VEGETATION SINCE SEVERE DAMAGE OR DESTRUCTION MAY RESULT.**

Repeat treatments may be necessary to control weeds regenerating from underground parts or seed.

Except as otherwise specified on this label, repeat treatments must be made before the crop emerges in accordance with the instructions of this label.

Except as otherwise specified in a crop section of this label, the combined total of all treatments must not exceed 8 quarts per acre of Sharda Glyphosate 41 SL per year.

For any crop not listed below, applications must be made at least 30 days prior to planting.

Do not harvest or feed treated vegetation for 8 weeks following application. Following spot treatment or selective equipment use, allow 14 days before grazing domestic livestock or harvesting forage grasses and legumes.

**Row Crops**

CORN (ALL)*	COTTON*	PEANUTS	SORGHUM (MILO)*
SOYBEANS*	SUGARCANE*		

**Cereal Grains**

BARLEY*	BUCKWHEAT*	MILLET (PEARL, PROSO)*	OATS*
RICE**	RYE*	TRITICALE*	WHEAT (ALL)*

420967

WILD RICE\*

**Citrus**

CITRUS	KUMQUAT	ORANGE (ALL)	TANGERINE
CALAMONDIN	LEMON	PUMMELO	TANGORS
CHIRONJA	LIME	TANGELO	
CITRON	MANDARIN ORANGE		
GRAPEFRUIT			

**Tree Nuts**

ALMOND	CASHEW	HICKORY NUT	PISTACHIO
BEECHNUT	CHESTNUT	MACADAMIA	WALNUT (BLACK, ENGLISH)
BRAZIL NUT	CHINQUAPIN	PECAN	
BUTTERNUT	FILBERT (HAZELNUT)		

**Small Fruits and Berries**

BLACKBERRY	CRANBERRY	ELDERBERRY	LOGANBERRY
BLUEBERRY	CURRENT	GOOSEBERRY	OLALLIEBERRY
BOYSENBERRY	DEWBERRY	HUCKLEBERRY	RASPBERRY (BLACK, RED)

**Tree Fruits**

APPLE	LOQUAT	OLIVE	PLUM/PRUNE (ALL)
APRICOTS	MAYHAW	PEACH	QUINCE
CHERRY (SWEET, SOUR)	NECTARINE	PEAR	

**Vegetables**

ARTICHOKE, JERUSALEM	CHARD, SWISS	LEEK	PUMPKIN***
ASPARAGUS*	CHICORY	LENTILS	RADISH
BEANS (ALL)	COLLARDS	LETTUCE	RAPE GREENS
BEET GREENS	CRENSHAW MELON***	MANGO MELON***	RHUBARB
BEETS (RED, SUGAR)	CUCUMBER***	MELONS (ALL)***	RUTABAGA
BROCCOLI (ALL)	EGGPLANT***	MUSKMELON***	SHALLOT
BRUSSELS SPROUTS	ENDIVE	MUSTARD GREENS	SPINACH (ALL)
CABBAGE (ALL)	GARLIC***	OKRA	SQUASH (SUMMER, WINTER)***
CABBAGE, CHINESE	GOURDS***	ONION	TOMATILLO***
CANTALOUPE***	GROUND CHERRY**	PARSLEY	TOMATO***†
CARROT	HONEYDEW MELON***	PARSNIPS	TURNIP
CAULIFLOWER	HONEY BALL MELON***	PEAS (ALL)	WATERCRESS***
CASABA MELON***	HORSERADISH	PEPPER (ALL)***	WATERMELON***
CELERIAC	KALE	PERSIAN MELON***	YAMS
CELERY	KOHLRABI	POTATO (IRISH, SWEET)	

**Vine Crops**

GRAPES	KIWI FRUIT
--------	------------

**Forage Crops and Legumes**

ALFALFA*	FORAGE GRASSES*	FORAGE LEGUMES*
----------	-----------------	-----------------

**Tropical Crops**

ACEROLA	COCOA BEANS	LONGAN	SAPODILLA
---------	-------------	--------	-----------

430967

ATEMOYA	COFFEE	LYCHEE	SAPOTE (BLACK
AVOCADO	DATES	MANGO	MAMEY, WHITE)
BANANA (PLANTAINS)	FIGS	PAPAYA	SOURSOP
BREADFRUIT	GENIP	PASSION FRUIT	SUGAR APPLE
CANISTEL	GUAVA	PERSIMMONS	TAMARIND
CARAMBOLA	JABOTICABA	PINEAPPLE****	TEA
CHERIMOYA	JACKFRUIT	POMEGRANATE	

- \* Spot treatments may be applied in these crops.
- \*\* Do not treat rice fields or levees when the fields contain flood water.
- \*\*\* Apply only prior to planting. Allow at least 3 days between application and planting.
- \*\*\*\*Do not feed or graze treated pineapple forage following application.
- † Use is restricted to direct seeded crops only.

When applying Sharda Glyphosate 41 SL prior to transplanting crops into plastic mulch, care must be taken to remove residues of Sharda Glyphosate 41 SL from the plastic prior to transplanting. Residues can be removed by 0.5 inch natural rainfall or by applying water via a sprinkler irrigation system.

**Spot Treatment:** (Only those crops with "\*" can be spot treated.) - Applications in growing crops must be made prior to heading of small grains and milo, initial pod set in soybeans, silking of corn, or boll opening on cotton.

For forage grasses and forage legumes see "SPOT TREATMENT" in the "PASTURES" section of "CROPPING SYSTEMS" in this label.

For dilution and rates of application using boom or hand-held equipment, see "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" and "WEEDS CONTROLLED" sections of this label.

**NOTE:** FOR FORAGE GRASSES AND FORAGE LEGUMES, NO MORE THAN ONE-TENTH OF ANY ACRE MUST BE TREATED AT ONE TIME. FOR ALL OTHER CROPS, DO NOT TREAT MORE THAN 10 PERCENT OF THE TOTAL FIELD AREA TO BE HARVESTED. THE CROP RECEIVING SPRAY IN TREATED AREA WILL BE KILLED. TAKE CARE TO AVOID DRIFT OR SPRAY OUTSIDE TARGET AREA FOR THE SAME REASON.

**Selective Equipment:** Sharda Glyphosate 41 SL may be applied through recirculating sprayers, shielded applicators or wiper applicators in cotton and soybeans. Shielded and wiper applicators may also be used in tree crops and grapes. Wiper applicators may be used in wheat, rutabagas, forage grasses and forage legumes, including pasture sites and grain sorghum (milo).

See the "SELECTIVE EQUIPMENT" part of the "APPLICATION EQUIPMENT and TECHNIQUES" section of this label for information on proper use and calibration of this equipment.

Allow at least the following time intervals between application and harvest:

Cotton, Soybeans	7 days
Apples, Citrus, Pear	1 day
Atemoya, Avocado, Breadfruit, Canistel, Carambola, Cherry, Grapes, Dates, Jaboticaba, Jackfruit, Longan, Lychee, Passion Fruit, Persimmons, Rutabagas, Sapodilla, Sapote, Soursop, Sugar Apple, Tamarind	14 days
Stone Fruit	17 days
Nut Crops	3 days
Wheat <sup>1</sup>	35 days
Sorghum (milo) <sup>1,2</sup>	40 days

<sup>1</sup> Do not use roller applicators.  
<sup>2</sup> Do not feed or graze treated milo fodder. Do not ensile treated vegetation.

440967

**ASPARAGUS**

When applied as directed for "CROPPING SYSTEMS" under the conditions described, Sharda Glyphosate 41 SL controls weeds listed on this label in asparagus. For specific rates of applications and instructions for control of various annual and perennial weeds, see the "WEEDS CONTROLLED" section of this label.

Prior to Crop Emergence - Apply Sharda Glyphosate 41 SL prior to crop emergence for the control of emerged labeled annual and perennial weeds. DO NOT APPLY WITHIN A WEEK BEFORE THE FIRST SPEARS EMERGE.

**Spot Treatment:** Apply Sharda Glyphosate 41 SL immediately after cutting, but prior to the emergence of new spears. Do not treat more than 10 percent of the total field area to be harvested. Do not harvest within 5 days of treatment.

**Postharvest:** Apply Sharda Glyphosate 41 SL after the last harvest and all spears have been removed. If spears are allowed to regrow, delay application until ferns have developed. Delayed treatments must be applied as directed or shielded spray in order to avoid contact of the spray with ferns, stems or spears. Direct contact of the spray with the asparagus may result in serious crop injury.

**NOTE:** Select and use specified types of spray equipment for postemergence postharvest applications. A directed spray is any application where the spray pattern is aligned in such a way as to avoid direct contact of the spray with the crop. A shielded spray is any application where a physical barrier is positioned and maintained between the spray and the crop to prevent contact of spray with the crop.

**BERRIES AND SMALL FRUITS**

Wiper applicators may be used in cranberries in accordance with instructions in this section.

For other berries, apply as a preplant broadcast application, or as a directed spray or wiper application post-planting. See "PRODUCT INFORMATION" and "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" sections of this label for essential product performance information. See the "SELECTIVE EQUIPMENT" part of the "APPLICATION EQUIPMENT and TECHNIQUES" section of this label for information on directed use and calibration of this equipment.

Allow a minimum of 30 days between last application and harvest of cranberries. For other small fruits and berries, allow a minimum of 14 days between last application and harvest.

**For Wick or other Wiper Applicators:** Mix 1 gallon of Sharda Glyphosate 41 SL in 4 gallons of water to prepare a 20 percent solution. In severe infestations, reduce equipment ground speed to ensure that adequate amounts of Sharda Glyphosate 41 SL are wiped on the weeds. A second treatment in the opposite direction may be beneficial.

Do not permit herbicide solution to contact desirable vegetation, including green shoots, canes or foliage.

**CORN (Field corn, seed corn, sweet corn and popcorn)**

**TYPES OF APPLICATIONS:** Preplant, preemergence, at-planting, hooded sprayers, spot treatment, preharvest, post-harvest.

Add an agriculturally approved nonionic surfactant at 0.375 percent by volume of spray solution. Adding 1 to 2 percent by weight of dry ammonium sulfate (or equivalent from other formulations) may increase the performance of Sharda Glyphosate 41 SL.

**Preplant, Preemergence and At-planting:** Sharda Glyphosate 41 SL may be applied before, during or after planting corn. Applications must be made prior to emergence of the crop.

Subject to any limitations stated on labeling of specific products, the following tank mixtures may be applied before, during or after planting in conventional tillage systems, into a cover crop, established sod or in previous crop residue. Apply these tank mixtures in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre.

ATRAZINE	DUAL®	HARNESS XTRA 5.6L®	PARTNER®
BANVEL®/Dicamba	DUAL® II	LARIAT®	PROWL®/Pendimethalin
BICEP®	EXTRAZINE®	LASSO®/Alachlor	SIMAZINE
BICEP® II	FRONTIER®	LINEX®	SURPASS®/Acetochlor

450f67

BLADEX®/Cyanazine	GUARDSMAN®	LOROX®/Linuron	SURPASS® 100
BROADSTRIKE®	HARNESS®/Acetochlor	MARKS MAN ®/Atrazine Dicamba	TOPNOTCH®
BULLET®	HARNESS XTRA®	MICRO-TECH®	

Sharda Glyphosate 41 SL may be tank-mixed with the listed herbicides provided the specific product is registered for use on these sites.

For Southern states, do not apply in nitrogen solutions to tough-to-control grasses such as barnyardgrass, fall panicum, broadleaf signalgrass, annual ryegrass and any perennial weeds.

For improved burndown, Sharda Glyphosate 41 SL may be tank mixed with 2,4-D (Weedar®, Weedone® and others) or dicamba.

For difficult-to-control annual weeds such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply Sharda Glyphosate 41 SL at 2 pints per acre in these tank mixtures. For other labeled annual weeds, apply 1 to 1-1/2 pints of Sharda Glyphosate 41 SL per acre when weeds are less than 6 inches tall, and 2 to 3 pints when weeds are over 6 inches tall.

Do not plant corn until at least 7 days after application of 2,4-D or dicamba.

The tank mix directions in this section are not registered in California.

**Hooded Sprayers:** Sharda Glyphosate 41 SL may be used through hooded sprayers for weed control between the rows of corn. Only hooded sprayers that completely enclose the spray pattern may be used. A hooded sprayer is a type of shielded applicator. The spray pattern is completely enclosed on the top and all 4 sides by a hood, thereby shielding the crop from the spray solution. This equipment must be set up and operated in a manner that avoids bouncing or raising the hoods off the ground in any way. If the hoods are raised, spray particles may escape and come into contact with the crop, causing damage or destruction of the crop. The spray hoods must be operated on the ground or skimming across the ground. Tractor speed must be adjusted to avoid bouncing of the spray hoods. Avoid operation on rough or sloping ground where the spray hoods might be raised off the ground. When applying to corn that is grown on raised beds, ensure that the hood is designed to completely enclose the spray solution. If necessary, extend the front and rear flaps of the hoods to reach the ground in deep furrows. Follow these requirements:

- The spray hoods must be operated on the ground or skimming across the ground.
- Do not apply more than 1 quart of Sharda Glyphosate 41 SL per acre per application.
- Corn must be at least 12 inches tall, measured without extending leaves.
- Leave at least an 8 inch untreated strip over the drill row. For example, if the crop row width is 38 inches, the maximum width of the spray hood must be 30 inches.
- Maximum tractor speed: 5 mph.
- Maximum wind speed: 10 mph.
- Use low-drift nozzles.

Crop injury may occur when the foliage of treated weeds comes into direct contact with leaves of the crop. Do not apply Sharda Glyphosate 41 SL when the leaves of the crop are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution may contact the crop and cause discoloration, stunting or destruction. Contact of Sharda Glyphosate 41 SL in any manner to any vegetation to which treatment is not intended may cause damage. Such damage shall be the sole responsibility of the applicator. For specific rates of application and instructions for control of various annual and perennial weeds, see the "WEEDS CONTROLLED" Section of the label booklet. Do not graze or feed corn forage or fodder following applications of Sharda Glyphosate 41 SL through hooded sprayers. Do not apply more than 3 quarts of Sharda Glyphosate 41 SL per acre per year for hooded sprayer applications.

**Spot treatment:** For spot treatments, apply Sharda Glyphosate 41 SL prior to silking of corn.

Do not treat more than 10 percent of the total field area to be harvested. The crop receiving spray in the treated area will be killed. Take care to avoid drift or spray outside target area for the same reason.

**Preharvest:** Make applications at 35 percent grain moisture or less. Ensure that maximum kernel fill is complete and the corn is physiologically mature (black layer formed). For ground applications, apply up to 3 quarts of Sharda Glyphosate 41 SL per acre. For aerial applications, apply up to 1 quart of Sharda Glyphosate 41 SL per acre.

460967

Allow a minimum of 7 days between application and harvest. It is not directed that corn grown for seed be treated preharvest because a reduction in germination or vigor may result.

**Post-harvest:** Sharda Glyphosate 41 SL may be applied after harvest of corn. Higher specified rates may be required for control of large weeds which were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used. Do not harvest or feed treated vegetation for 8 weeks following application.

**POSTEMERGENCE APPLICATIONS TO CORN WITH THE ROUNDUP READY® GENE  
PRODUCT INFORMATION**

SHARDA USA LLC DIRECTS USE OF SHARDA GLYPHOSATE 41 SL ONLY ON CORN HYBRIDS DESIGNATED AS CONTAINING THE ROUNDUP READY® GENE.

- Applying Sharda Glyphosate 41 SL to corn hybrids which are not designated as Roundup Ready® will result in severe crop injury and yield loss.
- The Roundup Ready® designation indicates that the corn contains a patented gene which provides tolerance to this herbicide. Information on Roundup Ready® corn may be obtained from your seed supplier or Sharda USA LLC representative.

**APPLICATION INSTRUCTIONS**

Sharda Glyphosate 41 SL may be applied postemergence to Roundup Ready® corn from emergence through the V8 stage (8 leaves with collars) or until corn height reaches 30 inches, whichever comes first. Single in-crop applications of Sharda Glyphosate 41 SL are not to exceed 1 quart per acre. Sequential in-crop applications of Sharda Glyphosate 41 SL from emergence through the V8 stage or 30 inches must not exceed 2 quarts per acre per growing season.

**Maximum Allowable Yearly Rates:**

- **Preplant:** Maximum amount of Sharda Glyphosate 41 SL which can be applied prior to crop emergence is 5 quarts per acre.
- **In-crop:** Maximum combined total of multiple in-crop applications from emergence through the V8 stage or 30 inches is 2 quarts per acre.
- **Preharvest:** Maximum amount of Sharda Glyphosate 41 SL that can be applied after maximum kernel fill is complete and the crop is physiologically mature (black layer formation) until 7 days before harvest is 1 quart per acre.
- **Cropping Season:** Combined total per year for all applications may not exceed 8 quarts per acre.

When applied as directed, Sharda Glyphosate 41 SL controls labeled annual grass and broadleaf weeds in Roundup Ready® corn. Many perennial grasses and broadleaf weeds will be controlled or suppressed with one or more applications of Sharda Glyphosate 41 SL. Applications must be made to actively growing weeds before they reach the maximum size listed in the label booklet. Refer to the label booklet for proper use instructions.

The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of Sharda Glyphosate 41 SL under hard water conditions, drought conditions or when tank mixed with Bullet®, Micro-Tech® or Partner® herbicides. Ensure that ammonium sulfate is completely dissolved in the spray tank before adding herbicides. Thoroughly rinse the spray system with clean water after use to reduce corrosion. The addition of other additives, including fertilizers and micronutrients is not directed with Sharda Glyphosate 41 SL since they may result in increased potential for crop injury.

Allow a minimum of 50 days between application of the product and harvest of corn forage and 7 days between application and harvest of corn grain. Allow a minimum of 10 days between in-crop applications of Sharda Glyphosate 41 SL. There are no rotational crop restrictions following applications of Sharda Glyphosate 41 SL.

**ATTENTION: AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING SHARDA GLYPHOSATE 41 SL TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN THE ROUNDUP READY® GENE. THOROUGHLY CLEAN THE SPRAY TANK AND ALL LINES AND FILTERS TO ELIMINATE POTENTIAL CONTAMINATION FROM OTHER HERBICIDES PRIOR TO MIXING AND APPLYING SHARDA GLYPHOSATE 41 SL.**

470767

**For ground applications:** Use the specified rates of Sharda Glyphosate 41 SL in 5 to 20 gallons of spray solution per acre as a broadcast spray. Carefully select correct nozzles and spray pressure to avoid spraying a fine mist. Check for even distribution of spray droplets.

**For aerial applications:** Use the specified rates of Sharda Glyphosate 41 SL in 3 to 15 gallons of spray solution per acre but do not exceed 1 quart of product per acre. Refer to label booklet for weeds controlled or suppressed. **AVOID DRIFT - DO NOT APPLY DURING INVERSION CONDITIONS, WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITIONS WHICH FAVOR DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.**

**Weed Control Directations**

Apply 24 to 32 fluid ounces of this herbicide per acre for control of labeled grasses and broadleaf weeds in conventional and no-till corn production systems. Refer to the label booklet for rate directations for specific annual weeds. Up to 1 quart per acre will control or suppress the growth of perennial weeds such as: bermudagrass, Canada thistle, common milkweed, field bindweed, hemp dogbane, horsenettle, nutsedge, quackgrass, rhizome johnsongrass, redvine, trumpetcreeper, swamp smartweed, and wirestem muhly. For additional information on perennial weeds, see the label booklet.

**Preemergence followed by Postemergence Weed Control Program:** Sharda Glyphosate 41 SL may be applied postemergence in-crop following any labeled preemergence herbicide application. The post application of Sharda Glyphosate 41 SL must be made before the weeds reach a height and/or density that the weeds become competitive with the crop. A single in-crop application of Sharda Glyphosate 41 SL at the specified rate will provide control of emerged weeds listed on the label. Sharda Glyphosate 41 SL may be applied postemergence to Roundup Ready® corn from emergence through the V8 stage (8 leaves with collars) or until corn height reaches 30 inches (free standing), whichever comes first.

**Postemergence followed by Postemergence Weed Control Program:** Sharda Glyphosate 41 SL may be applied alone as a postemergence in-crop application to provide control of emerged weeds listed on the label. The postemergence application of Sharda Glyphosate 41 SL must be made before the weeds reach a height and/or density that the weeds become competitive with the crop. If new flushes of weeds occur, a sequential application of Sharda Glyphosate 41 SL at 24 to 32 fluid ounces per acre will control the labeled grasses and broadleaf weeds. Sharda Glyphosate 41 SL may be applied postemergence to Roundup Ready® corn from emergence through the V8 stage or until corn height reaches 30 inches (free standing), whichever comes first.

Sharda Glyphosate 41 SL may be applied in a tank mixture with a labeled rate of Harness®, Harness Xtra®, Harness Xtra 5.6L®, Micro-Tech®, Bullet®, Partner®, Permit® or atrazine. Refer to the specific product label and observe all precautions and limitations on the labels for all products used in tank mixtures, including application timing restrictions, soil restrictions, minimum re-cropping interval and rotational guidelines - the more restrictive requirements apply. Tank mixtures with other products may result in increased potential for crop injury and/or weed antagonism. Refer to the table below for height limitation for tank mix partner.

Tank Mix Partner	Maximum Height Of Corn For Application
Harness® Harness Xtra®, Harness Xtra 5.6L®	11 inches
Bullet*® Micro-Tech®* Partner®*	5 inches
Permit®	24 inches
Atrazine	12 inches

\*Bullet®, Micro-Tech® and Partner® are not registered for use as a postemergence application in Texas.

**POSTEMERGENCE APPLICATIONS TO ROUNDUP READY® CORN 2  
PRODUCT INFORMATION**

Sharda USA LLC directs use of Sharda Glyphosate 41 SL only on corn hybrids designated as containing the Roundup Ready® gene.

480967

- Applying Sharda Glyphosate 41 SL to corn hybrids which are not designated as Roundup Ready® will result in severe crop injury and yield loss.
- The RoundUp Ready® designation indicates that the corn contains a patented gene, which provides tolerance to Sharda Glyphosate 41 SL.

**NOTE:** The instructions provided in this supplemental label are specific to, and must only be used with, Roundup Ready® Corn 2 hybrids. Do not combine the instructions in this supplemental with any other Roundup Ready® corn instructions on labeling for this or other glyphosate containing product. The use of the higher in-crop over the top rates described in this supplemental label on other than Roundup Ready® Corn 2 may cause crop injury and reduced yields.

**APPLICATION INSTRUCTIONS**

For Roundup Ready® Corn 2 from emergence through the V8 stage (8 leaves with collars) or until corn height reaches 30 inches, whichever comes first, Sharda Glyphosate 41 SL may be applied over-the-top broadcast or with drop nozzles. When corn height is 24 to 30 inches (free standing), for optimum spray coverage and weed control, drop nozzles are directed. For corn heights 30 to 48 inches (free standing) apply Sharda Glyphosate 41 SL only using ground application equipment with the drop nozzles adjusted to avoid spraying into the whorls of the corn plants. Single in-crop applications of Sharda Glyphosate 41 SL must not exceed 48 fluid ounces per acre.

**Maximum Allowable Combined Application Quantities Per Season**

**Preplant, At Planting, Preemergence:** Maximum amount of Sharda Glyphosate 41 SL which can be applied prior to crop emergence is 5.0 quarts per acre.

**Postemergence (in crop):** Maximum combined total of multiple in-crop applications from emergence through 48 inch stage is 3 quarts per acre.

**Preharvest:** Maximum preharvest rate after maximum kernel fill is complete and the crop is physiologically mature (black layer formation) with 35 percent grain moisture or less until 7 days before harvest is 1 quart per acre. See precautions and restrictions on preharvest applications.

**Cropping Season:** Combined total per year for all applications may not exceed 8.0 quarts per acre.

The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of Sharda Glyphosate 41 SL under hard water conditions, drought, or when tank mixed with "Bullet®" or "Micro-Tech®" herbicides. Ensure that ammonium sulfate is completely dissolved in the spray tank before adding herbicides. Thoroughly rinse the spray system with clean water after use to reduce corrosion. The addition of other additives, including surfactants, fertilizers and/or micro-nutrients are not directed with Sharda Glyphosate 41 SL since this may result in increased potential for crop injury.

**ATTENTION: AVOID DRIFT.** EXTREME CARE MUST BE USED WHEN APPLYING SHARDA GLYPHOSATE 41 SL TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN THE ROUNDUP READY® GENE. THOROUGHLY CLEAN THE SPRAY TANK AND ALL LINES AND FILTERS TO ELIMINATE POTENTIAL CONTAMINATION FROM OTHER HERBICIDES PRIOR TO MIXING AND APPLYING SHARDA GLYPHOSATE 41 SL.

**For ground applications:** Use the specified rates of Sharda Glyphosate 41 SL in 5 to 20 gallons of spray solution per acre as a broadcast spray. Carefully select correct nozzles and spray pressure to avoid spraying a fine mist. Check for even distribution of spray droplets.

**For aerial applications:** Use the specified rates of Sharda Glyphosate 41 SL in 3 to 15 gallons of spray solution per acre. Do not exceed 32 fluid ounces per acre. See "WEEDS CONTROLLED" section on this label. AVOID DRIFT - DO NOT APPLY DURING INVERSION CONDITIONS, WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITIONS WHICH FAVOR DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

**Weed Control Directations**

Apply 24 to 32 fluid ounces of Sharda Glyphosate 41 SL per acre for control of labeled grasses and broadleaf weeds in conventional and no-till corn production systems. Refer to the label booklet for Sharda Glyphosate 41 SL for rate directations for specific annual weeds. Sharda Glyphosate 41 SL applied at up to 48 fluid ounces per acre will control or suppress the growth of perennial weeds such as: bermudagrass,



49267

Canada thistle, common milkweed, field bindweed, hemp dogbane, horse nettle, nutsedge, quackgrass, rhizome johnsongrass, redvine, trumpet creeper, swamp smartweed, and wirestem muhly. For additional information on perennial weeds, see the "WEEDS CONTROLLED" section of the label booklet for Roundup Original herbicide.

**Preplant, Preemergence, At-Planting**

**USE INSTRUCTIONS:** Sharda Glyphosate 41 SL may be applied alone or in a tank-mixture before, during or after planting corn.

**TANK MIXTURES:** Sharda Glyphosate 41 SL may be tank mixed with Bullet®, Degree®, Degree® Xtra, Harness®, Harness Xtra®, Harness Xtra 5.6L®, Lariat®, Lasso® or Micro-Tech® at 50 to 100 percent of labeled rate. Refer to the specific product label and observe all precautions and limitations on the label for any preemergence herbicide application, including application timing restrictions, soil restrictions, minimum recropping interval and rotational guidelines - the more restrictive requirements apply.

**NOTE:** For maximum weed control, a postemergence (in-crop) application of Sharda Glyphosate 41 SL must be applied following the use of less than labeled rates of the preemergence residual products listed above.

**Preemergence followed by Postemergence Weed Control Program**

**USE INSTRUCTIONS:** Sharda Glyphosate 41 SL may be applied postemergence in-crop following any labeled preemergence herbicide application. The post application of Sharda Glyphosate 41 SL must be made before the weeds reach a height and/or density that the weeds become competitive with the crop. An in-crop application of Sharda Glyphosate 41 SL at the specified rate will provide control of emerged weeds listed on the label. Sharda Glyphosate 41 SL may be applied over-the-top broadcast or with drop nozzles postemergence to Roundup Ready® Corn 2 from emergence through the V8 state (8 leaves with collars) or until corn height reaches 30 inches (free standing), whichever comes first. When corn height is 24 to 30 inches drop nozzles are directed for optimum spray coverage and weed control. For corn heights 30 to 48 inches (free standing), apply Sharda Glyphosate 41 SL only using ground application equipment with drop nozzles and avoid spraying into the whorls of the corn plants.

**Postemergence Only Weed Control Program**

**USE INSTRUCTIONS:** Sharda Glyphosate 41 SL may be applied alone as a postemergence in-crop application to provide control of emerged weeds listed on the label. The postemergence application of Sharda Glyphosate 41 SL must be made before the weeds reach a height and/or density that the weeds become competitive with the crop. If new flushes of weeds occur, a sequential application of Sharda Glyphosate 41 SL at 24 to 32 fluid ounces per acre will control the labeled grasses and broadleaf weeds. Sharda Glyphosate 41 SL may be applied over-the-top broadcast or with drop nozzles postemergence to Roundup Ready® Corn 2 from emergence through the V8 stage or until corn height reaches 30 inches (free standing), whichever comes first. When corn height is 24 to 30 inches drop nozzles are directed for optimum spray coverage and weed control. For corn height 30 to 48 inches (free standing), apply Sharda Glyphosate 41 SL only using ground application equipment with drop nozzles and avoid spraying into the whorls of the corn plants.

**TANK MIXTURES:** Sharda Glyphosate 41 SL may be applied in tank mixture with a labeled rate of Degree®, Degree® Xtra, Harness®, Harness Xtra®, Harness Xtra 5.6L®, Micro-Tech® and Bullet® at 50 to 100 percent of labeled rate. Sharda Glyphosate 41 SL may be applied in tank mixture with Permit® and atrazine at labeled rates. Refer to the specific product label and observe all precautions and limitations on the label for all products used in tank mixtures, including application timing restrictions, soil restrictions, minimum recropping interval and rotational guidelines - the more restrictive requirements apply. Tank mixtures with other products may result in increased potential for crop injury and/or weed antagonism. Refer to the table below for height limitation for tank mix partner.

Tank Mix Partner	Maximum Height of Corn
Degree® Degree® Xtra Harness® Harness Xtra® Harness Xtra 5.6L®	11 inches

Bullet® Micro-Tech® *	5 inches
Permit®	30 inches
Atrazine	12 inches

\*Bullet® and Micro-Tech® are not registered for use as a postemergence application in Texas.

**PRECAUTIONS, RESTRICTIONS:**

Single in-crop applications of Sharda Glyphosate 41 SL must not exceed 48 fluid ounces per acre.

Allow a minimum of 10 days between in-crop applications of Sharda Glyphosate 41 SL. Allow a minimum of 50 days between application of Sharda Glyphosate 41 SL and harvest of corn forage or grain.

For applications at preharvest timing (see Preharvest section of this supplemental label), allow a minimum of 7 days between application and harvest or feeding of corn stover or grain.

There are no rotational crop restrictions following applications of Sharda Glyphosate 41 SL.

**PREHARVEST USE INSTRUCTIONS:** A single preharvest application of up to 32 fluid ounces per acre of Sharda Glyphosate 41 SL may be made, if no more than a total of 64 fluid ounces of Sharda Glyphosate 41 SL has been previously applied in over-the-top or drop nozzle applications. Make a preharvest application at 35 percent grain moisture or less. Ensure that maximum kernel fill is complete and the corn is physiologically mature (black layer formed).

**PRECAUTIONS, RESTRICTIONS:**

Do not make a preharvest application of Sharda Glyphosate 41 SL if more than a combined total of 64 fluid ounces of Sharda Glyphosate 41 SL has been previously applied in over-the-top or drop nozzle applications. Allow a minimum of 7 days between a preharvest application and harvest or feeding of corn stover or grain.

**POSTHARVEST USE INSTRUCTIONS:** Sharda Glyphosate 41 SL may be applied after harvest of corn. Higher rates may be required for control of large weeds that were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

**PRECAUTIONS, RESTRICTIONS:** Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

**POSTEMERGENCE APPLICATIONS TO COTTON WITH THE ROUNDUP READY® GENE**

SHARDA USA LLC DIRECTS SHARDA GLYPHOSATE 41 SL FOR USE ONLY OVER-THE-TOP OF OR DIRECTED ONTO IMPROVED COTTON VARIETIES THAT ARE DESIGNATED AS COTTON WITH THE ROUNDUP READY® GENE.

- Severe injury or death of cotton will result if any cotton varieties not properly designated as having the Roundup Ready® gene are sprayed with Sharda Glyphosate 41 SL. Avoid contact of herbicide with foliage, green stems, or fruit of crops, or any desirable plants and trees, other than crops with the Roundup Ready® gene, since severe injury or destruction will result.
- The Roundup Ready® designation indicates that the cotton contains a patented gene which provides tolerance to glyphosate herbicides. Information on Roundup Ready® cotton may be obtained from your seed supplier.

**APPLICATION INSTRUCTIONS**

Sharda Glyphosate 41 SL will control many troublesome weeds with over-the-top, post-directed, hooded sprayer, or preharvest applications in Roundup Ready® cotton.

**Maximum Allowable Yearly Rates:**

- Combined total per year for all applications: 8 quarts per acre
- Preplant, Preemergence applications: 5 quarts per acre
- Total in-crop applications from cracking to layby: 4 quarts per acre
- Maximum preharvest application rate: 2 quarts per acre

**For ground applications with broadcast equipment,** apply Sharda Glyphosate 41 SL in 5 to 20 gallons of spray solution per acre. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For

5/10/07

best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

**For aerial applications,** apply Sharda Glyphosate 41 SL in 3 to 15 gallons of water per acre.

DO NOT EXCEED A MAXIMUM RATE OF 1 QUART PER ACRE OF SHARDA GLYPHOSATE 41 SL WHEN MAKING APPLICATIONS BY AIR. AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING SHARDA GLYPHOSATE 41 SL TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN THE ROUNDUP READY® GENE. Do not apply during low-level inversion conditions, when winds are gusty or under any other conditions which favor drift. Drift may cause damage to any vegetation contacted to which treatment is not intended. To prevent injury to adjacent desirable vegetation, appropriate buffer zones must be maintained. There are no rotational crop restrictions following applications of Sharda Glyphosate 41 SL.

**Sprayer Preparation:** It is important that sprayer and mixing equipment be clean and free of pesticide residue before making applications of Sharda Glyphosate 41 SL to Roundup Ready® cotton. Follow the cleaning procedures specified on the label of the product(s) previously used. Cotton is very sensitive to many herbicides at extremely low concentrations and care must be taken to thoroughly clean all equipment prior to use.

**Types of Applications to Roundup Ready® Cotton:**

**Preplant Burndown:** Always plant into a weed free seedbed. In no-till and stale seedbed systems, always burn down existing weeds before cotton emerges. Apply a preplant burndown treatment of 16 to 48 fluid ounces per acre of Sharda Glyphosate 41 SL.

**Over-the-top applications:** Sharda Glyphosate 41 SL may be applied by aerial or ground application equipment postemergence to Roundup Ready® cotton from the ground cracking stage until the four leaf (node) stage of development (until the fifth true leaf reaches the size of a quarter). Over-the-top applications made after the four leaf (node) stage of development may result in boll loss, delayed maturity and/or yield loss. Any single over-the-top broadcast application must not exceed 1 quart per acre. No more than two over-the-top broadcast applications may be made from crop emergence through the four leaf (node) stage of development. Sequential over-the-top applications of Sharda Glyphosate 41 SL must be at least 10 days apart and cotton must have at least two nodes of incremental growth between applications.

**Post-directed or hooded applications:** Sharda Glyphosate 41 SL may be applied using precision post-directed or hooded sprayers to Roundup Ready® cotton through layby. At this stage, post-directed equipment must be used which directs the spray to the base of the cotton plants. Contact of the spray with cotton leaves must be avoided to the maximum extent possible. To minimize spray onto the leaves of the cotton plants, place nozzles in a low position directing a horizontal spray pattern under the cotton leaves to contact weeds in the row, and maintain low spray pressure (less than 30 PSI). For best results, make applications while weeds are small (less than 3 inches).

Any single post-directed application must not exceed 1 quart per acre of Sharda Glyphosate 41 SL. No more than two applications must be made from the fifth leaf through layby. Sequential in-crop applications of Sharda Glyphosate 41 SL must be at least 10 days apart and cotton must have at least two nodes of incremental growth between applications.

**ATTENTION:** USE OF SHARDA GLYPHOSATE 41 SL IN ACCORDANCE WITH LABEL DIRECTIONS IS EXPECTED TO RESULT IN NORMAL GROWTH OF ROUNDUP READY® COTTON. HOWEVER, VARIOUS ENVIRONMENTAL CONDITIONS, PRODUCT, EVEN WHEN APPLICATIONS ARE MADE IN CONFORMANCE WITH THE LABEL SPECIFICATIONS. IN SOME CASES, THESE FACTORS CAN RESULT IN BOLL LOSS, DELAYED MATURITY AND/OR YIELD LOSS.

**Salvage Treatment:** This treatment may be used after the four leaf stage of development and must only be used where weeds threaten to cause the loss of the crop. One quart per acre may be applied either as an over-the-top applications or as a post-directed treatments sprayed higher on the cotton plants and over the weeds.

**NOTE: SALVAGE TREATMENTS WILL RESULT IN SIGNIFICANT BOLL LOSS, DELAYED MATURITY AND/OR YIELD LOSS. NO MORE THAN ONE SALVAGE TREATMENT MUST BE USED PER GROWING SEASON.**

**Weeds controlled:** For specific rates of application and instructions for control of various annual and perennial weeds, refer to "ANNUAL WEEDS RATE TABLE" section of this booklet. Sharda Glyphosate 41 SL applied at 1 quart per acre will burndown or suppress the growth of the following perennial weeds and

520967

reduce crop competition from yellow and purple nutsedge rhizome johnsongrass, common bermudagrass, silverleaf nightshade, trumpet creeper, and redvine. Fall preharvest applications may be required for control of these perennial weeds. Tank mixtures with other herbicides may result in reduced weed control or crop injury and are not directed for over-the-top applications of Sharda Glyphosate 41 SL. Some weeds with multiple germination times or suppressed (stunted) weeds may require sequential applications of Sharda Glyphosate 41 SL for control.

**Preharvest applications:** Sharda Glyphosate 41 SL may be applied for preharvest annual and perennial weed control as a broadcast treatment to Roundup Ready® cotton after 20% boll crack. Allow a minimum of 7 days between application and harvest.

**NOTE:** Sharda Glyphosate 41 SL will not enhance performance of harvest aids when applied to Roundup Ready® cotton. DO NOT APPLY THIS HERBICIDE PREHARVEST TO CROPS GROWN FOR SEED.

Read the "Limit of Warranty and Liability" in this label booklet for Sharda Glyphosate 41 SL before using. For over-the-top uses on Roundup Ready® crop varieties, crop safety and weed control performance are not warranted by Sharda USA LLC when Sharda Glyphosate 41 SL is used in conjunction with "brown bag" or "bin-run" seed saved from previous year's production and replanted. If these terms are not acceptable, return the product unopened at once.

**FALLOW AND REDUCED TILLAGE SYSTEMS  
FOR AERIAL APPLICATION IN CALIFORNIA, REFER TO SUPPLEMENTAL LABEL.**

Use Sharda Glyphosate 41 SL in fallow and reduced tillage systems for control of annual weeds prior to emergence of crops listed in this label. Refer to the "WEEDS CONTROLLED" section of this label for specific rates and instructions. Sharda Glyphosate 41 SL may be applied using ground or aerial spray equipment. See the "APPLICATION EQUIPMENT and TECHNIQUES" section of this label for instructions.

**TANK MIXTURES**

- Sharda Glyphosate 41 SL plus BANVEL® or dicamba
- Sharda Glyphosate 41 SL plus 2,4-D
- Sharda Glyphosate 41 SL plus GOAL®\*

**DO NOT APPLY BANVEL®, DICAMBA OR 2,4-D TANK MIXTURES BY AIR IN CALIFORNIA.**

Applications of 2,4-D, Banvel® or dicamba must be made at least 7 days prior to planting corn.

The addition of Banvei or dicamba in a mixture with Sharda Glyphosate 41 SL may provide short-term residual control of selected weed species. Some crop injury may occur if Banvel® or dicamba is applied within 45 days of planting. Refer to the Banvel®, dicamba or 2,4-D labels for cropping restrictions and other use instructions.

Sharda Glyphosate 41 SL may be tank-mixed with the listed herbicides provided the specific product is registered for use on these sites.

**SHARDA GLYPHOSATE 41 SL plus Goal® Tank Mixtures**

Sharda Glyphosate 41 SL alone or in tank mixtures with Goal® will provide control of those weeds listed below.

Make applications when weeds are actively growing and at the specified stages of growth. Avoid spraying when weeds are subject to moisture stress, when dust is on the foliage or when straw canopy covers the weeds.

Sharda Glyphosate 41 SL 12 fluid oz./acre		Sharda Glyphosate 41 SL 16 fluid oz./acre-Annual grasses at left plus:			
Wheat	18"	Ryegrass, annual	6"	Crabgrass	12"
Barley	12"	Chickweed	6"	Johnsongrass, seedling	12"
Bluegrass, annual	6"	Groundsel	6"	Lambsquarters	12"
Barnyardgrass	6"	Marestail	6"	Oats, wild	12"
Rye	6"	Rocket, London	6"	Pigweed, redroot	12"
		Shepherdspurse	6"	Mustards	12"

53067

NOTE: Use 32 fluid ounces of Sharda Glyphosate 41 SL per acre where heavy weed densities exist.

Sharda Glyphosate 41 SL 12 fluid oz./acre plus GOAL®** 2 to 4 fluid oz./acre	Sharda Glyphosate 41 SL 16 fluid oz./acre plus GOAL®** 2 to 4 fluid oz./acre
Annual grasses above plus:	Annual weeds above plus:
Cheeseweed, common 3"	Cheeseweed, common 6"
Chickweed 3"	Groundsel 6"
Groundsel 3"	Chickweed 12"
Rocket, London 6"	Rocket, London 12"
Shepherdspurse 6"	Shepherdspurse 12"

NOTE: Use 32 fluid ounces of Sharda Glyphosate 41 SL per acre in mixtures with 2 to 4 fluid ounces of Goal® per acre where heavy weed densities exist.

\* Maximum height or length in inches.

\*\* Use the higher rate of Goal® when weeds approach maximum specified height or stands are dense.

These specified tank mixtures may be applied using ground or aerial spray equipment. Refer to the "WEEDS CONTROLLED" section of this label for specific rates and instructions.

**ECOFARMING SYSTEMS**

The directions made in this section are not registered for use in California.

The Ecofarming System consists of the following rotation: winter wheat, corn/sorghum, ecofallow.

Use the following tank mixtures for control of emerged annual weeds before planting corn or sorghum in the Ecofarming System.

- Sharda Glyphosate 41 SL at 16 to 20 fluid ounces per acre
- Sharda Glyphosate 41 SL plus 2,4-D at 0.375 to 0.5 pound a.i. per acre
- Sharda Glyphosate 41 SL plus Atrazine at 0.75 to 1 pound a.i. per acre
- Sharda Glyphosate 41 SL plus Lasso® at 2.5 to 3 quarts per acre

The above tank mixture must be applied in 28-0-0 or 32-0-0 liquid fertilizer carrier at 20 to 30 gallons per acre. The liquid fertilizer may be diluted with water to achieve the desired carrier volume. Sharda Glyphosate 41 SL may be tank-mixed with the listed herbicides provided the specific product is registered for use on these sites.

Brome, downy ( <i>Bromus tectorum</i> )	Foxtail, yellow ( <i>Setaria lutescens</i> )	Lettuce, prickly ( <i>Lactuca serriola</i> )	Thistle, Russian ( <i>Salsola kali</i> )
Cheat ( <i>Bromus secalinus</i> )	Kochia* ( <i>Kochia scoparia</i> )	Pigweed, redroot ( <i>Amaranthus retroflexus</i> )	Wheat, volunteer ( <i>Triticum aestivum</i> )

\*For improved control of kochia, add 4 fluid ounces per acre (0.125 pound a.i. per acre) of Banvel® or dicamba to the above tank mixture.

Risk of crop injury from 2,4-D, Banvel® or dicamba can be reduced by applying this treatment 7 to 14 days before planting. Refer to the label booklet for Lasso® herbicide for preemergence weed control achieved by this tank mixture. Refer to the specific product labels for crop rotation restrictions and cautionary statements for all products used in these tank mixtures.

**AID TO TILLAGE**

Sharda Glyphosate 41 SL, when used in conjunction with preplant tillage practices, will provide control of downy brome, cheat, volunteer wheat, tansy mustard and foxtail. Apply 8 fluid ounces of Sharda Glyphosate 41 SL in 3 to 10 gallons of water per acre. Make applications when weeds are actively growing and before they are 6 inches in height. Application must be followed by conventional tillage practices no later than 15 days after treatment and before regrowth occurs. Allow at least 1 day after application before tillage. Tank mixtures with residual herbicides may result in reduced performance.

**POSTHARVEST GRAIN SORGHUM, SORGHUM REGROWTH CONTROL**

Sharda Glyphosate 41 SL may be applied to grain sorghum (milo) stubble following harvest to suppress or control regrowth. Apply 1 quart of Sharda Glyphosate 41 SL per acre for control, or 1.5 pints of Sharda Glyphosate 41 SL per acre for suppression.

**PASTURES**

CANOPY®	LINURON®	PURSUIT®
COMMAND®	LOROX® PLUS	PURSUIT PLUS®
DUAL®	MICRO-TECH®	SCEPTER®
GEMINI®	PARTNER®	SENCOR®
LASSO®/ALACHLOR	PREVIEW®	SQUADRON®
LEXONE®	PROWL®	TURBO®

For improved burndown, Sharda Glyphosate 41 SL may be tank-mixed with the following herbicides:

Sharda Glyphosate 41 SL may be tank-mixed with the listed herbicides provided the specific product is registered for use on these sites.

2,4-DB

2,4-D\* (WEEDONE® 638, WEEDAR® 64, others)

\*See the label for 2,4-D for intervals between application and planting.

**CORN AND SOYBEANS**

**Annual Weeds:** For difficult-to-control weeds such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply Sharda Glyphosate 41 SL at 2 pints per acre in these tank mixtures. For other labeled annual weeds, apply 1 to 1.5 pints of Sharda Glyphosate 41 SL per acre when weeds are less than 6 inches tall, and 2 to 3 pints when weeds are over 6 inches tall. For a complete list of annual weeds controlled, see the "WEEDS CONTROLLED" section of this label.

**Perennial Weeds:** At normal application times in minimum tillage systems, perennial weeds may not be at the proper stage of growth for control. See the "WEEDS CONTROLLED" section of this label for the proper stage of growth for perennial weeds. Use of 2 to 4 quarts of Sharda Glyphosate 41 SL per acre in the tank mixtures mentioned above, under these conditions provides top kill and reduces competition from many emerged perennial grass and broadleaf weeds. For emerged perennial weeds controlled, see the "WEEDS CONTROLLED" section of this label. To obtain the desired stage of growth, it may be necessary to apply Sharda Glyphosate 41 SL alone in the late summer or fall and then follow with a label-approved, seedling weed-control program at planting.

**CORN**

For residual control, Sharda Glyphosate 41 SL may be tank-mixed with the following herbicides or combination of herbicides:

ATRAZINE	DUAL®	PARTNER®*
BICEP®	LARIAT®	PROWL® or pendimethalin
BULLET®	LASSO®/ALACHLOR	SIMAZINE
CYANAZINE	MICRO-TECH®	

For improved burndown, Sharda Glyphosate 41 SL may be tank-mixed with 2,4-D or dicamba. Applications of 2,4-D or dicamba must be made at least 7 days prior to planting corn. See the "WEEDS CONTROLLED" section for specific rate information.

\*Partner® herbicide is not registered in California.

USE OF THESE TANK MIXTURES FOR BERMUDAGRASS OR JOHNSONGRASS CONTROL IN MINIMUM TILLAGE SYSTEMS IS NOT DIRECTED. For bermudagrass control, follow the instructions under "CONTROL OF PERENNIAL WEEDS" section of this label and then use a label-approved, seedling weed-control program in a minimum tillage or conventional tillage system. For Johnsongrass control, follow instructions under "CONTROL OF PERENNIAL WEEDS" section of this label, and then use a label-approved, seedling weed-control program with conventional tillage.

**POSTEMERGENCE APPLICATIONS TO SOYBEANS WITH THE ROUNDUP READY® GENE**

**Product Information**

SHARDA USA LLC DIRECTS USE OF SHARDA GLYPHOSATE 41 SL FOR POSTEMERGENCE APPLICATION ONLY ON SOYBEAN VARIETIES DESIGNATED AS CONTAINING THE ROUNDUP READY® GENE.

- Applying Sharda Glyphosate 41 SL to soybean varieties which are not designated as Roundup Ready® will result in severe crop injury and yield loss. Avoid contact with foliage, green stems, or fruit of crops, or any desirable plants which do not contain the Roundup Ready® gene, since severe injury or destruction will result.
- The Roundup Ready® designation indicates that the soybean contains a patented gene which provides tolerance to glyphosate herbicides. Information on Roundup Ready® soybeans may be obtained from your seed supplier.

**Application Instructions**

Sharda Glyphosate 41 SL may be applied postemergence to Roundup Ready® soybeans from the cracking stage throughout flowering. Allow a minimum of 14 days between application and harvest of soybeans.

**Maximum Allowable Application Rates:**

- Combined total per year for all applications: 8 quarts per acre
- Preplant, Preemergence applications: 5 quarts per acre
- Total in-crop applications from cracking throughout flowering: 3 quarts per acre
- Maximum preharvest application rate: 1 quart per acre

When applied as directed, Sharda Glyphosate 41 SL will control labeled annual grasses and broadleaf weeds in Roundup Ready® soybeans. Many perennial grasses and broadleaf weeds will be controlled or suppressed with one or more applications of Sharda Glyphosate 41 SL.

**PRECAUTIONS/RESTRICTIONS:** The combined total application from crop emergence through harvest must not exceed 3 quarts per acre. The maximum rate for any single in crop application is 2 quarts per acre. The maximum combined total of Sharda Glyphosate 41 SL which can be applied during flowering is 2 quarts per acre. Allow a minimum of 14 days between final application and harvest of soybeans. There are no rotational crop restrictions following applications of Sharda Glyphosate 41 SL.

**For ground applications:** Use the specified rates of Sharda Glyphosate 41 SL in 5 to 20 gallons of spray solution per acre as a broadcast spray. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment, use nozzles which provide a flat fan pattern. Check for even distribution of spray droplets.

**For aerial applications:** Use the specified rates of Sharda Glyphosate 41 SL in 3 to 15 gallons of spray solution per acre. Do not exceed 1 quart of Sharda Glyphosate 41 SL per acre. DO NOT APPLY DURING LOW LEVEL INVERSION CONDITIONS. WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITIONS WHICH FAVOR DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. MAINTAIN APPROPRIATE BUFFER ZONES TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION.

AERIAL APPLICATIONS ON ROUNDUP READY® SOYBEANS, MAY BE MADE ONLY IN THE FOLLOWING STATES: ALABAMA, ARKANSAS, COLORADO, FLORIDA, GEORGIA, KANSAS, LOUISIANA, MISSISSIPPI, MISSOURI (BOOT HEEL ONLY), NEBRASKA, NORTH CAROLINA, NORTH DAKOTA, OKLAHOMA, SOUTH CAROLINA, SOUTH DAKOTA, TENNESSEE, TEXAS, VIRGINIA, AND WYOMING.

**ANNUAL WEED RATE DIRECTIONS**

The following rate directions will provide control of labeled grasses and broadleaf weeds in conventional and no-till soybean production systems. Refer to the rate directions for specific annual weeds in the "ANNUAL WEEDS" section of the label.

Sharda USA LLC will not warrant crop safety or weed control when Roundup Ready® soybeans are treated with herbicides not specified on this label. Because of the potential for; 1) crop injury, 2) poor weed control from antagonism, and/or 3) rotational crop restrictions; herbicides not specified on this label (or current supplemental label) ARE APPLIED AT THE SOLE RISK OF THE BUYER AND USER, whether applied preemergence or applied postemergence as a tank mixture with Sharda Glyphosate 41 SL. Sharda Glyphosate 41 SL may be used up to 2 quarts per acre in any single application for control of annual weeds, where heavy weed densities exist.

**Preplant Burndown:** The following directions are based on a clean start at planting by using a burndown application or tillage to control existing weeds before crop emergence, in no-till and stale seedbed systems,

50067

a preplant burn-down treatment of 16 to 64 fluid ounces per acre of Sharda Glyphosate 41 SL can be used to control existing weeds prior to crop emergence.

**MIDWEST/ MID-ATLANTIC DIRECTIONS**

**Narrow row or drilled soybeans:** A single in-crop application of Sharda Glyphosate 41 SL will provide effective control of labeled weeds. For best results, an initial application of 1 quart per acre on 4 to 8" weeds is directed. Weeds will generally be 4 to 8" tall 3 to 5 weeks after planting. If the initial application is delayed and weeds are 8 to 18" tall, use 48 ounces per acre for best results. Under adverse growing conditions such as drought, hail, wind damage or a poor soybean stand that slows or delays canopy closure, a sequential application of Sharda Glyphosate 41 SL at 24 to 32 fluid ounces per acre may be necessary to control late flushes of weeds. **Wide row soybeans:** An in-crop application of Sharda Glyphosate 41 SL will provide effective control of the initial stand of labeled weeds. For best results, an initial application of 1 quart per acre on 4 to 8" weeds is directed. Weeds will generally be 4 to 8" tall 3 to 5 weeks after planting. If new flushes of weeds occur, they can be controlled by sequential applications of Sharda Glyphosate 41 SL.

Initial and Sequential (if needed) Applications	
Weed Height (inches)	Rate (fluid ounces per acre)
1 – 3	24
4 – 8	32
8 – 18	48

**Giant ragweed:** Apply 1 quart per acre when the weed is 8 to 12" tall to avoid the need for sequential application.

**Black nightshade, Pennsylvania smartweed, Velvetleaf, and Waterhemp:** Apply 1 quart per acre to weeds 3 to 6" tall, and 48 fl oz per acre when weeds are up to 12" tall.

**Morning glory species:** Apply 1 quart when weeds are up to 4" tall and 48 fl. oz. per acre when weeds are up to 6" tall.

**Sequential Application for Certain Weeds:** Some weeds, such as black nightshade, woolly cupgrass, shattercane, wild proso millet, burcumber, and giant ragweed, with multiple germination times may require a sequential application of Sharda Glyphosate 41 SL. Suppressed or stunted weeds may also require sequential applications. Sequential applications must be made after some regrowth has occurred. Use a minimum of 24 fluid ounces of Sharda Glyphosate 41 SL per acre for sequential applications.

**SOUTHEAST DIRECTIONS**

**Narrow row, drilled, or wide-row soybeans:** An in-crop application of Sharda Glyphosate 41 SL will provide effective control of the initial stand of labeled weeds. For best results, an initial application of 1 quart per acre on 3 to 6" weeds is directed. Weeds will generally be 3 to 6" tall 2 to 3 weeks after planting.

Weed Height (inches)	Rate (fluid ounces per acre)
3 – 6	32
6 – 12	48

Under adverse growing conditions such as drought, hail, wind damage or a poor soybean stand that slows or delays canopy closure, a sequential application of Sharda Glyphosate 41 SL at 16 to 32 fluid ounces per acre may be necessary to control late flushes of weeds.

Sequential (if needed) Applications	
Weed Height (inches)	Rate (fluid ounces per acre)
2 – 3	16
3 – 6	24
6 – 12	32

**Florida pusley, hemp sesbania and spurred anoda:** Apply 1 quart per acre to weeds 2 to 4" tall for the initial application. Apply 1 quart per acre when these weeds are 3 to 6" tall if a sequential application is necessary.



570967

**Morning glory, black nightshade, groundcherry, and Pennsylvania smartweed:** Apply 24 fl. oz. per acre on 1 to 3" weeds, 32 fl. oz. per acre on 3 to 6" weeds, or 48 fl. oz. per acre on 6 to 12" weeds for the initial application.

**Sequential Application for Certain Weeds:** Some weeds, such as black nightshade, broadleaf signalgrass, Texas panicum, burcumber, and sicklepod, with multiple germination times may require a sequential application of Sharda Glyphosate 41 SL. Suppressed or stunted weeds may also require sequential applications. Sequential applications must be made after some regrowth has occurred. Use a minimum of 16 fluid ounces of Sharda Glyphosate 41 SL per acre for sequential applications. The combined total of all in-crop postemergence treatments must not exceed 96 fluid ounces per acre.

**DELTA/MID-SOUTH DIRECTIONS**

**Narrow row, drilled, or wide row soybeans:** An in-crop application of Sharda Glyphosate 41 SL will provide effective control of the initial stand of labeled weeds. A sequential application will be required to control new flushes of weeds. For best results, an initial application of 32 fluid ounces per acre on 2 to 4" weeds is directed. Weeds will generally be 2 to 4" tall 2 to 3 weeks after planting.

Initial Application	
Weed Height (inches)	Rate (fluid ounces per acre)
2 - 4	32
5 - 12	48

Sequential Applications	
Weed Height (inches)	Rate (fluid ounces per acre)
2 - 3	16
3 - 6	24
6 - 12	32

**Hemp sesbania and spurred anoda:** Apply a sequential treatment of 32 fl. oz. per acre on 3 to 6" weeds if necessary.

**Sequential Application for Certain Weeds:** Some weeds, such as black nightshade, broadleaf signalgrass, Texas panicum, burcumber, and sicklepod, with multiple germination times may require a sequential application of Sharda Glyphosate 41 SL. Suppressed or stunted weeds may also require sequential applications. Sequential applications must be made after some regrowth has occurred. Use a minimum of 16 fluid ounces of Sharda Glyphosate 41 SL per acre for sequential applications.

**PERENNIAL WEEDS RATE INSTRUCTIONS**

At the rate of 1 to 2 quarts per acre (single or multiple applications), Sharda Glyphosate 41 SL will control or suppress perennial weeds such as: bermudagrass, Canada thistle, common milkweed, field bindweed, hemp dogbane, horsenettle, maretail (horseweed), nutsedge, quackgrass, rhizome johnsongrass, redvine, trumpetcreeper, swamp smartweed, and wirestem muhly.

For best results, allow perennial weed species to achieve at least 6" of growth before spraying with this herbicide. For additional information on perennial weeds, see the "PERENNIAL WEEDS" section of this label. For some perennial species, repeat application may be required to eliminate crop competition throughout the growing season.

**PREHARVEST APPLICATIONS ON ALFALFA COTTON, GRAIN SORGHUM, SOYBEANS AND WHEAT**

When applied as directed under the conditions described, Sharda Glyphosate 41 SL controls annual and perennial weeds listed on this label prior to the harvest of cotton, grain sorghum, soybeans and wheat. For specific rates and application instructions for control of various annual and perennial weeds, see the "WEEDS CONTROLLED" section of this label. Sharda Glyphosate 41 SL may be applied by both ground and aerial application equipment. DO NOT APPLY MORE THAN 1 QUART PER ACRE OF SHARDA GLYPHOSATE 41 SL BY AIR. See the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for instructions for ground and aerial applications.

580A67

**NOTE:** Do not apply to crops grown for seed unless the likelihood of a reduction in germination and/or vigor is acceptable. Reduction in germination or vigor may occur.

**The use of Sharda Glyphosate 41 SL for preharvest grain sorghum (Milo) is not registered in California.**

### **SOYBEANS**

Apply after pods have set and lost all green color. Allow a minimum of 7 days between application and harvest of soybeans. Care must be taken to avoid excessive seed shatter loss due to ground application equipment. Do not graze or harvest treated crop for livestock feed within 25 days of last preharvest application. **DO NOT APPLY MORE THAN 6 QUARTS PER ACRE OF SHARDA GLYPHOSATE 41 SL FOR PREHARVEST APPLICATIONS.**

### **ALFALFA**

Sharda Glyphosate 41 SL may be used in declining alfalfa stands or any stand of alfalfa where crop destruction is acceptable. This application will severely injure or destroy the stand of alfalfa. The application rate of 1 quart per acre will control most annual and perennial weeds, including quackgrass, when applied prior to the harvest of alfalfa. The treated crop can be harvested and fed to livestock after 36 hours. Allow a minimum of 36 hours between application and harvest. For best results, harvest within 7 days of spraying.

Applications may be made at any time of year. Make only one preharvest application to an existing stand of alfalfa per year. For control of quackgrass, apply in the spring, late summer or fall when quackgrass is actively growing and at the proper growth stage (6 to 8 inches or more in height). Treatments for quackgrass must be followed by deep tillage for complete control.

**DO NOT APPLY MORE THAN 1 QUART OF SHARDA GLYPHOSATE 41 SL PER ACRE AS A PREHARVEST TREATMENT TO ALFALFA.**

### **COTTON**

**Broadcast Applications:** Sharda Glyphosate 41 SL may be applied using either aerial or ground spray equipment. For ground applications with broadcast equipment, apply Sharda Glyphosate 41 SL in 10 to 20 gallons of water per acre. For aerial applications, apply Sharda Glyphosate 41 SL in 3 to 10 gallons of water per acre. Sharda Glyphosate 41 SL provides weed control and cotton regrowth inhibition when applied prior to the harvest of cotton. Apply 1 to 2 quarts of Sharda Glyphosate 41 SL in 3 to 10 gallons of water per acre for cotton regrowth inhibition. Do not apply more than 2 quarts of Sharda Glyphosate 41 SL per acre for preharvest applications. **THE USE OF ADDITIVES FOR PREHARVEST APPLICATION TO COTTON IS PROHIBITED.**

Sharda Glyphosate 41 SL may be tank mixed with DEF® 6, Folex® or Prep® to provide additional enhancement of cotton leaf drop. Allow a minimum of 7 days between application and harvest of cotton.

Apply after sufficient bolls have developed to produce the desired yield of cotton. Applications made prior to this time could affect maximum yield potential. Do not feed or graze treated cotton forage or hay following preharvest applications.

### **GRAIN SORGHUM (MILO)**

Make applications at 30% grain moisture or less and at least 7 days prior to harvest. Apply up to 2 quarts of Sharda Glyphosate 41 SL per acre.

### **WHEAT**

Apply after the hard-dough stage of grain (30% or less grain moisture) and at least 7 days prior to harvest.

**DO NOT APPLY MORE THAN 1 QUART PER ACRE OF SHARDA GLYPHOSATE 41 SL FOR PREHARVEST APPLICATIONS TO WHEAT.**

### **PREHARVEST (FEED BARLEY AND WHEAT ONLY)**

**USE INSTRUCTIONS:** Sharda Glyphosate 41 SL provides weed control when applied prior to harvest of wheat or feed barley. For wheat, apply after the hard-dough stage of grain (30 percent or less grain moisture). For feed barley, apply after the hard dough-stage and when the grain contains 20 percent moisture or less. Stubble may be grazed immediately after harvest. Sharda Glyphosate 41 SL may be applied using either aerial or ground spray equipment. For ground applications, apply Sharda Glyphosate 41 SL in 3 to 10 gallons of water per acre.

**PRECAUTIONS, RESTRICTIONS:** Do not apply more than 1 quart of Sharda Glyphosate 41 SL per acre. Allow 7 days between application and harvest, feeding or grazing. Preharvest application is not directed for wheat or barley grown for seed, as a reduction in germination or vigor may occur.

**PREHARVEST AND SPOT TREATMENTS OF WEEDS IN DRY PEAS, LENTILS, AND CHICKPEAS BROADCAST SPRAY**

Sharda Glyphosate 41 SL may be applied as an over-the-top broadcast spray to control labeled weeds prior to the harvest of dry peas, lentils, and chickpeas. Apply up to 24 fluid ounces in 3 to 20 gallons of water per acre at the hard dough stage of the legume seed (30 percent grain moisture or less). Either ground broadcast or aerial applications may be made.

**PRECAUTIONS, RESTRICTIONS:**

- Apply at least 14 days before harvest.
- Only one application per year may be made; do not combine a preharvest spray with a spot treatment on the same crop area.
- Employ at least a 30 day plant-back interval between treatment and replanting for any crop not listed in Sharda Glyphosate 41 SL's label.
- Preharvest application is not directed for peas, lentils, or chickpeas grown for seed, as a reduction in germination or vigor may occur.
- Do not feed treated vines and hay from these crops to livestock.
- Do not apply Sharda Glyphosate 41 SL through any type of irrigation system.
- Do not treat field (feed) peas, since these are considered to be grown as livestock feed.

**SPOT TREATMENTS**

Sharda Glyphosate 41 SL may be applied as a spot treatment to control troublesome weeds such as Canada thistle, quackgrass, mayweed (dog fennel), and milkweed in dry peas, lentils, and chickpeas. Apply up to 20 fluid ounces in 10 to 20 gallons of water through ground spray equipment or use a 2 percent solution in a handheld sprayer. For best results, applications must be made at or beyond the bud stage of growth. The crop receiving spray in treated areas will be killed.

**PRECAUTIONS, RESTRICTIONS:**

- Apply at least 14 days before harvest.
- Only one application per year may be made; do not combine a preharvest spray with a spot treatment on the same crop area.
- Employ at least a 30 day plant-back interval between treatment and replanting for any crop not listed in this label.
- Do not feed treated vines and hay from these crops to livestock. Do not apply Sharda Glyphosate 41 SL through any type of irrigation system.
- Do not treat field (feed) peas, since these are considered to be grown as livestock feed.

**PREHARVEST AND SPOT TREATMENTS OF WEEDS IN DRY BEANS BROADCAST SPRAY**

Sharda Glyphosate 41 SL may be applied as an over-the-top broadcast spray to control labeled weeds prior to the harvest of dry beans. Apply up to 32 fluid ounces in 3 to 20 gallons of water per acre at the hard dough stage of the legume seed (30 percent grain moisture or less). Either ground broadcast or aerial applications may be made.

**PRECAUTIONS, RESTRICTIONS:**

- Apply at least 7 days before harvest.
- Only one application per year may be made; do not combine a preharvest spray with a spot treatment on the same crop area.
- Employ at least a 30 day plant-back interval between treatment and replanting for any crop not listed in this label.
- Preharvest application is not directed for dry beans grown for seed, as a reduction in germination or vigor may occur.
- Do not feed treated vines and hay from these crops to livestock. Do not apply Sharda Glyphosate 41 SL through any type of irrigation system.
- Do not treat field cowpeas, since these are considered to be grown as livestock feed.

60067

Apply Sharda Glyphosate 41 SL prior to planting forage grasses and legumes.

**Pasture or Hay Crop Renovation:** When applied as a broadcast spray, Sharda Glyphosate 41 SL controls the annual and perennial weeds listed in this label prior to planting forage grasses or legumes. Remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting.

**Spot Treatment:** When applied as a spot treatment as specified, Sharda Glyphosate 41 SL controls annual and perennial weeds listed in this label which are growing in pastures, forage grasses and forage legumes composed of bahiagrass, bermudagrass, bluegrass, brome, fescue, orchardgrass, ryegrass, timothy, wheatgrass, alfalfa or clover.

**Wiper Application:** When applied as directed, Sharda Glyphosate 41 SL controls or suppresses the weeds listed under "WIPER APPLICATORS" in the "SELECTIVE EQUIPMENT" section of this label.

For spot treatment and wiper application, apply in areas where the movement of domestic livestock can be controlled. No more than one-tenth of any acre must be treated at one time. Further applications may be made in the same area at 30-day intervals. Remove domestic livestock before application and wait 14 days after application before grazing livestock or harvesting.

**SUGARCANE**

When applied as directed for "CROPPING SYSTEMS", under the conditions described, Sharda Glyphosate 41 SL controls those emerged annual and perennial weeds listed on this label growing in or around sugarcane or in fields prior to the emergence of plant cane. Sharda Glyphosate 41 SL will also control undesirable sugarcane.

**NOTE:** Where repeat treatments are necessary, do not exceed a total of 10.6 quarts of Sharda Glyphosate 41 SL per acre per year. Do not apply to vegetation in or around ditches, canals or ponds containing water to be used for irrigation.

**Broadcast Treatment:** Apply Sharda Glyphosate 41 SL in 10 to 40 gallons of water per acre on emerged weeds prior to the emergence of plant cane. For specific rates of application and instructions for control of various annual and perennial weeds, see the "WEEDS CONTROLLED" section of this label. For removal of last stubble or ratoon cane, apply 4 to 5 quarts of Sharda Glyphosate 41 SL in 10 to 40 gallons of water per acre to new growth having at least 7 or more new leaves. Allow 7 or more days after application before tillage.

**Spot Treatment in or Around Sugarcane Fields:** For dilution and rates of application using hand-held equipment, see "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" and "WEEDS CONTROLLED" sections of this label. For control of volunteer or diseased sugarcane, make a 1 percent solution of Sharda Glyphosate 41 SL in water and spray to wet the foliage of vegetation to be controlled.

**NOTE:** When spraying volunteer or diseased sugarcane, the plants must have at least 7 new leaves. Avoid spray contact with healthy cane plants since severe damage or destruction may result. Do not feed or graze treated sugarcane forage following application.

**CONSERVATION TILLAGE, MINIMUM TILLAGE AND NO-TILL SYSTEMS  
CORN AND SOYBEANS TANK MIXTURES**

The directions made in this section are not registered for use in California.

When applied as specified under the conditions described, the tank mixtures listed in this section control many emerged weeds, and give preemergence control of many annual weeds where corn or soybeans will be planted directly into a cover crop, established sod or in previous crop residues.

Refer to specific product labels for crop rotation restrictions and precautionary statements of all products used in these tank mixtures. For mixing instructions, see the "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" section of this label. Apply these tank mixtures in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre before, during or after planting.

Do not apply these mixtures after crop emergence. The addition of 1 to 2 percent dry ammonium sulfate by weight may increase the performance of Sharda Glyphosate 41 SL.

**NOTE:** When using these tank mixtures, do not exceed 4 quarts of Sharda Glyphosate 41 SL per acre.

**SOYBEANS**

For residual control, Sharda Glyphosate 41 SL may be tank-mixed with the following herbicides or combination of herbicides:

**SPOT TREATMENTS**

Sharda Glyphosate 41 SL may be applied as a spot treatment to control troublesome weeds such as Canada thistle, quackgrass, mayweed (dog fennel), and milkweed in dry beans. Apply up to 26 fluid ounces in 10 to 20 gallons of water through ground spray equipment or use a 2 percent solution in a handheld sprayer. For best results, applications must be made at or beyond the bud stage of growth. The crop receiving spray in treated areas will be killed.

**PRECAUTIONS, RESTRICTIONS:**

- Apply at least 14 days before harvest.
- Only one application per year may be made; do not combine a preharvest spray with a spot treatment on the same crop area.
- Employ at least a 30 day plant-back interval between treatment and replanting for any crop not listed in Sharda Glyphosate 41 SL's label.
- Do not feed treated vines and hay from these crops to livestock. Do not apply Sharda Glyphosate 41 SL through any type of irrigation system.
- Do not treat field (feed) peas, since these are considered to be grown as livestock feed.

**TREE AND VINE CROPS**

Sharda Glyphosate 41 SL is specified for weed control in established groves, vineyards, and orchards, or for site preparation prior to transplanting crops listed in this section. Applications may be made with boom equipment, CDA, shielded sprayers, hand-held and high-volume wands, lances, orchard guns or with wiper applicator equipment, except as directed in this section. See the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for specific information on use of equipment. When applying Sharda Glyphosate 41 SL, refer to the "WEEDS CONTROLLED" section of this label and to specific directions in this section for rates to be used.

**NOTE:** Repeat treatments may be necessary to control weeds originating from underground parts of untreated weeds or from seeds. Sharda Glyphosate 41 SL does not provide residual weed control. For subsequent weed control, use repeated applications of Sharda Glyphosate 41 SL. Do not apply more than 10.6 quarts of Sharda Glyphosate 41 SL per acre per year.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE SOLUTION, SPRAY, DRIFT OR MIST WITH FOLIAGE OR GREEN BARK OF TRUNK, BRANCHES, SUCKERS, FRUIT OR OTHER PARTS OF TREES OR VINES. CONTACT OF SHARDA GLYPHOSATE 41 SL WITH OTHER THAN MATURED BROWN BARK CAN RESULT IN SERIOUS CROP DAMAGE. AVOID PAINTING OUT STUMPS WITH SHARDA GLYPHOSATE 41 SL AS INJURY RESULTING FROM ROOT GRAFTING MAY OCCUR IN ADJACENT TREES.

Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed or cut and have not been allowed to regrow to the specified stage for treatment.

For specific rates of applications and instructions, see the "WEEDS CONTROLLED" section of this label, and to specific directions which follow.

**MIDDLES MANAGEMENT**

**FOR ANNUAL WEEDS IN MIDDLES BETWEEN ROWS OF TREE AND VINE CROPS**

For citrus crops, treat uniformly between trees.

Sharda Glyphosate 41 SL alone or in mixtures with Goal® will control or suppress the annual weeds listed below.

Apply the specified rates of Sharda Glyphosate 41 SL, either alone or in mixtures with Goal® in 3 to 10 gallons of water per acre. Apply when weeds are actively growing and less than 6 inches in height or diameter. If weeds are under drought stress, irrigate prior to application. Reduced control may occur if weeds have been mowed prior to application. Up to 48 fluid ounces per acre of Sharda Glyphosate 41 SL may be used to control weeds, which have been mowed, are stressed or are growing in dense populations.

Weed	Maximum Height/Diameter (Inches)	Rate Per Acre Sharda Glyphosate 41 SL (fl. oz.)	Rate Per Acre Goal® (fl. oz.)
Barley ( <i>Hordeum vulgare</i> )	6	8	--

620967

Bluegrass, annual ( <i>Poa annua</i> )			
Barnyardgrass ( <i>Echinochloa crus-galii</i> ) Chickweed, common ( <i>Stellaria media</i> ) Red Maids ( <i>Calandrinia ciliata</i> )	6	12	--
Crabgrass ( <i>Digitaria spp.</i> ) Fleabane, hairy ( <i>Conyza bonariensis</i> ) Goundsel, common ( <i>Senecio vulgaris</i> ) Junglerice ( <i>Echinochloa colonum</i> ) Lambsquarters, common ( <i>Chenopodium album</i> ) Pigweed, redroot ( <i>Amaranthus retroflexus</i> ) Rocket, London ( <i>Sisymbrium irio</i> ) Ryegrass, common ( <i>Lolium multiflorum</i> )	6	16 or 16 to 32 plus	--  4 to 16**
Shepherdspurse ( <i>Capsella bursa-pastoris</i> ) Sowthistle, annual ( <i>Sonchus oleraceus</i> )	6	16 to 32 plus	4 to 16**
Cheesewood, common ( <i>Malva spp.</i> )	3	12 to 32 plus	4 to 16
Cheese weed, common ( <i>Malva spp.</i> ) Cheese weed, common ( <i>Malva spp.</i> ) Filaree* ( <i>Erodium spp.</i> ) Horseweed/Marestail ( <i>Conyza Canadensis</i> ) Nettle, stinging ( <i>Urtica dioica</i> ) Purselane, common* ( <i>Purtulaca oleracea</i> )	6	16 to 32 plus	4 to 16

\* Suppression only.

\*\*The mixture of Sharda Glyphosate 41 SL plus Goal® is directed when weeds are stressed or growing in dense populations.

**STRIPS**

**FOR ANNUAL AND PERENNIAL WEEDS IN STRIPS OF TREE AND VINE CROPS**

**TANK MIXTURES WITH RESIDUAL HERBICIDES**

When applied as a tank mixture, Sharda Glyphosate 41 SL provides control of the emerged annual weeds and control or suppression of emerged perennial weeds listed in this label. The following residual herbicides will provide preemergence control of those weeds listed in the individual product labels.

- Sharda Glyphosate 41 SL plus GOAL® 2XL
- Sharda Glyphosate 41 SL plus KARMEX® DF
- Sharda Glyphosate 41 SL plus KROVAR I®
- Sharda Glyphosate 41 SL plus KROVAR II®
- Sharda Glyphosate 41 SL plus PRINCEP® CALIBER® 90
- Sharda Glyphosate 41 SL plus SIMAZINE 4L, 80W or 90DF
- Sharda Glyphosate 41 SL plus SOLICAM® 80DF
- Sharda Glyphosate 41 SL plus SURFLAN® AS or 75W
- Sharda Glyphosate 41 SL plus PRINCEP® CALIBER® 90, SIMAZINE 4L, 80W or 90DF plus SURFLAN® AS or 75W
- Sharda Glyphosate 41 SL plus GOAL® 2XL plus SURFLAN® AS or 75W
- Sharda Glyphosate 41 SL plus GOAL® 2XL plus PRINCEP® CALIBER® 90, SIMAZINE 4L, 80W or 90DF
- Sharda Glyphosate 41 SL plus GOAL® 2XL plus SURFLAN® AS or 75W plus PRINCEP® CALIBER® 90, SIMAZINE 4L, 80W or 90DF

Do not apply these tank mixtures in Puerto Rico.

Refer to the individual product labels for specific crops, rates, geographical restrictions and precautionary statements.

Read and carefully observe the label claims, cautionary statements, rates and all other information on the labels of all products.

630f67

## SPECIFIED RATES

**Annual Weeds:** Apply 1 to 5 quarts per acre of Sharda Glyphosate 41 SL in these tank mixtures. Use rates at the higher end of the specified range when weeds are stressed, growing in dense populations or are greater than 12 inches tall.

**Perennial Weeds:** Apply 1 pint to 5 quarts per acre of Sharda Glyphosate 41 SL in these tank mixtures to control or suppress perennial weeds. Follow the directions in the "WEEDS CONTROLLED" section of this label for stage of growth and application rates for specific perennial weeds.

### Sharda Glyphosate 41 SL plus GOAL® plus SIMAZINE/SURFLAN®

Sharda Glyphosate 41 SL plus low rates of Goal® in 3-way or 4-way mixtures with simazine and/or Surflan® will provide postemergence control of the weeds listed below. Refer to the individual simazine and Surflan® labels for preemergence rates, weeds controlled, precautionary statements and other important information.

Apply these tank mixtures in 3 to 40 gallons of water. Apply 1 to 5 quarts per acre of Sharda Glyphosate 41 SL plus 4 to 48 fluid ounces per acre of Goal® plus labeled rates of simazine and/or Surflan® to control the following weeds:

Barley, wild ( <i>Hordeum leporinum</i> )	Filaree* ( <i>Erodium spp.</i> )	Nettle, stinging ( <i>Urtica dioica</i> )	Sowthistle, annual ( <i>Sonchus oleraceus</i> )
Bluegrass, annual ( <i>Poa annua</i> )	Fleabane, hairy ( <i>Conyza bonariensis</i> )	Pineappleweed ( <i>Matricaria matricarioides</i> )	
Cheeseweed, common ( <i>Malva spp.</i> )	Groundsel, common ( <i>Senecio vulgaris</i> )	Rocket, London ( <i>Sisymbrium irio</i> )	
Chickweed, common ( <i>Stellaria media</i> )	Horseweed/Marestail ( <i>Conyza Canadensis</i> )	Shepherdspurse ( <i>Capsella bursa-pastoris</i> )	

\*Use a minimum of 1.5 quarts of Sharda Glyphosate 41 SL in these mixtures.

**NOTE:** This direction does not preclude the use of Goal® in these mixtures at higher, labeled rates for preemergence weed control.

## PERENNIAL GRASS SUPPRESSION ON ORCHARD FLOORS

When applied as directed, Sharda Glyphosate 41 SL will suppress vegetative growth as indicated below.

**Bahiagrass:** Sharda Glyphosate 41 SL will provide significant inhibition of seedhead emergence and will suppress vegetative growth for a period of approximately 45 days with a single application and approximately 120 days with sequential applications. Apply Sharda Glyphosate 41 SL 1 to 2 weeks after full green-up or after mowing to a uniform height of 3 to 4 inches. Applications must be made prior to seedhead emergence. Apply 6 fluid ounces of Sharda Glyphosate 41 SL in 10 to 25 gallons of water per acre. Sequential applications of Sharda Glyphosate 41 SL may be made at approximately 45-day intervals to extend the period of seedhead and vegetative growth suppression. For continued seedhead suppression, sequential applications must be made prior to seedhead emergence. Apply no more than 2 sequential applications per year. As a first sequential application, apply 4 fluid ounces of Sharda Glyphosate 41 SL. A second sequential application of 2 to 4 fluid ounces may be made approximately 45 days after the last application.

**Bermudagrass:** For burndown, apply 1 to 2 quarts of Sharda Glyphosate 41 SL in 3 to 20 gallons of water per acre. Use 1 quart of Sharda Glyphosate 41 SL in 3 to 20 gallons of water per acre east of the Rocky Mountains. Use 1 to 2 quarts of Sharda Glyphosate 41 SL in 3 to 10 gallons of water per acre west of the Rocky Mountains. Use this treatment only if reduction of the bermudagrass stand can be tolerated. When burndown is required prior to harvest, allow at least 21 days to ensure sufficient time for burndown to occur.

**Suppression only (east of the Rocky Mountains):** Apply 6 to 16 fluid ounces of Sharda Glyphosate 41 SL in 3 to 20 gallons of water per acre no sooner than 1 to 2 weeks after full green-up. Mowing prior to application may occur provided a minimum height of 3 inches is maintained. Rates of 6 to 10 fluid ounces of Sharda Glyphosate 41 SL must be used in shaded conditions or where a lesser degree of suppression is desired. Sequential applications may be made when regrowth occurs and bermudagrass injury and stand reduction can be tolerated.

**Suppression only (west of the Rocky Mountains):** Apply 16 fluid ounces of Sharda Glyphosate 41 SL in 3 to 10 gallons of water per acre to bermudagrass up to 6 inches in height and no sooner than 1 to 2 weeks after full green-up. Mowing prior to application may occur provided a minimum height of 3 inches is

64067

maintained. Sequential applications may be made when regrowth occurs and bermudagrass injury and stand reduction can be tolerated.

**Cool Season Grass Covers**

For suppression of tall fescue, fine fescue, orchardgrass and quackgrass, apply 8 fluid ounces of Sharda Glyphosate 41 SL in 10 to 20 gallons of water per acre. For best suppression, add ammonium sulfate to the spray solution at a rate of 2 percent by weight or 17 pounds per 100 gallons of spray solution. For suppression of Kentucky bluegrass covers, apply 6 fluid ounces of Sharda Glyphosate 41 SL. Do not add ammonium sulfate.

For best results, mow cool-season grass covers in the spring to even their height and apply the specified rate of Sharda Glyphosate 41 SL 3 to 4 days after mowing. Avoid treating cool season grass covers under poor growing conditions, such as drought stress (drip irrigation), disease or insect damage.

**LOW VOLUME APPLICATION (FLORIDA AND TEXAS)**

For burndown or control of the weeds listed, apply the specified rates of Sharda Glyphosate 41 SL in 3 to 30 gallons of water per acre. Where weed foliage is dense, use 10 to 30 gallons of water per acre.

**Annual Weeds**

**Goatweed:** - Apply 2 to 3 quarts per acre of Sharda Glyphosate 41 SL plus 17 pounds of ammonium sulfate per 100 gallons of water. Apply in 20 to 30 gallons of water per acre when plants are actively growing. Use 2 quarts per acre when plants are less than 8 inches tall and 3 quarts per acre when plants are greater than 8 inches, if goatweed is greater than 8 inches tall, the addition of Krovar II® or Karmex® may improve control. Use labeled rates for these residual products. Read and carefully observe the label claims, cautionary statements, rates and all other information on the Krovar II® and Karmex® labels.

**Perennial Weeds**

Apply when weeds are actively growing and at the growth stages listed in the "PERENNIAL WEEDS CONTROLLED" section of this label. If perennial weeds are mowed, allow weeds to regrow to the specified stage of growth.

S = Suppression      B = Burndown      PC = Partial control      C = Control

Weed	Sharda Glyphosate 41 SL Rate Per Acre			
	1 qt.	2 qts.	3 qts.	5 qts.
Bermudagrass	B	--	PC	C
Guineagrass, Texas and Florida Ridge	B	C	C	C
Guineagrass, Florida Flatwoods	--	B	C	C
Paragrass	B	C	C	C
Torpedograss	S	--	PS	C

**TREE CROPS**

**Citrus:** calamondin, chironja, citron, grapefruit, kumquat, lemon, lime, mandarin orange, orange, pummelo, tangelo, tangerine, tangors. Allow a minimum of 1 day between last application and harvest.

**Nuts:** almond, beechnut, Brazil nut, butternut, cashew, chestnuts, chinquapin, filbert, hazel nut, hickory nut, macadamia, pecan, pistachio, walnut. Allow a minimum of 3 days between last application and harvest.

**Pome Fruit:** apple, loquat, mayhaw, pear, quince. Allow a minimum of 1 day between last application and harvest.

**Stone Fruit:** apricots, cherries, nectarines, olives, peaches, plums/prunes. Allow a minimum of 17 days between last application and harvest. For cherries, any application equipment listed in this section may be used in all states. For citron and olives, apply as a directed spray only.

Any application equipment listed in this section may be used in apricots, nectarines, peaches and plums/prunes growing in Arizona, California, Colorado, Idaho, Kansas, Kentucky, New Jersey, North Dakota, Oklahoma, Oregon, Texas, Utah and Washington, except for peaches grown in the states specified in the following paragraph. In all other states use wiper equipment only. For PEACHES grown in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina and Tennessee only,



apply with a shielded boom sprayer or shielded wiper applicator, which prevents any contact of Sharda Glyphosate 41 SL with the foliage or bark of trees. Apply no later than 90 days after first bloom. Applications made after this time may result in severe damage. Remove suckers and low-hanging limbs at least 10 days prior to application. Avoid applications near trees with recent pruning wounds or other mechanical injury. Apply only near trees which have been planted in the orchard for 2 or more years. **EXTREME CARE MUST BE TAKEN TO ENSURE NO PART OF THE PEACH TREE IS CONTACTED.**

### Tropical Fruit

**Allow a Minimum of 14 days between the last application and harvest for the following:**

Acerola, Atemoya, Avocado, Breadfruit, Canistel, Carambola, Cherimoya, Cocoa beans, dates, figs, genip, jaboticaba, jackfruit, longan, lychee, mango, mayhaw, passionfruit, persimmons, pomegranate, sapodilla, sapote, soursop, sugar apple, tamarind, tea

**Allow a Minimum of 28 days between the last application and harvest for the following:**

Plantains and Coffee

**Allow a Minimum of 1 day between the last application and harvest for the following:**

Banana, Guava, and Papaya

In coffee and bananas, delay applications 3 months after transplanting to allow the new coffee or bananas to become established.

### VEGETABLES: Peppers

**Hooded Sprayers:** Sharda Glyphosate 41 SL may be used through hooded sprayers for weed control between the rows of peppers. Only hooded sprayers that completely enclose the spray pattern may be used. A hooded sprayer is a type of shielded applicator. The spray pattern is completely enclosed on the top and all 4 sides by a hood, thereby shielding the crop from the spray solution. This equipment must be set up and operated in a manner that avoids bouncing or raising the hoods off the ground in any way. If the hoods are raised, spray particles may escape and come into contact with the crop, causing damage or destruction of the crop. The spray hoods must be operated on the ground or skimming across the ground. Tractor speed must be adjusted to avoid bouncing of the spray hoods. Avoid operation on rough or sloping ground where the spray hoods might be raised off the ground.

Follow these requirements:

- The spray hoods must be operated on the ground or skimming across the ground.
- Leave at least an 8 inch untreated strip over the drill row. For example, if the crop row width is 38 inches, the maximum width of the spray hood must be 30 inches.
- Maximum tractor speed: 5 mph.
- Maximum wind speed: 10 mph.
- Use low-drift nozzles.

### VINE CROPS - Kiwi Fruit, Grapes, Hops and Passion Fruit

Sharda Glyphosate 41 SL is directed for weed control in established groves, vineyards, and orchards, or for site preparation prior to transplanting crops listed in this section. Applications may be made with boom equipment, CDA, shielded sprayers, hand-held and high-volume wands, lances, orchard guns or with wiper applicator equipment, except as directed in this section. See the "APPLICATION EQUIPMENT AND TECHNIQUES" section for specific information on use of equipment and directions. When applying Sharda Glyphosate 41 SL, refer to the "WEEDS CONTROLLED" section and to specific directions in that section for rates to be used. Any variety of table, wine or raisin grape may be treated with any equipment listed in this section. Applications must not be made when green shoots, canes, or foliage are in the spray zone. Allow a minimum of 14 days between last application and harvest. In the northeast and Great Lakes regions, applications must be made prior to the end of bloom stage of grapes to avoid injury.

leaf 67

## STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

**PESTICIDE STORAGE:** STORE ABOVE 10°F (-12°C) TO KEEP PRODUCT FROM CRYSTALLIZING. Crystals will settle to the bottom. If allowed to crystallize, place in a warm room 68°F (20°C) for several days to redissolve and shake or roll to mix well before using.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of Sharda Glyphosate 41 SL that cannot be used or chemically reprocessed must be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures. Emptied container retains vapor and product residue. Observe all label safeguards until container is destroyed.

**NONREFILLABLE CONTAINERS:** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least on complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling, if available.

**RETURNABLE / REFILLABLE CONTAINERS:** Refill this container with Sharda Glyphosate 41% SL only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect thoroughly for damage such as cracks, punctures, abrasions and damaged or worn out threads on closure devices. Do not refill or transport damaged or leaking containers. Check for leaks after refilling and before transportation. If the container is not being refilled, return it to the point of purchase. Prior to disposal or refilling, triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least on complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

## LIMIT OF WARRANTY AND LIABILITY

Sharda USA LLC warrants that Sharda Glyphosate 41 SL conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE.

This warranty is also subject to the conditions and limitations stated herein. Buyer and all users shall promptly notify Sharda USA LLC of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

To the extent consistent with applicable law, buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of Sharda USA LLC, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF SHARDA USA LLC OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF

SHARDA GLYPHOSATE 41 SL (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF SHARDA GLYPHOSATE 41 SL INVOLVED, OR, AT THE ELECTION OF SHARDA USA LLC OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW IN NO EVENT SHALL SHARDA USA LLC OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

© Weedar® and Weedone® are registered trademarks of Nufarm Americas Inc.

© Bullet®, Degree®, Harness®, Lasso®, Micro-Tech®, Partner®, Roundup Ready® and Lariat® are registered trademarks of Monsanto Company.

© Bladex®, Oust®, Krovar®, Escort®, Extrazine®, Telar®, Canopy®, Gemini®, Lexone®, Lorox®, Preview® and Karmex® are registered trademarks of E.I. duPont de Nemours and Company.

© Dual, Bicep®, Princep® and Caliber® are registered trademarks of Syngenta Crop Protection.

© Broadstrike®, Surflan®, Surpass®, and Topnotch® are registered trademarks of Dow Agrosciences.

© Banvel®, Frontier®, Guardsman®, Prowl®, Pursuit®, Pursuit Plus®, Scepter®, Solicam®, and Squadron® are registered trademarks of BASF, Inc.

© Folex® and Prep® are registered trademarks of Bayer Crop Sciences.

© Goal® is a registered trademark of Rohm and Haas Company.

© DEF®, Sencor® and Turbo® are registered trademarks of Bayer AG.

© Command® is a registered trademark of FMC Corporation.

© Linex® is a registered trademark of Griffin LLC.

© Permit® is a registered trademark of Gowan Company.