



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

June 23, 2025

Freddy Shelley
freddy.shelley@syntechresearch.com
SHARDA USA LLC

Subject: Non-PRIA (Pesticide Registration Improvement Act) Labeling Amendment - Additional Sublabel to Add Uses, Additional ABN
Product Name: SHARDA THIAMETHOXAM 25% WDG
Admin Number: 83529-125
EPA Receipt Date: 05/19/2025
Action Case Number: 00659033

Dear Freddy Shelley:

The amended labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable.

The alternate brand name: Thelmi has been added to the registration. Our records have been updated accordingly. This approval does not affect any terms or conditions that were previously imposed on this registration. You continue to be subject to existing terms or conditions on your registration and any deadlines connected with them.

If, after formal consultation with the appropriate Service(s), additional modifications are identified in the Service's Biological Opinion(s) for thiamethoxam, EPA will notify Sharda USA LLC in writing consistent with the terms in the Biological Opinion of any necessary required changes. Sharda USA LLC must submit an application for amendment incorporating any required changes, including amended labels, consistent with the timeline specified in EPA's notification. If Sharda USA LLC fails to comply with this term, EPA may cancel the registration under an expedited process under FIFRA 6(e).

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release this product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR § 152.3.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains statements or claims substantially differing from statements or claims made in connection with obtaining a FIFRA section 3 registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6.

If you have questions, please contact Loren Lapointe via email at lapointe.loren@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Ralph Narain". The script is elegant and cursive, with the first letters of each name being capitalized and prominent.

Ralph Narain, Acting Product Manager 01
IVB3, RD
Office of Pesticide Programs

{MASTER LABEL}
{Sub-label A: Crop}
{Sub-label B: T&O}
{Sub-label C: Ornamental Plants & Christmas Trees}

THIAMETHOXAM	GROUP	4A	INSECTICIDE
--------------	-------	----	-------------

Sharda Thiamethoxam 25% WDG
[ABN: Artist] [Worth 25 WDG] [Thelmi]


For Control of Listed Insect Pests Infesting Artichoke (Globe), Barley, Bushberry and Caneberry, Low Growing Berry, Small Fruit Vine, Brassica (Cole) Leafy Vegetables, Citrus Fruit, Cranberry, Cucurbit Vegetables, Fruiting Vegetables, Leafy Vegetables, Mint, Pome Fruit, Root Vegetables, Stone Fruit, Tobacco, Tropical Fruit, and Tuberous and Corm Vegetables.

For Foliar and Systemic Control of Listed Insect Pests in Turfgrass, Sod Farms, Interiorscape and Landscape Plants.

For foliar and systemic control of insect pests in ornamental plants, fruit and nut trees (non-bearing), Christmas trees, forest seedlings and listed vegetables grown for transplant.

ACTIVE INGREDIENT:	WT. BY %
Thiamethoxam: 3-(2-Chloro-5-thiazolylmethyl)tetrahydro-5-methyl-N-nitro-4H-1,3,5-oxadiazin-4-imine	25.0%
OTHER INGREDIENTS:	75.0%
TOTAL:	100.0%

KEEP OUT OF REACH OF CHILDREN
CAUTION



Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID	
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">Take off contaminated clothing.Rinse skin immediately with plenty of water for 15-20 minutes.Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none">Move person to fresh air.If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none">Hold eye open and rinse slowly and gently with water for 15-20 minutes.Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none">Call a poison control center or doctor immediately for treatment advice.Have person sip a glass of water if able to swallow.DO NOT induce vomiting unless told to do so by a poison control center or doctor.DO NOT give anything by mouth to an unconscious person.
NOTE TO PHYSICIAN	
There is no specific antidote if ingested. Induce emesis or lavage stomach. Treat symptomatically.	
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222 .	

[Optional referral statements when booklets and container labels are used:
See Panel for First Aid Instructions and booklet for complete Precautionary Statements and Directions For Use.
See label booklet for complete Precautionary Statements, Directions For Use, and Storage and Disposal.
See label booklet for additional Precautionary Statements, Directions For Use, and Storage and Disposal.
See label booklet for complete Directions For Use.]

Manufactured for:

Sharda USA LLC



7217 Lancaster Pike, Suite A
Hockessin, Delaware 19707

Net Contents: _____ Lbs. [Kg.]

{Sub-label A: Crop}

THIAMETHOXAM GROUP 4A INSECTICIDE

Sharda Thiamethoxam 25% WDG

[ABN: Artist] [Worth 25 WDG]

For Control of Listed Insect Pests Infesting Artichoke (Globe), Barley, Bushberry and Caneberry, Low Growing Berry, Small Fruit Vine, Brassica (Cole) Leafy Vegetables, Citrus Fruit, Cranberry, Cucurbit Vegetables, Fruiting Vegetables, Leafy Vegetables, Mint, Pome Fruit, Root Vegetables, Stone Fruit, Tobacco, Tropical Fruit, and Tuberous and Corm Vegetables.

ACTIVE INGREDIENT:**WT. BY %**

Thiamethoxam: 3-(2-Chloro-5-thiazolylmethyl)tetrahydro-5-methyl-N-nitro-4H-1,3,5-oxadiazin-4-imine 25.0%

OTHER INGREDIENTS: 75.0%**TOTAL:** 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID	
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. DO NOT induce vomiting unless told to do so by a poison control center or doctor. DO NOT give anything by mouth to an unconscious person.
NOTE TO PHYSICIAN	
There is no specific antidote if ingested. Induce emesis or lavage stomach. Treat symptomatically.	
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222 .	

[Optional referral statements when booklets and container labels are used:

See Panel for First Aid Instructions and booklet for complete Precautionary Statements and Directions For Use.

See label booklet for complete Precautionary Statements, Directions For Use, and Storage and Disposal.

See label booklet for additional Precautionary Statements, Directions For Use, and Storage and Disposal.

See label booklet for complete Directions For Use.]

EPA Reg. No. 83529-125

EPA Est. No. XXXXX-XX-XXX

Batch Code: _____

Manufactured for:

Sharda USA LLC 7217 Lancaster Pike, Suite A
Hockessin, Delaware 19707

Net Contents: _____ Lbs. [Kg.]

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through skin, swallowed, or inhaled. Causes moderate eye irritation. Wear protective eyewear (goggles, face shield, safety glasses). Avoid contact with eyes, skin, or clothing. Avoid breathing dust or spray mist. Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton® ≥ 14 mils
- Shoes plus socks
- Wear protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607(d-f)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to wildlife and highly toxic to aquatic invertebrates. This pesticide is highly toxic to bees exposed to direct treatment on blooming crops/plants or weeds. **DO NOT** apply this product or allow it to drift to blooming crops/plants or weeds while bees are foraging in/or adjacent to the treatment area.

For Terrestrial Uses: DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. **DO NOT** contaminate water when cleaning equipment or disposing of equipment wash waters.

Groundwater Advisory

Thiamethoxam has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into the groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may contaminate water through drift of spray in wind. This product has a high potential for runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff (See manual at the following internet address: <http://www.wsi.nrcs.usda.gov/products/W2Q/pest/core4.html>).

Runoff of this product will be reduced by avoiding applications when rainfall is forecast to occur within 48 hours.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT use, pour, spill, or store near heat or open flame.

PROTECTION OF POLLINATORS



APPLICATION RESTRICTIONS EXIST FOR THIS PRODUCT BECAUSE OF RISK TO BEES AND OTHER INSECT POLLINATORS. FOLLOW APPLICATION RESTRICTIONS FOUND IN THE DIRECTIONS FOR USE TO PROTECT POLLINATORS.



Look for the bee hazard icon in the Directions for Use for each application site for specific use restrictions and instructions to protect bees and other insect pollinators.

This product can kill bees and other insect pollinators.

Bees and other insect pollinators will forage on plants when they flower, shed pollen, or produce nectar.

Bees and other insect pollinators can be exposed to this pesticide from:

- Direct contact during foliar applications, or contact with residues on plant surfaces after foliar applications.
- Ingestion of residues in nectar and pollen when the pesticide is applied as a seed treatment, soil, tree injection, as well as foliar applications.

When Using This Product Take Steps To:

- Minimize exposure of this product to bees and other insect pollinators when they are foraging on pollinator attractive plants around the application site.
- Minimize drift of this product on to beehives or to off-site pollinator attractive habitat. Drift of this product onto beehives or off-site to pollinator attractive habitat can result in bee kills.

Information on protecting bees and other insect pollinators may be found at the Pesticide Environmental Stewardship website at: <https://pesticidestewardship.org/pollinator-protection/>

Pesticide incidents (for example, bee kills) must immediately be reported to the State/Tribal lead agency. For contact information for your State, go to: <http://www.aapco.org>. Pesticide incidents must also be reported to the National Pesticide Information Center at: www.npic.orst.edu or directly to EPA at: beekill@epa.gov

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

See individual crops for specific pollinator protection application restrictions. If none exist under the specific crop, for foliar applications, follow these application directions for crops that are contracted to have pollinator services for food/feed crops and commercially grown ornamentals that are attractive to pollinators.

1. FOR CROPS UNDER CONTRACTED POLLINATION SERVICES



DO NOT apply this product while bees are foraging. **DO NOT** apply this product until flowering is complete and all petals have fallen unless the following condition has been met:

- If an application must be made when managed bees are at the treatment site, the beekeeper providing the pollination services must be notified no less than 48-hours prior to the time of the planned application so that the bees can be removed, covered, or otherwise protected prior to spraying.

2. FOR FOOD/FEED CROPS AND COMMERCIALY GROWN ORNAMENTALS NOT UNDER CONTRACT FOR POLLINATION SERVICES BUT ARE ATTRACTIVE TO POLLINATORS



DO NOT apply this product while bees are foraging. **DO NOT** apply this product until flowering is complete and all petals have fallen unless one of the following conditions is met:

- The application is made to the target site after sunset.
- The application is made to the target site when temperatures are below 55°F.
- The application is made in accordance with a government-initiated public health response.
- The application is made in accordance with an active state-administered apiary registry program where beekeepers are notified no less than 48-hours prior to the time of the planned application so that the bees can be removed, covered, or otherwise protected prior to spraying.
- The application is made due to an imminent threat of significant crop loss, and a documented determination consistent with an IPM plan or predetermined economic threshold is met. Every effort must be made to notify beekeepers no less than 48-hours prior to the time of the planned application so that the bees can be removed, covered, or otherwise protected prior to spraying.

This product can only be used in accordance with the Directions for Use on this label. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton® ≥ 14 mils
- Shoes plus socks

PRODUCT INFORMATION

Sharda Thiamethoxam 25% WDG is a selective water dispersible granule insecticide used to control many sucking and chewing insects through contact and ingestion on the crops listed on this label.

- Make application of **Sharda Thiamethoxam 25% WDG** when insect pest populations begin to build, but prior to populations reaching economically damaging levels. Economic thresholds for pests controlled by **Sharda Thiamethoxam 25% WDG** may be available from your local agricultural authorities.
- This products use is compatible with integrated pest management programs. However, **Sharda Thiamethoxam 25% WDG** is highly toxic to bees exposed to direct treatment on blooming crops or weeds.
- **Sharda Thiamethoxam 25% WDG** is rapidly taken up into foliage after being applied. Spray coverage is essential for optimal performance. To ensure good coverage, make application of **Sharda Thiamethoxam 25% WDG** in sufficient water. See specific treatment information in the **CROP USE DIRECTIONS** section of this label. The use of higher water volumes will generally result in better coverage, especially under adverse conditions (e.g., hot, dry) or where a dense plant canopy exists. The use of a spray adjuvant may improve spray coverage but is not required.
- **Sharda Thiamethoxam 25% WDG** is rainfast once the spray solution has dried on treated plants.
- **Sharda Thiamethoxam 25% WDG** may aid in the suppression of some pests. Suppression can mean either inconsistent control (good to poor), or consistent control at a level below that generally considered acceptable for commercial control.
- **Sharda Thiamethoxam 25% WDG** has a wide margin of plant safety when used in accordance with this label.

Use Restrictions

- **DO NOT** use **Sharda Thiamethoxam 25% WDG** in nurseries, greenhouses, plant propagation houses, or on any plants grown for use as transplants.
- **DO NOT** make application of **Sharda Thiamethoxam 25% WDG** to crops grown from seed treated with thiamethoxam.
- **DO NOT** apply this product, by any application method, to linden, basswood, or other *Tilia* species in the State of Oregon.
- **New York State:**
 - This product is classified as restricted use.
 - Not for Sale, Sale Into, Distribution and/or Use in Nassau and Suffolk Counties of New York State.
 - **DO NOT** exceed a total of 0.188 lb. a.i. of thiamethoxam-containing products per acre per growing season.
 - For Pome Fruit, **DO NOT** exceed a total of 0.188 lb. a.i. of thiamethoxam-containing products per acre per growing season.

THIAMETHOXAM	GROUP	4A	INSECTICIDE
--------------	-------	----	-------------

RESISTANCE MANAGEMENT

For resistance management, **Sharda Thiamethoxam 25% WDG** contains Thiamethoxam and is classified in the neonicotinoids chemical class as a Group 4A insecticide, nicotinic acetylcholine receptor (nAChR) competitive modulators.

Any insect population may contain individuals naturally resistant to **Sharda Thiamethoxam 25% WDG** and other Group 4A insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies must be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of **Sharda Thiamethoxam 25% WDG** or other Group 4A insecticides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. **DO NOT** rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues for the targeted pests between the individual components of a mixture.
- In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual insecticides selected for use in mixtures must be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide/acaricides use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological, and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance, contact Sharda USA LLC or representative.

MANDATORY SPRAY DRIFT MANAGEMENT

Aerial Applications:

- **DO NOT** release spray at a height greater than 10 ft. above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a Medium or coarser droplet size (ASABE S572.1).
- For aerial applications: **DO NOT** apply when wind speeds exceed 15 mph at the application site. If wind speed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters. Applicators must use ½ swath displacement upwind at the downwind edge of field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

Ground Boom Applications:

- Users must only apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy unless making a turf, pasture, or rangeland application, in which case applicators may apply with a nozzle height no more than 4 feet above the ground.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- **Adjust Nozzles** - Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles must be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom must remain level with the crop and have minimal bounce.

RELEASE HEIGHT – Aircraft

Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

APPLICATION INSTRUCTIONS

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Ground Application

Select spray nozzles which will provide accurate and uniform spray deposition. Use spray nozzles which provide medium-sized droplets and reduce drift. To help ensure accuracy, calibrate sprayer before each use. For information on spray equipment and calibration, consult nozzle manufacturers and/or State Extension Service specialists.

To provide thorough and uniform coverage, make applications of **Sharda Thiamethoxam 25% WDG** using sufficient water volume. Use greater water volumes where a dense canopy exists and/or pest pressure is high. The use of a spray adjuvant may improve spray coverage but is not required. **DO NOT** make applications under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur.

Aerial Application

Make applications of **Sharda Thiamethoxam 25% WDG** in water, using the minimum spray volume indicated in the **CROP USE DIRECTIONS** section of this label. Increase spray volume where practical to improve coverage. Avoid applying under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur.

Application Through Irrigation Systems (Chemigation) - Potatoes and Cranberry Only Cranberry - Solid Set Sprinkler System Only (See CROP USE DIRECTIONS)

Sharda Thiamethoxam 25% WDG alone or in combination with other products which are registered for treatment through sprinkler irrigation may be applied through irrigation systems. Make application of this product only through center pivot, solid set, hand move, or moving wheel irrigation systems. **DO NOT** make application of **Sharda Thiamethoxam 25% WDG** through any other type of irrigation system. Lack of effectiveness or illegal pesticide residues can result from non-uniform distribution of treated water. If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers, or other experts. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Using Water from Public Water Systems: DO NOT APPLY SHARDA THIAMETHOXAM 25% WDG THROUGH ANY IRRIGATION SYSTEM PHYSICALLY CONNECTED TO A PUBLIC WATER SYSTEM. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. **Sharda Thiamethoxam 25% WDG** may be applied through irrigation systems, which may be supplied by a public water system only if the water from the public water system is discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and to top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. Before beginning chemigation, always make sure that the air gap exists and that there is no blockage of the overflow of the reservoir tank.

Any irrigation system using water supplied from a public water system must also meet the following requirements.

Operating Instructions for All Specified Types of Irrigation Systems

1. The system must be calibrated to uniformly apply the rates specified. If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers, or other experts.
2. The system must contain a functional check-valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
6. The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
7. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
8. **DO NOT** apply when wind speed favors drift beyond the area intended.

Calibration and Application Instructions

Sharda Thiamethoxam 25% WDG must be applied under the schedule specified in the specific **CROP USE DIRECTIONS**, not according to the irrigation schedule unless the events coincide.

Set the equipment to make application to the minimum amount of water per acre. Run the system at 85 - 90% of the manufacturer's maximum rated travel speed.

The following calibration and treatment techniques are provided for user reference, but **DO NOT** constitute a warranty of fitness for treatment through sprinkler irrigation equipment. Check with State and local regulatory agencies for potential use restrictions before applying any agricultural chemical through sprinkler irrigation equipment.

Center Pivot Irrigation Equipment

Notes: (1) Use only drive systems that provide uniform water distribution. (2) **DO NOT** use end guns when chemigating **Sharda Thiamethoxam 25% WDG** through center pivot systems because of non-uniform application. (3) Plug the first nozzle closest to the well-head to protect the water source.

1. Determine the size of the area to be treated.
2. Determine the time required to apply 0.1 - 0.25 inch of water over the area to be treated when the system and injection equipment are operated at normal pressures as recommended by the equipment manufacturer. Run the system at 80 - 95% of the manufacturer's rated maximum travel speed.
3. Using water, determine the injection pump output when operated at normal line pressure.
4. Determine the amount of **Sharda Thiamethoxam 25% WDG**, and any tank mix partners, required to treat the area covered by the irrigation system.
5. Add the required amount of **Sharda Thiamethoxam 25% WDG**, any tank mix partners, and sufficient water to meet the injection time requirements to the solution tank. (See **MIXING PROCEDURES** section of this label.)
6. Make sure the system is fully charged with water before starting injection of the **Sharda Thiamethoxam 25% WDG** solution. Time the injection to last at least as long as it takes to bring the system to full pressure.
7. Maintain constant agitation in the solution tank during the injection period.
8. Inject the specified amount of **Sharda Thiamethoxam 25% WDG** per acre continuously for one complete revolution of the system.
9. Stop the injection equipment after treatment is completed. Continue to operate the system until the **Sharda Thiamethoxam 25% WDG** solution has cleared all of the sprinkler heads.
10. Allow time for all lines to flush the pesticide through all nozzles before turning off irrigation water.

Solid Set, Hand Move, and Moving Wheel Irrigation Equipment

1. Determine the acreage covered by the sprinklers.
2. Fill injector solution tank with plain water and calibrate the flow rate of the system to deliver the contents of the tank over a 20 - 40 minute time interval.
3. Determine the amount of **Sharda Thiamethoxam 25% WDG** required to treat the area covered by the irrigation system.
4. Add the required amount of **Sharda Thiamethoxam 25% WDG**, and any other tank mix partners, into the same quantity of water used to calibrate the injection period. (See **MIXING PROCEDURES** section of this label.)
5. Operate the system at the same pressure and time interval established during the calibration.
6. Inject specified amount of **Sharda Thiamethoxam 25% WDG** per acre for either a 20 - 40 minute period at the end of a regular irrigation set, or as a 20-40 minute injection as a separate application not associated with a regular irrigation to maximize retention of the insecticide by the foliage.
7. Stop injection equipment after treatment is completed. Continue to operate the system until the **Sharda Thiamethoxam 25% WDG** solution has cleared the last sprinkler head. To ensure lines are flushed and free from remaining pesticides, a dye indicator may be injected into the lines to mark the end of the application period.

MIXING PROCEDURES

Prepare no more spray mixture than is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Vigorous agitation is necessary for proper dispersal of the product. Maintain maximum agitation throughout the spraying operation. **DO NOT** let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area. Keep product container tightly closed when not in use.

Sharda Thiamethoxam 25% WDG Alone

Add ½ of the required amount of water to the mix tank. With the agitator running, add the desired amount of **Sharda Thiamethoxam 25% WDG** to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after **Sharda Thiamethoxam 25% WDG** has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

Sharda Thiamethoxam 25% WDG + Tank Mixtures

Add ½ of the required amount of water to the mix tank. Start the agitator running before adding any tank mix partners. Add tank mix partners in this order: products packaged in water-soluble packaging, wettable powders, wettable granules (dry flowables) such as **Sharda Thiamethoxam 25% WDG**, liquid flowables, liquids, emulsifiable concentrates, and surfactants/adjuvants. Always allow each tank mix partner to become fully dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation until all of the mixture has been applied.

Note: When using **Sharda Thiamethoxam 25% WDG** in tank mixtures, add all products in water-soluble packaging to the tank before any other tank mix partner, including **Sharda Thiamethoxam 25% WDG**. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank mix partner to the tank.

If using **Sharda Thiamethoxam 25% WDG** in a tank mixture, observe all directions for use, crop/sites, use rates, dilution ratios, precautions, and limitations, which appear on the tank mix product label. **DO NOT** exceed any label dosage rate, and follow the most restrictive label precautions and limitations. **DO NOT** mix this product with any product which prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are labeled.

Compatibility

Sharda Thiamethoxam 25% WDG is compatible with most commonly used pesticides, crop oils, adjuvants, and nutritional sprays. However, since it is not possible to test all possible mixtures, the user must pre-test to assure the physical compatibility and lack of phytotoxic effect of any proposed mixtures with **Sharda Thiamethoxam 25% WDG**. To determine the physical compatibility of **Sharda Thiamethoxam 25% WDG** with other products, use a jar test, as described below.

Using a quart jar, add the proportionate amounts of the products to 1 qt. of water. Add wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

The crop safety of all potential tank mixes on all crops has not been tested. Confirm the safety to the target crop before applying any tank mixture not specified on this label.

ROTATIONAL RESTRICTIONS

Treated areas may be replanted immediately following harvest, or as soon as practical following the last application, with any crop listed on this label or to barley, canola, cotton, corn, cucurbit vegetables, legume vegetables, oilseed crops (rapeseed, Indian rapeseed, Indian mustard seed, field mustard seed, black mustard seed, flax seed, safflower seed, crambe seed and borage seed), sorghum, sunflower and wheat. Any cover crop planted for erosion control or soil improvement may be planted as soon as practical following the last application. However, the cover crop may not be grazed or harvested for food or feed. For all other crops, a 120-day plant-back interval must be observed.


CROP USE DIRECTIONS




Pollinator Precautions

- **Sharda Thiamethoxam 25% WDG** is highly toxic to bees exposed to direct treatment on blooming crops/plants or weeds.
 - For apples, **DO NOT** make application of **Sharda Thiamethoxam 25% WDG** after pre-bloom (early pink growth stage) or before post-bloom (petal fall growth stage).
 - For citrus, **DO NOT** make application during pre-bloom or during bloom when bees are actively foraging.
 - For pears, **DO NOT** make application of **Sharda Thiamethoxam 25% WDG** after pre-bloom (green cluster stage) or before post-bloom (petal fall growth stage).
 - For stone fruit, **DO NOT** make application of **Sharda Thiamethoxam 25% WDG** between the pre-bloom (swollen bud) and post-bloom (petal fall) growth stages.
- **DO NOT** make application of **Sharda Thiamethoxam 25% WDG** or allow it to drift to blooming crops/plants or weeds if bees are foraging in or adjacent to the treatment area. This is especially critical if there are adjacent orchards that are blooming (refer to the **MANDATORY SPRAY DRIFT MANAGEMENT** section for additional information).
- After an **Sharda Thiamethoxam 25% WDG** application, wait at least 5 days before placing beehives in the treated field.
- If bees are foraging in the ground cover and it contains any blooming plants or weeds, always remove flowers before making an application. This may be accomplished by mowing, disking, mulching, flailing, or applying a labeled herbicide.
- Consult with your local cooperative extension service or State agency responsible for regulating pesticide use for additional pollinator safety practices.

Artichoke (Globe)


Pest	Rate per Acre per Application (Oz.)
Artichoke Aphid (<i>Capitophorus elaeagni</i>) Leafhoppers Proba Bug	3
<p>Make application prior to the pests reaching damaging levels. Scout fields and treat again if populations rebuild to potentially damaging levels. Make application at the higher rate within the listed rate range for heavy infestations. Use sufficient water volume to ensure thorough coverage of foliage.</p> <p>Restrictions:</p> <ul style="list-style-type: none"> • DO NOT apply more than a total of 6 oz. (0.094 lb. a.i.) of Sharda Thiamethoxam 25% WDG per acre per year. • DO NOT apply more than 2 applications per acre per year. • Pre-Harvest Interval (PHI): 4 days • Minimum Interval Between Applications: 7 days • DO NOT use less than 10 GPA for ground applications or 5 GPA for aerial applications. 	
 Refer to Pollinator Precautions and RESISTANCE MANAGEMENT sections.	

Barley


Pest	Rate per Acre per Application (Oz.)
Aphids	4
<p>Make application prior to pests reaching damaging levels. Scout fields and treat again if populations rebuild to potentially damaging levels. Use sufficient water volume to ensure thorough coverage of foliage.</p> <p>Restrictions:</p> <ul style="list-style-type: none"> • DO NOT apply more than a total of 8 oz. (0.125 lb. a.i.) of Sharda Thiamethoxam 25% WDG per acre per year. • DO NOT apply more than 2 applications per acre per year. • Pre-Harvest Interval (PHI): 21 days • Minimum Interval Between Applications: 7 days • DO NOT use less than 10 GPA for ground applications or 5 GPA for aerial applications. 	
 Refer to Pollinator Precautions and RESISTANCE MANAGEMENT sections.	

Bushberry Subgroup and Caneberry Subgroup

Crop	Pest	Rate per Acre per Application (Oz.)
Bushberry Subgroup	Aphids	3 - 4


Aronia berry, Black currant, Buffalo currant, Chilean guava, Edible honeysuckle, Elderberry, European barberry, Gooseberry, Highbush blueberry, Highbush cranberry, Huckleberry, Jostaberry, Juneberry, Native currant, Red currant, Salal, Sea buckthorn, and Cultivars, varieties and/or hybrids of these.	Cranberry Weevil Leafhoppers	
	Japanese Beetle Weevil (Adults)	4
Caneberry Subgroup Blackberry, Loganberry, Raspberry (black, red, and wild), and Cultivars, varieties and/or hybrids of these.	Aphids Leafhoppers	2 - 3
	Japanese Beetle Stinkbugs Tarnished Plant Bug Weevil (Adults) Whiteflies	3
<p>Make application prior to pests reaching damaging levels. Scout fields and treat again if populations rebuild to potentially damaging levels. Make application at the higher rate within the listed rate range for heavy infestations. Use sufficient water volume to ensure thorough coverage of foliage.</p> <p>Restrictions:</p> <ul style="list-style-type: none"> • Bushberry Subgroup - DO NOT apply more than a total of 12 oz. (0.188 lb. a.i.) of Sharda Thiamethoxam 25% WDG per acre per year. • Bushberry Subgroup – DO NOT apply more than 3 applications at 4 oz. per acre or 4 applications at 3 oz. per acre per year. • Caneberry Subgroup - DO NOT apply more than a total of 6 oz. (0.094 lb. a.i.) of Sharda Thiamethoxam 25% WDG per acre per year. • Caneberry Subgroup - DO NOT apply more than 2 applications at 3 oz. per acre or 3 applications at 2 oz. per acre per year. • Pre-Harvest Interval (PHI): 3 days • Minimum Interval Between Applications: 7 days • DO NOT use less than 10 GPA for ground applications or 5 GPA for aerial applications. <p> Refer to Pollinator Precautions and RESISTANCE MANAGEMENT sections.</p>		

Low Growing Berry Subgroup (except Cranberry)


Crop	Pest	Rate per Acre per Application (Oz.)
Bearberry, Bilberry, Cloudberry, Lingonberry, Lowbush blueberry, Muntries, Partridgeberry, and Strawberry	Aphids Leafhoppers	1.5 - 3
	Whiteflies	3 - 4
Refer to Cranberry section for use directions.	Lygus Bug (Suppression) Weevil (Adults)	4
<p>Make application prior to pests reaching damaging levels. Scout fields and treat again if populations rebuild to potentially damaging levels. Make application at the higher rate within the listed rate range for heavy infestations. Use sufficient water volume to ensure thorough coverage of foliage.</p> <p>Restrictions:</p> <ul style="list-style-type: none"> • DO NOT apply more than a total of 12 oz. (0.188 lb. a.i.) of Sharda Thiamethoxam 25% WDG per acre per year. • DO NOT apply more than 3 applications at 4 oz. per acre, 4 applications at 3 oz. per acre and 8 applications at 1.5 oz. per acre per year. • Pre-Harvest Interval (PHI): 3 days • Minimum Interval Between Applications: 10 days • DO NOT use less than 50 GPA for ground applications. • DO NOT apply by air. <p> Refer to Pollinator Precautions and RESISTANCE MANAGEMENT sections.</p>		

Small Fruit Vine Climbing Subgroup (except Fuzzy Kiwi Fruit and Gooseberry)

Crop	Pest	Rate per Acre per Application (Oz.)
Amur river grape, Grape, Kiwi fruit (hardy), Maypop, and Schisandra berry	Japanese Beetle Leafhoppers Mealybugs	1.75 - 3.5

Refer to Bushberry Subgroup section for use directions for gooseberry.	Sharpshooters	
<p>Make application prior to pests reaching damaging levels. Scout fields and treat again if populations rebuild to potentially damaging levels. Make application at the higher rate within the listed rate range for heavy infestations. Use sufficient water volume to ensure thorough coverage of foliage.</p> <p>Restrictions:</p> <ul style="list-style-type: none"> • DO NOT apply more than a total of 7 oz. (0.109 lb. a.i.) of Sharda Thiamethoxam 25% WDG per acre per year. • DO NOT apply more than 2 applications at 3.5 oz. per acre or 4 applications at 1.75 oz. per acre per year. • Pre-Harvest Interval (PHI): 5 days • Minimum Interval Between Applications: 14 days • DO NOT use less than 10 GPA for ground applications or 5 GPA for aerial applications. <p> Refer to Pollinator Precautions and RESISTANCE MANAGEMENT sections.</p>		

Brassica (Cole) Leafy Vegetables

Crop	Pest	Rate per Acre per Application (Oz.)
Head and Stem Brassica Broccoli, Broccoli (Chinese), Brussels sprouts, Cabbage, Cabbage (Chinese mustard and Napa), Cauliflower, Cavalo broccolo, and Kohlrabi	Aphids Flea Beetles	1.5 - 3
	Thrips Whiteflies	3 - 5.5
Leafy Brassica Greens Broccoli (raab), Chinese Cabbage (bok choy), Collards, Kale, Mizuna, Mustard greens, Mustard spinach, and Rape greens	Aphids Flea Beetles	1.5 - 3
	Thrips Whiteflies	3 - 5.5
<p>Make application prior to pests reaching damaging levels. Scout fields and treat again if populations rebuild to potentially damaging levels. Make application at the higher rate within the listed rate range for heavy infestations. Use sufficient water volume to ensure thorough coverage of foliage. Allow spray to dry before harvest.</p> <p>Restrictions:</p> <ul style="list-style-type: none"> • DO NOT apply more than a total of 11 oz. (0.172 lb. a.i.) of Sharda Thiamethoxam 25% WDG per acre per year. • DO NOT apply more than 2 applications at 5.5 oz. per acre per year. • Pre-Harvest Interval (PHI): 0 days (Head and Stem Brassica) and 7 days (Leafy Brassica Greens) • Minimum Interval Between Applications: 7 days • DO NOT use less than 10 GPA for ground applications or 5 GPA for aerial applications. <p> Refer to Pollinator Precautions and RESISTANCE MANAGEMENT sections.</p>		

Citrus Fruit Crop Group

Crop	Pest	Rate per Acre per Application (Oz.)
Calamondin, Citrus citron, Citrus hybrids (includes chironja, tangelo, and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Orange (sour and sweet), Pummelo, and Satsuma mandarin	Aphids Leafhoppers	3 - 4
	Asian Citrus Psyllid Citrus Black Fly Citrus Leafminer Mealybugs Sharpshooters Soft Scales Whiteflies	4 - 5.5
	Ants (except fire, harvester, carpenter, and Pharaoh ants) Armored Scales Citrus Root Weevil (Adults) Crickets Fruit Fly Grasshoppers Katydid Plant Bugs Stink Bugs Thrips (Foliage Feeding)	4.5 - 5.5

Make application prior to pests reaching damaging levels. For scales, time the treatments to coincide with the crawler stage. Scout fields and apply again if populations rebuild to potentially damaging levels. Make application at the higher rate within the listed rate range for heavy infestations. Use sufficient water volume to ensure thorough coverage of foliage. Aerial application may result in slower activity and reduced control compared to ground applications.

Restrictions:

- **DO NOT** apply more than a total of 11 oz. (0.172 lb. a.i.) of **Sharda Thiamethoxam 25% WDG** per acre per year.
- **DO NOT** apply more than 2 applications at 5.5 oz. per acre per year.
- Pre-Harvest Interval (PHI): 0 days
- Minimum Interval between Applications: 7 days
- **DO NOT** use less than 10 GPA for ground applications or 5 GPA for aerial applications.



Refer to **Pollinator Precautions** and **RESISTANCE MANAGEMENT** sections.

Cranberry

Pest	Rate per Acre per Application (Oz.)
Aphids Cranberry Flea Beetle Cranberry Weevil Leafhoppers	2 - 4
Japanese Beetle	4

Make application prior to pests reaching damaging levels. Scout fields and apply again if populations rebuild to potentially damaging levels. Make application at the higher rate within the listed rate range for heavy infestations. Use sufficient water volume to ensure thorough coverage of foliage. **Sharda Thiamethoxam 25% WDG** may be applied through a solid set sprinkler irrigation system at 200 - 650 gals. total volume per acre, and if applied during a regular irrigation set, only at the end of the irrigation set. For best results, it is recommended that 200 - 300 gals. total volume per acre be used for irrigation treatment.

Restrictions:

- **DO NOT** apply more than a total of 12 oz. (0.188 lb. a.i.) of **Sharda Thiamethoxam 25% WDG** per acre per year.
- **DO NOT** apply more than 3 applications at 4 oz. per acre and 6 applications at 2 oz. per acre per year.
- Pre-Harvest Interval (PHI): 30 days
- Minimum Interval Between Applications: 7 days
- **DO NOT** use less than 10 GPA for ground applications.
- **DO NOT** apply by air.
- **DO NOT** apply to flow-through bogs.
- **DO NOT** apply within 25 feet of bodies of water.
- **DO NOT** irrigate for the first 48 hours following application.
- **DO NOT** release water immediately following application; hold water within the bog system for 5 days following application prior to release.



Refer to **Pollinator Precautions** and **RESISTANCE MANAGEMENT** sections.

Cucurbit Vegetables

Crop	Pest	Rate per Acre per Application (Oz.)
Chayote, Chinese waxgourd, Citron melon, Cucumber, Edible gourd, Gherkin, <i>Momordica</i> spp., Muskmelon, Pumpkin, Squash (summer and winter), and Watermelon	Aphids Flea Beetles	1.5 - 3
	Cucumber Beetles (Suppression) Leafminers (Suppression) Whiteflies	3 - 5.5

Make application prior to pests reaching damaging levels. Scout fields and treat again if populations rebuild to potentially damaging levels. Make application at the higher rate within the listed rate range for heavy infestations. Use sufficient water volume to ensure thorough coverage of foliage. Allow spray to dry prior to harvest.

Restrictions:

- **DO NOT** apply more than a total of 11 oz. (0.172 lb. a.i.) of **Sharda Thiamethoxam 25% WDG** per acre per year.
- **DO NOT** apply more than 2 applications at 5.5 oz. per acre per year.
- Pre-Harvest Interval (PHI): 0 days
- Minimum Interval Between Applications: 5 days
- **DO NOT** use less than 10 GPA for ground applications or 5 GPA for aerial applications.



Refer to **Pollinator Precautions** and **RESISTANCE MANAGEMENT** sections.

Fruiting Vegetables

Crop	Pest	Rate per Acre per Application (Oz.)
Eggplant, Ground cherry, Pepino, Peppers (bell, chili, cooking, pimento, and sweet), Tomatillo, and Tomato	Aphids Colorado Potato Beetle Flea beetles Leafhoppers	2 - 3
	Pepper Weevil Stink Bugs Whiteflies	3 - 5.5

Make application prior to pests reaching damaging levels. Scout fields and apply again if populations rebuild to potentially damaging levels. Make application at the higher rate within the listed rate range for heavy infestations. Use sufficient water volume to ensure thorough coverage of foliage. Allow spray to dry before harvest.

Restrictions:

- **DO NOT** apply more than a total of 11 oz. (0.172 lb. a.i.) of **Sharda Thiamethoxam 25% WDG** per acre per year.
- **DO NOT** apply more than 2 applications at 5.5 oz. per acre per year.
- Pre-Harvest Interval (PHI): 0 days
- Minimum Interval Between Applications: 5 days
- **DO NOT** use less than 10 GPA for ground applications or 5 GPA for aerial applications.



Refer to **Pollinator Precautions** and **RESISTANCE MANAGEMENT** sections.

Leafy Vegetables (except Brassica)

Crop	Pest	Rate per Acre per Application (Oz.)
Amaranth, Arugula, Cardoon, Celery, Celery (Chinese), Celtuce, Chervil, Chrysanthemum (edible-leaved and garland), Corn salad, Cress (garden and upland), Dandelion, Dock, Endive, Fennel, Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Radicchio, Rhubarb, Spinach (New Zealand and vine), and Swiss chard	Aphids Flea Beetles Leafhoppers	1.5 - 3
	Whiteflies	3 - 5.5

Make application prior to pests reaching damaging levels. Scout fields and apply again if populations rebuild to potentially damaging levels. Make application at the higher rate within the listed rate range for heavy infestations. Use sufficient water volume to ensure thorough coverage of foliage.

Restrictions:

- **DO NOT** apply more than a total of 11 oz. (0.172 lb. a.i.) of **Sharda Thiamethoxam 25% WDG** per acre per year.
- **DO NOT** apply more than 2 applications at 5.5 oz. per acre per year.
- Pre-Harvest Interval (PHI): 7 days
- Minimum Interval Between Applications: 7 days
- **DO NOT** use less than 10 GPA for ground applications or 5 GPA for aerial applications.



Refer to **Pollinator Precautions** and **RESISTANCE MANAGEMENT** sections.

Mint

Crop	Pest	Rate per Acre per Application (Oz.)
Peppermint and Spearmint	Aphids Flea hoppers Leafhoppers Mint Flea Beetles	1.5 - 3
	Grasshoppers	3 - 4

Make application prior to pests reaching damaging levels. Scout fields and apply again if populations rebuild to potentially damaging levels. Make application at the higher rate within the listed rate range for heavy infestations. Use sufficient water volume to ensure thorough coverage of foliage.

Restrictions:

- **DO NOT** apply more than a total of 12 oz. (0.188 lb. a.i.) of **Sharda Thiamethoxam 25% WDG** per acre per year.
- **DO NOT** apply more than 3 applications at 4 oz. per acre, 4 applications at 3 oz. per acre and 8 applications at 1.5 oz. per acre per year.
- Pre-harvest Interval (PHI): 7 days
- Minimum Interval Between Applications: 14 days
- **DO NOT** use less than 10 GPA for ground applications or 5 GPA for aerial applications.



Refer to **Pollinator Precautions** and **RESISTANCE MANAGEMENT** sections.

Pome Fruit

Crop	Pest	Rate per Acre per Application (Oz.)	Application Directions
Apples, Crabapples, Loquat, Mayhaw, and Quince	Pre-bloom: Apple Aphid Apple Grain Aphid Green Peach Aphid Leafminers Mullein Bug (<i>Campylomma</i> spp.) Rosy Apple Aphid	4.5	Make application prior to pests reaching damaging levels. Rosy Apple Aphid: Make application when aphid colonies are first observed at the green tip through pink growth stage before leaf curling occurs. Leafminers: Make application when eggs are being deposited.
	Post-bloom: Leafhoppers	2 - 2.75	Make application prior to pests reaching damaging levels. Scout fields and apply again if populations rebuild to potentially damaging levels. Make application at the higher rate within the listed rate range for heavy infestations.
	Post-bloom: Apple Aphid Apple Grain Aphid European Apple Sawfly Green Peach Aphid Leafminers Plum Curculio	4.5 - 5.5	Make application prior to pests reaching damaging levels. Make application at the higher rate within the listed rate range for heavy infestations. Leafminers: To control first generation populations, apply immediately following petal fall. For control of second and third generations, apply to coincide with egg deposition. Make application at the higher rate within the listed rate range for heavy infestations. Plum Curculio: Apply immediately following petal fall. Additional treatments of a different insecticide may be necessary if pest pressure continues.
Pear and Oriental Pear (<i>Pyrus pyrifolia</i>)	Pre-bloom: Pear Psylla	5.5	Make application prior to pests reaching damaging levels.
	Pre-bloom: Apple Aphid	4.5 - 5.5	Make application prior to pests reaching damaging levels. Make application at the higher rate within the listed rate range for heavy infestations.
	Post-bloom: Pear Psylla	5.5	Make application prior to pests reaching damaging levels.
	Post-bloom: Apple Aphid Comstock Mealybug Plum Curculio	4.5 - 5.5	Make application prior to pests reaching damaging levels. Make application at the higher rate within the listed rate range for heavy infestations. Comstock Mealybug: Apply immediately following petal fall to control first generation crawlers. Plum Curculio: Apply immediately following petal fall. Make application at the higher rate within the listed rate range for heavy infestations. Additional treatments of a different insecticide may be necessary if pest pressure continues.
	Leafhoppers	2 - 2.75	Make application prior to pests reach damaging levels. Scout fields and apply again if populations rebuild to potentially damaging levels. Make application at the higher rate within the listed rate range for heavy infestations.

Restrictions:

- **DO NOT** apply more than a total of 16.5 oz. (0.258 lb. a.i.) of **Sharda Thiamethoxam 25% WDG** per acre per year.
- New York State: **DO NOT** apply more than a total of 0.172 lb. a.i. of thiamethoxam-containing products per acre per growing season on Pome Fruit.
- Pre-Harvest Interval (PHI): 35 days - Use rates exceeding 2.75 oz. per acre may not be applied closer than 35 days before harvest. 14 days - Application rates equal to or less than 2.75 oz. per acre may be applied up to 14 days before harvest.
- Minimum Interval Between Applications: 10 days
- **DO NOT** apply by air.
- Use a minimum of 50 GPA applied with ground equipment to ensure thorough coverage of foliage.



Refer to **Pollinator Precautions** and **RESISTANCE MANAGEMENT** sections.

Root Vegetables Subgroup (Except Sugarbeets)

Crop	Pest	Rate per Acre per Application (Oz.)
Radish	Aphids	1.5 - 3
	Flea Beetles	
	Leafhoppers	3 - 4
	Whiteflies	
Carrot, Celeriac, Chicory, Edible burdock, Garden beet, Ginseng, Horseradish, Oriental radish, Parsnip, Rutabaga, Salsify (black and Spanish), Skirret, Turnip, Turnip Rooted Chervil, and Turnip Rooted Parsley	Aphids	1.5 - 3
	Flea Beetles	
	Leafhoppers	3 - 4
	Whiteflies	

Make application prior to pests reaching damaging levels. Scout fields and apply again if populations rebuild to potentially damaging levels. Make application at the higher rate within the listed rate range for heavy infestations. Use sufficient water volume to ensure thorough coverage of foliage.

Restrictions:

- **Radish** - **DO NOT** apply more than a total of 4 oz. (0.063 lb. a.i.) of **Sharda Thiamethoxam 25% WDG** per acre per year.
- **Other Root Vegetables** - **DO NOT** apply more than a total of 8 oz. (0.125 lb. a.i.) of **Sharda Thiamethoxam 25% WDG** per acre per year.
- Pre-Harvest Interval (PHI): 7 days
- Minimum Interval Between Applications: 7 days
- **DO NOT** use less than 10 GPA for ground applications or 5 GPA for aerial applications.



Refer to **Pollinator Precautions** and **RESISTANCE MANAGEMENT** sections.

Stone Fruit

Crop	Pest	Rate per Acre per Application (Oz.)
Apricot, Cherry (sweet and tart), Chickasaw plum, Damson plum, Japanese plum, Nectarine, Peach, Plum, Plumcot, and Prune (fresh)	Aphids	3 - 4
	Leafhoppers	2 - 2.75
	Cherry Fruit Fly	4.5 - 5.5
	Plum Curculio	
	Stink Bugs	
	Tarnished Plant Bug	
	Thrips	

Make application prior to pests reaching damaging levels. Scout fields and treat again if populations rebuild to potentially damaging levels. Make application at the higher rate within the listed rate range for heavy infestations. Use sufficient water volume to ensure thorough coverage of foliage.

Restrictions:

- **DO NOT** apply more than a total of 11 oz. (0.172 lb. a.i.) of **Sharda Thiamethoxam 25% WDG** per acre per year.
- **DO NOT** apply more than 2 applications at 5.5 oz. per acre per year.
- Pre-Harvest Interval (PHI): 14 days
- Minimum Interval Between Applications: 7 days
- **DO NOT** use less than 50 GPA for ground applications.
- **DO NOT** apply by air.



Refer to **Pollinator Precautions** and **RESISTANCE MANAGEMENT** sections.

Tobacco

Pest	Rate per Acre per Application (Oz.)
Aphids Flea Beetles Japanese Beetles	2 - 3

Make application prior to pests reaching damaging levels. Make application at the higher rate within the listed rate range for heavy infestations. Use sufficient water volume to ensure thorough coverage of foliage.

Restrictions:

- **DO NOT** apply more than a total of 3 oz. (0.047 lb. a.i.) of **Sharda Thiamethoxam 25% WDG** per acre per year.
- **DO NOT** apply more than 1 application per acre per year.
- Pre-Harvest Interval: 14 days
- **DO NOT** use less than 20 GPA for ground applications.



Refer to **Pollinator Precautions** and **RESISTANCE MANAGEMENT** sections.

Tropical Fruit

Crop	Pest	Rate per Acre per Application (Oz.)
Avocado, Black sapote, Canistel, Mamey sapote, Mango, Papaya, Sapodilla, and Star apple	Aphids Leafhoppers Mealybugs Sharpshooters Thrips (Foliage Feeding) Whiteflies	4

Make application prior to pests reaching damaging levels. Scout fields and apply again if populations rebuild to potentially damaging levels. Use sufficient water volume to ensure thorough coverage of foliage.

Restrictions:

- **DO NOT** apply more than a total of 12 oz. (0.188 lb. a.i.) of **Sharda Thiamethoxam 25% WDG** per acre per growing season.
- **DO NOT** apply more than 3 applications per acre per year.
- Pre-Harvest Interval (PHI): 0 days
- Minimum Interval Between Applications: 7 days
- **DO NOT** use less than 50 GPA for ground application.



Refer to **Pollinator Precautions** and **RESISTANCE MANAGEMENT** sections.

Tuberous and Corm Vegetables

Crop	Pest	Rate per Acre per Application (Oz.)
Arracacha, Arrowroot, Canna, Cassava (bitter and sweet), Chayote (root), Chinese artichoke, Chufa, Dasheen, Ginger, Jerusalem artichoke, Leren, Potato, Sweet potato, Tanier, Turmeric, Yams, and Yam bean	Aphids	3
	Colorado Potato Beetle Flea Beetle Potato Leafhoppers	1.5 - 3

Make application prior to pests reaching damaging levels. Scout fields and apply again if populations rebuild to potentially damaging levels. Make application at the higher rate within the listed rate range for heavy infestations. Use sufficient water volume to ensure thorough coverage of foliage.

Restrictions:

- **DO NOT** apply more than a total of 6 oz. (0.094 lb. a.i.) of **Sharda Thiamethoxam 25% WDG** per acre per growing season.
- **DO NOT** apply more than 2 applications at 3 oz. per acre and 4 applications at 1.5 oz. per acre per year.
- Pre-harvest Interval: 14 days
- Minimum Interval Between Applications: 7 days
- **DO NOT** use less than 10 GPA for ground applications or 5 GPA for aerial applications.
- Chemigation: Use from 0.10 - 0.25" of water. (For more details - See application through irrigation systems for potatoes in the **APPLICATION INSTRUCTIONS** section.)



Refer to **Pollinator Precautions** and **RESISTANCE MANAGEMENT** sections.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal. For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup, procedures and disposal of wastes.

PESTICIDE STORAGE: Store in a cool, dry place.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of Federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

CONTAINER HANDLING:

[Nonrefillable Container (50 pounds or less):] Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by State and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Sharda USA LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Sharda USA LLC and Seller harmless for any claims relating to such factors.

Sharda USA LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or Sharda USA LLC and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, SHARDA USA LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Sharda USA LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SHARDA USA LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SHARDA USA LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Sharda USA LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Sharda USA LLC.

[All trademarks are the property of their respective owners.]

{Sub-label B: T&O}

THIAMETHOXAM GROUP 4A INSECTICIDE

Sharda Thiamethoxam 25% WDG

[ABN: Worth 25 WDG]

For Foliar and Systemic Control of Listed Insect Pests in Turfgrass, Sod Farms, Interiorscape and Landscape Plants.

ACTIVE INGREDIENT:	WT. BY %
Thiamethoxam: 3-(2-Chloro-5-thiazolylmethyl)tetrahydro-5-methyl-N-nitro-4H-1,3,5-oxadiazin-4-imine	25.0%
OTHER INGREDIENTS:	75.0%
TOTAL:	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID	
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. DO NOT induce vomiting unless told to do so by a poison control center or doctor. DO NOT give anything by mouth to an unconscious person.
NOTE TO PHYSICIAN	
There is no specific antidote if ingested. Induce emesis or lavage stomach. Treat symptomatically.	
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222 .	

[Optional referral statements when booklets and container labels are used:

See Panel for First Aid Instructions and booklet for complete Precautionary Statements and Directions For Use.

See label booklet for complete Precautionary Statements, Directions For Use, and Storage and Disposal.

See label booklet for additional Precautionary Statements, Directions For Use, and Storage and Disposal.

See label booklet for complete Directions For Use.]

EPA Reg. No. 83529-125

EPA Est. No. XXXXX-XX-XXX

Batch Code: _____

Manufactured for:

Sharda USA LLC


 7217 Lancaster Pike, Suite A
Hockessin, Delaware 19707

Net Contents: _____ Lbs. [Kg.]

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through skin, swallowed, or inhaled. Causes moderate eye irritation. Wear protective eyewear (goggles, face shield, safety glasses). Avoid contact with eyes, skin, or clothing. Avoid breathing dust or spray mist. Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton® ≥ 14 mils
- Shoes plus socks
- Wear protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607(d-f)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to wildlife and highly toxic to aquatic invertebrates. This pesticide is highly toxic to bees exposed to direct treatment on blooming crops/plants or weeds. **DO NOT** apply this product or allow it to drift to blooming crops/plants or weeds while bees are foraging in/or adjacent to the treatment area.

For Terrestrial Uses: DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. **DO NOT** contaminate water when cleaning equipment or disposing of equipment wash waters.

Groundwater Advisory

Thiamethoxam has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into the groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may contaminate water through drift of spray in wind. This product has a high potential for runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff (See manual at the following internet address: <http://www.wsi.nrcs.usda.gov/products/W2Q/pest/core4.html>).

PHYSICAL OR CHEMICAL HAZARDS

DO NOT use, pour, spill, or store near heat or open flame.

PROTECTION OF POLLINATORS



APPLICATION RESTRICTIONS EXIST FOR THIS PRODUCT BECAUSE OF RISK TO BEES AND OTHER INSECT POLLINATORS. FOLLOW APPLICATION RESTRICTIONS FOUND IN THE DIRECTIONS FOR USE TO PROTECT POLLINATORS.



Look for the bee hazard icon in the Directions for Use for each application site for specific use restrictions and instructions to protect bees and other insect pollinators.

This product can kill bees and other insect pollinators.

Bees and other insect pollinators will forage on plants when they flower, shed pollen, or produce nectar.

Bees and other insect pollinators can be exposed to this pesticide from:

- Direct contact during foliar applications, or contact with residues on plant surfaces after foliar applications.
- Ingestion of residues in nectar and pollen when the pesticide is applied as a seed treatment, soil, tree injection, as well as foliar applications.

When Using This Product Take Steps To:

- Minimize exposure of this product to bees and other insect pollinators when they are foraging on pollinator attractive plants around the application site.
- Minimize drift of this product on to beehives or to off-site pollinator attractive habitat. Drift of this product onto beehives or off-site to pollinator attractive habitat can result in bee kills.

Information on protecting bees and other insect pollinators may be found at the Pesticide Environmental Stewardship website at: <https://pesticidestewardship.org/pollinator-protection/>

Pesticide incidents (for example, bee kills) must immediately be reported to the State/Tribal lead agency. For contact information for your State, go to: <http://www.aapco.org>. Pesticide incidents must also be reported to the National Pesticide Information Center at: www.npic.orst.edu or directly to EPA at: beekill@epa.gov

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.



For outdoor applications, follow these non-agricultural application directions to protect pollinators.

NON-AGRICULTURAL USES

DO NOT apply **Sharda Thiamethoxam 25% WDG** while bees are foraging. **DO NOT** apply **Sharda Thiamethoxam 25% WDG** to plants that are flowering. Only apply after all flower petals have fallen off.

DO NOT apply this product, by any application method, to linden, basswood, or *Tilia* species.

The nonagricultural uses of this product are for application to ornamental trees, shrubs and non-food plants only.

New York State:

- This product is classified as restricted use.
- Not for Sale, Sale Into, Distribution and/or Use in Nassau and Suffolk Counties of New York State.

This product can only be used in accordance with the Directions for Use on this label. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also

contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton® ≥ 14 mils
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

DO NOT enter treated areas without protective clothing until sprays have dried.

PRODUCT INFORMATION

Sharda Thiamethoxam 25% WDG is a selective water dispersible granule insecticide that is effective when applied at label rates to turfgrass, plant foliage, and soil.

Sharda Thiamethoxam 25% WDG is active against listed sucking and chewing insect pests by contact and ingestion. **Sharda Thiamethoxam 25% WDG** is relatively short-lived on the surface of plant foliage, readily absorbed into plant tissues, and rainfast once it has dried. The rapid translaminar absorption and distribution within leaves provides excellent residual control of listed foliar-feeding insects.

In the soil, the active ingredient in **Sharda Thiamethoxam 25% WDG** will control listed soil pests upon contact or ingestion and is also readily taken up by plant roots. The active ingredient moves upwards in the plant to the site of pest infestation.

Through feeding on the plant, listed pests are exposed to the active ingredient in **Sharda Thiamethoxam 25% WDG**. Feeding will stop within minutes to hours of exposure and be followed by death of the pest. The moderate persistence of **Sharda Thiamethoxam 25% WDG** in the soil and foliage also provides residual control of labeled pests. **Sharda Thiamethoxam 25% WDG** is not active as an ovicide or as an insect growth regulator. Because residues on leaf surfaces are quickly degraded, **Sharda Thiamethoxam 25% WDG** is compatible with beneficial arthropods.

THIAMETHOXAM	GROUP	4A	INSECTICIDE
--------------	-------	----	-------------

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, **Sharda Thiamethoxam 25% WDG** contains Thiamethoxam and is classified in the neonicotinoids chemical class as a Group 4A insecticide, nicotinic acetylcholine receptor (nAChR) competitive modulators.

Any insect population may contain individuals naturally resistant to **Sharda Thiamethoxam 25% WDG** and other Group 4A insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies must be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of **Sharda Thiamethoxam 25% WDG** or other Group 4A insecticides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. **DO NOT** rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues for the targeted pests between the individual components of a mixture.
- In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual insecticides selected for use in mixtures must be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still

provide pest management benefits.

- The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide/acaricides use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological, and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance, contact Sharda USA LLC or go to <https://shardausa.com>.

MANDATORY SPRAY DRIFT MANAGEMENT

Ground Boom Applications:

- Users must only apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy unless making a turf, pasture, or rangeland application, in which case applicators may apply with a nozzle height no more than 4 feet above the ground.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

BOOM HEIGHT – Ground Boom

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom must remain level with the crop and have minimal bounce.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

APPLICATION INSTRUCTIONS

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Ground Application

Select spray nozzles which will provide accurate and uniform spray deposition. Use spray nozzles which provide medium-sized droplets and reduce drift. To help ensure accuracy, calibrate sprayer before each use. For information on spray equipment and calibration, consult nozzle manufacturers and/or State Extension Service specialists.

To provide thorough and uniform coverage, make applications of **Sharda Thiamethoxam 25% WDG** using sufficient water volume. Use greater water volumes where a dense canopy exists and/or pest pressure is high. The use of a spray adjuvant may improve spray coverage but is not required. **DO NOT** make applications under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur.

Using Water from Public Water Systems: DO NOT APPLY SHARDA THIAMETHOXAM 25% WDG THROUGH ANY IRRIGATION SYSTEM PHYSICALLY CONNECTED TO A PUBLIC WATER SYSTEM. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. **Sharda Thiamethoxam 25% WDG** may be applied through irrigation systems, which may be supplied by a public water system only if the water from the public water system is discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and to top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. Before beginning chemigation, always make sure that the air gap exists and that there is no blockage of the overflow of the reservoir tank.

Any irrigation system using water supplied from a public water system must also meet the following requirements.

Operating Instructions for All Specified Types of Irrigation Systems

9. The system must be calibrated to uniformly apply the rates specified. If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers, or other experts.
10. The system must contain a functional check-valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
11. The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
12. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
13. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
14. The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
15. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
16. **DO NOT** apply when wind speed favors drift beyond the area intended.

Calibration and Application Instructions

Sharda Thiamethoxam 25% WDG must be applied under the schedule specified in the specific **CROP USE DIRECTIONS**, not according to the irrigation schedule unless the events coincide.

Set the equipment to make application to the minimum amount of water per acre. Run the system at 85 - 90% of the manufacturer's maximum rated travel speed.

The following calibration and treatment techniques are provided for user reference, but **DO NOT** constitute a warranty of fitness for treatment through sprinkler irrigation equipment. Check with State and local regulatory agencies for potential use restrictions before applying any agricultural chemical through sprinkler irrigation equipment.

Center Pivot Irrigation Equipment

Notes: (1) Use only drive systems that provide uniform water distribution. (2) **DO NOT** use end guns when chemigating **Sharda Thiamethoxam 25% WDG** through center pivot systems because of non-uniform application. (3) Plug the first nozzle closest to the well-head to protect the water source.

11. Determine the size of the area to be treated.
12. Determine the time required to apply 0.1 - 0.25 inch of water over the area to be treated when the system and injection equipment are operated at normal pressures as recommended by the equipment manufacturer. Run the system at 80 - 95% of the manufacturer's rated maximum travel speed.
13. Using water, determine the injection pump output when operated at normal line pressure.
14. Determine the amount of **Sharda Thiamethoxam 25% WDG**, and any tank mix partners, required to treat the area covered by the irrigation system.
15. Add the required amount of **Sharda Thiamethoxam 25% WDG**, any tank mix partners, and sufficient water to meet the injection time requirements to the solution tank. (See **MIXING PROCEDURES** section of this label.)

16. Make sure the system is fully charged with water before starting injection of the **Sharda Thiamethoxam 25% WDG** solution. Time the injection to last at least as long as it takes to bring the system to full pressure.
17. Maintain constant agitation in the solution tank during the injection period.
18. Inject the specified amount of **Sharda Thiamethoxam 25% WDG** per acre continuously for one complete revolution of the system.
19. Stop the injection equipment after treatment is completed. Continue to operate the system until the **Sharda Thiamethoxam 25% WDG** solution has cleared all of the sprinkler heads.
20. Allow time for all lines to flush the pesticide through all nozzles before turning off irrigation water.

Solid Set, Hand Move, and Moving Wheel Irrigation Equipment

8. Determine the acreage covered by the sprinklers.
9. Fill injector solution tank with plain water and calibrate the flow rate of the system to deliver the contents of the tank over a 20 - 40 minute time interval.
10. Determine the amount of **Sharda Thiamethoxam 25% WDG** required to treat the area covered by the irrigation system.
11. Add the required amount of **Sharda Thiamethoxam 25% WDG**, and any other tank mix partners, into the same quantity of water used to calibrate the injection period. (See **MIXING PROCEDURES** section of this label.)
12. Operate the system at the same pressure and time interval established during the calibration.
13. Inject specified amount of **Sharda Thiamethoxam 25% WDG** per acre for either a 20 - 40 minute period at the end of a regular irrigation set, or as a 20-40 minute injection as a separate application not associated with a regular irrigation to maximize retention of the insecticide by the foliage.
14. Stop injection equipment after treatment is completed. Continue to operate the system until the **Sharda Thiamethoxam 25% WDG** solution has cleared the last sprinkler head. To ensure lines are flushed and free from remaining pesticides, a dye indicator may be injected into the lines to mark the end of the application period.

MIXING PROCEDURES

Prepare no more spray mixture than is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Vigorous agitation is necessary for proper dispersal of the product. Maintain maximum agitation throughout the spraying operation. **DO NOT** let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area. Keep product container tightly closed when not in use.

Sharda Thiamethoxam 25% WDG Alone

Add ½ of the required amount of water to the mix tank. With the agitator running, add the specified amount of **Sharda Thiamethoxam 25% WDG** to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after **Sharda Thiamethoxam 25% WDG** has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

Sharda Thiamethoxam 25% WDG + Tank Mixtures

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Sharda Thiamethoxam 25% WDG is compatible with many other commonly used insecticides, fungicides, and liquid fertilizers. Check compatibility before tank mixing.

Add ½ of the required amount of water to the mix tank. Start the agitator running before adding any tank mix partners. Add tank mix partners in this order: products packaged in water-soluble packaging, wettable powders, wettable granules (dry flowables) such as **Sharda Thiamethoxam 25% WDG**, liquid flowables, liquids, emulsifiable concentrates, and surfactants/adjuvants. Always allow each tank mix partner to become fully dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation until all of the mixture has been applied.

Important: When using **Sharda Thiamethoxam 25% WDG** in tank mixtures, add all products in water-soluble packaging to the tank before any other tank mix partner, including **Sharda Thiamethoxam 25% WDG**. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank mix partner to the tank.

Important: If using **Sharda Thiamethoxam 25% WDG** in a tank mixture, observe all directions for use, crop/sites, use rates, dilution ratios, precautions, and limitations, which appear on the tank mix product label. **DO NOT** exceed any label dosage rate, and follow the most restrictive label precautions and limitations. **DO NOT** mix this product with any product which prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are labeled.

Compatibility

Sharda Thiamethoxam 25% WDG is compatible with most commonly used pesticides, crop oils, adjuvants, and nutritional sprays. However, since it is not possible to test all possible mixtures, the user must pre-test to assure the physical compatibility and lack of phytotoxic effect of any proposed mixtures with **Sharda Thiamethoxam 25% WDG**. To determine the physical compatibility of **Sharda Thiamethoxam 25% WDG** with other products, use a jar test, as described below.

Using a quart jar, add the proportionate amounts of the products to 1 qt. of water. Add wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

The crop safety of all potential tank mixes on all crops has not been tested. Confirm the safety to the target crop before applying any tank mixture not specified on this label.

FIRE ANTS - INDIVIDUAL MOUND TREATMENTS (TURFGRASS AND SOIL OF ORNAMENTAL LANDSCAPE PLANTS)

Application Rate

For control of fire ants (*Solenopsis* species), prepare a drench solution at a concentration of 1 - 3 ounces of **Sharda Thiamethoxam 25% WDG** per 10 gallons of water. Thoroughly mix solution and apply directly to mounds. For optimum control of small ant mounds (< 6 inches in diameter at the surface), apply 1 gallon of the drench solution per mound.

For optimum control of larger ant mounds, apply 2 - 3 gallons of the drench solution per mound. Direct the drench application at the center of the mound and include a 6 inch diameter circle around the center of the mound. **DO NOT** apply less than 0.5 gallon or more than 3 gallons of drench solution per mound.

Application Timing

For control of fire ants, make a single drench application when mounds appear to be active.

Restrictions:

- **DO NOT** apply **Sharda Thiamethoxam 25% WDG** outdoors to plants that are flowering, only apply after all flower petals have fallen off.
- **DO NOT** exceed 17 oz. (0.266 lb. a.i.) of **Sharda Thiamethoxam 25% WDG** per acre per calendar year.

APPLICATION TO TURFGRASS

Sites of Application

Sharda Thiamethoxam 25% WDG can be applied to turfgrass on: golf courses, residential lawns, commercial grounds (i.e., office and shopping complexes, airports), parks, playgrounds, athletic fields, and sod farms.

Restrictions:

- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply more than 17 oz. of **Sharda Thiamethoxam 25% WDG** (0.266 pound thiamethoxam) per acre per calendar year.
- **DO NOT** use for seed production.
- **DO NOT** allow this product to contact plants in bloom while bees are foraging in the treatment area.
- **DO NOT** use in the commercial nursery production of ornamental plants.
- **DO NOT** allow children and pets to enter treated area until sprays have dried.

Broadcast Applications to Turfgrass

Apply **Sharda Thiamethoxam 25% WDG** as a broadcast application to turfgrass for control of targeted pests. Use sufficient water volume (1.5 - 5 gallons per 1,000 square feet) to uniformly distribute **Sharda Thiamethoxam 25% WDG** over the area being treated and to adequately move the active ingredient into the canopy and thatch layers.

To help prevent grubs, irrigate within 7 days of application to move **Sharda Thiamethoxam 25% WDG** into the root zone where grubs feed. To control listed grubs, irrigate within 1 day of application to move **Sharda Thiamethoxam 25% WDG** into the root zone where grubs feed. Maintain adequate soil moisture before and after application for optimum control and healthy turfgrass growth. Excessively wet or dry conditions may impact the performance of **Sharda Thiamethoxam 25% WDG** against white grubs and mole crickets. **DO NOT** mow turf until the treated area has been irrigated or rainfall has occurred to allow for maximum and uniform uptake into turfgrass.

Applications to Turfgrass

Use sites include turfgrass on golf courses, residential lawns, commercial grounds (i.e., office and shopping complexes, airports), parks, playgrounds, athletic fields, and sod farms. Use the higher rate in the specified rate range for heavy pest populations.

Use Pattern	Pest	Application Timing	Dosage of Sharda Thiamethoxam 25% WDG
Broadcast	Larvae (White Grubs) of: <i>Aphodius</i> species	Apply up to 45 days before the historical peak of adult flight to 2 nd instar grub of	3 - 4 oz. per 10,000 sq. ft.

	Asiatic Garden Beetles Billbugs (<i>Sphenophorus</i> species) Black Turfgrass Ataenius European Chafer Green June Beetles Japanese Beetles May or June Beetles (<i>Phyllophaga</i> species) Northern Masked Chafer Oriental Beetles Southern Masked Chafer	the species being targeted. For optimum control, treat from peak flight to peak egg hatch. To control grubs, make the application from egg hatch to second instar (grubs less than half their full size).	or 12.7 - 17 oz. per acre
	Craneflies (<i>Tipula</i> species)	For optimum control, apply at oviposition (egg lay)	
	Mole Crickets (Suppression) (<i>Scapteriscus</i> species)	To suppress damage, treat from first egg hatch to peak egg hatch. See note below.	
	Chinch Bugs (<i>Blissus</i> species)	Apply when young nymphs are first observed.	
	Flea Beetles Greenbugs Leafhoppers Sod Webworms Spittlebugs	For optimum control, make application when populations are first observed.	
	Ants (excluding Carpenter, Fire, Harvester, and Pharaoh Ants)	To control ants, treat when ant mounds are first observed. To achieve optimum control on green and tee surfaces, treat the affected area plus a surrounding 30 foot buffer. For fairways and roughs, treat at least 1.5 times the infested area. Water in to adequately move active ingredient to target area. For additional knockdown activity, utilize with Scimitar GC Insecticide in an insect control spray program. Consult the Scimitar GC Insecticide label for use.	
Individual Fire Ant Mounds	Fire Ants (<i>Solenopsis</i> species) Refer to FIRE ANTS - INDIVIDUAL MOUND TREATMENTS (TURFGRASS AND SOIL OF ORNAMENTAL LANDSCAPE PLANTS) section of label.		

For specific information about developmental stages of the target pest and optimal timing of applications, consult with your State Cooperative Extension Service.

For turf with heavy thatch (more than 0.75 inch), use the higher rates within the specified range. **Sharda Thiamethoxam 25% WDG** is not phytotoxic to any major turfgrass species.

Sharda Thiamethoxam 25% WDG provides suppression of mole crickets on turfgrass. Suppression can mean either erratic control, ranging from good to poor, or a consistent level of control below that generally acceptable for commercial purposes.

APPLICATION TO INTERIORESCAPE AND LANDSCAPE ORNAMENTAL PLANTS

Sites of Application

Sharda Thiamethoxam 25% WDG can be applied to ornamentals grown in residential and commercial landscapes, parks, golf courses and interiorscapes.

Restrictions:

- **DO NOT** apply **Sharda Thiamethoxam 25% WDG** outdoors to plants that are flowering. Only apply after all flower petals have fallen off.
- **DO NOT** exceed 17 oz. of **Sharda Thiamethoxam 25% WDG** (0.266 pounds Thiamethoxam) per acre per calendar year.

- **DO NOT** use on plants being grown for sale or other commercial use in greenhouses, or for commercial seed production.
- **DO NOT** apply this product, by any application method, to linden, basswood, or *Tilia* species.

APPLICATION METHODS

Sharda Thiamethoxam 25% WDG is an effective insecticide when applied to plant foliage, soil or soil-less media, and fire ant mounds.

Foliar Application

Make foliar applications in an adequate water volume to achieve thorough and uniform coverage without excessive runoff (to drip). On hard-to-wet plants, spray adjuvants that improve wetting of foliage are recommended. **DO NOT** use spray adjuvants that may bind **Sharda Thiamethoxam 25% WDG** to the leaf surface, which would limit absorption into the foliage.

Sharda Thiamethoxam 25% WDG can be applied in a range of spray volumes (concentrate to dilute spray volumes) provided thorough and uniform coverage is obtained. For concentrate sprays (i.e., ULV, foggers, air assist equipment), apply the same amount of **Sharda Thiamethoxam 25% WDG** per unit area as would be used in a dilute spray volume over the same area.

Begin applications when labeled pests first appear or when economic thresholds have been reached. Repeat applications as necessary to maintain control, but no sooner than every 7 days. Use the higher rate in the specified rate range for heavy pest populations.

Foliar Applications to Interiorscape and Landscape Ornamental Plants

Plant Type	Pest	Dosage of Sharda Thiamethoxam 25% WDG	Application Timing
Trees* Shrubs* Evergreens Flowers* Foliage Plants Groundcovers* Interior Plantscapes	Ants (excluding Carpenter, Fire, Harvester, and Pharaoh Ants) Aphids Lace Bugs Leafhoppers Leaf Beetles Tent Caterpillars	2 – 4 oz. per 100 gals.	Apply when listed pests are first observed. Reapply as needed but no sooner than every 7 days. For gall makers, apply during egg laying of the generation being targeted. For leafhoppers, plant bug, and aphids make the first application when populations are first observed. Repeat applications as necessary to maintain control, but no sooner than 7 days after the last application. For concentrated spray volumes, apply the same amount of product per unit area as would be applied with dilute spray volumes.
	Adelgids (including Hemlock Woolly Adelgid) Flea Beetles Japanese Beetles (Adults) Leaf-Gall-Forming Insects Leafminers Mealybugs Midges Plant Bug Psyllids (including Asian Citrus Psyllid) Sawflies Soft Scales Spittlebugs Thrips – Foliar Feeding (Suppression) Whiteflies	4 – 8.5 oz. per 100 gals.	Apply to landscape ornamental plants to reduce populations of aphids, whiteflies, and mealybugs, which produce honeydew which serves as a food source for some ant species. Use a 0.0038% - 0.01% liquid dilution (2 – 8 oz. of Sharda Thiamethoxam 25% WDG in 100 gals. water) as a foliar or banded application. The 0.01% dilution may be applied up to 2 gals./1,000 sq. ft. Re-treat as necessary to maintain control. DO NOT exceed 17 oz. (0.266 lb. a.i.) of Sharda Thiamethoxam 25% WDG per acre per calendar year.
	Black Vine Weevil (Adults) (Suppression)	4 oz. per 50 gals. or 8 oz. per 100 gals.	Apply to foliage of plants when adult feeding damage is first observed. Make only 1 application per generation. For concentrated spray volumes, apply the same amount of product per unit area as would be applied with dilute spray volumes. For best control, use in a rotational spray program with lambda-cyhalothrin. Consult lambda-cyhalothrin label for use.



- *DO NOT** apply Sharda Thiamethoxam 25% WDG outdoors to plants that are flowering. Only apply after all flower petals have fallen off.
- *DO NOT** exceed 17 oz. of Sharda Thiamethoxam 25% WDG (0.266 pounds Thiamethoxam) per acre per calendar year.

For specific information about developmental stages of the target pest, associated damage, and action thresholds to properly time applications consult with your State Cooperative Extension Service.

Application to Soil of Ornamental Plants

For plants grown in residential and commercial landscapes, parks, golf courses and interiorscapes.

Sharda Thiamethoxam 25% WDG is effective as a soil and systemic insecticide when applied to the soil using broadcast, soil injection, or soil drench applications.

As a broadcast application, use sufficient water volume (1.5 – 5 gals. per 1,000 square feet) to uniformly distribute **Sharda Thiamethoxam 25% WDG** at the labeled rate over the area being treated.

For soil injection around trees and shrubs, uniformly select sites and evenly distribute **Sharda Thiamethoxam 25% WDG** solution among injection sites.

For soil drench applications around the base of trees and shrubs, apply the proper rate uniformly in a minimum of 16 fluid ounces of solution per inch diameter of breast height (D.B.H.) or per foot of shrub height.

Sharda Thiamethoxam 25% WDG must move to the feeding site of the target pest, whether this is in the root zone or the above ground portion of the host plant. Therefore, control of pests from the systemic activity of **Sharda Thiamethoxam 25% WDG** may be delayed for 1 or more weeks depending on plant size, rate of uptake from the soil or rate of translocation within the plant. Systemic activity in large plants may require several weeks before control of stem and foliar pests is achieved. If economic thresholds for target pests are low, or plants are historically damaged, apply **Sharda Thiamethoxam 25% WDG** preventatively (before pests are evident) to the growing media.

Irrigate within 3 days of treatment (optimally within 24 hours) to move **Sharda Thiamethoxam 25% WDG** into the root zone of plants or to the location of soil-dwelling pests. **DO NOT** over irrigate during the first 7 days after treatment or reduced efficacy may occur. Use the higher rate in the specified rate range for heavy pest populations.

Soil Application to Soil-Inhabiting Pests of Interiorscape and Landscape Ornamental Plants

Plant Type	Pest	Dosage of Sharda Thiamethoxam 25% WDG	Application Timing
All Plant Types	Larvae (Grubs) of: <i>Aphodius</i> species Asiatic Garden Beetles Black Turfgrass Ataenius European Chafer Green June Beetles Japanese Beetles May or June Beetles (<i>Phyllophaga</i> species) Northern Masked Chafer Oriental Beetles Southern Masked Chafer	12.7 – 17 oz. per acre or 3 – 4 oz. per 10,000 sq. ft.	Application Timing for Grubs: Apply up to 45 days before the historical peak of adult flight to 2 nd instar grub of the species being targeted. For optimum control, treat from peak flight to peak egg hatch. To control grubs, make the application from egg hatch to second instar (grubs less than half their full size).
	Fire Ants (<i>Solenopsis</i> species) Refer to FIRE ANTS – INDIVIDUAL MOUND TREATMENTS (TURFGRASS AND SOIL OF ORNAMENTAL LANDSCAPE PLANTS) section of label.		

Application Instructions

- Apply to the soil using broadcast, soil injection, or soil drench applications. When using broadcast application, use the following water volume, 1.5 - 5 gals. per 1,000 square feet.

- **Important:** If leaching, wait a minimum of 7 days after application to allow for maximum uptake of **Sharda Thiamethoxam 25% WDG** by the plant's root system.
- Apply to the soil preventively in situations where tolerance of listed insects or damage is low. Allow a minimum of 1 week for smaller plants and at least 2 weeks for large plants to translocate **Sharda Thiamethoxam 25% WDG** to feeding sites of the target pest.
- Irrigate within 3 days of treatment (optimally within 24 hours) to move **Sharda Thiamethoxam 25% WDG** into the root zone of plants (unless the product is leaching).
- **DO NOT** exceed 17 oz. of **Sharda Thiamethoxam 25% WDG** (0.266 lb. thiamethoxam) per acre per calendar year when making applications to the soil for ornamentals.

For specific information about developmental stages of the target pest, associated damage, and action thresholds to properly time applications, consult with your State Cooperative Extension Service. Refer to additional comments under **PRODUCT INFORMATION** section relating to ornamental applications to soil and related irrigation events.

Soil Application to Above-Ground Pests of Interiorscape and Landscape Ornamental Plants

Pest	Plant Types	Dosage of Sharda Thiamethoxam 25% WDG	Application Instructions
Adelgids Aphids Lace Bugs Flea Beetles Japanese Beetle (Adults) Leaf Beetles Leafminers Mealybugs Root Aphids Root Weevil (Larvae) (including <i>Diaprepes abbreviatus</i>) Sawflies Soft Scales Spittlebugs Tent Caterpillars Thrips - Foliar Feeding (Suppression) Whiteflies	Trees	Single Tree 0.07 - 0.14 oz. (2 - 4 gm) per 1 inch of tree trunk diameter (D.B.H.)	Soil Injection Method for Trees Basal System: Apply the proper rate among 4 evenly spaced injection sites located within a foot of each trunk. Grid System: Apply the proper rate among injection sites on a grid pattern with 1.5 - 2.5 ft. centers extending to the drip line of the tree. Soil Drench Apply the proper rate uniformly around the base of the tree within 1 - 2 ft. of the trunk. Drench targeted area with the Sharda Thiamethoxam 25% WDG solution without runoff. Use a minimum of 16 fluid ounces solution per inch D.B.H. Irrigate after application for optimal performance.
	Shrubs	Single Shrub 0.07 - 0.14 oz. (2 - 4 gm) per foot of shrub height Hedge Row 0.07 - 0.14 oz. (2 - 4 gm) per foot of shrub height (<6 ft. tall) per plant	Soil Injection Method for Shrubs Within the canopy, apply the proper rate among a minimum of 4 evenly spaced injection sites per shrub. For multiple plants (hedge row), determine the number of plants per hedge row, multiply by the average height and rate to be applied per foot of height. Soil Drench Apply the proper rate uniformly around the shrub within 1 - 2 ft. of the base. Drench targeted area with the Sharda Thiamethoxam 25% WDG solution without runoff. Use a minimum of 16 fluid ounces solution per foot shrub height. For multiple plants (hedge row), determine the number of plants per hedge row, multiply by the average height and rate to be applied per foot of height. Irrigate after application for optimal performance.
	Annuals and Small Woody Landscape Plants	3 - 4 oz. per 10,000 sq. ft.	Apply as a broadcast treatment to the soil directed to the plant root zone. Irrigate after application for optimal performance.

Application Timing for Above-Ground Pests

To Help Prevent Pests: Apply 1 - 4 weeks before pest has historically been observed.

To Control Pests: Allow 1 - 4 weeks for reduction in pest density to occur.

- **Important:** If leaching, wait a minimum of 7 days after application to allow for maximum uptake of **Sharda Thiamethoxam 25% WDG** by the plant's root system.
- Apply to the soil preventively in situations where tolerance of insects or damage is low. Allow a minimum of 1 week for smaller plants and at least 2 weeks for large plants to translocate **Sharda Thiamethoxam 25% WDG** to feeding sites of the target pest.
- **DO NOT** exceed 17 oz. per acre per calendar year when making applications to the soil for ornamentals.

For specific information about developmental stages of the target pest, associated damage, and action thresholds to properly time applications, consult with your State Cooperative Extension Service. Refer to additional comments under Use Information section relating to ornamental applications to soil and related irrigation events.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal. For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup, procedures and disposal of wastes.

PESTICIDE STORAGE: Store in a cool, dry place.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of Federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

CONTAINER HANDLING:

[Nonrefillable Container (50 pounds or less):] Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by State and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Sharda USA LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Sharda USA LLC and Seller harmless for any claims relating to such factors.

Sharda USA LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or Sharda USA LLC and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, SHARDA USA LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Sharda USA LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SHARDA USA LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SHARDA USA LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Sharda USA LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Sharda USA LLC.

[All trademarks are the property of their respective owners.]

{Sub-label C: Ornamental Plants & Christmas Trees}

THIAMETHOXAM GROUP 4A INSECTICIDE

Sharda Thiamethoxam 25% WDG

[ABN: Thelmi]

For foliar and systemic control of insect pests in ornamental plants, fruit and nut trees (non-bearing), Christmas trees, forest seedlings and listed vegetables grown for transplant.

ACTIVE INGREDIENT:

Thiamethoxam: 3-(2-Chloro-5-thiazolylmethyl)tetrahydro-5-methyl-N-nitro-4H-1,3,5-oxadiazin-4-imine 25.0%

OTHER INGREDIENTS: 75.0%**TOTAL:** 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID	
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">Take off contaminated clothing.Rinse skin immediately with plenty of water for 15-20 minutes.Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none">Move person to fresh air.If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none">Hold eye open and rinse slowly and gently with water for 15-20 minutes.Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none">Call a poison control center or doctor immediately for treatment advice.Have person sip a glass of water if able to swallow.DO NOT induce vomiting unless told to do so by a poison control center or doctor.DO NOT give anything by mouth to an unconscious person.
NOTE TO PHYSICIAN	
There is no specific antidote if ingested. Induce emesis or lavage stomach. Treat symptomatically.	
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222 .	

[Optional referral statements when booklets and container labels are used:

See Panel for First Aid Instructions and booklet for complete Precautionary Statements and Directions For Use.

See label booklet for complete Precautionary Statements, Directions For Use, and Storage and Disposal.

See label booklet for additional Precautionary Statements, Directions For Use, and Storage and Disposal.

See label booklet for complete Directions For Use.]

EPA Reg. No. 83529-125

EPA Est. No. XXXXX-XX-XXX

Batch Code: _____

Manufactured for:

Sharda USA LLC 7217 Lancaster Pike, Suite A
Hockessin, Delaware 19707

Net Contents: _____ Lbs. [Kg.]

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through skin, swallowed, or inhaled. Causes moderate eye irritation. Wear protective eyewear (goggles, face shield, safety glasses). Avoid contact with eyes, skin, or clothing. Avoid breathing dust or spray mist. Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton® ≥ 14 mils
- Shoes plus socks
- Wear protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607(d-f)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to wildlife and highly toxic to aquatic invertebrates. This pesticide is highly toxic to bees exposed to direct treatment on blooming crops/plants or weeds. **DO NOT** apply this product or allow it to drift to blooming crops/plants or weeds while bees are foraging in/or adjacent to the treatment area.

For Terrestrial Uses: DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. **DO NOT** contaminate water when cleaning equipment or disposing of equipment wash waters.

Groundwater Advisory

Thiamethoxam has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into the groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may contaminate water through drift of spray in wind. This product has a high potential for runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff (See manual at the following internet address: <http://www.wsi.nrcs.usda.gov/products/W2Q/pest/core4.html>).

Runoff of this product will be reduced by avoiding applications when rainfall is forecast to occur within 48 hours.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT use, pour, spill, or store near heat or open flame.

PROTECTION OF POLLINATORS



APPLICATION RESTRICTIONS EXIST FOR THIS PRODUCT BECAUSE OF RISK TO BEES AND OTHER INSECT POLLINATORS. FOLLOW APPLICATION RESTRICTIONS FOUND IN THE DIRECTIONS FOR USE TO PROTECT POLLINATORS.



Look for the bee hazard icon in the Directions for Use for each application site for specific use restrictions and instructions to protect bees and other insect pollinators.

This product can kill bees and other insect pollinators.

Bees and other insect pollinators will forage on plants when they flower, shed pollen, or produce nectar.

Bees and other insect pollinators can be exposed to this pesticide from:

- Direct contact during foliar applications, or contact with residues on plant surfaces after foliar applications.
- Ingestion of residues in nectar and pollen when the pesticide is applied as a seed treatment, soil, tree injection, as well as foliar applications.

When Using This Product Take Steps To:

- Minimize exposure of this product to bees and other insect pollinators when they are foraging on pollinator attractive plants around the application site.
- Minimize drift of this product on to beehives or to off-site pollinator attractive habitat. Drift of this product onto beehives or off-site to pollinator attractive habitat can result in bee kills.

Information on protecting bees and other insect pollinators may be found at the Pesticide Environmental Stewardship website at: <https://pesticidestewardship.org/pollinator-protection/>

Pesticide incidents (for example, bee kills) must immediately be reported to the State/Tribal lead agency. For contact information for your State, go to: <http://www.aapco.org>. Pesticide incidents must also be reported to the National Pesticide Information Center at: www.npic.orst.edu or directly to EPA at: beekill@epa.gov

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

See individual use sites for specific pollinator protection application restrictions. If none exist, follow these directions for foliar applications to commercially-grown plants and ornamentals that are attractive to pollinators and non-agricultural use sites:

1. FOR FOOD/FEED CROPS AND COMMERCIALY GROWN ORNAMENTALS NOT UNDER CONTRACT FOR POLLINATION SERVICES BUT ARE ATTRACTIVE TO POLLINATORS



DO NOT apply this product while bees are foraging. **DO NOT** apply this product until flowering is complete and all petals have fallen unless one of the following conditions has been met:

- The application is made to the target site after sunset.
- The application is made to the target site when temperatures are below 55°F.
- The application is made in accordance with a government-initiated public health response.
- The application is made in accordance with an active state-administered apiary registry program where beekeepers are notified no less than 48-hours prior to the time of the planned application so that the bees can be removed, covered or otherwise protected prior to spraying.
- The application is made due to an imminent threat of significant crop loss, and a documented determination consistent with an IPM plan or predetermined economic threshold is met. Every effort should be made to notify beekeepers no less than 48-hours prior to the time of the planned application so that the bees can be removed, covered or otherwise protected prior to spraying.

2. FOR NON-AGRICULTURAL USES



DO NOT apply this product while bees are foraging. **DO NOT** apply this product to plants that are flowering. Only apply after all flower petals have fallen off.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

RESTRICTIONS

- This product is classified as restricted use in New York State.
- Sale, use and distribution of this product in Nassau and Suffolk counties in the state of New York is prohibited.
- In New York State, use is limited to Christmas trees and indoor greenhouse use only.
- **DO NOT** apply this product, by any application method, to linden, basswood, or *Tilia* species.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours except for automatic fogger use in greenhouses (See "Automatic Fogger Use in Greenhouses" below).

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of barrier laminate, butyl rubber >14 mils, nitrile rubber >14 mils, neoprene rubber >14 mils, polyvinyl chloride (PVC) >14 mils, or Viton® >14 mils
- Shoes plus socks

Automatic Cold Fogger Use in Greenhouses

- If there is no ventilation, the REI is **24 hours**.

Early Entry based upon Ventilation Criteria

- Early entry is permitted (with the PPE listed above) and respiratory protection consisting of a protection factor (PF) 10 respirator fitted with an organic-vapor removing filter after **four (4) hours** to perform short duration activities such as opening windows or turning on fans or ventilation systems.
- Early entry is permitted (with the PPE listed above) after **two (2) hours** of ventilation using fans or other mechanical ventilation systems.
- Early entry is permitted (with the PPE listed above) after **four (4) hours** of ventilation using vents, windows or other passive ventilation.
- Early entry is permitted (with the PPE listed above) after **ten (10) complete air exchanges** have been completed.

After 12 hours, if the greenhouse has been ventilated according to the above criteria, no PPE is required to enter the treated area.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

DO NOT enter treated areas without protective clothing until sprays have dried.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR INSECT CONTROL, AND/OR ILLEGAL RESIDUES.

PRODUCT INFORMATION

Sharda Thiamethoxam 25% WDG is a broad-spectrum insecticide that is effective when applied at label rates to the foliage and soil. **Sharda Thiamethoxam 25% WDG** is active against many sucking and chewing insect pests by contact and ingestion.

Sharda Thiamethoxam 25% WDG is relatively short-lived on the surface of plant foliage, readily absorbed into plant tissues, and rainfast once it has dried. The rapid translaminar absorption and distribution within leaves provides excellent knockdown and residual control of foliar feeding insects.

In the soil, the active ingredient in **Sharda Thiamethoxam 25% WDG** will control soil pests upon contact or ingestion and is also readily taken up by plant roots. The active ingredient moves upwards in the plant to the site of pest infestation. Through feeding on the plant, pests are exposed to the active ingredient in **Sharda Thiamethoxam 25% WDG**. Feeding will stop within minutes to hours of exposure,

followed by death of the pest. The moderate persistence of **Sharda Thiamethoxam 25% WDG** in the soil and foliage also provides residual control of labeled pests.

THIAMETHOXAM	GROUP	4A	INSECTICIDE
--------------	-------	----	-------------

RESISTANCE MANAGEMENT

For resistance management, **Sharda Thiamethoxam 25% WDG** contains Thiamethoxam and is classified in the neonicotinoids chemical class as a Group 4A insecticide, nicotinic acetylcholine receptor (nAChR) competitive modulators.

Any insect population may contain individuals naturally resistant to **Sharda Thiamethoxam 25% WDG** and other Group 4A insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies must be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of **Sharda Thiamethoxam 25% WDG** or other Group 4A insecticides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. **DO NOT** rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues for the targeted pests between the individual components of a mixture.
- In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual insecticides selected for use in mixtures must be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide/acaricides use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological, and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance, contact Sharda USA LLC or go to <https://shardausa.com>.

Plant Safety

Sharda Thiamethoxam 25% WDG is safe on most ornamental plants. To determine the plant safety of mixtures of **Sharda Thiamethoxam 25% WDG** with other products, or if you do not have experience with a particular tank mix or with **Sharda Thiamethoxam 25% WDG** applied alone to a particular plant, test **Sharda Thiamethoxam 25% WDG** or the mixture on a small number of plants to determine any adverse plant safety effects before making large scale applications to a greater number of plants.

Among container-grown ornamentals and ornamentals grown in planting or liner beds, some plant species may be sensitive to fertilizers. **DO NOT** tank mix **Sharda Thiamethoxam 25% WDG** with fertilizers for treatment of these plants grown under these conditions.

Applications of **Sharda Thiamethoxam 25% WDG** to varieties of Yellow Honey Locust may result in leaf chlorosis and leaf abscission. Although the effects are temporary, do not use **Sharda Thiamethoxam 25% WDG** on Yellow Honey Locust varieties.

MIXING PROCEDURES

Prepare only the amount of spray mixture that is needed for the immediate application. Thoroughly clean spray equipment before using this product. Vigorous agitation is necessary for proper dispersal of the product. Maintain maximum agitation throughout the spraying operation.

DO NOT let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area.

Sharda Thiamethoxam 25% WDG Alone

Add 1/2 of the required amount of water to the mix tank. With the agitator running, add the **Sharda Thiamethoxam 25% WDG** to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after the **Sharda Thiamethoxam 25% WDG** has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

Tank Mix Compatibility

Sharda Thiamethoxam 25% WDG is compatible with most insecticide, fungicide, and foliar nutrients products. However, test the physical compatibility of **Sharda Thiamethoxam 25% WDG** with tank-mix partners before use. To determine the physical compatibility of **Sharda Thiamethoxam 25% WDG** with other products, use a jar test, as described below.

Using a quart jar, add the proportionate amounts of the products to 1 qt of water. Add wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

Sharda Thiamethoxam 25% WDG + Tank Mixtures

Add 1/2 of the required amount of water to the mix tank. Start the agitator running before adding any tank mix partners. Add the tank mix partners in this order: products packaged in water-soluble packaging, wettable powders, wettable granules (dry flowables) such as **Sharda Thiamethoxam 25% WDG**, liquid flowables, liquids and emulsifiable concentrates. Always allow each tank mix partner to become fully dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation until all the mixture has been applied.

Note: When using **Sharda Thiamethoxam 25% WDG** in tank mixtures, add all products in water-soluble packaging to the tank before any other tank mix partner, including **Sharda Thiamethoxam 25% WDG**. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank mix partner to the tank.

If using **Sharda Thiamethoxam 25% WDG** in a tank mixture, observe all directions for use, crop/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank mix product label. **DO NOT** exceed any label dosage rate, and follow the most restrictive label precautions and limitations. **DO NOT** mix this product with any product, which prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are labeled.

APPLICATION PROCEDURES

DO NOT apply this product, by any application method, to linden, basswood, or *Tilia* species.

Foliar Applications

Make foliar applications in an adequate water volume to achieve thorough and uniform coverage without excessive runoff (to drip). On hard to wet plants, spray adjuvants that improve wetting of foliage are recommended. **DO NOT** use spray adjuvants that may bind **Sharda Thiamethoxam 25% WDG** to the leaf surface, which would limit absorption into the foliage. **Sharda Thiamethoxam 25% WDG** can be applied in a range of spray volumes (concentrate to dilute spray volumes), provided thorough and uniform coverage is obtained.

Handheld Application Equipment

Applications can be made through most types of low pressure handheld application equipment.

DO NOT apply through any type of handheld fogger equipment.

Automatic Cold Fogger Applications – Greenhouses

Applications can be made in greenhouses with automatic cold fogger equipment (such as the Damm AutoFog). Apply the same amount of **Sharda Thiamethoxam 25% WDG** per unit area as would be applied in a dilute spray volume over the same area.

DO NOT apply through fogger equipment requiring workers to be present in the greenhouse during the fogger application.

Important: Read and follow the REI and PPE requirements for “**Automatic Cold Fogger Use in Greenhouses**” in the Worker Protection Standard **AGRICULTURAL USE REQUIREMENTS** box.

Soil or Soil-less Medium Applications

Sharda Thiamethoxam 25% WDG is effective as a soil and systemic insecticide when applied to the soil using band, broadcast or drench applications. When applying **Sharda Thiamethoxam 25% WDG** as a drench, apply enough solution to sufficiently wet ½ – ¾ of the root zone without allowing runoff out of the container. With irrigation or rainfall, move **Sharda Thiamethoxam 25% WDG** into the root zone of plants or the location of soil-dwelling pests, preferably within 3 days of treatment and, optimally, within 24 hours.

Excessive watering or heavy rainfall may reduce the effectiveness of **Sharda Thiamethoxam 25% WDG**. Allow a minimum of 7 days after watering-in for maximum uptake of **Sharda Thiamethoxam 25% WDG** by the plant's root system.

DO NOT apply to soils or growing media above field or pot capacity. **DO NOT** apply to waterlogged soil or growing media.

Application Timing

Begin foliar applications when labeled pests first appear or when economic thresholds have been reached.

For applications made to growing media (soil or soil-less media), **Sharda Thiamethoxam 25% WDG** must move to the feeding site of the target pest, whether this is in the root zone or the above ground portion of the host plant. Therefore, control of pests from the systemic activity of **Sharda Thiamethoxam 25% WDG** may be delayed for 1 or more weeks depending on plant size, rate of uptake from the soil, or rate of translocation within the plant, etc.

Systemic activity in large plants such as those in field nurseries or Christmas tree plantations may require several weeks to months before control of stem and foliar pests is achieved. If economic thresholds for target pests are low, or plants are historically damaged, apply **Sharda Thiamethoxam 25% WDG** to the growing media (soil or soil-less media) to help prevent pests.

Application Through Irrigation Systems

Apply **Sharda Thiamethoxam 25% WDG** only through micro-irrigation systems (for example, drip, trickle, spaghetti tubes, and micro-sprinklers), overhead irrigation, ebb and flood irrigation, or hand held or motorized calibrated irrigation equipment. **DO NOT** apply this product through any other type of irrigation equipment. Allow a minimum of 7 days after watering-in for maximum uptake of **Sharda Thiamethoxam 25% WDG** by the plant's root system. See additional precautions regarding irrigation system application below.

Chemigation

Uniform Water Distribution and System Calibration: The irrigation system must provide uniform distribution of treated water. Plant injury, lack of effectiveness, or illegal pesticide residues in the plant can result from non-uniform distribution of treated water.

The system must be calibrated to uniformly apply the rates specified. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

DO NOT connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

Chemigation Monitoring: A person knowledgeable of the Chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Required System Safety Devices: The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

DO NOT apply when wind speed favors drift beyond the area intended for treatment.

Specific Instructions for Public Water Systems

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back-flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the

water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

DO NOT apply when wind speed favors drift beyond the area intended for treatment.

Ebb and Flood Application

To determine the water volume uptake of **Sharda Thiamethoxam 25% WDG** in ebb and flood applications, bring a minimum of 10 plants up to a known field capacity and allow them to dry out for one to two days.

Re-wet these plants to determine how much water on average each plant will absorb to bring it back to field capacity. Use the water volume absorbed per plant, multiply by the number of pots of the same size being treated, and add the minimum water volume to irrigate the area. This should minimize the return back to the storage tank. Re-use the returned volume with subsequent irrigation or nutrients on the same plants.

Aerial Application

For application to ornamental field nurseries, forest seed/seedling field nurseries or Christmas trees.

Apply **Sharda Thiamethoxam 25% WDG** in water. Increase spray volume where practical to improve coverage. **DO NOT** make applications under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur.

MANDATORY SPRAY DRIFT MANAGEMENT

Aerial Applications:

- **DO NOT** release spray at a height greater than 10 ft. above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a Medium or coarser droplet size (ASABE S572.1).
- For aerial applications: **DO NOT** apply when wind speeds exceed 15 mph at the application site. If wind speed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters. Applicators must use ½ swath displacement upwind at the downwind edge of field. Mount the spray boom on the aircraft so as to minimize drift caused by wing tip vortices.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

Ground Boom Applications:

- Users must only apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy unless making a turf, pasture, or rangeland application, in which case applicators may apply with a nozzle height no more than 4 feet above the ground.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- **Adjust Nozzles** - Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles must be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom must remain level with the crop and have minimal bounce.

RELEASE HEIGHT – Aircraft

Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift. To reduce risk of exposure to sensitive aquatic areas, **DO NOT** make applications when wind direction is toward the aquatic area. **DO NOT** apply when wind velocity exceeds 10 mph. Make applications when wind velocity favors on-target product deposition (approximately 3–10 mph). **DO NOT** apply when wind gusts approach 10 mph.

DO NOT cultivate or plant crops within 25 feet of the aquatic area as to allow growth of a vegetative filter strip.

SPECIFIC USE DIRECTIONS

Foliar Applications

- Ornamental plants, fruit and nut trees (non-bearing), Christmas trees, and forest seedlings grown in greenhouses, lath and shadehouses, containers, field nurseries and interiorscapes.
- In New York State, use is limited to Christmas trees and indoor greenhouse use only.

Pests	Rate	Use Restrictions
Ants (excluding harvester, carpenter and Pharaoh ants)	2–4 oz./100 gal	Make the first application when economic action levels for the target pest are reached. For crops without action levels, make the first application before pest populations cause significant damage to the host plant. Re-apply as needed, but no sooner than every 7 days.
Aphids	or 4–8 oz./A	
Adelgids (including Hemlock Woolly Adelgid)	4–8.5 oz./100 gal	
Lace Bugs	or 8–17 oz./A	

Leaf Beetles Leafhoppers Leafminer Leaf gall forming insects Mealybugs Midges Plant Bug Psyllids (including Asian Citrus Psyllid) Sawflies Soft Scales Thrips – foliar feeding Whiteflies		Apply higher use rates in rate range for longer control or high insect populations. For leaf gall forming insects, apply during egg laying of the generation being targeted. For black vine weevil, apply to foliage of plants when adult feeding damage is first observed. Make only one application per generation. For concentrated spray volumes, apply the same amount of product per unit area as would be applied with dilute spray volumes.
Black vine weevil (adult)	8 oz./100 gal or 17 oz./A or 2 oz./25 gal	
<ul style="list-style-type: none"> DO NOT exceed 17 oz./A per crop or year. For specific information about developmental stages of the target pest, associated damage and action thresholds to properly time applications, consult with your State Cooperative Extension Service. 		

Applications to Soil or Other Growing Media

- Ornamental plants, fruit and nut trees (non-bearing), Christmas trees, and forest seedlings grown in greenhouses, lath and shadehouses, containers, field nurseries and interiorscapes.
- In New York State, use is limited to Christmas trees and indoor greenhouse use only.

Pests	Rate	Use Restrictions
Aphids Fungus gnats Leaf Beetles Leafminers Mealybugs Root Aphid Root Weevil larvae (including <i>Diaprepes abbreviatus</i>) Soft Scales Thrips – foliar feeding (suppression) Whiteflies	4–8.5 oz./100 gal or 8–17 oz./A	Apply to the soil or growing media to help prevent pests in situations where tolerance of insects or damage is low. Allow a minimum of 1 week for smaller plants and at least 2 weeks for large plants (typically found in nurseries, Christmas tree plantations) to translocate Sharda Thiamethoxam 25% WDG to feeding sites of the target pest. Re-apply as needed, but no sooner than every 7 days. Apply as a broadcast, band or drench. Refer to the Application Procedures and Banded Application sections for additional direction.
Larvae (Grubs) of: <i>Aphodius</i> spp. Asiatic garden beetle Billbugs Black turfgrass atenioides Dung beetle European chafer Green June beetle Japanese beetle May or June beetle Northern masked chafer Oriental beetle Scarab beetle Southern masked chafer	6–8.5 oz./100 gal or 12.7–17 oz./A	Apply from adult flight through peak egg hatch of the targeted species. Apply as a broadcast, band or drench treatment over the active root zone. Refer to the Application Procedures and Banded Application sections for additional direction. For drench and chemigation applications made to containerized plant material, apply sufficient volume to moisten the media around the root zone without loss from the bottom of the container.
Fire Ants	1–3 oz./10 gal	Individual Mound Treatment: For small mounds (<6 inch in diameter) apply 1 gal of water per mound and for larger mounds apply 2–3 gal of dilute solution per mound for optimum control. Direct the application to the center of the mound, including a 6-inch diameter circle.
<ul style="list-style-type: none"> For broadcast applications, typical water volume is 1.5–5 gal/1,000 sq. ft.. 		

- For band applications, minimum water volume is 5 gal/1,000 linear feet.
- **DO NOT** exceed 17 oz./A per crop or year.
- **DO NOT** over-water or leach for a minimum of 7 days after watering in for maximum uptake of **Sharda Thiamethoxam 25% WDG** by the plant root system.
- For specific information about developmental stages of the target pest, associated damage, and action thresholds to properly time applications, consult with your State Cooperative Extension Service.

Banded Application

For banded applications, apply uniformly over the row or active root zone in a minimum water volume of 5 gallons per 1,000 linear feet. For optimum activity, rainfall or irrigation is necessary within 24 hours of application.

Sharda Thiamethoxam 25% WDG rate (ounces of product) per 1,000 linear feet of row for particular row spacing.

Rate/A	lb. a.i./A	Row Spacing										
		3 ft	4 ft	4.5 ft	5 ft	6 ft	7 ft	8 ft	9 ft	10 ft	11 ft	12 ft
8 oz./A	0.125	0.55	0.73	0.83	0.92	1.10	1.29	1.47	1.65	1.84	2.02	2.20
12.7 oz./A	0.186	0.88	1.16	1.32	1.46	1.76	2.05	2.33	2.65	2.93	3.23	3.51
17oz./A	0.265	1.17	1.55	1.76	1.95	2.34	2.74	3.11	3.53	3.91	4.30	4.68

Drench Application

When applying **Sharda Thiamethoxam 25% WDG** as a drench, apply enough solution to sufficiently wet $\frac{1}{2}$ – $\frac{3}{4}$ of the root zone without allowing runoff out of the container. Use the following chart as a guideline for the number of pots that this volume will treat based on the size of the pot and drench volume. Use higher drench volumes for woody ornamentals, containers having multiple plants, and those crops with longer production times or needing longer protection.

Note: Drench volumes will vary based on the porosity of the growing media. Conduct your own tests to determine correct drench volumes for your growing media.

Pot Size	Suggested Drench Volume fl. oz./pot	Pots Treated per 100 gal
4 inch standard	2 – 3	6,400 – 4,267
6 inch standard	3 – 4	4,267 – 3,200
8 inch standard	5 – 10	2,560 – 1,280
10 inch standard	12.5 – 25	1,024 – 512

Applications to Grass and Turf Areas In and Around Ornamental Production Facilities

- Grass and turf around container grown plants and areas surrounding field nurseries and greenhouses.
- In New York State, use is limited to Christmas trees and indoor greenhouse use only.
- Not for use on residential lawns or other grass/turf in residential areas.
- Apply **Sharda Thiamethoxam 25% WDG** as a broadcast application to turfgrass for control of targeted pests. To help prevent pests, irrigate within seven (7) days of application to move **Sharda Thiamethoxam 25% WDG** into the root zone where grubs feed. To control pests, irrigate within one day (1) of application to move **Sharda Thiamethoxam 25% WDG** into the root zone where grubs feed. Maintain adequate soil moisture before and after application for optimum control and healthy turfgrass growth. Excessively wet or dry conditions may negatively impact the performance of **Sharda Thiamethoxam 25% WDG** against white grubs and mole crickets. **DO NOT** mow turf until the treated area has been irrigated or rainfall has occurred to allow for maximum and uniform uptake into turfgrass.

Pests	Rate	Use Restrictions
Larvae (Grubs) of: <i>Aphodius</i> spp. Asiatic garden beetle Billbugs Black turfgrass atenioides Dung beetle European chafer Green June beetle Japanese beetle May or June beetle Northern masked chafer Southern masked chafer Oriental beetle Scarab beetle	12.7–17 oz./A or 1.5–1.95 oz./5,000 sq. ft.	Apply up to 45 days before the historical peak of adult flight to 2nd instar grub of the species targeted. For optimum control, treat from peak flight to peak egg hatch.

Mole Crickets (suppression)*	12.7–17 oz./A or 1.5–1.95 oz./5,000 sq. ft.	To suppress damage, treat from first egg hatch to peak egg hatch.
Chinch bugs (suppression)*	12.7–17 oz./A or 1.5–1.95 oz./5,000 sq. ft.	Apply when young nymphs are first observed.
Fire Ants	1–3 oz./10 gal	Individual Mound Treatment: For small mounds (<6 inch in diameter), apply 1 gal of dilute solution per mound. For larger mounds, apply 2–3 gal per mound for optimum control. Direct the application to the center of the mound, including a 6-inch diameter circle.
<p>* Suppression can mean either erratic control, ranging from good to poor, or a consistent level of control below that generally acceptable for commercial purposes.</p> <ul style="list-style-type: none"> Use sufficient water volume (1.5–5 gal/1,000 sq. ft.) to uniformly distribute Sharda Thiamethoxam 25% WDG over the area being treated and to adequately move the active ingredient into the canopy and thatch layers. For turf with heavy thatch (more than ¾ inches), use the higher rates within the rate range. DO NOT exceed 17 oz./A per year. For specific information about developmental stages of the target pest, associated damage and action thresholds to properly time applications, consult with your State Cooperative Extension Service. 		

Foliar Application to Vegetable Plants Grown for Re-sale to Consumers

- For tomatoes, peppers, eggplant, and cucurbit plants grown for re-sale to consumers.
- In New York State, use is limited to Christmas trees and indoor greenhouse use only.

Crop	Pests	Rate	Use Restrictions
Fruiting Vegetables Eggplant Ground-cherry Pepino Peppers (bell, chili, cooking, pimento, and sweet) Tomatillo Tomato	Aphids Colorado Potato Beetle Flea beetles Leafhoppers Whiteflies Pepper Weevil Stink Bugs	2–3 oz./A or 0.23–0.34 oz./ 5,000 sq. ft. 3–5.5 oz./A or 0.34–0.63 oz./ 5,000 sq. ft.	Apply before pests reach damaging levels. Apply higher rates within the listed rate range for heavy infestations. Apply before pests reach damaging levels. Apply higher rates within the listed rate range for heavy infestations.
<ul style="list-style-type: none">• DO NOT exceed 11 oz./A (1.26 oz./5,000 sq. ft.) of Sharda Thiamethoxam 25% WDG per crop.• Allow at least 5 days between applications.			
Cucurbit Vegetables Chayote Chinese waxgourd Citron melon Cucumber Edible gourd Gherkin <i>Momordica</i> species Muskmelon Pumpkin Squash: summer and winter Watermelon	Aphids Flea Beetles Cucumber Beetles (suppression) Leafminers (suppression) Whiteflies	1.5–3 oz./A or 0.17–0.34 oz./ 5,000 sq. ft. 3–5.5 oz./A or 0.34–0.63 oz./ 5,000 sq. ft.	Apply before pests reach damaging levels. Apply higher rates within the listed rate range for heavy infestations.
<ul style="list-style-type: none">• DO NOT exceed 11 oz./A (1.26 oz./5,000 sq. ft.) of Sharda Thiamethoxam 25% WDG per crop.• Allow at least 5 days between applications.			

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal. For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup, procedures and disposal of wastes.

PESTICIDE STORAGE: Store in a cool, dry place.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of Federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control

Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

CONTAINER HANDLING:

[Nonrefillable Container (50 pounds or less):] Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by State and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Sharda USA LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Sharda USA LLC and Seller harmless for any claims relating to such factors.

Sharda USA LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or Sharda USA LLC and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, SHARDA USA LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Sharda USA LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SHARDA USA LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SHARDA USA LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Sharda USA LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Sharda USA LLC.

[All trademarks are the property of their respective owners.]