U.S. ENVI	RONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505P) 1200 Pennsylvania Ave., N.W. Washington, D.C. 20460	EPA Reg. Number: 83529-247	Date of Issuance: 5/9/2023
NOTICE OF P <u>X</u> Regis	ESTICIDE: tration	Term of Issuance: Unconditional	
(under FIFRA, as am	FRA, as amended)	Name of Pesticide Product:	
Name and Address of Registrant (include ZI Sharda USA LLC c/o Wagner Regulatory Associ P.O. Box 640 Hockessin, DE 19707	P Code): ates, Inc.		
<b>Note:</b> Changes in labeling differing in substanc Registration Division prior to use of the label in	e from that accepted in connection with this regist a commerce. In any correspondence on this produ	tration must be submitted to ar act always refer to the above E	nd accepted by the PA registration number.
On the basis of information furnished by the registrant, the above-named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others. This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you: 1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.			
Signature of Approving Official:		Date:	
Heather & McFarley Heather McFarley, Acting Produce Fungicide and Herbicide Branch, Office of Pesticide Programs	ct Manager 24 Registration Division (7505P)	5/9	9/2023
Registration Notice Unconditional v.20150320			

Page 2 of 2 EPA Reg. No. 83529-247 Decision No. 587435

- 2. Make the following label changes before you release the product for shipment:
  - Revise the EPA Registration Number to read, "EPA Reg. No. 83529-247."
- 3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

• Basic CSF dated 5/02/2023

If you have any questions, please contact Sayed Islam by phone at 202-566-2796, or via email at islam.sayed@epa.gov

Enclosure:

• Stamped label

PENOXSULAM GROUP 2 HERBICIDE

# Sharda Penoxsulam 0.24% GR

# **ABN: Marble**

# For Selective Weed Control in Water-Seeded Rice in California.

ACTIVE	<b>INGREDIENT:</b>
--------	--------------------

Penoxsulam: 2-(2,2-difluoroethoxy)-N-(5,8-dimethoxy[1,2,4] triazolo[1,5c]pyrimidin-2-yl)-6-	
(trifluoromethyl)benzenesulfonamide	
OTHER INGREDIENTS:	
TOTAL:	

Contains 0.24 lb. of active ingredient per 100 lbs. bag.

# **KEEP OUT OF REACH OF CHILDREN**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you **DO NOT** understand this label, find someone to explain it to you in detail.)

FIRST AID		
IF IN EYES:	<ul> <li>Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li> </ul>	
	<ul> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> </ul>	
	Call a poison control center or doctor for treatment advice.	
HOTLINE NUMBER		
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at <b>1-800-222-1222</b> . For non-emergency resource information concerning this product, call the National Pesticides Information Center (NPIC) at <b>1-800-858-7378</b> Monday – Friday 8 am – Noon Pacific Time, (NPIC Web site: www.npic.orst.edu). For Chemical Spill, Leak, Fire or Exposure, call CHEMTREC <b>800-424-9300</b> .		
Optional referral stat	ements when booklets and container labels are used:]	

[See label booklet for [complete] [additional] [First Aid,] [Precautionary Statements], [Directions For Use], and [Storage and Disposal].]



7217 Lancaster Pike, Suite A Hockessin, Delaware 19707 EPA Reg. No. 83529-XXX EPA Est. No. XXXXX-XX-XXX

Net Contents: \_\_\_\_\_ [Lbs./Kgs.]



# PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

# PERSONAL PROTECTIVE EQUIPMENT (PPE)

# Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

#### USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

## ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

#### Users should:

• Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

## **ENVIRONMENTAL HAZARDS**

Except when treating rice fields as specified in this product label, **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

# **Non-Target Organism Advisory Statement**

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

## **Groundwater Advisory Statement**

This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

# Surface Water Advisory Statement

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of penoxsulam from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

# **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Read all Directions for Use carefully before applying.

# AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

# DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material

Shoes plus socks

# PRODUCT INFORMATION

Sharda Penoxsulam 0.24% GR is a post-flood, post-emergence herbicide for selective control of susceptible grass, broadleaf, and sedge weeds in California rice. Susceptible weeds at the 1- to 3-leaf stage emerged at the time of application will be controlled (refer to Application Rates and Weeds Controlled table). Apply Sharda Penoxsulam 0.24% GR directly into flooded fields when rice is at a minimum 2 ½-leaf stage of growth. At this stage, the second leaf is fully expanded or unfurled with the third leaf emerged but not expanded (refer to Application Timing section). Average water depth at application must be maintained at 4 - 5 inches, with a minimum water depth of 3 inches in the shallowest part of the field with completely submerged weeds and no exposed soil (refer to Water Management section).

Rice crops grown under adverse environmental conditions, including extreme cold or heat, may express temporary crop injury when **Sharda Penoxsulam 0.24% GR** is applied including slight height reduction or root stunting. Any crop stress or environmental factors which decrease plant metabolism and growth may reduce weed control efficacy and crop tolerance. Such effects are transient and **DO NOT** affect yield. Fertilizer top dressing will speed crop recovery.

# **RESTRICTIONS:**

- Apply Sharda Penoxsulam 0.24% GR by aerial application only.
- Pre-Harvest Interval: DO NOT apply Sharda Penoxsulam 0.24% GR within 60 days of rice harvest.
- DO NOT apply Sharda Penoxsulam 0.24% GR following rainfall until the water depth in checks or basins has been adjusted to meet the correct water depth requirements.
- DO NOT apply where runoff or irrigation water may flow directly onto agricultural land other than rice fields.
- **DO NOT** make more than 1 application per year.
- DO NOT apply more than 18.5 lbs. (0.044 lb. a.i. penoxsulam) of Sharda Penoxsulam 0.24% GR per acre per year.
- **DO NOT** rotate treated land to crops other than rice for 3 months following application.
- DO NOT use on wild rice.
- DO NOT fish or commercially grow fish, shellfish or crustaceans on treated acres during the year of treatment.
- Chemigation: **DO NOT** apply this product through any type of irrigation system.

# PRECAUTIONS:

- Sharda Penoxsulam 0.24% GR may not reliably control known ALS-resistant weed biotypes.
- Applications made to greater than 3 leaf weeds, dense weed populations, and/or average water depth greater than 6 inches may result in reduced weed control.
- Static water conditions for 7 days following an application of **Sharda Penoxsulam 0.24% GR** are advised for best results. It is important not to allow the minimum water depth to drop below 3 inches; therefore, adding water may be required. Water must be trickled in or added very slowly.
- For best results, it is advised that water not be released for 14 days following an application of **Sharda Penoxsulam 0.24% GR**. During this time water depth must be maintained or can be increased as rice grows. Early release of water may result in reduced weed control.
- Shallow water conditions can result in reduced weed control as it may promote additional weed flushes and can increase the potential for crop injury.
- Reduced weed control can be expected at field water inlets due to dilution and movement of water.
- The use of **Sharda Penoxsulam 0.24% GR** on rice varieties with known reduced seedling vigor when established and grown under adverse weather conditions is not advised. Use care when using **Sharda Penoxsulam 0.24% GR** on newly released rice varieties until variety tolerance can be determined.
- Application of **Sharda Penoxsulam 0.24% GR** to fields that have been leveled within 12 months prior to application may result in serious rice injury in areas that have been cut or filled. This does not apply to normal annual land planning activities.
- Under poor field preparation practices (i.e., lack of soil aeration, unleveled ground, large clods, excessive crop residue, weeds emerged prior to flooding, etc.), reduced weed control may occur when using Sharda Penoxsulam 0.24% GR.
- Cold and/or alternating weather conditions during the planting season can result in prolonged weed seed germination resulting in reduced weed control.

# MANDATORY SPRAY DRIFT MANAGEMENT

# Aerial Applications:

- **DO NOT** release spray at a height greater than 10 ft above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- Applicators must use 1/2 swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply when wind speeds exceed 10 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor
- diameter for helicopters.
- **DO NOT** apply during temperature inversions.

# SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

# IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

#### **Controlling Droplet Size – Aircraft**

Adjust Nozzles - Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

#### **RELEASE HEIGHT - Aircraft**

Higher release heights increase the potential for spray drift.

# SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

# **TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

## **TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

## WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

## Handheld Technology Applications

Take precautions to minimize spray drift.

# WEED RESISTANCE MANAGEMENT

For resistance management, **Sharda Penoxsulam 0.24% GR** is a Group 2 (acetolactate synthase (ALS) enzyme inhibitor) herbicide. Any weed population may contain or develop plants naturally resistant to **Sharda Penoxsulam 0.24% GR** and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies must be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of **Sharda Penoxsulam 0.24% GR** or other Group 2 (acetolactate synthase (ALS) enzyme inhibitor) herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method including hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Sharda USA, LLC [at www.shardausa.com].

In addition to the guidance above, registrants are encouraged to incorporate the appropriate elements of Best Management Practices from HRAC and WSSA on the label.

#### APPLICATION INSTRUCTIONS

## **Best Management Practices for Aerial Application**

Equipment used in the application of **Sharda Penoxsulam 0.24% GR** must be carefully calibrated before use and checked frequently during application to be sure it is working properly and delivering a uniform distribution pattern. Avoid overlaps that will increase **Sharda Penoxsulam 0.24% GR** dosage above specified limits.

## **Application Timing**

Application timing is dependent upon the proper rice growth stage and optimum timing for weed species present. Sharda **Penoxsulam 0.24% GR** must be applied to rice after it reaches a minimum of 2 ½-leaf stage (Figure 1); at this stage the second rice leaf is fully expanded or unfurled with third leaf emerged. Sharda Penoxsulam 0.24% GR must be applied when susceptible weeds are from the 1- to 3-leaf stage up to 60 days before harvest (refer to the Application Rates and Weeds Controlled table).

It is important that the stage of rice be closely monitored prior to application to determine the correct timing for an application of **Sharda Penoxsulam 0.24% GR**. Under cool growing conditions, rice development can be delayed; therefore, close crop development monitoring is recommended prior to application.



# Water Management

Immediately prior to application:

- Average water depth at application must be maintained at 4 5 inches, with a minimum water depth of 3 inches in the shallowest part of the field, with completely submerged weeds and no exposed soil.
- If rainfall occurs prior to application, **DO NOT** apply **Sharda Penoxsulam 0.24% GR** until water depth in checks or basins can be adjusted to meet the correct water depth requirements.

First week following application:

- For best results, maintain a static flood for 7 days following application.
- During the static flood period it is important to maintain a minimum water depth of 3 inches. If necessary, water can be slowly added or trickled in to maintain water depth.
- If water is lost from a basin or check, re-flood immediately. Early release of water may result in reduced weed control.
- Failure to maintain the recommended flood depth after an application of Sharda Penoxsulam 0.24% GR may result in reduced weed control or crop tolerance.

Second week following application:

- Following the 7-day static flood period it is advised that water not be released for an additional 7 days.
- During this time, water depth must be maintained or increased as rice grows.
- Shallow water conditions can result in reduced weed control as it may promote additional weed flushes.
- If water is lost from a basin or check, re-flood immediately. Early release of water may result in reduced weed control.

At the conclusion of the 2-week treatment period, it is important to manage water flood depth to prevent secondary weed flushes which can occur if water is released or lowered.

	-0
Weeds Controlled <sup>1</sup> Common Name ( <i>Scientific Name</i> )	Application Rate and Stage of Weed Development
Barnyardgrass (Echinochloa crus-galli) California Arrowhead (Sagittaria montevidensis spp., Caclycina)	
Common Waterplantain (Alisma plantago-aquatica)	45  Hz = 6  and  4  (0.026 Hz = 1/4)
Ducksalad (Heteranthera spp.)	15 lbs. of product/A (0.036 lb. a.i/A)
Watergrass (early and late) (Echinochlog spn.)	un to 3-leaf
Weeds Suppressed	up to 5-lear
Common Name (Scientific Name)	
Gregg's Arrowhead (Sagittaria longiloba)	
Redstem (Ammannia spp.) <sup>3</sup>	
Rice Mimic (Echinochloa spp.)	
Ricefield Bulrush (Scirpus mucronatus) <sup>3</sup>	
Smallflower Umbrellaplant (Cyperus difformis)	
<sup>1</sup> Sharda Penoxsulam 0.24% GR may not reliably control known	AIS-resistant weed histories. Weeds that emerge after the

<sup>1</sup>Sharda Penoxsulam 0.24% GR may not reliably control known ALS-resistant weed biotypes. Weeds that emerge after the treatment period will not be controlled.

<sup>2</sup>Under conditions of late application when weeds are greater than 3-leaf stage, dense weed populations, or average water depth greater than 6 inches deep, 18.5 lbs. (0.044 lb. a.i. penoxsulam) of **Sharda Penoxsulam 0.24% GR** per acre may be applied. Application of **Sharda Penoxsulam 0.24% GR** under these conditions may result in reduced weed control.

<sup>3</sup>Ricefield bulrush (*Scirpus mucronatus*) and redstem (*Ammannia* spp.) are effectively controlled if emerged at the time of treatment with **Sharda Penoxsulam 0.24% GR**.

# STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in the original container. **DO NOT** store in a manner where cross-contamination with other pesticides, fertilizers, food, or feed could occur. In the event of a spill during handling or storage, absorb with sand or other inert material and dispose of absorbent in accordance with the Pesticide Disposal instructions listed below.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

# CONTAINER HANDLING:

[Non-refillable Plastic Containers (Capacity Equal to or Less than 50 Pounds): DO NOT reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two or more times. Offer for recycling, or puncture and dispose of in a sanitary landfill, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.]

[Non-refillable Plastic Containers (Capacity Equal to or more than 50 Pounds): DO NOT reuse or refill this container. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank. Hold container upside down over application equipment or a mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. DO NOT burn, unless allowed by State and local ordinances.] CONTAINER IS NOT SAFE FOR FOOD. FEED. OR DRINKING WATER!

# CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Sharda USA LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Sharda USA LLC and Seller harmless for any claims relating to such factors.

Sharda USA LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or Sharda USA LLC and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, SHARDA USA LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Sharda USA LLC nor Seller shall be liable for any incidental, consequential, or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SHARDA USA LLC AND SELLER FOR ANY AND ALL

CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SHARDA USA LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

Sharda USA LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Sharda USA LLC.

[All trademarks are the property of their respective owners.]

[OPTIONAL MARKETING LANGUAGE]

	[www.shardausa.com]	
1		
2	[Handle with Care]	
3	[This side Up]	
	{The below graphic to be added to box if formulated in the United States}	
4	Proudly Formulated & Packaged In The U.S.A.	