



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505T)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

83529-261

Date of Issuance:

5/10/24

NOTICE OF PESTICIDE:

☒ Registration
☐ Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Sharda Spiromesifen 45.2% SC II

Name and Address of Registrant (include ZIP Code):

Wagner Regulatory Associates, Inc.
P.O. Box 640
7217 Lancaster Pike, Suite A
Hockessin, DE 19707

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.


On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

Continues page 2

Signature of Approving Official:  Melissa Bridges, Product Manager 07 IVB3, Registration Division (7505T)	Date: 5/10/24
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------

EPA Form 8570-6

2. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 83529-261.”
3. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The record for this product currently contains the following CSF(s):

- Basic CSF dated 11/01/2022

If you have any questions, please contact Joseph Belsky at 202-566-2495 or at belsky.joseph@epa.gov.

Enclosure

[MASTER LABEL]

SPIROMESIFEN GROUP 23 INSECTICIDE/ACARICIDE

Sharda Spiromesifen 45.2% SC II

ABN: Speech 4 SC

Intended for professional use. For control of mites and whiteflies on ornamental plants in and around areas such as parks, golf courses, recreational areas, and residential or commercial buildings.

ACTIVE INGREDIENT:	WT. BY %
Spiromesifen: 2-oxo-3-(2,4,6-trimethylphenyl)-1-oxaspiro[4.4]non-3-en-4-yl 3,3-dimethylbutanoate*	45.2%
OTHER INGREDIENTS:	54.8%
TOTAL:	100.0%

*Contains 4 pounds of spiromesifen per U.S. gallon (480 grams per liter).

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you **DO NOT** understand this label, find someone to explain it to you in detail.)


FIRST AID	
IF INHALED:	<ul style="list-style-type: none">Move the person to fresh airIf person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possibleCall a poison control center or doctor for further treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none">Call a poison control center or doctor immediately for treatment advice.Have person sip a glass of water if able to swallow.DO NOT induce vomiting unless told to do so by a poison control center or doctor.DO NOT give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">Take off contaminated clothingRinse skin immediately with plenty of water for 15-20 minutes.Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222 .	
NOTE TO PHYSICIAN	
No specific antidote is known. Treat symptomatically.	

[Optional referral statements when booklets and container labels are used:]
[See label booklet for [complete] [additional] [First Aid,] [Precautionary Statements], [Directions For Use], and [Storage and Disposal].]

Manufactured for:
Sharda USA LLC 
7217 Lancaster Pike, Suite A
Hockessin, Delaware 19707

EPA Reg. No. 83529-EAR
EPA Est. No. XXXXX-XX-XXX

Net Contents: _____ Gals. [L.]



PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, loaders and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves (barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, natural rubber \geq 14 mils, polyethylene, polyvinyl chloride (PVC) \geq 14 mils, viton \geq 14 mils)
- Shoes plus socks

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- shoes plus socks.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove Personal Protective Equipment immediately after handling this product.
- Wash the outside of gloves before removing.
- As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. **DO NOT** contaminate surface water through spray drift. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. **DO NOT** contaminate water when cleaning equipment or disposing of equipment washwater.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of spiromesifen from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours. Sound erosion control practices will reduce this product's potential to reach aquatic sediment via runoff.

Groundwater Advisory

Degradates of spiromesifen have properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Endangered Species Advisory

The use of any pesticide in a manner that may kill or otherwise harm an endangered species or adversely modify their habitat is a violation of Federal law.

Physical or Chemical Hazards

DO NOT mix or allow to come in contact with oxidizing or reducing agents. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire label before using this product. For use on plants intended for aesthetic purposes or climatic modification in areas such as landscapes, interior plantscapes, campgrounds, golf courses, athletic fields, parks, gardens, rights of way and other easements, and around residential, institutional, public, commercial, and industrial buildings..

Not for use on plants being grown for sale or commercial production, or for commercial seed production, or for research purposes.

PRODUCT INFORMATION

Sharda Spiromesifen 45.2% SC II is a Flowable (Suspension Concentrate) formulation that is intended for control of mites and whiteflies on ornamental plants, flowers and foliage plants. The active ingredient contained in **Sharda Spiromesifen 45.2% SC II** is active on all mite development stages. However, mite juvenile stages are often more susceptible than adults. Application should be timed to coincide with early threshold levels in a developing mite population.

Sharda Spiromesifen 45.2% SC II is also highly effective against whitefly nymphs, plus it has a significant effect on the otherwise difficult to control pupal stage. For best results against whitefly, applications should target these life stages. **Sharda Spiromesifen 45.2% SC II** can be applied by ground equipment. However, thorough coverage of all plant parts is required for optimum performance. Performance evaluations should be made 4 - 10 days after application. Apply specific dosage of **Sharda Spiromesifen 45.2% SC II** as needed for control. For best results, uniform coverage of the upper and lower leaf surfaces is recommended. Avoid excess application that results in runoff from plant foliage. An adjuvant may be used to improve coverage on hard to wet foliage. For best results, the treatment should be made when whitefly or mite population begins to build and before a damaging population becomes established. Rate range is provided and is generally dependent on size of the plant and density of the foliage.

Product Features

Controls all stages of whiteflies. Provides knockdown and residual control of all developmental stages of mites.

{Only Necessary if Product Intend to be Applied with Ground Boom}

[MANDATORY SPRAY DRIFT MANAGEMENT

Ground Boom Applications:

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators must select nozzle and pressure that deliver medium or courser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 572 (ASABE S572).
- **DO NOT** apply when wind speeds exceed 15 mph at the application site.
- **DO NOT** apply during temperature inversions.

Wind Speed Restrictions:

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size, canopy and equipment specifications determine drift potential at any given wind speed. **DO NOT** apply when winds are greater than 15 mph or when gusty and windless conditions are present. Risk of exposure to sensitive aquatic areas can be reduced by not making applications when wind direction is toward the aquatic area.

Restrictions During Temperature Inversions:

DO NOT make ground applications during temperature inversions. Drift potential is high during temperature inversions. Temperature inversions restrict vertical air mixing, which causes small, suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical mixing.]

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.

BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

Importance Of Droplet Size

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

[Controlling Droplet Size – Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.]

[BOOM HEIGHT – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.]

Shielded Sprayers

Shielding the individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

Temperature And Humidity

When making applications in hot and dry conditions, use larger droplets to reduce the effects of evaporation.

Temperature Inversions

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Wind

Drift potential generally increases with wind speed. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Boomless Ground Applications:

- Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications:

- Take precautions to minimize spray drift.

RESISTANCE MANAGEMENT

For resistance management, **Sharda Spiromesifen 45.2% SC II** contains a Group 23, lipid biosynthesis inhibitor (LBI) insecticide/acaricide. Any insect/mite population may contain individuals naturally resistant to **Sharda Spiromesifen 45.2% SC II** and other Group 23 insecticides/acaricides. The resistant individuals may dominate the insect/mite population if this group of insecticides/acaricides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay insecticide/acaricide resistance, take the following steps:

- Rotate the use of **Sharda Spiromesifen 45.2% SC II** or other Group 23 insecticides/acaricides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides/acaricides from a different group that are equally effective on the target pest when such use is permitted. **DO NOT** rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues for the targeted pests between the individual components of a mixture.
- In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide/acaricides use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.

- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact your local Sharda USA representative at [1-(910) 859-3090 Monday through Friday, 9AM to 4 PM EST, or] shardausa.com.

APPLICATION INSTRUCTIONS

Apply to shrubs, trees (including non-bearing fruit and nut trees), flowers and foliage plants in outdoor landscapes and interior plantscapes.

Apply specific dosage of **Sharda Spiromesifen 45.2% SC II** as needed for control. Good coverage of the foliage is recommended. An adjuvant may be used to improve coverage on hard to wet foliage. For best results the treatment should be made when the whitefly or mite population begins to build and before a damaging population becomes established. **Sharda Spiromesifen 45.2% SC II** is highly effective against the egg and nymphal stages of whiteflies and mites. For best results, applications should target these stages.

For maximum efficacy against adult whiteflies, use the 4 fl. oz. rate of **Sharda Spiromesifen 45.2% SC II**. For heavy infestations, two applications may be necessary to control adult whiteflies. Rate range is provided and is generally dependent on size of the plant and density of the foliage. Apply when pests first appear and prior to leaf damage or discoloration. Apply with ground application equipment including [ground boom], boomless, or handheld sprayers, in adequate water for uniform coverage.

Pests Controlled	Rate Per Application
Spider mites (including twospotted spider mite, southern red mite, maple spider mite, spruce spider mite, honeylocust spider mite, Euonymus mite, boxwood spider mite, tumid mite and Lewis mite) False spider mites Rust and blister mites (family <i>Eriophyidae</i>) Tarsonemid mites (including broad mites and cyclamen mites)	1.4-4 Fl. Oz. (40-120 mL)/100 gallons of spray solution or 2-6 mL/5 gallons of spray solution
Whiteflies (Including silverleaf, greenhouse, and sweet potato whitefly)	2-4 Fl. Oz. (60-120 mL)/100 gallons of spray solution or 3-6 mL/5 gallons of spray solution
Restrictions: <ul style="list-style-type: none"> • Maximum Application Rate for Residential Trees and Greenhouse Ornamentals: 0.125 lbs. a.i./A. • Maximum Application Rate for Ornamental Gardens: 0.0625 lbs. a.i./A • DO NOT use on vegetable gardens. • DO NOT use on plants intended for use as feed or forage. • DO NOT apply via chemigation. • DO NOT use in commercial greenhouses or nurseries or on bearing fruit or nut trees. • DO NOT apply Sharda Spiromesifen 45.2% SC II more than three times per calendar year. • For all crops and use sites DO NOT use more than 0.35 lbs. spiromesifen per acre per year. 	

Plant Tolerance

Sharda Spiromesifen 45.2% SC II has been evaluated for phytotoxicity on a wide range of ornamental plants. However, due to the large number of species and varieties of ornamental plants, it is impossible to test every one for tolerance to **Sharda Spiromesifen 45.2% SC II**. The professional user should determine if **Sharda Spiromesifen 45.2% SC II** could be used safely prior to widespread use. In a small area, test the recommended rates on a small number of plants for phytotoxicity prior to widespread use. Before using **Sharda Spiromesifen 45.2% SC II** in tank mixture with other products and adjuvants, test the mixture on a small number of plants for phytotoxicity prior to widespread use.

Use of **Sharda Spiromesifen 45.2% SC II** on ferns, English ivy (*Hedera helix*), lily-of-the-Incas (*Alstroemeria*), *Dracaena*, *Peperomia*, croton, *Cordyline*, Mexican heather, fuschia, geraniums (*Pelargonium* spp.), *Schefflera*, and *Impatiens* spp. is not recommended.

Irrigation Timing

If irrigation is used, conduct irrigations efficiently to prevent excessive loss of irrigation waters through runoff. Time the applications to allow sprays to dry prior to rain or sprinkler irrigations. Allow at least 24 hours between application of product and any irrigation that results in surface runoff into lakes, reservoirs, rivers, permanent streams, marshes, potholes, vernal pools, natural ponds, estuaries, or commercial fish farm ponds.

ADDITIONAL DIRECTIONS FOR USE WHEN USED IN CALIFORNIA

Allow growth of a vegetative filter strip within 25 feet (on which the product should not be applied) along lakes, reservoirs, rivers, permanent streams, marshes, potholes, vernal pools, natural ponds, estuaries, or commercial fish farm ponds.

MIXING INSTRUCTIONS

Mix pesticides in areas not prone to runoff such as concrete mixing/loading pads, disked soil in flat terrain or graveled mix pads, or use a suitable method to contain spills and/or rinsate. Properly empty and triple-rinse pesticide containers at time of use as stated in the **STORAGE AND DISPOSAL** section.

Mixing and Loading Requirements

To help prevent potential contamination of groundwater, use of a properly designed and maintained containment pad for mixing and loading of any pesticide into application equipment. If containment pad is not used, maintain a minimum distance of 25 feet between mixing and loading area and potential surface to groundwater conduits such as field sumps, uncased well heads, sink-holes, or field drains.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

COMPATIBILITY

Sharda Spiromesifen 45.2% SC II is physically and biologically compatible with many registered pesticides and fertilizers or micronutrients. When considering mixing **Sharda Spiromesifen 45.2% SC II** with other pesticides, or other additives, first contact your supplier for advice. For further information, contact your local Sharda Representative. If your supplier has no experience with the combination you are considering, conduct a test to determine physical compatibility. To determine physical compatibility, add the required proportions of each chemical with the same proportion of water, as will be present in the chemical supply tank, into a suitable container, mix thoroughly and allow to stand for five minutes. If the combination remains mixed, or can be readily re-mixed, the mixture is considered physically compatible.

ORDER-OF-MIXING

Sharda Spiromesifen 45.2% SC II may be used with other recommended pesticides, fertilizers and micronutrients. Mix pesticides in areas not prone to runoff such as concrete mixing/loading pads, disked soil in flat terrain or graveled mix pads, or use a suitable method to contain spills and/or rinsate. Properly empty and triple-rinse pesticide containers at the time of use.

The proper mixing procedure for **Sharda Spiromesifen 45.2% SC II** alone or in tank mix combinations with other pesticides is:

1. Fill the spray tank $\frac{1}{4}$ to $\frac{1}{2}$ full with clean water;
2. While recirculating and with the agitator running, add any products in PVA bags. Allow time for thorough mixing;
3. Continue to fill spray tank with water until $\frac{1}{2}$ full;
4. Add any other wettable powder (WP) or wettable granules (WG) products;
5. Add the required amount of **Sharda Spiromesifen 45.2% SC II**, and any other "flowable" (FL or SC) type products;
6. Allow enough time for thorough mixing of each product added to tank;
7. If applicable, add any remaining tank mix components: emulsifiable concentrates (EC), fertilizers and micronutrients.
8. Fill spray tank to desired level and maintain constant agitation to ensure uniformity of spray mixture.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store this product in a cool, dry place in its original container only. **DO NOT** store this product near fertilizers, seeds, or other pesticides. If this product is spilled, sweep up the spillage and dispose pursuant to the below Pesticide Disposal instructions. **DO NOT** allow product or containers to freeze.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

[Less Than or Equal to 5 Gallons] {HDPE} [Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or

reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration.]

[Greater Than 5 Gallons] {HDPE} [Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system Repeat this rinsing procedure two more times.]

[For Bulk and Mini-Bulk Containers] {HDPE} [Refillable container. Refill this container with pesticide only. **DO NOT** use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by State and local authorities.]

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Sharda USA LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Sharda USA LLC and Seller harmless for any claims relating to such factors.


Sharda USA LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or Sharda USA LLC and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, SHARDA USA LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Sharda USA LLC nor Seller shall be liable for any incidental, consequential, or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SHARDA USA LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SHARDA USA LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Sharda USA LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Sharda USA LLC.

[All trademarks are the property of their respective owners.]

[OPTIONAL MARKETING LANGUAGE]

1	[shardausa.com] 
2	[Handle with Care]
3	[This side Up]
4	{The below graphic to be added to box if formulated in the United States} 