



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

July 24, 2025

SENT BY EMAIL

Freddy Shelley
freddy.shelley@syntechresearch.com
SHARDA USA LLC

Subject: Labeling Notification per Pesticide Registration Notice (PRN) 98-10 - correct misspellings and aerial restriction language.
Product Name: Sharda Trifloxystrobin 50% WDG
Admin Number: 83529-297
EPA Receipt Date: 07/10/2025
Action Case Number: 00661660

Dear Freddy Shelley:

The U.S. Environmental Protection Agency is in receipt of your application for notification under Pesticide Registration Notice 98-10 for the above referenced product. The EPA has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The labeling submitted with this application has been stamped "Notification" and will be placed in our records.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains statements or claims substantially differing from statements or claims made in connection with obtaining a FIFRA section 3 registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

If you have questions, please contact Thomas Harty via email at harty.thomas@epa.gov.

Sincerely,

Thomas Harty, Risk Manager
FB, RD Office of Pesticide Programs

NOTIFICATION[MASTER LABEL]
83529-297

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

07/24/2025

TRIFLOXYSTROBIN	GROUP	11	FUNGICIDE
-----------------	-------	----	-----------

Sharda Trifloxystrobin 50% WDG**[ABN: Globe WDG]**

For Control of Certain Foliar, Stem, and Root Diseases of Turfgrass Including Golf Courses, Institutional, Commercial and Residential Lawns, Sod Farms, Sports Fields, Parks, Municipal Grounds and Cemeteries, and of Ornamentals Grown in Interiorscapes, Field Nursery Plantings, Forest Nurseries, Residential and Commercial Landscapes, Greenhouses, Lath and Shadehouses, Containers, and Other Enclosed Structures.

ACTIVE INGREDIENT:

Trifloxystrobin: (E,E)-alpha-(methoxyimino)-2-[[[1-[3-(trifluoromethyl)phenyl]ethylidene]amino]oxy]methyl]-methylester

WT. BY %

50.0%

OTHER INGREDIENTS: 50.0%

TOTAL: 100.0%

[Water-dispersible granule]

KEEP OUT OF REACH OF CHILDREN**CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you **DO NOT** understand the label, find someone to explain it to you in detail.)

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.
HOTLINE NUMBERS	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222 . For general information about this product, contact the National Pesticides Information Center (NPIC) at 1-800-858-7378 , Monday through Friday, 8 AM to 12 PM PST, or at http://npic.orst.edu .	
NOTE TO PHYSICIAN	
If ingested, induce emesis or lavage stomach. Treat symptomatically.	

{Optional referral statements when booklets and container labels are used:}

[See label booklet for [complete] [additional] [First Aid,] [Precautionary Statements], [Directions For Use], and [Storage and Disposal].]

EPA Reg. No.: 83529-297

EPA Est. No.: _____

Manufactured for:

Sharda USA LLC

7217 Lancaster Pike, Suite A
Hockessin, Delaware 19707

Net Contents: _____ Lbs. [Kg]
[Batch Code/Lot No.: _____]

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Harmful if absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material (nitrile rubber ≥ 14 mils, butyl rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber ≥ 14 mils, polyethylene, polyvinyl chloride (PVC) ≥ 14 mils, ~~viton~~Viton ≥ 14 mils, and/or barrier laminate)
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high-water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. **DO NOT** contaminate water when disposing of equipment wash water or rinsate. Applying this product when rain is not predicted for the next 24 hours will help reduce potential risk to aquatic invertebrates by reducing pesticide runoff from the treatment area into water bodies. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark.

Groundwater Advisory

Several trifloxystrobin degradates have properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT use, pour, spill, or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Intended for use by professional applicators.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **12 hours**.

Exception: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material (nitrile rubber ≥ 14 mils, butyl rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber ≥ 14 mils, polyethylene, polyvinyl chloride (PVC) ≥ 14 mils, ~~viton~~Viton ≥ 14 mils, and/or barrier laminate)
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

DO NOT enter treated areas without protective clothing until sprays have dried.

PRODUCT INFORMATION

Sharda Trifloxystrobin 50% WDG is a site-specific fungicide for use on turfgrass and ornamentals with protective and curative activity. **Sharda Trifloxystrobin 50% WDG** penetrates the plant and provides trans laminar activity via a high affinity for the waxy layer of the plant surface, localized vapor movement and re-deposition on the plant.

Use Restrictions:

- ~~Aerial Application: DO NOT apply this product by aerial application.~~
- ~~Chemigation: DO NOT apply this product through any type of irrigation system.~~

Maximum Use Rates:

- For turfgrass, up to 34.5 oz. of **Sharda Trifloxystrobin 50% WDG** can be applied per acre per year.
- For plants grown in outdoor nurseries, outdoor seedbeds, field plantings, and landscapes, up to 34.5 oz. of **Sharda Trifloxystrobin 50% WDG** per acre of production or acre of landscape per year or crop cycle can be used.
- For seedlings and plants grown in greenhouses, containers, and other enclosed structures, up to 120 oz. of **Sharda Trifloxystrobin 50% WDG** per acre per year or crop cycle can be used. In California only, **DO NOT** apply more than 34.5 oz. of **Sharda Trifloxystrobin 50% WDG** per acre per year or crop cycle to seedlings or plants grown in greenhouses, containers, and other enclosed structures.

Resistance Management

Sharda Trifloxystrobin 50% WDG belongs to the strobilurin class of chemistry which exhibits no known cross-resistance to other chemical classes including sterol inhibitors, dicarboximides, benzimidazoles, anilinopyrimidines, or phenylamides. However, certain fungal pathogens are known to develop resistance to products used repeatedly. Because resistance development cannot be predicted, the use of this product must conform to resistance management strategies. Such strategies may include rotating and/or tank mixing with products having different modes of action; or limiting the total number of applications per season. Sharda USA LLC encourages responsible product stewardship to ensure effective long-term control of the fungal diseases on this label. See specific directions in the **TURFGRASS DISEASE CONTROL** and **ORNAMENTAL DISEASE CONTROL** sections.

TANK MIXING PROCEDURES

Water Dispersible Granules

Prepare no more spray mixture than is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Agitation is necessary for proper dispersal of the product. Maintain agitation throughout the spraying operation. **DO NOT** let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area.

Sharda Trifloxystrobin 50% WDG Alone: Add half of the required amount of water to the mix tank. With the agitator running, add the **Sharda Trifloxystrobin 50% WDG** to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after **Sharda Trifloxystrobin 50% WDG** has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

Sharda Trifloxystrobin 50% WDG Tank Mixtures: Add half of the required amount of water to the mix tank. Start the agitator running before adding any tank mix partners. Add tank mix partners in this order: (1) products packaged in water-soluble packaging, wettable powders, wettable granules (dry flowables) such as **Sharda Trifloxystrobin 50% WDG**; (2) liquid flowables, liquids; and (3) emulsifiable concentrates. Always allow each tank mix partner to become fully dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation until all of the mixture has been applied.

NOTE: When using **Sharda Trifloxystrobin 50% WDG** in tank mixtures, add all products in water-soluble packaging to the tank before any other tank mix partner, including **Sharda Trifloxystrobin 50% WDG**. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank mix partner to the tank.

If using **Sharda Trifloxystrobin 50% WDG** in a tank mixture, observe all directions for use, crop/sites, use rates, dilution ratios, precautions, and limitations that appear on the tank mix product label. Label dosage rate must not be exceeded, and the most restrictive label precautions and limitations must be followed. **DO NOT** mix this product with any product that prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are labeled.

Sharda Trifloxystrobin 50% WDG is compatible with most insecticide, fungicide, and foliar nutrient products. However, the compatibility of **Sharda Trifloxystrobin 50% WDG** with tank mix partners must be tested before use.

To determine biological compatibility with other products, mix the products in the desired proportions, spray on target plants, and observe for phytotoxicity 7 days after the application.

To determine the physical compatibility of **Sharda Trifloxystrobin 50% WDG** with other products, use a jar test, as described below. Using a quart jar, add the proportionate amounts of the products to 1 qt. of water. Add wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

Observe all directions, precautions, and limitations on labeling of all products used in tank mixes. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are registered.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Use with Additives: Use of spray additives are not required. Evaluate any spray additive prior to use. **DO NOT** use in conjunction with organosilicate-based products, or plant injury may occur. Label directions are based on data with no additives.

Aerial Application: DO NOT apply this product by aerial application.

Chemigation: DO NOT apply this product through any type of irrigation system.

[Water Soluble Packets]

The enclosed packets containing **Sharda Trifloxystrobin 50% WDG** are water soluble. **DO NOT** allow packets to become wet prior to adding to the spray tank. **DO NOT** handle with wet hands. Reseal outer carton/container to protect remaining packets.

To prepare the spray mixture, drop the required number of unopened packets into the spray tank while filling the water to the desired level (refer to the **Conversion Table for Sharda Trifloxystrobin 50% WDG in Water-Soluble Packets** section). Operate the agitator while mixing. Depending on the water temperature and the degree of agitation, the packets must be completely dissolved within a few minutes from the time they were added to the water.

ATTENTION: DO NOT use **Sharda Trifloxystrobin 50% WDG** water-soluble packets in a tank-mix with products that contain boron or release free chlorine. The resultant reaction of PVA and boron or free chlorine is a plastic which is not soluble in water or solvents such as diesel oils, kerosene, gasoline, or alcohol. Use of chlorinated water is acceptable. **DO NOT** attempt to use the PVA packets directly in liquid fertilizer, diesel oils, or summer spray type oils as in ULV or LV uses. PVA packets are water soluble not fertilizer or oil soluble. If liquid fertilizer or oils are to be added to the mix water, add **Sharda Trifloxystrobin 50% WDG** packets first and wait until they are dissolved before adding liquid fertilizer or oil.

Sharda Trifloxystrobin 50% WDG Alone: Add half of the required water to the mix tank. With the agitator running, add the required number of **Sharda Trifloxystrobin 50% WDG** packets to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after the **Sharda Trifloxystrobin 50% WDG** packets have completely dissolved and dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

Sharda Trifloxystrobin 50% WDG plus Tank Mixtures: Add half of the required amount of water to the tank mix. Start the agitation running before adding **Sharda Trifloxystrobin 50% WDG** water-soluble packets and any tank mix partner(s). Add tank mix partners in this order: (1) products packaged in water-soluble packaging (such as **Sharda Trifloxystrobin 50% WDG**), wettable powders, wettable granules (dry flowables); (2) liquid flowables, liquids; and (3) emulsifiable concentrates. Always allow each tank mix partner to become fully dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation

until all of the mixture has been applied.

NOTE: When using **Sharda Trifloxystrobin 50% WDG** in tank mixtures, all products in water-soluble packaging must be added to the tank before any other tank mix partner, including **Sharda Trifloxystrobin 50% WDG**. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank mix partner to the tank.

If using **Sharda Trifloxystrobin 50% WDG** in a tank mixture, observe all directions for use, crop/sites, use rates, dilution ratios, precautions, and limitations, which appear on the tank mix product label. **DO NOT** exceed labeled rates and observe the most restrictive of the labeling limitations and precautions of all products used in mixtures. This product must not be mixed with any product, which prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are registered.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Sharda Trifloxystrobin 50% WDG is compatible with most insecticide, fungicide, and foliar nutrient products. However, the compatibility of **Sharda Trifloxystrobin 50% WDG** with tank mix partners must be tested before use.

To determine biological compatibility with other products, mix the products in the desired proportions, spray on target plants and observe for phytotoxicity 7 days after the application.

Observe all directions, precautions, and limitations on labeling of all products used in tank mixes. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are registered.

Use with Additives: Use of spray additives are not required. Evaluate any spray additive prior to use. **DO NOT** use in conjunction with organosilicate-based products, or plant injury may occur. Label directions are based on data with no additives.]

MANDATORY SPRAY DRIFT MANAGEMENT

Aerial Applications:

- For aerial applications, do not apply when wind speeds exceed 15 mph at the application site. If the wind speed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- **DO NOT** release spray at a height greater than 10 ft above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators must use 1/2 swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply during temperature inversions.

Airblast Applications:

- Sprays must be directed into the canopy.
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- User must turn off outward pointing nozzles at row ends and when spraying outer row.
- **DO NOT** apply during temperature inversions.

Ground Boom Applications:

- Apply with the nozzle height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size - Ground Boom

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

~~Controlling Droplet Size—Aircraft~~

- ~~Adjust Nozzles—Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.~~

BOOM HEIGHT - Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

~~RELEASE HEIGHT—Aircraft~~

~~Higher release heights increase the potential for spray drift.~~

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Handheld Technology Applications:

- Take precautions to minimize spray drift.

TURFGRASS DISEASE CONTROL

Sharda Trifloxystrobin 50% WDG is a preventive and curative fungicide that may be applied to turf sites including golf courses, institutional, commercial and residential lawns, sod farms, sports fields, parks, municipal grounds and cemeteries.

Sharda Trifloxystrobin 50% WDG is a broad-spectrum fungicide for the control of brown patch (*Rhizoctonia solani*), anthracnose (*Colletotrichum graminicola*), red thread (*Corticium fusiforme*), pink patch (*Limonomyces roseipellis*), Leaf spot (*Bipolaris* spp., *Drechslera* spp.), gray leaf spot (*Pyricularia grisea*), rust (*Puccinia* spp.), pink snow mold (*Microdochium nivale*), fusarium patch (*Fusarium nivale*), rapid blight, and summer patch (*Magnaporthe poae*). For optimum control of pink snow mold, fusarium patch, and summer patch, mix or rotate **Sharda Trifloxystrobin 50% WDG** with Triadimefon. When conditions are favorable for Pythium blight (*Pythium* spp.), mix a Pythium control fungicide with **Sharda Trifloxystrobin 50% WDG**.

Sharda Trifloxystrobin 50% WDG will provide suppression of dollar spot (*Sclerotinia homoeocarpa*) and will not flare this disease.

However, during periods of dollar spot pressure, mix **Sharda Trifloxystrobin 50% WDG** with Triadimefon or other dollar spot control fungicide.

1. STROBILURIN FUNGICIDES SUCH AS THIS PRODUCT ARE BEST USED IN A PREVENTIVE DISEASE CONTROL PROGRAM. CURATIVE APPLICATIONS CAN BE MADE FOR CERTAIN DISEASES.
2. Apply in 1 - 2 gals. of water per 1,000 sq. ft. to ensure thorough coverage. For soil-borne diseases, use sufficient water to reach the crown and upper root zone.
3. Apply after mowing OR allow sprayed area to completely dry before mowing.
4. For control of both foliar and soil-borne diseases, allow sprayed area to completely dry before irrigation.
5. Under conditions optimum for high disease pressure, use the higher rate and the shorter interval.
6. For optimum turf quality and disease control, use **Sharda Trifloxystrobin 50% WDG** in conjunction with turf management practices that promote good health and optimum disease control.
7. Before use of any fungicide, proper diagnosis of the organism causing the disease is important. Use of diagnostic kits or other means of identification of the disease organism is essential to determine the best control measures.

Tank Mixes

For longer and more broad-spectrum control including dollar spot, **Sharda Trifloxystrobin 50% WDG** can be tank mixed with other fungicides. Check compatibility before tank mixing.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

SHARDA TRIFLOXYSTROBIN 50% WDG CAN BE APPLIED TO ALL MAJOR TURFGRASS SPECIES. RATES ARE DEPENDENT UPON DISEASE AND THE ORGANISM THAT IT IS INCITED BY.

Resistance Management for Turfgrass

DO NOT apply more than 2 sequential **Sharda Trifloxystrobin 50% WDG** applications for gray leaf spot control. **DO NOT** apply more than 3 sequential applications of **Sharda Trifloxystrobin 50% WDG** for all other diseases. Alternate with a fungicide having a different mode of action.

Turfgrass Diseases Controlled with Sharda Trifloxystrobin 50% WDG			
Disease	Fungicide(s)	Rate of Product per 1,000 Sq. Ft.	Application Interval/Timing
Brown Patch	Sharda Trifloxystrobin 50% WDG	0.1 - 0.2 oz. (0.003-0.006 lb. a.i.)	14 days
		Apply when conditions are favorable for disease development.	
		0.15 - 0.25 oz. (0.004-0.007 lb. a.i.)	21 days
		For curative applications, apply 0.2 oz. of Sharda Trifloxystrobin 50% WDG and repeat applications on a 21-day interval.	
	Sharda Trifloxystrobin 50% WDG + Triadimefon	0.1 - 0.15 oz. (0.003-0.004 lb. a.i.) + Refer to product label.	14 - 21 days
		For longer and more broad-spectrum disease control including dollar spot, mix Sharda Trifloxystrobin 50% WDG with Triadimefon.	
Leaf Spot	Sharda Trifloxystrobin 50% WDG	0.1 - 0.15 oz. (0.003-0.004 lb. a.i.)	14 days
		Apply when conditions are favorable for disease development.	
		0.15 - 0.25 oz. (0.004-0.007 lb. a.i.)	21 - 28 days
		For curative applications, apply Sharda Trifloxystrobin 50% WDG at 0.15 - 0.25 oz. and repeat applications according to the application interval timing.	
Anthracnose	Sharda Trifloxystrobin 50% WDG	0.15 - 0.2 oz. (0.004-0.006 lb. a.i.)	14 days
		0.25 oz. (0.007 lb. a.i.)	21 days
		Apply when conditions are favorable for disease development.	
	Sharda Trifloxystrobin 50% WDG + Triadimefon	0.1 - 0.15 oz. (0.003-0.004 lb. a.i.) + Refer to triadimefon product label for use information.	14 - 21 days
		For longer and more broad-spectrum disease control including dollar spot, mix Sharda Trifloxystrobin 50% WDG with Triadimefon.	
Summer Patch	Sharda Trifloxystrobin 50% WDG and/or Triadimefon	0.2 - 0.25 oz. (0.006-0.007 lb. a.i.) and/or Refer to triadimefon product label for use information.	21 - 28 days
		Under heavy disease pressure, apply Triadimefon when the soil temperatures reach 65°F.	
		AFTER 21 - 28 days, apply either Sharda Trifloxystrobin 50% WDG at 0.2 oz. every 14 days OR a combination of Triadimefon + Sharda Trifloxystrobin 50% WDG at 0.2 - 0.25 oz. every 21 - 28 days.	
		DO NOT make more than 3 sequential applications of Sharda Trifloxystrobin 50% WDG before rotating back to Triadimefon.	
Gray Leaf Spot Rapid Blight	Sharda Trifloxystrobin 50% WDG	0.2 - 0.25 oz. (0.006-0.007 lb. a.i.) and/or Refer to triadimefon product label for use information.	21 - 28 days
		Under light disease pressure, make 2 - 3 applications of Sharda Trifloxystrobin 50% WDG at 0.2 - 0.25 oz. + Triadimefon every 21 - 28 days.	
	Sharda Trifloxystrobin 50% WDG	0.15 - 0.2 oz. (0.004-0.006 lb. a.i.)	14 days
		0.25 oz. (0.007 lb. a.i.)	21 days
	Sharda Trifloxystrobin 50% WDG + Triadimefon	0.1 - 0.2 oz. (0.003-0.006 lb. a.i.) +	14 - 21 days

	WDG + Triadimefon	Refer to triadimefon product label for use information.	
		Apply when conditions are favorable for disease development. For best results on Rapid Blight apply Sharda Trifloxystrobin 50% WDG at the highest rate.	
		For longer and more broad-spectrum disease control including dollar spot, mix Sharda Trifloxystrobin 50% WDG with Triadimefon.	
Red Thread, Pink Patch	Sharda Trifloxystrobin 50% WDG	0.1 - 0.15 oz. (0.003-0.004 lb. a.i.)	14 days
		Apply when conditions are favorable for disease development.	
		0.2 - 0.25 oz. (0.006-0.007 lb. a.i.)	21 days
		For curative applications, apply Sharda Trifloxystrobin 50% WDG at 0.15 - 0.25 oz. and repeat applications according to the application interval timing.	
	Sharda Trifloxystrobin 50% WDG + Triadimefon	0.1 - 0.15 oz. (0.003-0.004 lb. a.i.) +	14 - 21 days
		Refer to triadimefon product label for use information.	
Rust	Sharda Trifloxystrobin 50% WDG	0.1 - 0.15 oz. (0.003-0.004 lb. a.i.)	14 days
		Apply when conditions are favorable for disease development.	
		0.2 - 0.25 oz. (0.006-0.007 lb. a.i.)	21 days
		For curative applications, apply Sharda Trifloxystrobin 50% WDG at 0.15 - 0.25 oz. and repeat applications according to the application interval timing.	
	Sharda Trifloxystrobin 50% WDG + Triadimefon	0.1 - 0.15 oz. (0.003-0.004 lb. a.i.) +	14 - 21 days
		Refer to triadimefon product label for use information.	
Fusarium Patch	Sharda Trifloxystrobin 50% WDG	0.2 - 0.25 oz. (0.006-0.007 lb. a.i.)	Fall-Early Spring
		Apply when conditions are favorable for disease development.	
	Sharda Trifloxystrobin 50% WDG + Triadimefon	0.15 - 0.25 oz. (0.004-0.007 lb. a.i.) +	Fall-Early Spring
		Refer to triadimefon product label for use information.	
		Under conditions of heavy disease pressure, tank mix with Triadimefon.	
Pink Snow Mold	Sharda Trifloxystrobin 50% WDG	0.25 oz. (0.007 lb. a.i.)	Late Fall
		Apply 1 application in late fall before snow cover or early spring after snow melts. DO NOT apply on top of snow.	
	Sharda Trifloxystrobin 50% WDG + Triadimefon	0.15 - 0.25 oz. (0.004-0.007 lb. a.i.) +	Late Fall
		Refer to triadimefon product label for use information.	
		Under conditions of heavy disease pressure, tank mix with Triadimefon.	

Conversion Table for Sharda Trifloxystrobin 50% WDG in Water-Soluble Packets

Oz. per 1,000 Sq. Ft.	Oz. per Acre	Each 8 oz. Soluble Packet Treats	Number of 8 Oz. Soluble Packets needed to Treat 20 Acres of Turfgrass
0.1 (0.003 lb. a.i.)	4.4 (0.137 lb. a.i.)	1.82 acres	11
0.15 (0.004 lb. a.i.)	6.5 (0.203 lb. a.i.)	1.22 acres	16
0.185 (0.005 lb. a.i.)	8 (0.25 lb. a.i.)	1 acre	20
0.2 (0.006 lb. a.i.)	8.7 (0.272 lb. a.i.)	0.92 acre	22
0.25 (0.007 lb. a.i.)	10.9 (0.340 lb. a.i.)	0.73 acre	27

ORNAMENTAL DISEASE CONTROL

Sharda Trifloxystrobin 50% WDG is a broad-spectrum fungicide for the control of certain foliar, stem, and root diseases of ornamentals grown in interiorscapes, field nursery plantings, forest nurseries, residential and commercial landscapes, greenhouses, lath and shadehouses, containers, and other enclosed structures.

Foliar Diseases: **Sharda Trifloxystrobin 50% WDG** will control foliar diseases of ornamentals when applied as a foliar spray. Apply **Sharda Trifloxystrobin 50% WDG** at 1 - 4 oz. (0.031-0.125 lb. a.i.) per 100 gals. to the point of drip and repeat at 7- to 14-day intervals until the threat of disease is over. Start applications when conditions are favorable for disease development and continue until the threat of disease is over.

Damping off of New Seedlings: **Sharda Trifloxystrobin 50% WDG** will control damping off of new seedlings caused by *Rhizoctonia solani* when applied as a drench to seedlings and transplants. Drench the growth media at a rate of 0.5 oz. (0.015 lb. a.i.) per 100 gals.

Repeat every 21 - 28 days. If *Pythium* spp. are also present, mix **Sharda Trifloxystrobin 50% WDG** with a Pythium control fungicide.

The plants that **Sharda Trifloxystrobin 50% WDG** has been tested on, diseases that are controlled, and specific directions for use are listed in **Tables 1, 2, and 3**. Refer to **Table 1** for information on ornamentals and diseases that have been evaluated, **Table 2** for specific pathogens controlled, and **Table 3** for specific guidelines on the rates and timing of application.

NOTICE TO USER: Plant tolerance to **Sharda Trifloxystrobin 50% WDG** has been found to be acceptable on all ornamentals that it has been tested on. Due to the large number of species and varieties of ornamentals and nursery plants, it is impossible to test every one for tolerance to **Sharda Trifloxystrobin 50% WDG**. Neither the Manufacturer nor the Seller has determined whether or not **Sharda Trifloxystrobin 50% WDG** can be used safely on ornamental plants not specified on this label. The professional user must determine if **Sharda Trifloxystrobin 50% WDG** can be used safely prior to commercial use. In a small area test the prescribed rates on a small number of plants for phytotoxicity prior to widespread use.

Before using **Sharda Trifloxystrobin 50% WDG** in tank mixture with other products, test the mixture on a small number of plants for phytotoxicity prior to widespread use.

Before using **Sharda Trifloxystrobin 50% WDG** on plants for diseases that are not listed in the Directions for Use, test **Sharda Trifloxystrobin 50% WDG** on a small scale first.

Resistance Management for Ornamentals

Sharda Trifloxystrobin 50% WDG is a site-specific fungicide belonging to the strobilurin class of chemistry. Fungal pathogens are known to develop resistance to fungicides with a specific mode of action. When site-specific fungicides are introduced without a clear resistance management strategy, resistance development may be rapid, particularly with greenhouse use.

Sharda Trifloxystrobin 50% WDG exhibits cross-resistance to other strobilurins and fungicides within the Strobilurin Type Action and Resistance group (STAR compounds), but there is no known cross-resistance to fungicides of other classes including sterol inhibitors, dicarboximides, benzimidazoles, anilinopyrimidines, phenylpyrroles, or phenylamides.

Many fungi which attack ornamentals and flowering plants including Botrytis and powdery mildews have a history of fungicide resistance development. Because resistance development cannot be predicted, implementation of suitable strategies to manage the resistance risk to **Sharda Trifloxystrobin 50% WDG** is needed. To minimize the risk of resistance development to **Sharda Trifloxystrobin 50% WDG**, the following practices are advised:

1. Use **Sharda Trifloxystrobin 50% WDG** preventively.
2. For Leaf Spots and diseases other than Powdery Mildew, Downy Mildew, and Botrytis:
 - a) Use no more than 2 applications of **Sharda Trifloxystrobin 50% WDG** before rotating to another effective product that is not in the strobilurin class of chemistry for 2 applications before rotating back to **Sharda Trifloxystrobin 50% WDG**.
-OR-
 - b) Rotate to another fungicide of nonstrobilurin chemistry after each **Sharda Trifloxystrobin 50% WDG** application.
3. For Powdery Mildew, Downy Mildew, and Botrytis:
 - a) Between each **Sharda Trifloxystrobin 50% WDG** application, make 2 applications of a fungicide of nonstrobilurin chemistry before rotating back to **Sharda Trifloxystrobin 50% WDG**.
-OR-
 - b) Rotate to another fungicide of nonstrobilurin chemistry after each **Sharda Trifloxystrobin 50% WDG** application.
4. Make no more than 4 foliar applications of **Sharda Trifloxystrobin 50% WDG** per crop cycle or season for each at risk pathogen. Soil applications are independent of this limit.
5. **DO NOT** use **Sharda Trifloxystrobin 50% WDG** for disease control in vegetables grown in greenhouses for crop production or in vegetable production of transplants for outdoor use.

Restrictions:

- For ground application, use a minimum of 50 gals. per acre.
- To avoid spray drift, **DO NOT** apply when conditions favor drift beyond the target area. Avoid spray overlap.
- For information on spray equipment and calibration, consult sprayer manufacturers and State recommendations. For specific local directions and spray schedules, consult the current State Agricultural Experiment Station recommendations.
- Use of spray additives are not required. Evaluate any spray additive prior to use. **DO NOT** use in conjunction with organosilicate-based products, or plant injury may occur. Label directions are based on data with no additives.

Maximum Use Rates in Ornamentals:

- For plants grown in outdoor nurseries, outdoor seedbeds, field plantings, and landscapes, up to 34.5 oz. (1.07 lb. a.i.) of **Sharda Trifloxystrobin 50% WDG** per acre of production or acre of landscape per year or crop cycle can be used.
- For seedlings and plants grown in greenhouses, containers, and other enclosed structures, up to 120 oz. (3.75 lb. a.i.) of **Sharda Trifloxystrobin 50% WDG** per acre per year or crop cycle can be used. In California only, **DO NOT** apply more than 34.5 oz. (1.07 lb. a.i.) of **Sharda Trifloxystrobin 50% WDG** per acre per year or crop cycle to seedlings and plants grown in greenhouses,

containers, and other enclosed structures.

- For foliar applications, **DO NOT** apply more than 8 oz. (0.25 lb. a.i.) of **Sharda Trifloxystrobin 50% WDG** per acre per application.

Table 1

Sharda Trifloxystrobin 50% WDG has been tested for phytotoxicity and been found safe to the following plants*. The numbers in () indicate the diseases listed in Table 2. For plants not listed and for use of Sharda Trifloxystrobin 50% WDG in tank mixtures, refer to the NOTICE TO USER section.

Ajuga (2,11,12)	Dogwood (<i>Cornus</i> spp.) (2,11)	Ligustrum (1,2,8,12)
Aloe Vera (12)	Dracaena (8)	Lilac (5,11,12)
Alyssum (7,12)	Dusty Miller (1,12)	Liriope (12)
Apple, Non-Bearing** (11,13,14, 17)	Dwarf Ivy (12)	Lupines (2,5,11,12)
Aptenia (12)	Euonymus (2,11)	Marigold (1,5,11,12)
Aster (11,12,13)	Exacum (5)	Mint (11,13)
Azalea (2,5,11,12,13)	Ficus (2,12)	Monarda (Bee Balm) (5,11)
Azalea (Rhododendron) (2,10,11,12,13)	Geranium (1,5,7,11,13)	Moonflower (12)
Bamboo (12,14)	Gerbera (Caution on open flower) (2,5,11)	Nandina (2,11)
Barberry, Japanese (4)	Grasses: (2,8,11,13)	Nectarine, Non-Bearing** (8,11,14,17)
Begonia (5,11,12)	<i>Avena</i> : Blue Grass	Pansy* (1,2,5,7,8,11,12)
Bellis (1,2,5)	<i>Festuca glauca</i> : Blue Fescue	Peach, Non-Bearing** (5,12, 17)
Betula (11,13)	<i>Festuca glauca</i> : Sea Urchin	Penstemon (2,8,11)
Blue Daze (12)	<i>Festuca glauca</i> : Elijah Blue	Petunia (5,12)
Bottle Brush (14)	<i>Lagurus</i> : Rabbit Tail	Phlox (5,7,11)
Brachycome (12)	Pampas Grass: Pink	Photinia (4,8,11)
Buddleia (Butterfly Bush) (8)	Pampas Grass: White	Pittosporum (1,8,12)
Camellia (2,5,12)	<i>Pennisetum rubrum</i> (Crimson Fountain Grass)	Plum, Non-Bearing** (5,11,14)
Campanula (Bell Flower) (11,12,13)	<i>Phalaris picta</i> (Varigated Ribbon Grass)	Poinsettia (1,5,11,12,14)
Caladium (12)	Silver Banner Grass	Poppy (5)
Cast Iron Plant (12)	Hawthorn (5,8,14)	Primula (Primrose) (5)
Catnip (5,8,12)	Hawthorn, Indian (8,12)	Prunus (2,5)
Cedar (8,13)	Heather, Mexican (12)	Rabbit's Foot Fern*** (5,12)
Celosia (5,12)	<i>Hedera</i> spp. (2,5,10,12)	Ranunculus (7,11)
Cherry, Non-Bearing** (11,14,17)	Hen and Chickens, Flowering (12)	Photinia (Red Tip) (8)
Chrysanthemum (5,8,12,13)	Hibiscus (2,5,10,12)	Rose (3,5,6,7,11,12,13,14)
Citrus, Non-Bearing** (11)	Holly (Ilex) (4)	Pothos (9,10,12)
Coleus (5,12)	Hosta (5,12)	Salvia (1,5,7,11,12,13)
Coontie Palm (12)	Hydrangea (11)	Snapdragon (2,5,7,8,11,12,13)
Coreopsis (1,5,7,11,12)	Hypericum (13)	Spathiphyllum (6,9,10)
Cosmos (11,12)	Hypoestes (12)	Spirea (11)
Crabapple, Non-Bearing** (8,11,13,14)	Impatiens (1,7,12)	Syngonium (9,12)
Daisy (12)	Iris (Dwarf, Japanese, Siberian) (8,12)	Verbena (5,7,11,12)
Dianthus (1,5,8,12,13)	Jasmine (2,12)	Veronica (11)
Day Lily (8,12,13)	<i>Juniperus tortuosum</i> (12)	<i>Viburnum</i> spp. (2,8,12)
Delphinium (5,8,11,12)	Lantana (12,13)	Vinca (Catharanthus) (1,4,5,10,12)
<i>Dieffenbachia</i> spp. (9,10,12)	Lagerstroemia (Crape Myrtle) (11)	<i>Vinca Minor</i> (2,10)
Digitalis (Foxglove) (7)		Zinnia (1,11)

***DO NOT** drench pansy

****DO NOT** apply **Sharda Trifloxystrobin 50% WDG** to fruit trees that will bear harvestable fruit within 12 months of the last application.

*****DO NOT** use **Sharda Trifloxystrobin 50% WDG** on leatherleaf fern.

Table 2 - Diseases Controlled by Sharda Trifloxystrobin 50% WDG

Common Name	Scientific Name
1. Alternaria (B)	<i>Alternaria</i> spp.
2. Anthracnose (B)	<i>Colletotrichum</i> spp.
3. Black Root (D)	<i>Thielaviopsis</i> spp.
4. Black Spot (B)	<i>Diplocarpon rosae</i>
5. Botrytis (B)	<i>Botrytis</i> spp.
6. Cythrodcladium (D)	<i>Cylindrocladium</i> spp.
7. Downy Mildew (A)	<i>Peronospora</i> spp.
8. Leaf Spot (B)	<i>Septoria</i> spp.
9. Myrothecium (A)	<i>Myrothecium</i> spp.
10. Phytophthora Aerial (A) Phytophthora Root (D)	<i>Phytophthora parasitica</i>

11. Powdery Mildew (A)	<i>Erysiphe</i> spp., <i>Microsphaera</i> spp., <i>Oidium</i> spp., <i>Podosphaera</i> spp., <i>Sphaerotheca</i> spp.
12 Rhizoctonia Stem/Root Rot	<i>Rhizoctonia solani</i>
13. Rust (B)	<i>Gymnosporangium</i> spp., <i>Phragmidium</i> spp., <i>Puccinia</i> spp., <i>Uromyces</i> spp.
14. Scab (B)	<i>Cladosporium</i> spp., <i>Sphaceloma</i> , <i>Venturia inaequalis</i>
15. Myrothecium (A)	<i>Myrothecium</i> spp.
16. Rhizoctonia Root Rot (C)	<i>Rhizoctonia solani</i>
17. Blossom Blight (E)	<i>Monilinia</i> spp.

Table 3 - Specific Use Directions for Selected Pathogens

- A. Apply **Sharda Trifloxystrobin 50% WDG** as a foliar spray at 1 - 2 oz. (0.031-0.062 lb. a.i.) per 100 gals. to the point of drip before disease is detected or when conditions are favorable for disease development. Continue at 7- to 14-day intervals until the threat of disease is over.
- B. Apply **Sharda Trifloxystrobin 50% WDG** as a foliar spray at 2 - 4 oz. (0.062-0.125 lb. a.i.) per 100 gals. to the point of drip before disease is detected or when conditions are favorable for disease development. Continue at 7- to 14-day intervals until the threat of disease is over. Under heavy pressure, use the highest rate and the shortest interval. Under light disease pressure, the application interval may be extended.
- C. Apply 0.5 oz. (0.015 lb. a.i.) per 100 gals. as a drench to wet the upper half of the growing media. Start the application at the time of seeding, again at transplanting and at 21- to 28-day intervals thereafter.
- D. Apply 1 - 2 oz. (0.031-0.062 lb. a.i.) product per 100 gals. as a drench to wet the upper half of the growing media. Start the application at the time of planting and at 14 - 28 days depending on disease pressure.
- E. Apply 2 - 3 oz. (0.062-0.094 lb. a.i.) per 100 gals. starting at bud break on non-bearing stone and pome fruit listed in **Table 1**. If conditions are favorable for disease development, apply again at full bloom and at petal fall or on a 14- to 21-day spray schedule.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container and keep tightly closed when not in use. Store in a cool dry place. Avoid cross-contamination with other pesticides.

PESTICIDE DISPOSAL: Pesticides wastes are toxic. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of Federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

[[Nonrefillable Outer Bag:] Do not reuse or refill the outer bag. Completely empty bag into application equipment. Offer bag for recycling if available. If recycling is not available, then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.]

[[Nonrefillable Plastic Containers (Capacity Equal to or Less Than 50 Pounds):] Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by State and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.]

[[Nonrefillable Plastic Containers (Capacity Greater Than 50 Pounds):] Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by State and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.]

[[Non-Seed Treatment Products in Non-Refillable Foil outer pouches of Water-Soluble Packets (WSP):] Offer foil pouch for recycling if available or dispose of empty pouch in the trash as long as WSP is unbroken.]

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Sharda USA LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Sharda USA LLC and Seller harmless for any claims relating to such factors.

Sharda USA LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent consistent with applicable law, this warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or Sharda USA LLC and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, SHARDA USA LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Sharda USA LLC nor Seller shall be liable for any incidental, consequential, or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SHARDA USA LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SHARDA USA LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Sharda USA LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Sharda USA LLC.

[All trademarks are the property of their respective owners.]

[OPTIONAL MARKETING LANGUAGE]

1	<div><div>[https://www.shardausa.com/]</div><div></div></div>
2	[Handle with Care]
3	[This side Up]
4	<div><div>{The below graphic to be added to box if formulated in the United States}</div><div></div></div>