



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505T)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

83529-409

Date of Issuance:

9/17/26

NOTICE OF PESTICIDE:

☒ Registration
☐ Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Sharda Glyphosate 30% +
Oxyfluorfen 3.75% SC

Name and Address of Registrant (include ZIP Code):

Sharda USA LLC
c/o Syntech Research, Inc.
P.O. Box 640
Hockessin, DE 19707

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. Submit one copy of the final printed label for the record before you release the product for shipment.

Continues page 2

Signature of Approving Official:

Emily Schmid

Emily Schmid, Product Manager 25
Herbicide Branch, Registration Division (7505T)

Date:

9/17/26

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

Please note that the alternate brand name, "PAMIRA SC" has been added to the product record.

The record for this product currently contains the following CSF:

- Basic CSF dated 04/24/2026

If you have any questions, please contact Jamie Millard at (202) 566-2726 or by email at millard.jamie@epa.gov.

Enclosure

{MASTER LABEL}

GLYPHOSATE	GROUP	9	HERBICIDE
OXYFLUORFEN	GROUP	14	HERBICIDE

Sharda Glyphosate 30% + Oxyfluorfen 3.75% SC

FOR CONTROL OR SUPPRESSION OF WEEDS IN DORMANT TREE AND VINE CROPS, FALLOW SYSTEMS, AND
COTTON (INCLUDING ROUNDUP® READY COTTON VARIETIES)

ACTIVE INGREDIENTS:	WT. BY %
Glyphosate; N-(phosphonomethyl)glycine, in the form of its isopropylamine salt	30.00%
Oxyfluorfen; 2-chloro-1-(3-ethoxy-4-nitrophenoxy)-4-(trifluoromethyl)benzene	3.75%
OTHER INGREDIENTS:	66.25%
TOTAL:	100.0%

*Contains 443 grams per liter or 3.7 lbs. per U.S. gallon of the active ingredient glyphosate, in the form of its isopropyl-amine salt, equivalent to 328.5 grams per liter or 2.74 lbs. per U.S. gallon of the acid, glyphosate and 41 grams per liter or 0.34 lb. per U.S. gallon of the active ingredient, oxyfluorfen.

KEEP OUT OF REACH OF CHILDREN CAUTION / PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID	
IF ON SKIN OR CLOTHING:	- Take off contaminated clothing. - Rinse skin immediately with plenty of water for 15 - 20 minutes. - Call a poison control center or doctor for treatment advice.
IF IN EYES:	- Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. - Call a poison control center or doctor for treatment advice.
HOTLINE NUMBERS	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call your poison control center at 1-800-222-1222 . For general information about this product, contact the National Pesticides Information Center (NPIC) at 1-800-858-7378 , Monday through Friday, 8 AM to 12 PM PST, or at http://npic.orst.edu .	
NOTE TO PHYSICIAN	
Domestic Animals: This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.	

{Optional referral statements when booklets and container labels are used:}

[See label booklet for [additional OR complete] [{include applicable section(s) of} First Aid, Precautionary Statements, Directions For Use, and/or Storage and Disposal.]

EPA Reg. No.: 83529-409

Manufactured For [By]:

Manufactured for:
Sharda USA LLC 
7217 Lancaster Pike, Suite A
Hockessin, Delaware 19707

EPA Est. No.: XXXXX-XX-XXX

Net Contents: _____ [Gals./L/OZ.]

ACCEPTED

9/17/2026

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No. 83529-409

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION / PRECAUCIÓN

Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash

contaminated clothing before reuse. Wear protective clothing specified in the **Personal Protective Equipment (PPE)** section below.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber ≥ 14 mils, polyethylene, polyvinyl chloride (PVC) ≥ 14 mils, or Viton ≥ 14 mils

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides 40 CFR 170.607(d-f), the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment wash waters.

This product is highly toxic to aquatic invertebrates, aquatic plants, wildlife, and fish. Use with care when applying in areas frequented by wildlife or adjacent to any body of water or wetland area. **DO NOT** apply when weather conditions favor drift or erosion from target areas. Runoff may be hazardous to aquatic organisms in neighboring areas.

NON-TARGET ORGANISM ADVISORY: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

ENDANGERED AND THREATENED SPECIES PROTECTION REQUIREMENTS: To determine if your county or parish has a Bulletin, and to obtain that Bulletin, consult <https://www.epa.gov/endangered-species/endangered-species-protection-bulletins>, or call 1-844-447-3813 no more than 6 months before using this product. Applicators must use Bulletins that are in effect in the month in which the pesticide will be applied. New Bulletins will generally be available from the above sources 6 months before their effective dates.

PHYSICAL AND CHEMICAL HAZARDS

Spray solutions of this product should be mixed, stored, and applied only in stainless steel, aluminum, fiberglass, plastic, and plastic-lined steel containers.

DO NOT MIX, STORE, OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.

This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette, or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the **restricted-entry interval (REI) of 24 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves
- Chemical-resistant gloves made of barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, natural rubber ≥14 mils, polyethylene, polyvinyl chloride (PVC) ≥14 mils, or Viton ≥14 mils

PRODUCT INFORMATION

This product is a postemergent, systemic herbicide for control or suppression of emerged annual broadleaf weeds and grasses in fallow systems and dormant tree and vine crops. This product is also for use as a post directed/hooded sprayer application for broadleaf weed control and suppression in cotton and Roundup Ready cotton. It is formulated as a water-based suspension concentrate. It may be applied through most standard industrial or field type sprayers after dilution and through mixing with water or other carriers according to label instructions.

How this product works: This product moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days, but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay development of visual symptoms. Visible effects are a gradual wilting and yellowing of the plant which advances to complete browning of above-ground growth and deterioration of underground plant parts.

Stage of Weeds: Annual weeds are easiest to control when they are small. Refer to the “**Annual Weeds Rate Table**” for recommendations for specific weeds.

Always use the higher rate of this product per acre within the recommended range when weed growth is heavy or dense or weeds are growing in an undisturbed (non-cultivated) area.

DO NOT treat weeds under poor growing conditions such as drought stress, disease, or insect damage, as reduced weed control may result. Reduced results may also occur when treating weeds heavily covered with dust.

Cultural Considerations: Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed, or cut and have not been allowed to regrow to the recommended stage for treatment.

Rainfastness: Heavy rainfall soon after application may wash this product off of the foliage and a repeat application may be required for adequate control.

Spray Coverage: For best results, spray coverage should be uniform and complete. **DO NOT** spray weed foliage to the point of runoff.

Tank Mixing: At recommended use rates, this product does not provide residual weed control. For subsequent residual weed control, follow a label-approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used. Use according to the most restrictive label directions for each product in the mixture. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions, limitations, and directions for use on all product labels involved in the tank mixture. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

To the extent consistent with applicable law buyer and all users are responsible for loss or damage in connection with the use or handling of mixtures of the product with herbicides, or other materials that are not expressly recommended in this labeling. Mixing this product with herbicides or other materials not recommended on this label may result in reduced performance.

Annual Maximum Use Rate: Except as otherwise specified in a crop section of this label, the combined total of all treatments must not exceed 8.76 quarts of this product per acre per year.

APPLICATION RESTRICTIONS

Restrictions:

- **DO NOT** use any plants treated with **Sharda Glyphosate 30% + Oxyfluorfen 3.75% SC** for feed or forage.
- **DO NOT** feed or allow animals to graze on any areas treated with **Sharda Glyphosate 30% + Oxyfluorfen 3.75% SC**.
- Avoid direct application to any body of water.
- **DO NOT** apply this product through any type of irrigation or chemigation system.
- **DO NOT** contaminate irrigation water or water used for domestic purposes.
- **DO NOT** apply during rain.

Rotation Crop Restrictions:

- **DO NOT** rotate to small-grain crops (includes barley, buckwheat, corn, oats, pearl millet, proso millet, popcorn, rice, rye, sorghum, triticale, wheat, wild rice) within 10 months following a **Sharda Glyphosate 30% + Oxyfluorfen 3.75% SC** treatment.
- **DO NOT** direct seed any crop, other than **Sharda Glyphosate 30% + Oxyfluorfen 3.75% SC** labeled crops, within 60 days following the **Sharda Glyphosate 30% + Oxyfluorfen 3.75% SC** treatment.
- **DO NOT** transplant seedling crops, other than **Sharda Glyphosate 30% + Oxyfluorfen 3.75% SC** labeled crops, within 30 days following the **Sharda Glyphosate 30% + Oxyfluorfen 3.75% SC** treatment.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS (EXCEPT AS SPECIFIED FOR INDIVIDUAL ROUNDUP READY CROPS), DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

Precaution: Use of this product in any manner not consistent with this label *may* result in injury to persons, animals or crops, or other unintended consequences.

MIXING

Clean sprayer parts immediately after using this product by thoroughly flushing with water.

NOTE: REDUCED RESULTS MAY OCCUR IF WATER CONTAINING SOIL IS USED, SUCH AS VISIBLY MUDDY WATER OR WATER FROM PONDS AND DITCHES THAT IS NOT CLEAR.

SHAKE PRODUCT CONTAINER WELL BEFORE USE.

Mixing with Water: This product mixes readily with water. Mix spray solutions of this product as follows:

Fill the mixing or spray tank with the required amount of water. Add the recommended amount of this product near the end of the filling process and mix well. Use caution to avoid siphoning back into the carrier source. Use approved anti-back-siphoning devices where required by state or local regulations. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, avoid the use of mechanical agitators, terminate by-pass and return lines at the bottom of the tank and, if needed, use an approved anti-foam or defoaming agent.

Tank Mixing Procedure: Mix labeled tank mixtures of this product with water as follows:

1. Place a 20 to 35 mesh screen or wetting basket over filling port.
2. Through the screen, fill the spray tank one-half full with water and start agitation.
3. If a wettable powder is used, make a slurry with the water carrier, and add it SLOWLY through the screen into the tank. Continue agitation.
4. If a flowable formulation is used, premix one part flowable with one part water. Add diluted mixture SLOWLY through the screen into the tank. Continue agitation.
5. If an emulsifiable concentrate formulation is used, premix one part emulsifiable concentrate with two parts water. Add diluted mixture slowly through the screen into the tank. Continue agitation.
6. Continue filling the spray tank with water and add the required amount of this product near the end of the filling process.
7. Add individual formulations to the spray tank as follows: wettable powder, flowable, emulsifiable concentrate, drift control additive and water soluble liquid.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to re-suspend the mixture before spraying is resumed.

Keep by-pass line on or near the bottom of the tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50 mesh.

Always predetermine the compatibility of labeled tank mixtures of this product with water carrier by mixing small proportional quantities in advance.

Refer to the "Tank Mixing" section of for additional precautions.

Mixing for Hand-Held Sprayers: Prepare the desired volume of spray solution by mixing the amount of this product in water as shown in the following table:

Spray Solution

Amount of Sharda Glyphosate 30% + Oxyfluorfen 3.75% SC Herbicide

Desired Volume	1/2%	1%	1 1/2%	2%	5%	10%
1 Gal	2/3 oz	1 1/3 oz	2 oz	2 2/3 oz	6 1/2 oz	13 oz
25 Gal	1 pt	1 qt	1 1/2 qt	2 qt	5 qt	10 qt

100 Gal	2 qt	1 gal	1 1/2 gal	2 gal	5 gal	10 gal
---------	------	-------	-----------	-------	-------	--------

2 tablespoons = 1 fluid ounce

For use in knapsack sprayers, it is suggested that the recommended amount of this product be mixed with water in a large container. Fill sprayer with the mixed solution.

Surfactants: Nonionic surfactants which are labeled for use with herbicides may be used. **DO NOT** reduce rates of this product when adding surfactant. When adding additional surfactant use 0.5 percent surfactant concentration (2 quarts per 100 gallons of spray solution) when using surfactants with at least 70 percent active ingredient or a 1 percent surfactant concentration (4 quarts per 100 gallons spray solution) for those surfactants containing less than 70 percent active ingredient. Read and carefully observe surfactant cautionary statements and other information appearing on the surfactant label.

Ammonium Sulfate: The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of this product, particularly when tank mixed with certain residual herbicides on annual and perennial weeds. The equivalent rate of ammonium sulfate in a liquid formulation may also be used. Ensure that ammonium sulfate is completely dissolved in the spray tank before adding herbicides. Thoroughly rinse the spray system with clean water after use to reduce corrosion.

Note: When using ammonium sulfate, apply this product at rates recommended in this label. Lower rates will result in reduced performance.

Colorants and Dyes: Agriculturally approved colorants or marking dyes may be added to this product. Colorants or dyes used in spray solutions of this product may reduce performance, especially at lower rates or dilutions. Use colorants or dyes according to the manufacturer's recommendations.

Drift Control Additives: Drift control additives may be used with all equipment types, except wiper applicators, sponge bars and CDA equipment. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

APPLICATION EQUIPMENT AND TECHNIQUES

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

DO NOT allow the herbicide solution to mist, drip, drift, or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of injury, occurring from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing, or when there are other meteorological conditions that favor spray drift. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) which are likely to drift. AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.

This product may be applied with the following application equipment:

Aerial: Fixed Wing and Helicopter.

Ground Broadcast Spray: Boom or boomless systems, pull type sprayer, floaters, pick-up sprayers, spray coupes, and other ground broadcast equipment.

Hand-Held and High-Volume Spray Equipment: Knapsack and backpack sprayers, pump-up pressure sprayers, handguns, hand wands, mist blowers*, lances and other handheld and motorized spray equipment used to direct the spray onto weed foliage.

*This product is not registered in California or Arizona for use in mist blowers.

Selective Equipment: Recirculating sprayers, shielded and hooded sprayers, wiper applicators, and sponge bars.

Controlled Droplet Applicator (CDA): Hand-held or boom-mounted applicators which produce a spray consisting of a narrow range of droplet sizes.

APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES.

Aerial Equipment

DO NOT APPLY THIS PRODUCT USING AERIAL SPRAY EQUIPMENT EXCEPT UNDER CONDITIONS AS SPECIFIED WITHIN THIS LABEL.

Use the recommended rates of this herbicide in 3 to 15 gallons of water per acre unless otherwise specified on this label. Unless otherwise specified, **DO NOT** exceed 2.19 quarts per acre (1.5 lbs.-Glyphosate, 0.2 lbs.-Oxyfluorfen/acre). Aerial applications of this product may be made in annual cropping conventional tillage systems, fallow and reduced tillage systems. Refer to the individual use area section of this label for recommended volumes, application rates and further instructions.

FOR AERIAL APPLICATION IN CALIFORNIA, REFER TO THE FEDERAL SUPPLEMENTAL LABEL FOR AERIAL APPLICATIONS IN THAT STATE FOR SPECIFIC INSTRUCTIONS, RESTRICTIONS, AND REQUIREMENTS. FOR AERIAL APPLICATIONS, CONSULT WITH STATE OR LOCAL AUTHORITIES REGARDING ANY ADDITIONAL REQUIREMENTS FOR AERIAL TREATMENTS.

Dicamba tank mixtures may not be applied by air in California.

Ensure uniform application: To avoid streaked, uneven, or overlapped application, use appropriate marking devices.

Aerial Spray Drift Management: The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops.

1. The distance of the outermost nozzles on the boom must not exceed $\frac{3}{4}$ the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they should be observed.

Importance of Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions section of this label).

Controlling Droplet Size

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is released backwards parallel to the airstream will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.
- **Boom Length** - For some use patterns, reducing the effective boom length to less than $\frac{3}{4}$ of the wingspan or rotor length may further reduce drift without reducing swath width.
- **Application Height** - Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

Wind

Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. **NOTE:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small, suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a connected cloud (under low wind conditions) indicates an inversion, while smoke that moves upwards and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

Equipment care and Maintenance: Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR IS MORE SUSCEPTIBLE. The maintenance of an organic coating (paint), which meets aerospace specification MIL-C-38413, may prevent corrosion.

Ground Broadcast Equipment

Use the recommended rates of this product in 3 to 40 gallons of water per acre as a broadcast spray unless otherwise specified. As density of weeds increases, spray volume should be increased within the recommended range to ensure complete coverage. Carefully select proper nozzles to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

Hand-Held and High Volume Equipment

Apply to foliage of vegetation to be controlled. For applications made on a spray-to-wet basis, spray coverage should be uniform and complete. **DO NOT** spray to the point of runoff. Use coarse sprays only.

For control of weeds listed in the Annual Weeds Rate Table, apply a 0.5 percent solution of this product to weeds less than 6" in height or runner length. Apply prior to seedhead formation in grass or bud formation in broadleaf weeds. For annual weeds over 6 inches tall, or unless otherwise specified, use a 1 percent solution.

For best results, use a 2 percent solution on harder to control perennials, such as Bermuda grass, and field bindweed.

When using application methods which result in less than complete coverage, use a 5 percent solution for annual and perennial weeds.

Selective Equipment

This product may be applied through recirculating spray systems, shielded applicators, hooded sprayers, wiper applicators or sponge bars after dilution and thorough mixing with water to listed weeds growing in any noncrop site specified on this label and only when specifically recommended in cropping systems.

A recirculating spray system directs the spray solution onto weeds growing above desirable vegetation while spray solution not intercepted by weeds is collected and returned to the spray tank for reuse.

A shielded or hooded applicator directs the herbicide solution onto weeds, while shielding desirable vegetation from the herbicide.

A wiper or sponge applicator applies the herbicide solution onto weeds by rubbing the weed with an absorbent material containing the herbicide solution.

AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION.

Contact of the herbicide solution with desirable vegetation may result in damage or destruction. Applicators used above desirable vegetation should be adjusted so that the lowest spray stream or wiper contact point is at least 2 inches above the desirable vegetation. Droplets, mist, foam, or splatter of the herbicide solution setting on desirable vegetation may result in discoloration, stunting or destruction.

Applications made above the crops should be made when the weeds are a minimum of 6 inches above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations or when the height of the weeds varies so that not all weeds are contacted. In these instances, repeat treatment may be necessary.

Shielded and Hooded Applicators: Use nozzles that provide uniform coverage within the treated area. Keep shields on these sprayers adjusted to protect desirable vegetation. EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE WITH DESIRABLE

VEGETATION.

A hooded sprayer is a type of shielded applicator. The spray pattern is completely enclosed on the top and all 4 sides by a hood, thereby shielding the crop from the spray solution. This equipment must be set up and operated in a manner that avoids bouncing or raising the hoods off the ground in any way. If the hoods are raised, spray particles may escape and come into contact with the crop, causing damage or destruction of the crop. The spray hoods must be operated on the ground or skimming across the ground. Tractor speed must be adjusted to avoid bouncing of the spray hoods. Avoid operation on rough or sloping ground where the spray hoods might be raised off the ground.

Wiper Applicators and Sponge Bars: Wiper applicators are devices that physically wipe appropriate amount of this product directly onto the weed.

Equipment must be designed, maintained and operated to prevent the herbicide solution from contacting desirable vegetation. Operate this equipment at ground speeds no greater than 5 mph. Performance may be improved by reducing speed in areas of heavy weed infestations to ensure adequate wiper saturation. Better results may be obtained if 2 applications are made in opposite directions.

Avoid leakage or dripping onto desirable vegetation. Adjust height of applicator to ensure adequate contact with weeds. Keep wiping surfaces clean. Be aware that, on sloping ground, the herbicide solution may migrate, causing dripping on the lower end and drying of the wicks on the upper end of a wiper applicator.

Restriction: DO NOT use wiper equipment when weeds are wet.

Mix only the amount of solution to be used during a 1-day period, as reduced activity may result from use of leftover solutions. Clean wiper parts immediately after using this product by thoroughly flushing with water.

For Rope or Sponge Wick Applicators: Mix 1 gallon of this product in 2 gallons of water to prepare a 33 percent solution. Apply this solution to weeds listed in this section.

For Porous-Plastic Applicators: Solutions ranging from 33 to 100 percent of this product in water may be used in porous-plastic wiper applicators.

CDA Equipment

The rate of this product applied per acre by vehicle mount CDA equipment must not be less than the amount recommended in this label when applied by conventional broadcast equipment. For vehicle-mounted CDA equipment, apply 3 to 15 gallons of water per acre.

For the control of annual weeds with hand-held CDA units, apply a 20 percent solution of this product at a flow rate of 2 fluid ounces per minute and a walking speed of 1.5 mph (1 quart per acre). For the control of perennial weeds, apply a 20 to 40 percent solution of this product at a flow rate of 2 fluid ounces per minute and a walking speed of 0.75 mph (2.19 to 4.38 quarts per acre).

Controlled droplet application equipment produces a spray pattern which is not easily visible. Extreme care must be exercised to avoid spray or drift contacting the foliage or any other green tissue of desirable vegetation, as damage or destruction may result.

CROPS

This section is organized alphabetically by crop category. There may be several labeled crops listed in a crop category.

Timing of Application: This product should be applied postemergence to vigorously growing weeds when they have reached the recommended size given in the "**Annual Weeds Rate Table**" or "**Perennial Weeds Rate Table**" of this label. Application should be delayed until maximum emergence of the target weeds, but before weeds exceed the maximum size recommended. For annual weeds, allow 1 day after treatment before tillage.

Reduced control may result if treatments are made during poor growing conditions such as drought stress, disease or insect damage or if weeds have been mowed, grazed or cut. Heavy dust on foliage or an overstory canopy covering targeted weeds may also reduce control.

Heavy rainfall or irrigation soon after application may wash this product off the foliage and a repeat treatment may be required for adequate control.

Citrus Crops (Nonbearing only)

Labeled Crops: Calamondin, Chironia, Citron, Citrus Hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Orange (All), Pummelo, Tangelo, Tangor

FOR USE ONLY IN PERMANENTLY ESTABLISHED GROVES IN ARIZONA AND CALIFORNIA.

NOTE: FOR GENERAL USE DIRECTIONS, SEE THE "**TREE AND VINE CROPS**" SECTION. THE FOLLOWING DIRECTIONS ARE SPECIFIC TO CITRUS CROPS.

TYPES OF APPLICATIONS: General weed control, middles (between rows of trees), strips (in row of trees), selective equipment. For rates, refer to the "**Annual Weeds Rate Table**". Single applications cannot exceed 4.38 quarts of **Sharda Glyphosate 30% lbs. + Oxyfluorfen 3.75% SC lbs.** per acre (3lbs. ai Glyphosate, 0.4 lbs. ai Oxyfluorfen/Acre). Repeat applications may be made up to a maximum of 8.76 quarts per acre per year (6lbs. ai Glyphosate, 0.8 lbs. ai Oxyfluorfen).

PRECAUTIONS: Applications may be made to newly planted trees or to young trees that will not bear fruit for one year.

RESTRICTIONS: **DO NOT** apply this product during periods of new foliage growth. Applications should be made after foliage has fully expanded and hardened off. Direct spray toward the base of trees. Avoid direct spray contact on the citrus foliage.

Cotton (Including Glyphosate-Resistant Cotton Varieties)

TYPES OF APPLICATIONS: Preplant, hooded sprayer, selective equipment, and precision post-directed equipment. PRECISION POST-DIRECTED EQUIPMENT IS PERMITTED ONLY IN COTTON WITH THE ROUND UP READY GENE.

PRECAUTION: IN-CROP APPLICATIONS OF THIS PRODUCT THROUGH PRECISION POST-DIRECTED EQUIPMENT DIRECTED ONTO COTTON IS RECOMMENDED ONLY ON IMPROVED COTTON VARIETIES THAT ARE DESIGNATED AS COTTON WITH THE ROUNDUP READY GENE. SEVERE INJURY OR DEATH OF COTTON WILL RESULT IF ANY COTTON VARIETIES NOT PROPERLY DESIGNATED AS HAVING THE ROUNDUP READY GENE ARE SPRAYED WITH THIS PRODUCT. AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, OR ANY DESIRABLE PLANTS AND TREES SINCE SEVERE INJURY OR DESTRUCTION WILL RESULT. GLYPHOSATE- RESISTANT COTTON VARIETIES MUST BE PURCHASED FROM AN AUTHORIZED LICENSED SEED SUPPLIER. THE DESIGNATION "GLYPHOSATE-RESISTANT Y" INDICATES THE COTTON VARIETY CONTAINS A PATENTED PROPRIETARY TRAIT.

Preplant: Use Instructions: This product may be applied before planting cotton. **DO NOT** apply **Sharda Glyphosate 30% + Oxyfluorfen 3.75% SC** within 7 days prior to planting. The fallow beds should be worked thoroughly to a depth of at least 2.5 inches prior to planting.

Selective Equipment: Use Instructions: This product may be applied through shielded or hooded sprayers which **DO NOT** allow contact of the spray with cotton. Applications may be made from 6-inch tall cotton through layby. CONTACT OF THE SPRAY SOLUTION WITH COTTON FOLIAGE, GREEN STEMS, OR FRUIT MAY RESULT IN SEVERE CROP INJURY OR DESTRUCTION.

GLYPHOSATE-RESISTANT COTTON: In addition to preplant and selective equipment applications, this product may be applied using precision post-directed or hooded sprayers to Glyphosate-Resistant cotton after the fifth leaf (node) stage of development through layby. Cotton must be at least 6-8 inches tall. Post-directed equipment that directs the spray to the base of the cotton plants should be used. Contact of the spray with cotton leaves should be avoided to the maximum extent possible. To minimize the spray onto the leaves of the cotton plants, place nozzles in a low position directing a horizontal spray pattern under the cotton leaves to contact in the row, and maintain low spray pressure (less than 30 PSI).

For best results, make applications while weeds are small (less than 3 inches). Any single post-directed application should not exceed 1 quart per acre of this product. No more than 2 applications should be made from the fifth leaf through layby. Sequential in-crop applications of this product must be at least 10 days apart and cotton must have at least two nodes of incremental growth between applications. **DO NOT** apply within 75 days of harvest.

PRECAUTION: USE OF SHARDA GLYPHOSATE 30% + OXYFLUORFEN 3.75% SC IN ACCORDANCE WITH LABEL DIRECTIONS IS EXPECTED TO RESULT IN NORMAL GROWTH OF GLYPHOSATE-RESISTANT COTTON. HOWEVER, VARIOUS ENVIRONMENTAL CONDITIONS, AGRONOMIC PRACTICES AND OTHER FACTORS MAKE IT IMPOSSIBLE TO ELIMINATE ALL RISKS ASSOCIATED WITH THIS PRODUCT, EVEN WHEN APPLICATIONS ARE MADE IN CONFORMANCE WITH THE LABEL SPECIFICATIONS. IN SOME CASES, THESE FACTORS CAN RESULT IN BOLL LOSS, DELAYED MATURITY, AND/OR YIELD LOSS.

NOTE: Total in-crop applications of herbicides containing glyphosate as an active ingredient, whether applied as a mixture or separately, may not exceed 3 pounds of glyphosate active ingredient from the cracking stage to layby in Glyphosate-Resistant cotton. When using this product as part of an overall weed control program that includes other herbicides containing glyphosate as an active ingredient, calculate the application rates and ensure that the total use of this and other glyphosate containing products does not exceed stated maximum use rates.

Fallow Systems

TYPES OF APPLICATION: Chemical fallow, preplant fallow beds.

NOT FOR USE ON FALLOW BEDS TO BE PLANTED TO SOYBEANS IN CALIFORNIA.

USE INSTRUCTIONS: This product may be applied during the fallow period prior to planting or emergence of any crop listed on this label. For any crop not listed on this label, applications must be made at least 60 days prior to direct seeding and 30 days prior to transplanting crops. For rates refer to the **"Annual Weeds Rate Table"**. Single applications cannot exceed 4.38 quarts of this product per acre (3 lbs. ai/acre-Glyphosate; 0.4 lbs. ai/acre-Oxyfluorfen). Repeat applications may be made up to a maximum of 8.76 quarts per acre (6 lbs. ai/acre-Glyphosate; 0.8 lbs. ai/acre-Oxyfluorfen) per year. This product may be used as a substitute for tillage to control annual weeds in fallow fields. Ground or aerial application equipment may be used.

Minimum Treatment-Planting Interval (DAYS)

DIRECT SEEDED CROPS	SHARDA GLYPHOSATE 30% + OXYFLUORFEN 3.75% SC USE RATE	
	Up to 4.38 quarts/acre (3lbs. ai Glyphosate, 0.4 lbs. ai Oxyfluorfen/Acre)	Up to 8.76 quarts/acre (6lbs. ai Glyphosate, 0.8 lbs. ai Oxyfluorfen/Acre)
Carrot	90 Days	90 Days
Potato	60 Days	60 Days
Sugarbeet	60 Days	90 Days
Other root/tuber crops	90 Days	90 Days
Onions	180 Days	180 Days
Other bulb vegetables	180 Days	180 Days
Cabbage	90 Days	90 Days
Cauliflower	90 Days	90 Days
Other brassica crops	120 Days	120 Days
Lettuce	90 Days	120 Days
Other leafy vegetables (except brassica crops)	120 Days	120 Days
Pepper	90 Days	120 Days
Tomato	60 Days	120 Days
Other fruiting vegetables	120 Days	120 Days
Cantaloupe	60 Days	90 Days
Squash	90 Days	120 Days
Watermelon	60 Days	60 Days
Other cucurbits	90 Days	120 Days
Dry beans	60 Days	60 Days
Peanut	60 Days	60 Days
Other legume vegetables	60 Days	60 Days
Safflower	60 Days	60 Days
Cereal grains (includes barley, buckwheat, com, proso millet, pearl millet, oats, popcorn, rice, rye, sorghum, triticale, wheat, wild rice)	10 Months	10 Months
Cotton and soybean	7 Days	7 Days

TRANSPLANTED CROPS	SHARDA GLYPHOSATE 30% + OXYFLUORFEN 3.75% SC USE RATE	
	Up to 4.38 Quarts/acre ((3lbs. ai Glyphosate, 0.4 lbs. ai Oxyfluorfen/Acre)	Up to 8.76 quarts/acre (6lbs. ai Glyphosate, 0.8 lbs. ai Oxyfluorfen/Acre)
Broccoli	0 Days	30 Days
Cabbage	0 Days	30 Days
Cauliflower	0 Days	30 Days
Celery	30 Days	30 Days
Conifer	0 Days	0 Days
Garlic	0 Days	30 Days
Grape/Kiwi	3 Days	3 Days
Onion	0 Days	30 Days
Pepper	30 Days	30 Days
Strawberries	30 Days	30 Days
Tomato	30 Days	30 Days
Treefruit/Nut/Citrus	3 Days	3 Days

IMPORTANT: The fallow beds should be worked thoroughly to a depth of at least 2.5 inches prior to planting. FAILURE TO ACHIEVE

THOROUGH AND COMPELTE INCORPORATED, OR TO FOLLOW THE RECOMMENDED TREATMENT PLANTING INTERVAL MAY RESULT IN STAND REDUCTION AND/OR VIGOR REDUCTION OF THE PLANTED CROP.

For best results, apply this product after most weed seeds have germinated but before seedhead formation in grasses or flower bud formation in broadleaves.

When applied as directed, this product will provide control or suppression of weeds listed in the annual and perennial weed tables.

Tree and Vine Crops (General)

TYPES OF APPLICATIONS: General weed control, middles (between rows of trees), strips (in row of trees), selective equipment (except kiwi) in DORMANT tree and vine crops.

NOTE: THIS SECTION GIVES GENERAL DIRECTIONS THAT APPLY TO ALL CITRUS CROPS (NON BEARING ONLY), TREE FRUITS, TREE NUTS AND VINE CROPS. SEE THE INDIVIDUAL CROP SECTION AND RESTRICTIONS FOR SPECIFIC CROPS.

This product may be applied in middles, strips, and for general weed control in established citrus groves, tree fruit and tree nut orchards, and vineyards. This product may also be used for site preparation and prior to transplanting these crops. For rates refer to the **"ANNUAL WEEDS RATE TABLE"**. Single applications cannot exceed 4.38 quarts of **Sharda Glyphosate 30% + Oxyfluorfen 3.75% SC** per acre. Repeat applications may be made up to a maximum of 8.76 quarts per acre per year. Allow a minimum of 3 days between application and transplanting. Applications may be made with boom equipment, CDA, shielded sprayers, handheld and high-volume wands, lances, orchard guns or with wiper applicator equipment, except as directed.

Middles (between rows): USE INSTRUCTIONS: This product will control or suppress annual weeds growing between the rows of labeled tree and vine crops. If weeds are under drought stress, irrigate prior to application. Reduced control may result if weeds have been mowed prior to application.

Strips (in rows): USE INSTRUCTIONS: This product may be applied in rows of tree or vine crops and may also be tank mixed with the following products:

- Bromacil
- Diuron
- Napropamide
- Norflurazon
- Oryzalin
- Propyzamide
- Simazine

Refer to the individual product labels for specific crops, rates, geographic restrictions and precautionary statements.

Selective equipment: Shielded and wiper applicators may be used in tree crops and grapes. Refer to the individual crop sections for time interval between application and harvest.

GENERAL PRECAUTIONS: **Sharda Glyphosate 30% + Oxyfluorfen 3.75% SC** or any of the combinations recommended on this label should be applied only to healthy growing trees and vines. Direct spray toward the base of tree or vines. Avoid direct plant contact.

GENERAL RESTRICTIONS:

For citron and olives, apply as a post-directed spray only.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE SOLUTION, SPRAY, DRIFT OR MIST WITH FOLIAGE OR GREEN BARK OF TRUNK, BRANCHES, SUCKERS, FRUIT OR OTHER PARTS OF TREES AND VINES. CONTACT OF THIS PRODUCT WITH OTHER THAN MATURED BROWN BARK CAN RESULT IN SERIOUS CROP DAMAGE.

AVOID PAINTING CUT STUMPS WITH THIS PRODUCT AS INJURY RESULTING FROM ROOT GRAFTING MAY OCCUR IN ADJACENT TREES.

THE FOLLOWING RESTRICTIONS APPLY ONLY TO TREE FRUITS, NUT AND VINE CROPS.

DO NOT apply **Sharda Glyphosate 30% + Oxyfluorfen 3.75% SC** during the period between bud swell and completion of final harvest or when fruit or nuts are present. **Sharda Glyphosate 30% + Oxyfluorfen 3.75% SC** can be applied upon completion of final harvest.

In Arizona and California: **Sharda Glyphosate 30% + Oxyfluorfen 3.75% SC** can be applied during the period following completion of final harvest up to February 15 (February 1st in Coachella Valley, CA). Applications made after the calendar dates above, but prior to bud swell, may result in significant crop injury and are the responsibility of the user.

Tree Fruits (Including Tropical)

LABELED CROPS: Apple, Apricot, Cherry (Sweet, Sour), Crabapple, Date, Fig, Loquat, Mayhaw, Nectarine, Olive, Peach, Pear, Persimmon, Plum/Prune (All), Pomegranates, Quince.

TYPES OF APPLICATIONS: General weed control, middles (between rows of trees), strips (in rows of trees), selective equipment. For rates refer to the **"Annual Weeds Rate Table"**. Single applications cannot exceed 4.38 quarts of **Sharda Glyphosate 30% + Oxyfluorfen 3.75% SC** per acre. Repeat applications may be made up to a maximum of 8.76 quarts per acre per year.

NOTE: FOR GENERAL USE DIRECTIONS, SEE THE **"TREE AND VINE CROP "** SECTION. THE FOLLOWING DIRECTIONS ARE SPECIFIC TO TREE FRUITS.

Restrictions on application equipment: For cherries, any application equipment listed in this section may be used in all states.

Any application equipment listed in this section may be used in apricots, nectarines, peaches and plums/prunes growing in Arizona, California, Oregon, and Washington.

Tree Nuts

LABELED CROPS: Almond, Beechnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (Hazelnut), Hickory nut, Macadamia, Pecan, Pistachio, Walnut (Black, English).

TYPES OF APPLICATIONS: General weed control, middles (between rows of trees), strips (in row of trees), selective equipment.

NOTE: FOR GENERAL USE DIRECTIONS, SEE THE **"TREE AND VINE CROPS "** SECTION FOLLOWING DIRECTIONS THAT ARE SPECIFIC TO TREE NUTS.

For rates refer to the **"Annual Weeds Rate Table"**. Single applications cannot exceed 4.38 quarts of **Sharda Glyphosate 30% + Oxyfluorfen 3.75% SC** per acre. Repeat applications may be made up to a maximum of 8.76 quarts per acre per year.

Vegetable Crops

LABELED CROPS: Broccoli (All), Cabbage (All), Cauliflower, Garlic, Horseradish*, Onion.

USE INSTRUCTIONS: This product may be applied up to 4.38 quarts/acre prior to transplanting these listed vegetables. For rates between 4.38 and 8.76 quarts/acre, **DO NOT** transplant within 30 days following the **Sharda Glyphosate 30% + Oxyfluorfen 3.75% SC** treatment.

*For horseradish, applications must be made after the horseradish roots have been planted and prior to plant emergence.

RESTRICTIONS: When applying this product prior to transplanting crops into plastic mulch, care must be taken to remove residues of this product which could cause crop injury from the plastic prior to transplanting. Residues can be removed by a single 0.5 inch application of water, either by natural rainfall or via a sprinkler system. Applications made at emergence will result in injury or death to emerged seedlings. **DO NOT** apply this product preemergence to direct-seeded broccoli, cabbage or cauliflower. **DO NOT** apply this product post transplant or postemergence to broccoli, cabbage or cauliflower. **DO NOT** apply this product in an enclosed greenhouse structure, as injury to plant foliage may result.

Vine Crops

LABELED CROPS: Grapes (raisin, table, wine, juice), Kiwi fruit.

TYPES OF APPLICATIONS: General weed control, middles (between rows), strips (in row), selective equipment (except kiwi).

NOTE: FOR GENERAL USE DIRECTIONS, SEE THE **"TREE AND VINE CROPS (GENERAL)"** SECTION. THE FOLLOWING DIRECTIONS ARE SPECIFIC TO VINE CROPS.

Direct spray toward the base of vines.

For rates refer to the **"Annual Weeds Rate Table"**. Single applications cannot exceed 4.38 quarts **Sharda Glyphosate 30% + Oxyfluorfen 3.75% SC** per acre. Repeat applications may be made up to a maximum of 8.76 quarts per acre per year.

PRECAUTIONS, RESTRICTIONS: Applications should not be made when green shoots, canes or foliage are in the spray zone. **DO NOT** apply to grapes or kiwi that are not staked or trellised unless vines are free-standing.

Annual Weeds Rate Table

Water carrier volumes of 3 to 10 gallons per acre for ground applications and 3 to 5 gallons per acre for aerial applications are recommended.

Apply to actively growing annual weeds. Annual weeds are easiest to control when they are small.

Always use the higher rate within the provided rate range when applications are made to larger weeds within the range. For weeds that have been mowed, grazed or cut, allow regrowth to occur prior to treatment.

For those rates less than 52.5 fluid ounces per acre, this product may be used up to 52.5 (1.12 lbs. ai/A-Glyphosate, 0.14 lbs. ai/A-Oxyfluorfen) fluid ounces per acre where heavy weed densities exist.

WEED SPECIES	RATE (FLUID OUNCES PER ACRE)	
	26.28-35 (0.56-0.75 lbs. ai/A)- Glyphosate (0.07-0.09 lbs. ai/A)- Oxyfluorfen	43.8-52.5 (0.94- 1.12 lbs. ai/A)- Glyphosate (0.12-0.14 lbs. ai/A)- Oxyfluorfen
	MAXIMUM HEIGHT/LENGTH	
Barnyardgrass	3" - 8"	9" - 12"
Bluegrass, annual	8" - 12"	- -
Buttercup, smallflower	12" - 24"	- -
Carolina geranium	2" - 4"	5" - 9"
Carpetweed	6" - 12"	- -
Cheeseweed (<i>Malva parviflora</i>)	3" - 6"	6" - 9"
Chickweed	12" - 24"	- -
Cocklebur	12" - 24"	- -
Crabgrass	6" - 24"	- -
Cutleaf evening primrose	2" - 3"	4" - 6"
Fiddleneck, Coast	4" - 6"	7" - 12"
Filaree, broadleaf	- - 3"	6" - 12"
Filaree, whitestem	- - 3"	6" - 12"
Filaree, redstem	- - 3"	6" - 12"
Fleabane, hairy (<i>Conyza bonariensis</i>)	- - 3"	6" - 9"
Fleabane, rough	3" - 12"	- -
Florida, pusley	2" - 4"	5" - 6"
Goosegrass	3" - 8"	9" - 18"
Groundcherry, cutleaf	6" - 12"	- -
Groundcherry, Wright	6" - 12"	- -
Groundsel, common	6" - 12"	- -
Hemp, sesbania	2" - 4"	5" - 8"
Henbit	4" - 6"	7" - 20"
Horseweed/Marestail (<i>Conyza Canadensis</i>)	6" - 12"	- -
Jimsonweed	4" - 6"	7" - 12"
Johnsongrass, seedling	12" - 24"	- -
Junglerice	3" - 8"	9" - 12"
Knotweed	3" - 12"	13" - 20"
Lambsquarters	6" - 12"	13" - 20"
London rocket	6" - 12"	- -
Morningglory (<i>Ipomoea</i> spp.)	2" - 4"	5" - 6"
Mustard, blue	6" - 18"	- -
Mustard, tansy	6" - 18"	- -
Mustard, tumble	6" - 18"	- -
Mustard, wild	6" - 18"	- -
Nightshade, black	6" - 18"	- -
Nightshade, hairy	6" - 18"	- -
Pigweed	12" - 24"	- -
Prickly lettuce	6" - 20"	- -
Purslane	4" - 6"	7" - 12"
Redmaids	6" - 12"	- -
Ryegrass, common	4" - 6"	7" - 18"
Shepherd's purse	6" - 12"	- -
Sicklepod	2" - 4"	5" - 8"
Smartweed, ladysthumb	4" - 8"	9" - 12"
Smartweed, Pennsylvania	4" - 8"	9" - 12"

Sowthistle, annual	4" - 6"	7" - 12"
Spurge, prostrate	6" - 20"	- -
Spurge, spotted	6" - 20"	- -
Teaweed/Prickly sida	1" - 3"	4" - 6"
Velvetleaf South	2" - 4"	5" - 8"
Velvetleaf North	3" - 12"	- -
Virginia pepperweed	12" - 18"	- -

Perennial Weeds Rate Table

Apply to actively growing perennial weeds.

RESTRICTION: If weeds have been mowed, grazed, cut or tilled, **DO NOT** treat until plants have resumed active growth and have reached the recommended stages.

Best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seed. Repeat treatments must be made prior to crop emergence.

Unless otherwise stated, allow 7 or more days after application before tillage. Best results are obtained when soil moisture is adequate for active weed growth.

WEED SPECIES	RATE (QT/A)	WATER VOLUME (GPA)	HAND-HELD % SOLUTION
Alfalfa	1.095 – 2.19	3 - 10	2%
After fall cutting and 6 to 8" regrowth, Deep till 7 days after application.			
Alligatorweed*	4.38	3 - 20	1.5%
Broom stage. Repeat applications will be required to maintain control.			
Anise (fennel)	-	-	1 – 2%
Bud to full-bloom stage.			
Bahiagrass	3.285 – 5.475	3 - 20	2%
Early head stage.			
Bentgrass*	1.642	10 - 20	2%
Resumed growth of crown area to at least 3" in height.			
Bermudagrass	3.285 – 5.475	3 - 20	2%
Actively growing and seedheads present.			
Bermudagrass, water (knotgrass)	1.095 - 1.64	5 - 10	2%
12 to 18" weeds, 7 days before tilling. This product is not registered in California for use on water bermudagrass.			
Bindweed, field	0.547 – 5.475	3 - 20	2%
At or beyond full bloom in late summer or fall. Also for control, apply 2 quarts of this product plus 0.5 pound a.i. of Dicamba in 10 to 20 gallons of water per acre.			
Restrictions: DO NOT apply by air. DO NOT treat when weeds are under drought stress.			
Bluegrass, Kentucky	2.19	3 - 30	2%
Boot to early seedhead stage.			
Blueweed, Texas	3.285 – 5.475	3 - 30	2%
At or beyond full bloom in late summer or fall.			
Brackenfern	3.285 – 4.38	3 - 30	1 - 1.5%
Fully expanded fronds which are at least 18 inches long.			
Bromegrass, smooth	2.19	3 - 30	2%
Boot-to-early seedhead stage.			
Bursage, woolly-leaf	-	3 - 20	2%
For control, apply 2 quarts plus 1 pint of Dicamba per acre. At or beyond flowering and new active growth.			
Canary grass, reed	2.19 – 3.285	3 - 30	2%
Boot-to-head stage.			
Cattail	3.285 – 5.475	3 - 30	2%
Early head stage.			
Clover; red, white	3.285 – 5.475	3 – 20	2%
Early head stage.			
Cogongrass	3.285 – 5.475	10 - 30	2%
At least 18 inches tall in late summer or fall.			

Dallisgrass	3.285 – 5.475	3 - 20	2%
Early head stage.			
Dandelion	3.285 -- 5.475	3 - 30	2%
Early bud stage. Also for control, apply 16 fluid ounces plus 0.5 pound a.i. 2,4-D in 3 to 10 gallons of water per acre.			
Dock, curly	3.285 -- 5.475	3 - 30	2%
Early bud stage. Also for control, apply 16 fluid ounces plus 0.5 pound a.i. 2,4-D in 3 to 10 gallons of water per acre.			
Dogbane, hemp	4.38	3 - 30	2%
Late bud to flower stage in late summer or fall. For suppression, apply 16 fluid ounces plus 0.5 pound a.i. of 2,4-D in 3 to 10 GPA by ground and 3 to 5 GPA by air.			
Fescue (except fall)	3.285 -- 5.475	3 - 20	2%
Early head stage.			
Boot-to-early seedhead stage. Fall applications only; apply 1 quart in 3 to 10 GPA to fescue with 6 to 12" of new growth.			
Guineagrass	3.285	3 - 30	1%
7-leaf stage.			
Horsenettle	3.285 -- 5.475	3 - 20	2%
Early bud stage.			
Horseradish	4.38	3 - 30	2%
Late bud to flower stage in late summer or fall.			
Iceplant	-	-	1.5 - 2%
At or beyond the early bud stage. Thorough coverage is necessary for best control.			
Jerusalem artichoke	3.285 -- 5.475	3 - 20	2%
Early bud stage.			
Johnsongrass	1.095-3.285	3 - 30	1%
Boot-to-head stage of growth or in the fall prior to frost.			
Kikuyugrass	2.19-3.285	3 - 30	2%
At least 8 inches in height (3- or 4-leaf stage of growth).			
Knapweed	4.38	3 - 30	2%
Late bud to flower stage in late summer or fall.			
Lantana	-	-	1 – 1.25%
At or beyond the bloom stage.			
Lespedeza	3.285 -- 5.475	3 - 20	2%
Early bud stage.			
Milkweed, common	3.285	3 - 30	2%
Late bud to flower stage.			
Muhly, wirestem	1.095-2.19	3 - 30	2%
8" or greater.			
Mullein, common	3.285 -- 5.475	3 - 20	2%
Early bud stage.			
Napiergrass	3.285 -- 5.475	3 - 20	2%
Early head stage.			
Nightshade, silverleaf	2.19	3 - 10	2%
Apply when at least 60 percent of the plants have berries.			
Nutsedge; purple, yellow	1.095-3.285	3 - 30	1 - 2%
Apply 3 quarts when plants are in flower or when the new nutlets can be found at rhizome tips. Sequential applications: 1 to 2 quarts when plants are in the 3- to 5-leaf stage (less than 6 inches tall).			
Orchardgrass	2.19	3 - 30	2%
Boot-to-early seedhead stage, actively growing plants.			
Pampasgrass	-	-	1.5 - 2%
At or beyond the boot stage of growth.			
Paragrass	3.285 -- 5.475	3 - 20	2%
Early head stage.			
Phragmites*	3.285 -- 5.475	10 - 30	1 - 2%
Actively growing and in full bloom in late summer or fall.			
Poison hemlock	-	-	1 - 2%
Apply as a spray to wet treatment. Optimum results are obtained when plants are treated at the bud to full bloom stage of growth.			
Pokeweed, common	1.095	3 - 30	2%
Apply to actively growing plants up to 24 inches tall.			
Quackgrass	1.095-3.285	3 - 30	2%
6 to 8" tall followed by deep tillage.			
Redvine*	2.19	5 - 10	2%
At least 18" tall in late September or early October.			

Reed, giant	-	-	2%
Apply in late summer to fall for best results.			
Ryegrass, perennial	1.095-3.285	3 - 30	1%
Boot-to-head stage or in the fall prior to frost. DO NOT tank-mix with residual herbicide when using the 1-quart per acre rate.			
Smartweed, swamp	3.285 -- 5.475	3 - 30	2%
Early bud stage. Also for control, apply 16 fluid ounces plus 0.5 pound a.i. of 2,4-D in 3 to 10 GPA in the late summer or fall.			
Sowthistle, perennial	2.19-3.285	3 - 30	2%
At or beyond bud stage.			
Spurge, leafy*	-	3 - 10	2%
Apply 16 fluid ounces plus 0.5 pound a.i. 2,4-D in 3 to 10 gallons of water per acre in the late summer or fall.			
Starthistle, yellow	2.19	10 - 30	2%
Rosette, bolting and early flower stages.			
Sweet potato, wild*	-	-	2%
At or beyond the bloom stage.			
Thistle, artichoke*	-	-	2%
At or beyond the bloom stage of growth.			
Thistle, Canada	2.19-3.285	3 - 30	2%
At or beyond the bud stage, actively growing.			
Timothy	2.19-3.285	3 - 30	2%
Boot-to-head stage.			
Torpedograss*	4.38-5.475	3 - 30	2%
At or beyond the seedhead stage.			
Trumpetcreeper*	2.19	5 - 10	2%
At least 18" tall in late September and October.			
Vaseygrass	3.285 -- 5.475	3 - 20	2%
Early head stage.			
Velvetgrass	3.285 -- 5.475	3 - 20	2%
Early head stage.			
Wheatgrass, western	2.19-3.285	3 - 30	2%
Boot-to-head stage.			

*Partial Control.

For management of burning nettle in vine crops, tree fruits, tree nuts, citrus crops (nonbearing only), fallow systems, and vegetable crops.

See the "**crops**" section of this label for specific instructions on applications in Vine Crops, Tree Fruits (Including Tropical), Tree Nuts, Citrus Crops (Nonbearing Only), Fallow Systems, and Vegetable Crops.

Application Information

Apply up to 4.38 quarts per acre of **Sharda Glyphosate 30% + Oxyfluorfen 3.75% SC** per single application. Repeat applications may be made up to a maximum of 8.76 quarts per acre. For best results, apply this product after most weeds have germinated but before weeds exceed the maximum size recommended.

Weed Species	Rate (fluid ounces per acre)
Burning Nettle	26.28-35
	Maximum Height/Length (in inches)
	3-12

Restrictions

The maximum amount of **Sharda Glyphosate 30% + Oxyfluorfen 3.75% SC** applied must not exceed 8.76 quarts per acre per year.

Precautions

Sharda Glyphosate 30% + Oxyfluorfen 3.75% SC should be applied only around healthy growing trees.

Direct spray toward base of tree. Avoid direct herbicide contact with foliage or nuts.

STORAGE AND DISPOSAL

STORAGE: DO NOT contaminate water, food, or feed by storage or disposal. Store at temperatures above 3° F. If allowed to freeze, remix before using

PESTICIDE STORAGE: DO NOT use or store near heat or open flame. Keep the container tightly closed and dry in a cool, well-ventilated place. Storage temperature should not exceed 125°F. Protect against direct sunlight.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

[Less Than 5 Gallons]

[Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or mix tank. Drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.]

[Greater Than 5 Gallons]

[Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration.]

[For Bulk and Mini-Bulk Containers]

[Refillable container. Refill this container with pesticide only. **DO NOT** use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by State and local authorities.]

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

WARRANTY AND DISCLAIMER STATEMENT

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Sharda USA, LLC. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SHARDA USA LLC MAKES NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Sharda USA, LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SHARDA USA LLC DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT SHARDA USA LLC'S ELECTION, THE REPLACEMENT OF PRODUCT.

[OPTIONAL MARKETING LANGUAGE]

	 []
2	[Handle with Care]
3	[This side Up]