



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505T)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

83772-15

Date of Issuance:

2/27/25

NOTICE OF PESTICIDE:

☒ Registration
☐ Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

AgSaver™ Glyphosate 5.4 Plus

Name and Address of Registrant (include ZIP Code):

AgSaver, LLC
203 East Ash Street
McGehee, AR 71654

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

Continues page 2

Signature of Approving Official:

Emily Schmid

Emily Schmid, Product Manager 25
Herbicide Branch, Registration Division (7505P)

Date:

2/27/25

2. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 83772-15."
3. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The record for this product currently contains the following CSF(s):

- Basic CSF dated 2/23/2023
- Alternate CSF 1-3 dated 2/23/2023

If you have any questions, please contact Lydia Crawford at 202-566-2575 or at crawford.lydia@epa.gov.

Enclosure

{Note to reviewer: [Text] in brackets denotes optional text. {Text} in braces denotes where in the final label text will appear and notes to reviewer.}

{SUB-LABEL A - For non-aquatic terrestrial uses only}

GLYPHOSATE	GROUP	9	HERBICIDE
------------	-------	---	-----------

AgSaver™ Glyphosate 5.4 Plus
{ABN:} [CropSmart Glyphosate 5.4 Plus]

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

1.0 Ingredients

% BY WT.

ACTIVE INGREDIENT: Glyphosate*, N-(phosphonomethyl)glycine, in the form of its isopropylamine salt ... 53.8%

OTHER INGREDIENTS: 46.2%

TOTAL..... 100.0%

*Contains 648 grams per liter or 5.4 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its isopropylamine salt. Equivalent to 480 grams per liter or 4 pounds per U.S. gallon of the acid, glyphosate.

2.0 Emergency Phone Numbers

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE CALL CHEMTREC (800) 424-9300

KEEP OUT OF REACH OF CHILDREN
CAUTION

3.0 PRECAUTIONARY STATEMENTS

3.1 HAZARDS TO HUMANS AND DOMESTIC ANIMALS

ACCEPTED

2/27/2025

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No. 83772-15

DOMESTIC ANIMALS: This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

3.1.1 PHYSICAL OR CHEMICAL HAZARDS

Spray solutions of this product must be mixed, stored, and applied using only stainless steel, aluminum, fiberglass, plastic, or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette, or other ignition source.

EPA Reg. No. 83772-[xx]

EPA Est. No.

[[[(When more than one EPA Est. No. is used this may appear:)] Last letter in lot number corresponds to the EPA Est. No. used.]

Manufactured by:

AgSaver, LLC 203 East Ash Street McGehee, AR 71654

[Net Contents:]

[Batch Code:]

{Net Contents: May appear on label and/or container.} {Batch Code: May appear on label and container.}

[AgSaver is a trademark of AgSaver, LLC]

Section	DESCRIPTION
1.0	INGREDIENTS
2.0	EMERGENCY PHONE NUMBERS
3.0	PRECAUTIONARY STATEMENTS
3.1	Hazards To Humans & Domestic Animals
3.1.1	Physical & Chemical Hazards
3.2	Personal Protective Equipment (PPE)
3.3	User Safety Recommendations
3.4	Environmental Hazards
3.5	Directions for Use
3.6	Agricultural Use Requirements
3.7	Non-agricultural Use Requirements
3.8	Seed Potato Precautions
4.0	STORAGE & DISPOSAL
5.0	USE INFORMATION (Mode of Action)
6.0	WEED RESISTANCE MANAGEMENT
6.1	Weed Management Recommendations
6.2	Management Recommendations For Glyphosate Resistant Biotypes
7.0	MIXING
7.1	Mixing with Water
7.2	Surfactants
7.3	Tank Mixing Procedure
7.4	Mixing For Hand-Held Sprayers
7.5	Ammonium Sulfate
7.6	Colorants or Dyes
7.7	Drift Control Additives
8.0	APPLICATION EQUIPMENT & TECHNIQUES
8.1	Aerial Equipment
8.2	Aerial Spray Drift Management
8.3	Ground Broadcast Equipment
8.4	Hand-Held Or High-Volume Equipment
8.5	Selective Equipment
8.6	Injection Equipment
8.7	CDA Equipment
9.0	ANNUAL & PERRENIAL CROPS (Alphabetical)
9.1	Cereal & Grain Crops
9.2	Corn
9.3	Cotton
9.4	Fallow Systems
9.5	Grain Sorghum (Milo)
9.6	Herbs & Spices
9.7	Oil Seed Crops
9.8	Soybeans (Non-Roundup Ready)
9.9	Sugarcane
9.10	Vegetable Crops

Section	DESCRIPTION
9.11	Miscellaneous Crops
10.0	TREE, VINE & SHRUB CROPS (ALPHABETICAL)
10.1	Berry Crops
10.2	Citrus
10.3	Miscellaneous Tree Food Crops
10.4	Non-Food Tree Crops
10.5	Pome Fruit
10.6	Stone Fruit
10.7	Tree Nuts
10.8	Tropical & Subtropical Trees & Fruits
10.9	Vine Crops
11.0	PASTURE GRASSES, FORAGE LEGUMES & RANGELANDS
11.1	Alfalfa, Clover & Other Forage Legumes
11.2	Conservation Reserve Program Acres (CRP)
11.3	Grass or Turfgrass Seed Production
11.4	Pastures
11.5	Rangelands
11.6	Turf Grass Sod Production
11.7	Release of Bermuda Grass & Bahia Grass
12.0	ROUNDUP READY CROPS
12.1	Roundup Ready Canola (Spring Varieties)
12.2	Roundup Ready Canola (Fall Varieties)
12.3	Roundup Ready Corn
12.4	Roundup Ready Cotton
12.5	Roundup Ready Flex Cotton
12.6	Roundup Ready Soybeans
12.7	Roundup Ready Sugar Beet
12.8	Roundup Ready Alfalfa
13.0	NON-CROP USES AROUND THE FARMSTEAD
13.1	Weed Control, Trim-and-Edge & Bare Ground
13.2	Greenhouse/Shadehouse
13.3	Chemical Mowing
13.4	Cut Stumps
13.5	Habitat Management
14.0	FORESTRY, INDUSTRIAL, TURF & ORNAMENTAL
14.1	Forestry Site Preparation
14.2	Noncrop Areas & Industrial Sites
14.3	Injection & Frill (Woody Brush & Trees)
14.4	Hollow-Stem Injection
14.5	Ornamentals, Plant Nurseries & Christmas Trees
14.6	Parks, Recreational & Residential Areas
14.7	Railroads
14.8	Roadsides
14.9	Utility Sites
15.0	ANNUAL WEEDS RATE TABLE (ALPHABETICAL BY SPECIES)
15.1	Annual Weeds – Rates For 10 To 40 Gpa

Section	DESCRIPTION
15.2	Annual Weeds – Tank Mixtures With 2,4-D, Dicamba Or Picloram
16.0	PERENNIAL WEEDS RATE TABLE (ALPHABETICAL BY SPECIES)
17.0	WOODY BRUSH & TREES RATE TABLE (ALPHABETICAL BY SPECIES)
18.0	WARRANTY DISCLAIMER, INHERENT RISKS OF USE, LIMITATION OF REMEDIES

{Note to Reviewer: Table of Contents Optional on Final Label; section and page numbers may be adjusted as applicable for accuracy.}

3.2 PERSONAL PROTECTIVE EQUIPMENT: (PPE)

Applicators and other handlers must wear:

1. long-sleeved shirt and long pants,
2. shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE (Personal Protective Equipment). If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

3.3 USER SAFETY RECOMMENDATIONS:

Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing or PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

3.4 ENVIRONMENTAL HAZARDS

FOR TERRESTRIAL USE ONLY: Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

3.5

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

3.6 AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, are:

1. coveralls,
2. waterproof gloves,
3. shoes plus socks.

3.7 NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep people and pets off treated areas until spray solution has dried to prevent transfer of this product onto desirable vegetation.

3.8 Seed Potato Precautions

Potatoes grown for seed are very sensitive to glyphosate at extremely low concentrations. Exposure of the seed potato crop can cause germination failure or deformities. Daughter tuber damage may occur at levels where mother crop symptoms are not viable. Multiple sprouting from eyes, weak and distorted stems, "little potato syndrome", cauliflower sprouts, root distortions, excessive root growth, suppressed tuber initiation and bulking failure or delay in opening of eyes and rotting of tubers in the field or store can result. Subsequent plantings of seed pieces from the exposed mother crop can result in delayed or no emergence or produce lower than normal yields.

Glyphosate can contaminate seed potato crops through carryover residue in application equipment or drift from applying glyphosate to nearby crops.

Always follow good wash-out procedures using detergents or other suitable cleaning agents to remove all residual traces of glyphosate from application equipment that may be used to apply other products to seed potato crops.

To avoid contamination from spray drift, follow the precautions in the "Spray Drift Management" section of the label.

4.0 STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

STORAGE: STORE ABOVE 10°F (-12°C) TO KEEP FROM CRYSTALIZING. Crystals will settle to the bottom. If crystals form, allow product to warm above 50°F (10°C) and mix well or recirculate to redissolve.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed must be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state, or local procedures. Emptied container retains vapor and product residue. Observe all labeled safeguards until container is cleaned, reconditioned, or destroyed.

CONTAINER HANDLING: (See the Net Contents section on the container to determine if it non-refillable or refillable.) APPROPRIATE BOX MUST BE CHECKED.

Non-refillable containers (1 and 2.5 gallon): Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Non-refillable containers (>5 gallon): Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Refillable containers: Refillable container. Refill this container with glyphosate only. Do not reuse this container for any other purpose.

When this container is empty, replace the cap and seal all openings that have been made during usage and return the container to the point of purchase, or to an alternate location designated by the manufacturer at the time of purchase of this product. If not returned, clean container the empty container and offer for recycling, if available.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the re-filler.

To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or re-circulate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing process two more times.

If the container cannot be refilled, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Read the entire label before using this product. Use only according to label instructions.

Read the "CONDITIONS OF SALE AND WARRANTY" statement at the end of the label before buying or using. If terms are not acceptable, return at once unopened.

5.0 USE INFORMATION

Product Description: This product is a postemergent, systemic herbicide with no soil residual activity. It is generally non-selective and gives broad-spectrum control of many annual weeds, perennial weeds, woody brush, and trees. It is formulated as a water-soluble liquid. It may be applied through most standard industrial or field-type sprayers after dilution and thorough mixing with water or other carriers according to label instructions.

Surfactant may be included in the tank mixture if desired and should only be done so based on field experience or further instructions from your local extension service, crop consultant or field representative.

Time to Symptoms: This product moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days, but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay development of visual symptoms. Visible effects are a gradual wilting and yellowing of the plant which advances to complete browning of aboveground growth and deterioration of underground plant parts.

Stage of Weeds: Annual weeds are easiest to control when they are small. Best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity. Refer to the annual, perennial, woody brush, and trees rate tables for rates of specific weeds.

Always use the higher rate of this product per acre within the labeled range when weed growth is heavy or dense or weeds are growing in an undisturbed (non-cultivated) area.

Do not treat weeds under poor growing conditions such as drought stress, disease or insect damage, as reduced weed control may result. Reduced results may also occur when treating weeds heavily covered with dust.

Cultural Considerations: Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed, or cut, and have not been allowed to regrow to the labeled stage for treatment.

Rainfastness: Heavy rainfall soon after application may wash this product off of the foliage and a repeat application may be required for adequate control.

Spray Coverage: For best results, spray coverage should be uniform and complete. Do not spray weed foliage to the point of runoff.

Mode of Action: The active ingredient in this product inhibits an enzyme found only in plants and microorganisms that is essential to formation of specific amino acids.

No Soil Activity: Weeds must be emerged at the time of application to be controlled by this product. Weeds germinating from seed after application will not be controlled. Unemerged plants arising from unattached underground rhizomes or rootstocks of perennials will not be affected by the herbicide and will continue to grow.

When this product comes in contact with soil, it is bound to soil particles. Under labeled use situations, once this product is bound to soil particles, it is not available for plant uptake and will not harm off-site vegetation where roots grow into the treated area or if the soil is transported off-site. The strong affinity of this product to soil particles prevents this product from leaching out of the soil profile and entering ground water.

Biological Degradation: Degradation of this product is primarily a biological process carried out by soil microbes.

Tank Mixing: This product does not provide residual weed control. For subsequent residual weed control, follow a label-approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used. Use according to the most restrictive label directions for each product in the mixture.

To the extent consistent with applicable law, buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressly allowed

in this labeling. Mixing this product with herbicides or other materials not labeled on this label may result in reduced performance.

Annual Maximum Use Rate:

- Except as otherwise specified in a crop section of this label, the combined total of all treatments must not exceed 4.5 quarts (4.5 lbs ae) of this product per acre per year.
- For tree, vine, or shrub crops, the combined total of all treatments must not exceed 7.8 quarts (7.8 lbs ae) per acre per year, unless otherwise specified.
- For noncrop uses, the combined total of all treatments must not exceed 6.0 quarts (6 lbs ae) of this product per acre per year, unless otherwise specified.

ATTENTION

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift, or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of injury occurring from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing or when there are other meteorological conditions that favor spray drift. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) which are likely to drift.

Refer to the Spray Drift Management guidance in Section 8.0.

NOTE: Keep container closed to prevent spills and contamination.

6.0 WEED RESISTANCE MANAGEMENT

For resistance management, AgSaver™ Glyphosate 5.4 Plus is a Group 9 herbicide. Any weed population may contain or develop plants naturally resistant to AgSaver™ Glyphosate 5.4 Plus and other Group 9 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistant management strategies should be followed.

6.1 WEED MANAGEMENT DIRECTIONS

For resistance management, AgSaver™ Glyphosate 5.4 Plus is a Group 9 herbicide. Any weed population may contain or develop plants naturally resistant to AgSaver™ Glyphosate 5.4 Plus and other Group 9 herbicides. Weed species with acquired resistance to Group 9 herbicides may eventually dominate the weed population if Group 9 herbicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by AgSaver™ Glyphosate 5.4 Plus or other Group 9 herbicides. Users should scout before and after application.

Suspected herbicide-resistant weeds may be identified by these indicators:

- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
- A spreading patch of non-controlled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.

To delay herbicide resistance, take one or more of the following steps:

- Avoid the consecutive use of AgSaver™ Glyphosate 5.4 Plus or other target site of action Group 9 herbicides that might have a similar target site of action, on the same weed species.

- Use tank mixtures or premixes with herbicides from different target site of action Groups as long as the involved products are all registered for the same use, have different sites of action and are both effective at the tank mix or prepack rate on the weed(s) of concern (an herbicide mode of action classification by itself may not adequately address specific weeds that are resistant to specific herbicides).
- Base herbicide use on a comprehensive Integrated Pest Management (IPM) program.
- Scout fields prior to application to identify the weed species present and their growth state to determine if the intended application will be effective.
- Scout fields after application to verify that the treatment was effective.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- Report any incidence of repeated non-performance of this product on a particular weed to your AgSaver™, LLC representative, local retailer, or county extension agent.

6.2 MANAGEMENT DIRECTIONS FOR GLYPHOSATE RESISTANCE BIOTYPES

Note: Appropriate testing is critical in order to determine if a weed is resistant to glyphosate. Contact your AgSaver™, LLC representative to determine if resistance has been confirmed to any particular weed biotype in your area, or visit on the internet www.weedscience.org. For more information see the “ANNUAL WEEDS RATE SECTION” and “PERENNIAL WEEDS RATE SECTION” of this label.

Control directions for biotypes confirmed as resistant to glyphosate are made available on separately published supplemental labeling or fact sheets for this product and can be obtained from your local retailer or AgSaver™, LLC representative.

Since the occurrence of new glyphosate resistant weeds cannot be determined until after product use and scientific confirmation, to the extent consistent with applicable law, AgSaver™, LLC is not responsible for any losses that may result from the failure of this product to control glyphosate-resistant weed biotypes.

The following good agronomic practices are recommended to reduce the spread of confirmed glyphosate resistant biotypes:

- If a naturally occurring resistant biotype is present in your field, this product should be tank-mixed or applied sequentially with an appropriately labeled herbicide with a different mode of action to achieve control
- Cultural and mechanical control practices (e.g., crop rotation or tillage) may also be used as appropriate.
- One method for adding other herbicides into a continuous Roundup Ready® system is to rotate to other Roundup Ready crops.
- Scout treated fields after herbicide applications and control escaping weeds including resistant biotypes before they set seed.
- Thoroughly clean equipment before leaving fields known to contain resistant biotypes.

7.0 MIXING

Clean sprayer parts immediately after using this product by thoroughly flushing with water.

Precaution: Reduced results may occur if water containing soil is used, such as visibly muddy water or water that is not clear from ponds and ditches.

7.1 Mixing with Water

This product mixes readily with water. Mix spray solutions of this product as follows: Fill the mixing or spray tank with the required amount of water. Add the labeled amount of this product near the end of the filling process and mix well. Use caution to avoid siphoning back into the carrier source. Use approved anti-back-siphoning devices where required by state or local regulations. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, avoid the use of mechanical agitators, terminate by-pass, and return lines at the bottom of the tank and, if needed, use an approved anti-foam or defoaming agent.

7.2

Surfactant

Surfactant may be included in the tank mixture if desired and should only be done so based on field experience or further recommendation of your local extension service, crop consultant or field representative.

7.3

Tank Mixing Procedure

Mix labeled tank mixtures of this product with water as follows:

1. Place a 20 to 35-mesh screen or wetting basket over filling port.
2. Through the screen, fill the spray tank one-half full with water and start agitation.
3. If a wettable powder is used, make a slurry with the water carrier, and add it SLOWLY through the screen into the tank. Continue agitation.
4. If a flowable formulation is used, premix one part flowable with one part water. Add diluted mixture SLOWLY through the screen into the tank. Continue agitation.
5. If an emulsifiable concentrate formulation is used, premix one part emulsifiable concentrate with two parts water. Add diluted mixture slowly through the screen into the tank. Continue agitation.
6. Continue filling the spray tank with water and add the required amount of this product near the end of the filling process.
7. When using nonionic surfactant add it to the spray tank before completing the filling process.
8. Add individual formulations to the spray tank as follows: wettable powder, flowable, emulsifiable concentrate, drift control additive, water-soluble liquid followed by surfactant.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed. Keep by-pass line on or near the bottom of the tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50 mesh.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Always predetermine the compatibility of labeled tank mixtures of this product with water carrier by mixing small proportional quantities in advance.

Refer to the "TANK MIXING" section of "USE INFORMATION" for additional precautions.

7.4

Mixing for Hand-held Sprayers

Prepare the desired volume of spray solution by mixing the amount of this product in water as shown in the following table:

Spray Solution

AMOUNT OF PRODUCT

Desired Volume	3/4%	1%	1 1/2%	2%	5%	10%
1 Gal	1 fl oz (0.03 lb ae)	1 1/3 fl oz (0.04 lb ae)	2 fl oz (0.06 lb ae)	2 2/3 fl oz (0.08 lb ae)	6 1/2 fl oz (0.2 lb ae)	13 fl oz (0.41 lb ae)
25 Gal	1 1/2 pt (0.75 lb ae)	1 qt (1 lb ae)	1 1/2 qt (1.5 lbs ae)	2 qt (2 lbs ae)	5 qt (5 lbs ae)	2 1/2 gal (10 lbs ae)
100 Gal	3 qt (3 lbs ae)	1 gal (4 lbs ae)	1 1/2 gal (6 lbs ae)	2 gal (8 lbs ae)	5 gal (20 lbs ae)	10 gal (40 lbs ae)

2 tablespoons = 1 fluid ounce

For use in knapsack sprayers, it is suggested that the labeled amount of this product be mixed with water in a larger container. Fill sprayer with the mixed solution.

7.5

Ammonium Sulfate

The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of this product when tank mixed with certain residual herbicides on annual and perennial weeds. The equivalent rate of ammonium sulfate in a liquid formulation may also be used. Ensure that ammonium sulfate is completely dissolved in the spray tank before adding herbicides. Thoroughly rinse the spray system with clean water after use to reduce corrosion.

NOTE: When using ammonium sulfate, apply this product at rates labeled in this label. Lower rates will result in reduced performance.

7.6

Colorants or Dyes

Agriculturally approved colorants or marking dyes may be added to this product. Colorants or dyes used in spray solutions of this product may reduce performance, especially at lower rates or dilutions. Use colorants or dyes according to the manufacturer's directions.

7.7

Drift Control Additives

Drift control additives may be used with all equipment types, except wiper applicators, sponge bars and CDA equipment. When a drift control additive is used, read, and carefully observe the cautionary statements and all other information appearing on the additive label.

8.0

APPLICATION EQUIPMENT AND TECHNIQUES

Do not apply this product through any type of irrigation system.

This product may be applied with the following application equipment:

Aerial – Fixed Wing and Helicopter

Ground Broadcast Spray – Boom or boomless systems, pull-type sprayer, floaters, pick-up sprayers, spray coupes and other ground broadcast equipment.

Hand-held and High-Volume Spray Equipment – Knapsack and backpack sprayers, pump-up pressure sprayers, handguns, handwands, mistblowers*, lances and other hand-held and motorized spray equipment used to direct the spray onto weed foliage.

*Not Registered for Use in mistblowers by California and Arizona

Selective Equipment – Recirculating sprayers, shielded and hooded sprayers, wiper applicators and sponge bars.

Injection Systems – Aerial or ground injection sprayers.

Controlled Droplet Applicator (CDA) – Hand-held or boom-mounted applicators which produce a spray consisting of a narrow range of droplet sizes.

Apply these spray solutions in properly maintained and calibrated equipment capable of delivering desired volumes.

8.1

Aerial Equipment

DO NOT APPLY THIS PRODUCT USING AERIAL SPRAY EQUIPMENT EXCEPT UNDER CONDITIONS AS SPECIFIED WITHIN THIS LABEL.

Use the labeled rates of this herbicide in 3 to 15 gallons of water per acre unless otherwise specified on this label. Unless otherwise specified, do not exceed 24 fl oz (0.75 lb ae) per acre. Aerial applications of this product may be

made in annual cropping conventional tillage systems, fallow, and reduced tillage systems and preharvest applications. Refer to the individual use area sections of this label for labeled volumes and application rates.

NOTE: For aerial application in California or Arkansas, refer to the Federal supplemental label for aerial applications in that state for specific instructions, restrictions, and requirements. For aerial applications, consult with state or local authorities regarding any additional requirements for aerial treatments.

Not Registered for Use by California for dicamba tank mixtures applied by air. Avoid direct application to any body of water.

AVOID DRIFT – DO NOT APPLY DURING LOW-LEVEL INVERSION CONDITIONS, WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION WHICH FAVORS DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine spray droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure.

Drift control additives may be used. When a drift control additive is used, read, and carefully observe the cautionary statements and all other information appearing on the additive label.

Ensure uniform application – To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills.

Prolonged exposure of this product to uncoated steel surfaces may result in corrosion and possible failure of the part. Landing gear are most susceptible.

The maintenance of an organic coating (paint), which meets aerospace specification MIL-C-38413, may prevent corrosion.

8.2 AERIAL SPRAY DRIFT MANAGEMENT

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed $\frac{3}{4}$ the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they must be observed.

The applicator must be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory.

Aerial Drift Reduction Advisory

This section is advisory in nature and does not supersede the mandatory label requirements.

INFORMATION ON DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (See Wind, Temperature and Humidity, and Temperature Inversions).

CONTROLLING DROPLET SIZE

- Volume – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure – Do not exceed the nozzle manufacturer's specified pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of nozzles – Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation – Orient nozzles so that the spray is released parallel to the airstream which produces larger droplets than other orientations. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- Nozzle Type – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

BOOM LENGTH

For some use patterns, reducing the effective boom length to less than $\frac{3}{4}$ of the wingspan or rotor length may further reduce drift without reducing swath width.

APPLICATION HEIGHT

Applications must not be made at a height greater than 10 feet above the top of the target plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

SWATH ADJUSTMENT

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance must increase with increasing drift potential (higher wind, smaller drops, etc.)

WIND

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application must be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator must be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

TEMPERATURE INVERSIONS

Applications must not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small, suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SENSITIVE AREAS

The pesticide must only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

8.3 Ground Broadcast Equipment

Use the labeled rates of this product in 3 to 40 gallons of water per acre as a broadcast spray unless otherwise specified. As density of weeds increases, spray volume should be increased within the labeled range to ensure complete coverage. Carefully select proper nozzles to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

8.4 Hand-Held and High-Volume Equipment

Apply to foliage of vegetation to be controlled. For applications made on a spray-to-wet basis, spray coverage must be uniform and complete. Do not spray to the point of runoff. Use coarse sprays only. For control of weeds listed in the annual weeds rate tables, apply a 1/2 percent solution of this product to weeds less than 6 inches in height or runner length. Apply prior to seedhead formation in grass or bud formation in broadleaf weeds. For annual weeds over 6 inches tall, or unless otherwise specified, use a 1 percent solution.

For best results, use a 1 1/2 percent solution on harder-to-control perennials, such as bermudagrass, dock, field bindweed, hemp dogbane, milkweed, and Canada thistle.

When using application methods which result in less than complete coverage, use a 3.75 percent solution for annual and perennial weeds and a 3.75 to 5 percent solution for woody brush and trees.

8.5 Selective Equipment

This product may be applied through recirculating spray systems, shielded applicators, hooded sprayers, wiper applicators or sponge bars after dilution and thorough mixing with water to listed weeds growing in any noncrop site specified on this label and only when specifically labeled in cropping systems.

A recirculating spray system directs the spray solution onto weeds growing above desirable vegetation, while spray solution not intercepted by weeds is collected and returned to the spray tank for reuse.

A shielded or hooded applicator directs the herbicide solution onto weeds, while shielding desirable vegetation from the herbicide.

A wiper or sponge applicator applies the herbicide solution onto weeds by rubbing the weed with an absorbent material containing the herbicide solution.

AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION.

Contact of the herbicide solution with desirable vegetation may result in damage or destruction. Applicators used above desirable vegetation must be adjusted so that the lowest spray stream or wiper contact point is at least 2 inches above the desirable vegetation. Droplets, mist, foam, or splatter of the herbicide solution settling on desirable vegetation may result in discoloration, stunting or destruction.

Applications made above the crops should be made when the weeds are a minimum of 6 inches above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations or when the height of the weeds varies so that not all weeds are contacted. In these instances, repeat applications can be made up to the maximum number of applications for that use site.

Shielded and hooded applicators

Use nozzles that provide uniform coverage within the treated area. Keep shields on these sprayers adjusted to protect desirable vegetation. Extreme care must be exercised to avoid contact of herbicide with desirable vegetation.

A hooded sprayer is a type of shielded applicator. The spray pattern is completely enclosed on the top and all 4 sides by a hood, thereby shielding the crop from the spray solution. This equipment must be set up and operated in a manner that avoids bouncing or raising the hoods off the ground in any way. If the hoods are raised, spray particles may escape and come into contact with the crop, causing damage or destruction of the crop. The spray hoods must be operated on the ground or skimming across the ground. Tractor speed must be adjusted to avoid bouncing of the spray hoods. Avoid operation on rough or sloping ground where the spray hoods might be raised off the ground.

Wiper applicators and sponge bars

Wiper applicators are devices that physically wipe appropriate amounts of this product directly onto the weed.

Equipment must be designed, maintained, and operated to prevent the herbicide solution from contacting desirable vegetation. Operate this equipment at ground speeds no greater than 5 mph. Performance may be improved by reducing speed in areas of heavy weed infestations to ensure adequate wiper saturation. Better results may be obtained if 2 applications are made in opposite directions.

Avoid leakage or dripping onto desirable vegetation. Adjust height of applicator to ensure adequate contact with weeds. Keep wiping surfaces clean. Be aware that, on sloping ground, the herbicide solution may migrate, causing dripping on the lower end and drying of the wicks on the upper end of a wiper applicator.

Do not use wiper equipment when weeds are wet.

Mix only the amount of solution to be used during a 1-day period, as reduced activity may result from use of leftover solutions. Clean wiper parts immediately after using this product by thoroughly flushing with water.

Include a nonionic surfactant at a rate of 10 percent by volume of total herbicide solution with all wiper applications.

For Rope or Sponge Wick Applicators – Mix 3 quarts (3 lbs ae) of this product in 2 gallons of water to prepare a 25 percent solution. Apply this solution to weeds listed in this section.

For Porous-Plastic Applicators – Solutions ranging from 25 to 100 percent of this product in water may be used in porous-plastic wiper applicators.

When applied as labeled, this product CONTROLS the following weeds:

Corn, volunteer	Sicklepod	Shattercane
Panicum, Texas	Spanishneedles	
Rye, common	Starbur, bristly	

When applied as labeled, this product SUPPRESSES the following weeds:

Beggarweed, Florida	Ragweed, common
Bermudagrass	Ragweed, giant
Dogbane, hemp	Smutgrass
Dogfennel	Sunflower
Guineagrass	Thistle, Canada
Johnsongrass	Thistle, musk
Milkweed	Vaseygrass
Nightshade, silverleaf	Velvetleaf
Pigweed, redroot	

8.6

Injection Systems

This product may be used in aerial or ground injection spray systems. It may be used as a liquid concentrate or diluted prior to injecting into the spray stream.

RESTRICTION:

- Do not mix this product with the concentrate of other products when using injection systems.

8.7

CDA Equipment

The rate of this product applied per acre by vehicle mounted CDA equipment must not be less than the amount labeled in this label when applied by conventional broadcast equipment. For vehicle mounted CDA equipment, apply 3 to 15 gallons of water per acre.

For the control of annual weeds with hand-held CDA units, apply a 20 percent solution of this product at a flow rate of 2 fl oz (0.06 lb ae) per minute and a walking speed of 1.5 mph (1.5 pints (0.75 lb ae) per acre). For the control of perennial weeds, apply a 20 to 40 percent solution of this product at a flow rate of 2 fl oz (0.06 lb ae) per minute and a walking speed of 0.75 mph (3 to 6 pints (1.5-3 lbs ae) per acre).

Controlled droplet application equipment produces a spray pattern which is not easily visible. Extreme care must be exercised to avoid spray or drift contacting the foliage or any other green tissue of desirable vegetation, as damage or destruction may result.

9.0 - ANNUAL & PERENNIAL CROPS (Alphabetical)

This section is organized alphabetically by crop category. There may be several labeled crops listed in a crop category.

See the individual crop categories for specific instructions, preharvest intervals, and additional precautions and restrictions.

See the "ROUNDUP READY CROPS" section of this label for instructions for treating Roundup Ready crops.

TYPES OF APPLICATIONS

Chemical fallow, Pre-plant fallow beds, Pre-plant, Pre-emergence, At Planting, Hooded Sprayers in Row-Middles, Shielded Sprayers in Row-Middles, Wiper Applications in Row-Middles, and Post-Harvest Treatments.

Additional application types may be specified or allowed in individual Crop Categories.

USE DIRECTIONS

Apply this product during fallow intervals preceding planting, prior to planting or transplanting, at planting, or pre-emergent to annual and perennial crops listed in this label, except where specifically limited. For any crop NOT listed in this label, applications must be made at least 30 days prior to planting.

Unless otherwise specified weed control applications must be made according to the rates listed in the “Annual Weeds,” “Perennial Weeds,” and “Woody Brush & Trees” rate tables in this label.

Post-directed hooded sprayers and wiper equipment capable of preventing all crop contact with herbicide solutions may be used in mulched or un-mulched row middles after crop establishment. Where specifically noted below, wipers may also be used above certain crops to control tall weeds. Refer to the “SELECTIVE EQUIPMENT” section of this label for essential precautions when using hooded sprayers or wipers to avoid crop injury caused by leakage of spray mists or dripping onto crops. Crop injury is possible with these methods of application.

The maximum use rates stated throughout this product’s labeling apply to this product combined with the use of all other herbicides containing glyphosate or sulfosate as the active ingredient, whether applied as mixtures or separately. Calculate the application rates and ensure that the total use of this and other glyphosate or sulfosate containing products does not exceed stated maximum use rate.

PRECAUTIONS

- Avoid contact of herbicide with foliage, green shoots or stems, bark, exposed roots (including those emerging from plastic mulch), or fruit of crops because severe injury or destruction may result.
- Apply before seed germination in coarse sandy soils to further minimize the risk of injury.
- When making pre-emergence and at planting applications, applications must be made before crop emergence to avoid severe crop injury. Broadcast applications made at emergence will result in injury or death to emerged seedlings.

RESTRICTIONS

- **DO NOT** apply more than 7.5 pints (3.75 lbs ae) per acre per application, unless otherwise specified for an individual use.
- **DO NOT** apply more than 4.5 quarts (4.5 lbs ae) per acre per year as a combined total of all treatments, except for applications in non-agricultural sites or in tree, vine, or shrub crops and unless otherwise specified for an individual use.
- Pre-harvest Interval (PHI): Treatments with selective equipment including wipers and hooded sprayers must be made at least 14 days prior to harvest, unless otherwise specified for an individual use.
- Pre-harvest Interval (PHI): For broadcast post-emergent treatments, **DO NOT** harvest or feed treated vegetation for 8 weeks following application, unless otherwise specified for an individual use.
- In crops where spot treatments are allowed, **DO NOT** treat more than 10 percent of the total field to be harvested. The crop receiving spray in treated area will be killed. Take care to avoid drift or spray outside the target area for the same reason.
- Post-harvest or fallow applications must be made at least 30 days prior to planting any non-labeled crop.

9.1 - CEREAL CROPS		
LABELED CROPS: Barley, Buckwheat, Millet (Pearl & Proso), Oats, Rice, Rye, Quinoa, Teff, Teosinte, Triticale, Wheat (All), Wild rice.		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
See Section 9.0	See Use Directions in Section 9.0	See Section 9.0
Pre-Plant, Pre-Emergence, At-Planting	This product may be applied before, during or after planting of cereal crops. Applications must be made prior to emergence of the crop.	Do not treat rice fields or levees when the field contains floodwater.

9.1 - CEREAL CROPS		
LABELED CROPS: Barley, Buckwheat, Millet (Pearl & Proso), Oats, Rice, Rye, Quinoa, Teff, Teosinte, Triticale, Wheat (All), Wild rice.		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
See Section 9.0	See Use Directions in Section 9.0	See Section 9.0
Red Rice Control (prior to planting rice)	Apply 2.25 pints (1.13 lbs ae) of this product in 5 to 10 gallons of water per acre. Flush fields prior to application to obtain uniform germination and stand of red rice. Make application when the majority of the red rice plants are in the 2-leaf stage and no more than 4 inches tall. Red rice plants with less than 2 true leaves may only be partially controlled. Avoid spraying during low humidity conditions, as reduced control may result.	Do not treat rice field or levees when the fields contain flood water. Do not re-flood treated fields for 8 days following application.
Spot treatment (except rice)	This product may be applied as a spot treatment in cereal crops. Apply this product before heading in small grains.	Do not treat more than 10 percent of the total field area to be harvested. The crop receiving spray in the treated area will be killed. Do not spray or allow drift outside target area for the same reason.
Over the Top Wiper applications (Feed barley & wheat only)	Wiper applications may be used in wheat. To control common rye or cereal rye, apply after the weeds have headed and achieved maximum growth, when the rye is at least 6 inches above the wheat crop.	Pre-harvest Interval (PHI): Allow at least 35 days between application and harvest. Do not use roller applicators.
Pre-harvest (Feed barley & wheat only)	This product provides weed control when applied prior to harvest of wheat. Apply after the hard-dough stage of grain (30% or less grain moisture) and at least 7 days prior to harvest. Wheat stubble may be grazed immediately after harvest. This product may be applied using either aerial or ground spray equipment. For ground applications, apply this product in 10 to 20 gallons of water per acre. For aerial applications, apply this product in 3 to 10 gallons of water per acre.	Do not apply more than 1.5 pints (0.75 lb ae) per acre per application. Do not apply to wheat or barley grown for seed. Pre-harvest Interval (PHI): Allow 7 days between application and harvest or grazing.
Post-harvest	This product may be applied after harvest of cereal crops. Higher rates may be required for control of large weeds which were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.	For any crop not listed on this label, applications must be made at least 30 days prior to planting the next crop. Pre-harvest Interval (PHI): Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

9.2 – CORN (Non-Roundup Ready)		
LABELED CROPS: Field corn, Seed corn, Silage corn, Sweet corn and Popcorn		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
See Section 9.0	See Use Directions in Section 9.0	See Section 9.0
Pre-plant, Pre-emergence, At planting	This product may be applied before, during or after planting corn. Applications must be made prior to emergence of the crop. TANK MIXTURES: Apply these tank mixtures in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.	Do not apply in nitrogen solutions to tough-to-control grasses such as barnyardgrass, fall panicum, broadleaf signalgrass, annual ryegrass and any perennial weeds in the following area: From Route 50 South in Illinois and Indiana and the following states: Alabama, Arkansas, Delaware, Florida, Georgia, Kentucky, Louisiana, Maryland, Mississippi, New Jersey, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia, and West Virginia.

9.2 – CORN (Non-Roundup Ready)		
LABELED CROPS: Field corn, Seed corn, Silage corn, Sweet corn and Popcorn		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
	<p>acetochlor dimethenamid pendimethalin acetochlor + atrazine dimethenamid + atrazine rimsulfuron + atrazine flufenacet + isoxaflutole thifensulfuron- atrazine + s-metolachlor flufenacet + metribuzin methyl carfentrazone-ethyl flumetsulam simazine dicamba + atrazine isoxaflutole S-metolachlor diflufenzopyr + dicamba linuron 2,4-D</p> <p>For difficult to control annual weeds such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signal grass up to 2 inches tall and Pennsylvania smartweed up to 6 inches tall, apply this product at 1.5 pints (0.75 lb ae). per acre in these tank mixtures. For other labeled weeds, apply 1-1.5 pints (0.5-0.75 lb ae) of this product per acre when weeds are less than 6 inches tall, 1.5 – 2.25 pints (0.75-1.13 lbs ae) when weeds are over 6 inches tall. When using nitrogen solutions as the carrier, use rate may need to be increased to the 2.25 pints (1.13 lbs ae) rate for acceptable weed control.</p>	
Spot treatment	For spot treatments, apply this product prior to silking of corn.	Do not treat more than 10 percent of the total field area to be harvested. The crop receiving spray in the treated area will be killed. Do not spray or allow drift outside target area for the same reason.
Hooded sprayers	<p>This product may be used through hooded sprayers for weed control between the rows of corn.</p> <p>Only hooded sprayers that completely enclose the spray pattern may be used.</p> <p>See additional instructions for the use of hooded sprayers in the "Application Equipment and Techniques" section of this label.</p> <p>Precaution: Contact of this product in any manner to any vegetation to which treatment is not intended may cause damage. To the extent consistent with applicable law, such damage shall be the sole responsibility of the applicator.</p>	<p>Corn must be at least 12 inches tall, measured without extending leaves.</p> <p>Do not apply more than 1.5 pints (0.75 lb ae) per acre per application.</p> <p>Do not apply more than 2.25 quarts (2.25 lbs ae). per acre per year.</p>
Pre-harvest	<p>Make applications at 35 percent grain moisture or less. Ensure that maximum kernel fill is complete, and the corn is physiologically mature (black layer formed).</p> <p>For ground applications, apply up to 2.25 quarts (2.25 lbs ae) of this product per acre.</p> <p>For aerial applications, apply up to 3 pints (1.5 lbs ae) of this product per acre.</p>	<p>Do not apply more than 2.25 quarts (2.25 lbs ae) per acre per ground application.</p> <p>Do not apply more than 3 pints (1.5 lbs ae) per acre per aerial application.</p> <p>Pre-harvest Interval (PHI): Allow a minimum of 7 days between application and harvest.</p> <p>Do not make applications to corn grown for seed.</p>
Post-harvest	This product may be applied after harvest of corn. Higher rates may be required for control of large weeds which were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.	Pre-harvest Interval (PHI): Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

9.3 – COTTON		
LABELED CROPS: Cotton (non-Roundup Ready)		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
See Section 9.0	See Use Directions in Section 9.0	See Section 9.0
Pre-plant, Pre-emergence, At-planting	This product may be applied before, during or after planting cotton.	Applications must be made prior to emergence of the crop.
Hooded sprayer, Selective equipment	This product may be applied through hooded sprayers, shielded applicators or wiper applicators in cotton.	<p>See the "Selective Equipment" part of the "Application Equipment and Techniques" section of this label for information on proper use and calibration of this equipment.</p> <p>Pre-harvest Interval (PHI): Allow at least 7 days between application and harvest.</p>

9.3 – COTTON		
LABELED CROPS: Cotton (non-Roundup Ready)		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
Spot treatment	For spot treatments, apply this product prior to boll opening of cotton.	Do not treat more than 10 percent of the total field area to be harvested. The crop receiving spray in the treated area will be killed. Do not spray or allow drift outside target area for the same reason.
Pre-harvest	<p>This product provides weed control and cotton regrowth inhibition when applied prior to harvest of cotton. For weed control, apply at rates given in the annual, perennial, and woody brush tables. Apply 12 – 48 fl oz (0.38-1.5 lbs ae) of this product per acre for cotton regrowth inhibition.</p> <p>Apply up to 3 pints (1.5 lbs ae) of this product using either aerial or ground spray equipment. Apply after sufficient bolls have developed to produce the desired yield of cotton. Applications made prior to this time could affect maximum yield potential.</p> <p>TANK MIXTURES:</p> <p>This product may be tank mixed with tribufos, diuron + thidiazuron, or ethephon to provide additional enhancement of cotton leaf drop. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.</p>	<p>Do not apply more than 3 pints (1.5 lbs ae) per acre per application.</p> <p>Pre-harvest Interval (PHI): Allow at least 7 days between application and harvest.</p> <p>Do not apply to cotton grown for seed, as a reduction in germination or vigor may occur.</p> <p>The use of additives other than those listed on this label, for preharvest application to cotton is prohibited.</p>

9.4 - FALLOW SYSTEMS		
LABELED CROPS: This product may be applied during the fallow period prior to planting or emergence of any crop on this label.		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
	See Use Directions in Section 9.0	See Section 9.0
Chemical Fallow	<p>This product may be applied during the fallow period prior to planting or emergence of any crop listed on this label.</p> <p>This product may be used as a substitute for tillage to control annual weeds in fallow fields. Also, broadcast or spot treatments will control or suppress many perennial weeds in fallow fields. Ground or aerial application equipment may be used. Tank mixtures with 2,4-D and dicamba may be used. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.</p> <p>Applications up to 3 pints (1.5 lbs ae). per acre may be made by aerial application in fallow sites where there is sufficient buffer to prevent injury due to drift onto adjacent crops.</p>	<p>Do not apply more than 3 pints (1.5 lbs ae) per acre per aerial application.</p> <p>For any crop not listed on this label, applications must be made at least 30 days prior to planting.</p> <p>Not Registered for Use by California for dicamba tank mixtures applied by air.</p> <p>Refer to the specific product labels for crop rotation restrictions and cautionary statements of all products used in tank mixtures.</p>
Pre-plant Fallow Beds	<p>This product may be applied to fallow beds prior to planting or emergence of any crop listed on this label. This product will control weeds listed in the annual, perennial, and woody brush tables.</p> <p>TANK MIXTURES:</p> <p>It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.</p> <p>In addition, 9 fl oz (0.28 lb ae) of this product plus the labeled rate of oxyfluorfen per acre will control the following weeds with the maximum height or length indicated: 3" – common cheeseweed, chickweed, groundsel; 6" – London rocket, shepherdspurse.</p>	

9.4 - FALLOW SYSTEMS		
LABELED CROPS: This product may be applied during the fallow period prior to planting or emergence of any crop on this label.		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
	<p>12 fl oz (0.38 lb ae) of this product plus the labeled rate of oxyfluorfen per acre will control the following weeds with the maximum height or length indicated: 6" – common cheeseweed, groundsel, marestail (<i>Conyza canadensis</i>), 12" – chickweed, London rocket, shepherdspurse.</p> <p>PRECAUTION: Some crop injury may occur if dicamba is applied within 45 days of planting.</p>	
Aid-to-Tillage	<p>This product may be used in conjunction with tillage practices in fallow systems or pre-plant to labeled crops to control downy brome, cheat, volunteer wheat, tansy mustard and foxtail. Apply 9 fl oz (0.28 lb ae) of this product in 3 to 10 gallons of water per acre. Make applications before weeds are 6 inches in height. Application must be followed by conventional tillage practices no later than 15 days after treatment and before regrowth occurs.</p> <p>PRECAUTION: Tank mixtures with residual herbicides may result in reduced performance. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.</p>	Allow at least 1 day after application before tillage.

9.5 - GRAIN SORGHM (Milo)		
LABELED CROPS: Grain Sorghum (Milo)		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
See Section 9.0	See Use Directions in Section 9.0	See Section 9.0
Pre-Plant, Pre-Emergence, At-Planting	<p>This product may be applied alone or in tank mixture before, during or after planting grain sorghum. Applications must be made prior to emergence of the crop.</p> <p>TANK MIXTURES: Apply these tank mixtures in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.</p> <p style="text-align: center;">S-metolachlor</p> <p>atrazine atrazine + S-metolachlor</p> <p>For difficult-to-control annual weeds such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply this product at 1.5 pints (0.75 lb ae) per acre in these tank mixtures. For other labeled annual weeds, apply 18-24 fl oz (0.56-0.75 lb ae) of this product per acre when weeds are less than 6 inches tall, and 1.5 – 2.25 pints (0.75-1.13 lbs ae) when weeds are over 6 inches tall. When using nitrogen solutions as the carrier, the 2.25 pints (1.13 lbs ae) use rate may need to be increased for acceptable weed control.</p>	<p>For spot treatment, do not treat more than 10 percent of the total field area to be harvested.</p> <p>The crop receiving spray in the treated area will be killed. Do not spray or allow drift outside target area for the same reason.</p> <p>Pre-harvest Interval (PHI): For wiper applicators, allow at least 40 days between application and harvest.</p> <p>Do not use roller applicators.</p> <p>Do not feed or graze treated milo fodder.</p> <p>Do not ensile treated vegetation.</p>
Spot Treatment, Over-the-Top Wiper Applications	<p>This product may be applied as a spot treatment in grain sorghum. Make spot treatments before heading of milo.</p> <p>This product may be applied with wiper applicators to control or suppress the weeds listed under "WIPER APPLICATORS" in the "SELECTIVE EQUIPMENT" section of this label.</p>	
Hooded Sprayers	<p>This product may be used through hooded sprayers for weed control between the rows of milo. Only hooded sprayers that completely enclose the spray pattern may be used. See additional instruction for the use of hooded sprayers in the "APPLICATIONS EQUIPMENT AND TECHNIQUES" section of this label.</p> <p>Treat before milo sends tillers between the drill rows. If such tillers are contacted with the spray solution, the main plant may be killed. Contact of this product in any manner to any vegetation to which treatment is not</p>	<p>Milo must be at least 12 inches tall, measured without extending leaves.</p> <p>Do not graze or feed milo forage or fodder following applications of this product through hooded sprayers.</p> <p>Do not apply more than 1.5 pints (0.75 lb ae) per acre per application.</p> <p>Do not apply more than 2.25 quarts (2.25</p>

9.5 - GRAIN SORGHM (Milo)		
LABELED CROPS: Grain Sorghum (Milo)		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
	intended may cause damage. To the extent consistent with applicable law, such damage is the responsibility of the applicator. Crop injury may occur when the foliage of treated weeds comes into direct contact with leaves of the crop. Droplets, mist, foam, or splatter of the herbicide solution may contact the crop and cause discoloration, stunting or destruction.	lbs ae) per acre per year. Do not apply this product when the leaves of the crop are growing in direct contact with weeds to be treated.
Pre-harvest	Make applications at 30% grain moisture or less. As with other herbicides that cause sudden plant death, avoid pre-harvest applications of this product to milo infected with charcoal rot as lodging can occur.	Do not apply more than 3 pints (1.5 lbs ae) per acre per application. Pre-harvest Interval (PHI): Allow a minimum of 7 days between application and harvest of sorghum. Do not make applications to sorghum grown for seed. Not Registered for Use on pre-harvest grain sorghum (milo) by California..
Post-harvest	This product may be applied after harvest of grain sorghum. A 2.25 pints (1.13 lbs ae) rate may be required for control of large weeds which were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used. This product may be applied to grain sorghum (milo) stubble following harvest to suppress or control regrowth. Apply 1.5 pints (0.75 lb ae) of this product per acre for control, or 18 fl oz (0.56 lb ae) of this product per acre for suppression.	Pre-harvest Interval (PHI): Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

9.6 - HERBS AND SPICES		
LABELED CROPS: Allspice, Angelica, Star anise, Annatto (seed), Balm, Basil, Borage, Burnet, Chamomile, Caper buds, Caraway, Black caraway, Cardamom, Cassia bark, Cassia buds, Catnip, Celery seed, Chervil (dried), Chive, Chinese chive, Cilantro (seed), Cinnamon, Clary, Clove buds, Coriander leaf (cilantro or Chinese parsley), Coriander seed (cilantro), Costmary, Culantro (leaf), Cumin, Curry (leaf), Dill (dillweed), Dill (seed), Epazote, Fennel seed (common and Florence), Fenugreek, White ginger flower, Grains of paradise, Horehound, Hyssop, Juniper berry, Lavender, Lemongrass, Lovage (leaf and seed), Mace, Marigold, Marjoram (including oregano), Mexican oregano, Miaga flower, Mustard (seed), Nasturtium, Nutmeg, Parsley (dried), Pennyroyal, Pepper (black and white), Pepper leaves, Peppermint, Perilla, Poppy (seed), Rosemary, Rue, Saffron, Sage, Savory (summer and winter), Spearmint, Stevia leaves, Sweet bay, Tansy, Tarragon, Thyme, Vanilla, Wintergreen, Woodruff, Wormwood.		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
See Section 9.0	See Use Directions in Section 9.0 PRECAUTION: This product could cause crop injury. When applying this product prior to transplanting or direct-seeding crops into plastic mulch, care must be taken to remove product residues from the plastic prior to planting. Residual product can be removed by a single 0.5-inch application of water, either by natural rainfall or via a sprinkler system. Care must be taken to ensure that the washwater flushes off the plastic mulch and does not enter transplant holes. Applications made at emergence will result in injury or death to emerged seedlings. For some crops below, applications must be made 3 days before transplanting or planting.	See Section 9.0
Over-the-Top Wiper Application, Spot Treatment (Peppermint and Spearmint only)	This product may be applied as a spot treatment or over the top of peppermint or spearmint with wiper applications in spearmint and peppermint. Apply spot treatments on a spray-to-wet basis with hand-held equipment, such as backpack sprayers, pump-up pressure sprayers, hand-guns, hand-wands or any other hand-held or motorized spray equipment	Pre-harvest Interval (PHI): Allow at least 7 days between application and harvest. In spot treatment applications, no more than 10 percent of the total field area to be harvested can be treated at one time. The crop receiving spray in the treated area will be killed. Do not spray or allow drift outside target area

9.6 - HERBS AND SPICES

<p>LABELED CROPS: Allspice, Angelica, Star anise, Annatto (seed), Balm, Basil, Borage, Burnet, Chamomile, Caper buds, Caraway, Black caraway, Cardamom, Cassia bark, Cassia buds, Catnip, Celery seed, Chervil (dried), Chive, Chinese chive, Cilantro (seed), Cinnamon, Clary, Clove buds, Coriander leaf (cilantro or Chinese parsley), Coriander seed (cilantro), Costmary, Culantro (leaf), Cumin, Curry (leaf), Dill (dillweed), Dill (seed), Epazote, Fennel seed (common and Florence), Fenugreek, White ginger flower, Grains of paradise, Horehound, Hyssop, Juniper berry, Lavender, Lemongrass, Lovage (leaf and seed), Mace, Marigold, Marjoram (including oregano), Mexican oregano, Miaga flower, Mustard (seed), Nasturtium, Nutmeg, Parsley (dried), Pennyroyal, Pepper (black and white), Pepper leaves, Peppermint, Perilla, Poppy (seed), Rosemary, Rue, Saffron, Sage, Savory (summer and winter), Spearmint, Stevia leaves, Sweet bay, Tansy, Tarragon, Thyme, Vanilla, Wintergreen, Woodruff, Wormwood.</p>		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
	<p>used to direct the spray solution to a limited area.</p> <p>In wiper applications, the applicator must be adjusted so that the wiper contact point is at least 2 inches above the crop. Weeds should be a minimum of 6 inches taller than the crop.</p> <p>PRECAUTION: Contact of the herbicide solution with the crop may result in discoloration, stunting, or destruction.</p> <p>Further applications may be made in the same area at 30-day intervals.</p>	for the same reason.

9.7 OIL SEED CROPS

<p>LABELED CROPS: Borage, Buffalo gourd (seed), Canola (non-Roundup Ready), Crambe, Flax, Jojoba, Lesquerella, Meadowfoam, Mustard (seed), Rape, Safflower, Sesame, Sunflower.</p>		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
See Section 9.0	<p>See Use Directions in Section 9.0</p> <p>This product may be applied before, during or after planting oil seed crops listed in this section. Broadcast applications must be made prior to crop emergence. Wiper applications or hooded sprayers may be used between the rows once the crop is established.</p> <p>TANK MIXTURES: For sunflowers, a tank mixture with pendimethalin may be applied according to the labeled directions for that product.</p>	<p>See Section 9.0</p> <p>For use with canola, do not apply more than 1.5 quarts (1.5 lbs ae) per acre per application.</p> <p>For use with sunflowers, do not apply more than 1.5 pints (0.75 lb ae) per acre as a single pre-plant or pre-emergence application.</p> <p>Do not feed or graze sunflower forage following application of this product.</p>
Pre-Harvest (Sunflower & safflower)	<p>This product provides weed control when applied as a harvest aid to a physiologically mature crop prior to harvest of sunflower or safflower.</p> <p>For safflower, apply when seed has lost its opaque character, approximately 20 to 30 days after the end of flowering of the secondary branches.</p> <p>For sunflower, apply when the backsides of sunflower heads are yellow and bracts are turning brown and seed moisture content is less than 35%.</p>	<p>Pre-harvest Interval (PHI): Allow a minimum of 7 days between treatment and harvest or livestock feeding.</p> <p>Do not apply more than 72 fl oz (2.25 lbs ae) per acre at a pre-harvest timing to safflower.</p> <p>Do not apply more than 1.5 pints (0.75 lb ae) per acre at a pre-harvest timing to sunflower.</p>
Post-Harvest (Sunflower & safflower)	<p>This product may be applied after harvest of safflower or sunflower.</p> <p>A 2.25 pints (1.13 lbs ae) per acre rate may be required for control of large weeds, which are growing in the crops at the time of harvest.</p> <p>Tank mixtures with 2,4-D or dicamba may be used. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.</p>	<p>Pre-harvest Interval (PHI): Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.</p> <p>Applications must be made at least 30 days prior to planting any crop not listed on the AgSaver™ Glyphosate 5.4 Plus label booklet.</p>

9.8 - SOYBEANS																													
LABELED CROPS: Soybeans (non-Roundup Ready)																													
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS																											
See Section 9.0	See Use Directions in Section 9.0	See Section 9.0																											
Pre-Plant, Pre-Emergence, At-Planting	<p>This product may be applied before, during or after planting soybeans. Applications must be made prior to emergence of the crop.</p> <p>Products that can be tank mixed with AgSaver™ Glyphosate 5.4 Plus are in the table below. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.</p> <p>This product may be tank-mixed with 2,4-D or 2,4-DB. See the 2,4-D label for intervals between application and planting.</p> <p>For difficult-to-control weeds such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply this product at 1.5 pints (0.75 lb ae) per acre in these tank mixtures. For other labeled annual weeds, apply 18-24 fl oz (0.56-0.75 lb ae) of this product per acre when weeds are less than 6 inches tall, and 1.5-2.25 pints (0.75-1.13 lbs ae) when weeds are over 6 inches tall.</p> <p>TANK MIXES:</p> <table> <tr> <td>carfentrazone-ethyl</td><td>fomesafen</td><td>metribuzin + S-</td></tr> <tr> <td>clomazone</td><td>imazaquin</td><td>metolachlor</td></tr> <tr> <td>cloransulam</td><td>imazethapyr</td><td>pendimethalin</td></tr> <tr> <td>dimethenamid</td><td>imazethapyr +</td><td>quizalofop-P-ethyl</td></tr> <tr> <td>fluzifop-P-butyl +</td><td>pendimethalin</td><td>S-metolachlor</td></tr> <tr> <td>fenoxaprop</td><td>linuron</td><td>sulfentrazone</td></tr> <tr> <td>flufenacet +</td><td>metribuzin</td><td>sulfentrazone +</td></tr> <tr> <td>metribuzin</td><td>metribuzin +</td><td>cloransulam-methyl</td></tr> <tr> <td>flumioxazin</td><td>chlorimuron-ethyl</td><td></td></tr> </table>	carfentrazone-ethyl	fomesafen	metribuzin + S-	clomazone	imazaquin	metolachlor	cloransulam	imazethapyr	pendimethalin	dimethenamid	imazethapyr +	quizalofop-P-ethyl	fluzifop-P-butyl +	pendimethalin	S-metolachlor	fenoxaprop	linuron	sulfentrazone	flufenacet +	metribuzin	sulfentrazone +	metribuzin	metribuzin +	cloransulam-methyl	flumioxazin	chlorimuron-ethyl		Not Registered for Use in tank mixtures on soybeans by California.
carfentrazone-ethyl	fomesafen	metribuzin + S-																											
clomazone	imazaquin	metolachlor																											
cloransulam	imazethapyr	pendimethalin																											
dimethenamid	imazethapyr +	quizalofop-P-ethyl																											
fluzifop-P-butyl +	pendimethalin	S-metolachlor																											
fenoxaprop	linuron	sulfentrazone																											
flufenacet +	metribuzin	sulfentrazone +																											
metribuzin	metribuzin +	cloransulam-methyl																											
flumioxazin	chlorimuron-ethyl																												
Spot treatment	<p>For spot treatments, apply this product prior to initial pod set in soybeans.</p> <p>The crop receiving spray in the treated area will be killed.</p>	<p>Do not treat more than 10 percent of the total field area to be harvested.</p> <p>Do not spray or allow drift outside target area for the same reason.</p>																											
Pre-harvest	<p>This product provides weed control when applied prior to harvest of soybeans.</p> <p>Apply at rates given in the annual, perennial, and woody brush tables.</p> <p>This product may be applied using either aerial or ground spray equipment.</p> <p>Apply after pods have set and lost all green color. Care should be taken to avoid excessive seed shatter loss due to ground application equipment.</p>	<p>Do not apply more than 3.75 quarts (3.75 lbs ae) per acre for pre-harvest applications.</p> <p>Do not apply more than 3 pints (1.5 lb ae) per acre per application of this product by air.</p> <p>Pre-harvest Interval (PHI): Allow a minimum of 7 days between application and harvest of soybeans.</p> <p>Do not graze or harvest treated hay or fodder for livestock feed within 25 days of last pre-harvest application. (If the application rate is 1.5 pints (0.75 lb ae) per acre or lower, the grazing restriction is reduced to 14 days after the last pre-harvest application.)</p> <p>Do not apply to soybeans grown for seed.</p>																											
Selective equipment	<p>This product may be applied through shielded applicators, hooded sprayers, wiper applicators or sponge bars in soybeans.</p> <p>See the "Selective Equipment" part of the "Application and Equipment Techniques" section of this label for information on proper use and calibration of this equipment.</p>	Pre-harvest Interval (PHI): Allow at least 7 days between application and harvest.																											

9.9 – SUGARCANE		
LABELED CROPS: Sugarcane		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
See Section 9.0	See Use Directions in Section 9.0	See Section 9.0
Pre-plant, Pre-emergence, At-planting	This product may be applied in or around sugarcane fields or in fields prior to the emergence of plant cane.	Do not apply to vegetation in or around ditches, canals or ponds containing water to be used for irrigation.
Spot Treatment	This product may be applied as a spot treatment in sugarcane. For control of volunteer or diseased sugarcane, make a 1 percent solution of this product in water and spray to wet the foliage of vegetation to be controlled. Volunteer or diseased sugarcane should have at least 7 new leaves. Avoid spray contact with healthy cane plants since severe damage or destruction may result.	Do not feed or graze treated sugarcane foliage following application.
Fallow treatments	This product may be used as a replacement for tillage in fields that are lying fallow between sugarcane crops. This product may also be used to remove the last stubble of ratoon cane. For removal of last stubble of ratoon cane, apply 3 to 3.75 quarts (3-3.75 lbs ae) of this product in 10 to 40 gallons of water per acre to new growth having at least 7 new leaves. Ground or aerial application equipment may be used. Applications up to 4.5 pints (2.25 lbs ae) per acre may be made by aerial application in fallow sites where there is sufficient buffer to prevent injury due to drift onto adjacent crops. Tank mixtures with 2,4-D and dicamba may be used. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.	Do not apply more than 4.5 pints (2.25 lbs ae) per acre per aerial application. Allow 7 or more days after application before tillage.
Hooded sprayers	This product may be used through hooded sprayers for weed control between the rows of sugarcane. See Section 8.0 for "Application Equipment and Techniques" for additional USE DIRECTIONS. Minimize the potential for spray particles to escape from under the hood by operating the sprayer at appropriate ground speeds, nozzle pressures and wind speeds. Operation on rough or sloping ground may result in spray particles escaping from the hood. When applying to sugarcane that is grown on raised beds, ensure that the hood is designed to completely enclose the spray. If necessary, extend the front and rear flaps of the hoods to reach the ground in furrows between the rows. PRECAUTION: Droplets, mist, foam, or splatter of the herbicide solution settling on the crop may result in discoloration, stunting or destruction. To the extent consistent with applicable law, such damage shall be the sole responsibility of the applicator.	Do not allow treated weeds to come into contact with the crop.
FOR AID IN SUGARCANE RIPENING (FLORIDA, HAWAII, LOUISIANA, PUERTO RICO AND TEXAS)	This product is a foliar-applied plant growth regulator to hasten ripening and increase the level of sucrose in sugarcane. It is effective in both low and high-tonnage sugarcane. When applied as directed under the conditions described, this product will hasten ripening and extend the period of high sucrose level in sugarcane. As a result of leaf desiccation, improved trash burn can be expected. Most of the sucrose increase is concentrated in the top nodes of the treated cane stalk. In order to recover the maximum sugar where topping is practiced during harvest, top at the base of the fourth leaf. Prior to application, consult your state sugarcane authority or local AgSaver™, LLC representative regarding the degree of sucrose response anticipated from the variety of sugarcane to be treated. APPLICATION RATES: Use the following application rates and timing instructions according to the State in which the sugarcane is grown. PRECAUTION: Application of this product may initiate development of shooting eyes. This product may not increase the sucrose content of sugarcane under conditions of good nature ripening. Within 2 to 3 weeks after application, this product may produce a slight yellowing to pronounced browning and drying of leaves, and a shortening of upper internodes. Spindle death may occur.	Do not apply more than 9 fl oz (0.28 lb ae) per acre per application in Florida, Louisiana, and Texas. Do not apply more than 15.75 fl oz (0.49 lb ae) per acre per application in Hawaii. Do not apply more than 3.75 fl oz (0.12 lb ae) per acre per application in Puerto Rico. Do not make application to sugarcane grown for seed. Do not feed or graze treated sugarcane forage following application. Do not apply for enhanced ripening to any crops other than sugarcane. Do not plant subsequent crops in treated fields other than the following for 30 days after application: alfalfa or other forage

9.9 – SUGARCANE		
LABELED CROPS: Sugarcane		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
	<p>Rainfall within 6 hours after application may reduce effectiveness.</p> <p>NOTE: Use the higher rate within the labeled range when treating sugarcane under adverse ripening conditions or when less responsive varieties are to be treated.</p> <p>FLORIDA – Apply 3.75-9 fl oz (0.12-0.28 lb ae) of this product per acre 3 to 6 weeks before harvest of LAST RATTON CANE ONLY.</p> <p>HAWAII – Apply 6.75-15.75 fl oz (0.21-0.49 lb ae) of this product per acre 4 to 10 weeks before harvest.</p> <p>LOUISIANA – Apply 2.6-9 fl oz (0.08-0.28 lb ae) of this product per acre 3 to 7 weeks before harvest of RATOON CANE ONLY.</p> <p>PUERTO RICO – Apply 3.75 fl oz (0.12 lb ae) of this product per acre 3 to 5 weeks before harvest of RATOON CANE ONLY.</p> <p>TEXAS – Apply 3.75-9 fl oz (0.12-0.28 lb ae) of this product per acre 3 to 5 weeks before harvest of RATOON CANE ONLY.</p>	legumes, beans (all types), corn (all types), cotton, melons (all types), pasture grasses, peanuts, potatoes (Irish or sweet), sorghum (milo), soybeans, squash (all types) or wheat.

9.10 - VEGETABLE CROPS

This “VEGETABLE CROPS” section gives directions that apply to all listed vegetable crops within section 9.10 grouped alphabetically below. See the individual crop categories for specific instructions, preharvest intervals, precautions, and restrictions.

TYPES OF APPLICATIONS: Chemical Fallow, Preplant Fallow Beds, Preplant, Preemergence, Prior to Transplanting Vegetables, At-Planting, Hooded Sprayers in Row Middles, Shielded Sprayers in Row Middles, Wiper Applications in Row Middles, and Post Harvest, Directed Applications (Non-Bearing Ginseng), Over-the-top Wiper Applications (Rutabagas Only).

PRECAUTIONS:

- When applying this product prior to transplanting or direct-seeding crops into plastic mulch, care must be taken to remove residues of this product, which could cause crop injury, from the plastic prior to planting.
- Residues can be removed by single 0.5-inch application of water, either by natural rainfall or via a sprinkler system. Care must be taken to ensure that the wash water flushed off the plastic mulch and does not enter transplant holes.
- Applications made at emergence will result in injury or death to emerged seedlings.
- Apply before seed germination in coarse sandy soils to further minimize the risk of injury.
- Avoid contact of herbicide with foliage, green shoots or stems, bark, exposed roots (including those emerging from plastic mulch), or fruit of crops because severe injury or destruction may result.

RESTRICTIONS:

- When making pre-emergence and at planting applications, applications must be made before crop emergence to avoid severe crop injury.
- In crops with vines, hooded sprayer, shielded sprayer and wiper applications to row middles must be made prior to vine development to prevent severe injury or destruction.
- Pre-Harvest Interval: Unless otherwise specified in this product’s labeling, treatments with selective equipment including wipers and hooded sprayers must be made at least 14 days prior to harvest.
- Post-harvest or fallow applications must be made at least 30 days prior to planting any non-labeled crop.

See “Application Equipment and Techniques” section of this label for additional information.

9.10.1 – BRASSICA VEGETABLES		
LABELED CROPS: Broccoli, Broccoli (raab), Brussels sprouts, Cabbage, Cabbage (Chinese), Cabbage (Chinese mustard), Cauliflower, Cavalo broccolo, Chinese broccoli (gai lon), Chinese cabbage (bok choy & napa), Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
See Section 9.10	See Use Directions under Section 9.0	See Sections 9.0 and 9.10

9.10.2 – BULB VEGETABLES		
LABELED CROPS: Garlic, Great-headed garlic, Leek, Onion (dry bulb & green), Shallot, Welsh onion, Shallot		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
See Section 9.10	See Use Directions under Section 9.0	See Sections 9.0 and 9.10

9.10.3 – CUCURBIT VEGETABLES & FRUITS		
LABELED CROPS: Chayote (fruit), Chinese waxgourd, Citron melon, Cucumber, Gherkin, Gourds, Gourds (edible including hyotan, cucuzza, hechima, Chinese okra), Melons (All), <i>Momordica</i> spp. (includes balsam apple, balsam pear, bittermelon, Chinese cucumber), Muskmelon (cantaloupe, casaba, crenshaw, golden pershaw, honeydew, honey ball, mango melon & Persion, pineapple, Santa Claus, snake), Pumpkin, Summer Squash (including crookneck, scallop, straightneck, vegetable marrow, zucchini) Winter squash (including butternut, calabaza, hubbard, acorn, spaghetti), Watermelon		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
See Section 9.10	See Use Directions under Section 9.0	See Sections 9.0 and 9.10 For Cantaloupe, Casaba melon, Crenshaw melon, Cucumber, Gherkin, Gourds, Honeydew melon, Honey ball melon, Mango melon, Melons (all) Muskmelon, Persian melon, Pumpkin, Squash (summer & winter), AND Watermelon, allow at least 3 days between application and planting.

9.10.4 – LEAFY VEGETABLES		
LABELED CROPS: Amaranth (Chinese spinach), Arrugula (roquette), Beet greens, Cardoon, Celery, Celery (Chinese), Celtuce, Chaya, Chervil, Chrysanthemum (edible leaved), Chrysanthemum (Garland), Corn salad, Cress (garden & upland), Dandelion, Dock (sorrel), Dokudami, Endive (escarole), Fennel (Florence), Gow kee, Lettuce (head & leaf), Orach, Parsley, Purslane (garden & winter), Radicchio (red chicory), Rhubarb, Spinach (All), Swiss Chard, Watercress (upland), Water Spinach		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
See Section 9.10	See Use Directions under Section 9.0 For Watercress, avoid application within 3 days prior to seeding and during the period between seeding and emergence to minimize the risk of injury.	See Sections 9.0 and 9.10

9.10.5 - FRUITING VEGETABLES		
LABELED CROPS: Eggplant, Ground cherry (<i>Physalis</i> spp.), Pepino, Pepper (includes bell, chili, cooking, pimento, sweet), Tomatillo, Tomato		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
See Section 9.10	See Use Directions under Section 9.0	See Sections 9.0 and 9.10 For Eggplant, Ground cherry, Pepino, Pepper (all), Tomatillo and Tomato, allow at least 3 days between application and planting. Do not use hooded or shielded sprayer applications in row middles of tomatoes.

9.10.6 – LEGUME VEGETABLES (succulent or dried)		
<p>LABELED CROPS: Bean (Lupinus: includes grain lupin, sweet lupin, white lupin, and white sweet lupin), Bean (Phaseolus: includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean), Bean (Vigna: includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean), Broad bean (fava), Chickpea (garbanzo), Guar, Jackbean, Lablab bean, Lentil, Pea (Pisum: includes dwarf pea, edible podded pea, English pea, field pea, garden pea, green pea, snowpea, sugar snap pea), Pigeon pea, Soybean (immature seed), Sword bean</p>		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
See Section 9.10	See Use Directions under Section 9.0	See Sections 9.0 and 9.10
Pre-harvest broadcast spray (Dry beans)	This product may be applied as an over the top broadcast spray to control labeled weeds prior to the harvest of dry beans. Apply up to 1.5 pints (0.75 lb ae) stage of the legume seed (30 percent grain moisture or less). Either ground broadcast or aerial applications may be made.	Do not apply more than 1.5 pints (0.75 lb ae) per acre per application for preharvest broadcast spray on dry beans. Do not apply more than 3 pints (1.5 lbs ae) per acre per application for pre-harvest broadcast spray on dry peas, lentils, and chickpeas.
Pre-harvest broadcast spray (Dry Peas, Lentils & Chickpeas)	This product may be applied as an over the top broadcast spray to control labeled weeds prior to the harvest of dry peas, lentils, and chickpeas. Apply up to 3 pints (1.5 lbs ae) in 3 to 20 gallons of water per acre at the hard dough stage of the legume seed (30 percent grain moisture or less). Either ground broadcast or aerial applications may be made.	Do not apply more than 1.5 pints (0.75 lb ae) per acre per year for preharvest broadcast spray on dry beans. Do not apply more than 3 pints (1.5 lbs ae) per acre per year for pre-harvest broadcast spray on dry peas, lentils, and chickpeas. Pre-harvest Interval (PHI): Apply at least 7 days before harvest for Dry Beans, Dry Peas, Lentils & Chickpeas. Do not combine a pre-harvest spray with a spot treatment on the same crop area. Do not feed treated vines and hay from these crops to livestock. Do not apply this product through any type of irrigation system. Do not treat field (feed) peas since these are considered to be grown as livestock feed. Do not make pre-harvest applications to Dry Beans, Dry Peas, Lentils & Chickpeas grown for seed.
Spot treatment (Dry beans, Dry peas, Lentils, Chickpeas)	This product may be applied as spot treatment to control troublesome weeds such as Canada thistle, quackgrass, mayweed (dog fennel), and milkweed in dry beans. Apply up to 19.5 fl oz (0.61 lb ae) in 10 to 20 gallons of water through ground spray equipment or use a 2 percent solution in a handheld sprayer. For best results, applications should be made at or beyond the bud stage of growth. The crop receiving spray in treated areas will be killed.	Do not apply more than 19.5 fl oz (0.61 lb ae) per acre per year. Do not apply more than 19.5 fl oz (0.61 lb ae) per acre per single application. Pre-harvest Interval (PHI): Apply at least 14 days before harvest. Only one application per year may be made. Do not combine a pre-harvest spray with a spot treatment on the same crop area. Do not feed treated vines and hay from these crops to livestock. Do not apply this product through any type of irrigation system. Do not treat field cowpeas, since these are considered to be grown as livestock feed.

9.10.7 – ROOT & TUBER VEGETABLES		
<p>LABELED CROPS: Arracacha, Arrowroot, Artichoke (Chinese & Jerusalem), Beet (garden), Burdock, Canna, Carrot, Cassava (bitter & sweet), Celeriac, Chayote (root), Chervil, Chicory, Chufa, Dasheen, Galangal, Ginger, Ginseng, Horseradish, Leren, Kava, Parsley, Parsnips, Potato (Irish), Radish, Radish (Oriental), Rutabaga, Salsify, Salsify (Black & Spanish), Skirret, Sweet potato, Tanier, Tumeric, Turnip, Wasabi, Yacon, Yams, Yam bean, Yam (True)</p>		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
See Section 9.10	See Use Directions under Section 9.0	See Sections 9.0 and 9.10
Direct Application (Non-bearing Ginseng)	<p>This product may be used for weed control in established non-bearing ginseng. Applications may be made with boom equipment, CDA, shielded sprayers, hand-held and high volume wands, lances, and orchard guns or with wiper application equipment.</p> <p>PRECAUTION: Extreme care must be exercised to avoid contact of herbicide solution, spray, drift or mist with foliage or green bark of trunk, branches, suckers, fruit or other parts of desirable plants. Contact of this product with other than matured</p>	Pre-harvest Interval (PHI): Applications must be made at least one year prior to harvest.

9.10.7 – ROOT & TUBER VEGETABLES

Labeled Crops: Arracacha, Arrowroot, Artichoke (Chinese & Jerusalem), Beet (garden), Burdock, Canna, Carrot, Cassava (bitter & sweet), Celeriac, Chayote (root), Chervil, Chicory, Chufa, Dasheen, Galangal, Ginger, Ginseng, Horseradish, Leren, Kava, Parsley, Parsnips, Potato (Irish), Radish, Radish (Oriental), Rutabaga, Salsify, Salsify (Black & Spanish), Skirret, Sweet potato, Tanier, Tumeric, Turnip, Wasabi, Yacon, Yams, Yam bean, Yam (True)		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
	brown bark can result in serious crop damage.	
Over-the-Top Wiper Application (Rutabaga Only)	Wiper applicators may be used over-the-top of rutabagas.	Pre-harvest Interval (PHI): Allow at least 14 days between application and harvest of rutabagas.

9.11 - MISCELLANEOUS CROPS

Labeled Crops: Aloe vera, Asparagus, Bamboo shoots, Globe artichoke, Okra, Peanut (ground nut), Pineapple, Strawberry, Sugar Beet (non-Roundup Ready)		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
See Section 9.10	See Use Directions under Section 9.0	See Sections 9.0 and 9.10
	Avoid contact of herbicide with foliage, green shoots, or stems. Bark, exposed roots (including those emerging from plastic mulch), or fruit of crops because severe injury or destruction may result.	When making pre-emergence and at planting applications, applications must be made before crop emergence to avoid serious crop injury. Apply before seed germination in coarse sandy soils to further minimize the risk of injury. In crops with vines, hooded sprayer, shielded sprayer and wiper applications to row middles must be made prior to vine development. Treatments with selective equipment including wipers and hooded sprayers must be made at least 14 days prior to harvest. Post harvest or fallow applications must be made at least 30 days prior to planting any non-labeled crop.
Weed control, Site preparation	This product may be applied for weed control or for site preparation prior to planting or transplanting crops listed in this section. When applying this product prior to transplanting or direct seeding crops into plastic mulch, care must be taken to remove residues of this product from the plastic prior to transplanting. Residues can be removed by 0.5 inch natural rainfall or by applying water via a sprinkler system. Care must be taken to ensure that the wash water flushes off the plastic mulch and does not enter transplant holes. Injury made at emergence will result in injury or death to emerged seedlings.	Do not apply within a week before the first asparagus spears emerge. Do not feed or graze treated pineapple forage following application.
Spot treatment (Asparagus)	This product may be applied immediately after cutting, but prior to the emergence of new spears.	Do not treat more than 10 percent of the total field area to be harvested. Pre-harvest Interval (PHI): Do not harvest within 5 days of treatment.
Post-harvest (Asparagus)	This product may be applied after the last harvest and all spears have been removed. If spears are allowed to regrow, delay application until ferns have developed. Delayed treatments should be applied as a directed or shielded spray in order to avoid contact of the spray with ferns, stems or spears. Select and use recommended types of spray equipment for post-emergence postharvest applications. A directed spray is any application where the spray pattern is aligned in such a way as to avoid direct contact of the spray with the crop. A shielded spray is any application where a physical barrier is positioned and maintained between the spray	Do not allow direct contact of the spray with the asparagus which will result in serious crop injury.

9.11 - MISCELLANEOUS CROPS

LABELED CROPS: Aloe vera, Asparagus, Bamboo shoots, Globe artichoke, Okra, Peanut (ground nut), Pineapple, Strawberry, Sugar Beet (non-Roundup Ready)

and the crop to prevent contact of spray with the crop.

10.0 - TREE, VINE, & SHRUB CROPS (Alphabetical)

This section gives directions that apply to all listed tree, vine & shrub crops within section 10 crop groups. Individual crops may have more specific instructions, pre-harvest intervals, precautions, and restrictions.

TYPES OF APPLICATIONS: Pre-plant (Site Preparation) Broadcast Sprays, Weed control, Middles (between rows of trees, vines or shrubs), Strips (within rows of trees, vines or shrubs), Selective Equipment (shielded sprayers, wiper treatments), Directed Sprays, Spot Treatments, Perennial Grass Suppression, Cut Stump.

Applications may be made with boom equipment, CDA, shielded sprayers, hand-held and high-volume wands, lances, orchard guns or with wiper applicator equipment, except as directed.

USE DIRECTIONS:

This product may be applied in middles (between rows of trees or vines), strips (within rows of trees or vines), and for weed control or perennial grass suppression in established tree fruit and tree nut groves, orchards, berries, and vineyards. This product may also be used for site preparation prior to planting or transplanting these crops. Apply at 12 – 120 fl oz (0.38-3.75 lbs ae) per acre according to the “Annual Weeds” and “Perennial Weeds” rate tables sections of this label. Utilize rates at the higher end of the labeled rate range when weeds are stressed, growing in dense populations or are greater than 12 inches tall. Repeat applications may be made up to a maximum of 7.8 quarts (7.8 lbs ae) per acre per year.

The maximum use rates stated throughout this product’s labeling apply to this product combined with the use of all other herbicides containing glyphosate or sulfosate as the active ingredient, whether applied as mixtures or separately. Calculate the application rates and ensure that the total use of this and other glyphosate or sulfosate containing products does not exceed stated maximum use rate.

PRECAUTIONS:

- Extreme care must be exercised to avoid contact of herbicide solution, spray, drift or mist with foliage or green bark of trunk, branches, suckers, fruit or other part of the trees, canes, and vines.
- Avoid applications when recent pruning wounds or other mechanical injury has occurred.
- Contact of this product other than matures brown bark can result in serious crop damage or destruction.
- For applications in strips (within rows of trees), only selective equipment (directed sprays, hooded sprayers, shielded applicators, or wipers) must be used to minimize the potential for leakage or drift of herbicide sprays onto crop.

See “APPLICATION EQUIPMENT AND TECHNIQUES” section of this label for additional directions and precautions.

RESTRICTIONS:

- **DO NOT** apply more than 7.5 pints (3.75 lbs ae) per acre per application, unless otherwise specified for an individual use.
- **DO NOT** apply more than 7.8 quarts (7.8 lbs ae) per acre per year as a combined total of all treatments for applications in tree, vine, or shrub crops, unless otherwise specified for an individual use.
- For suppression of vegetative growth and seedhead inhibition of bahiagrass up to 120 days, do not apply more than 3 fl oz (0.09 lb ae) per acre per application.
- For suppression of vegetative growth and seedhead inhibition of bahiagrass up to 120 days, do not apply more than 6 fl oz (0.19 lb ae) per acre per application.
- Only wipers or shielded applicators capable of preventing all contact with crop may be used.

- Only shielded or directed sprayers may be used in crops with potential for crop contact, and then only where there is sufficient clearance.
- For berry crops, hooded or shielded sprayers must be fully enclosed including top, sides, front and back.
- Allow a minimum of 3 days between applications and transplanting.

Middles (between rows of trees, vines, or bushes)

USE DIRECTIONS: This product will control or suppress annual and perennial weeds and ground covers growing between the rows of labeled tree and vine crops. If weeds are under drought stress, irrigate prior to application. Reduced control may result if weeds have been mowed prior to application.

TANK MIXTURES: It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

A tank mixture of this product plus oxyfluorfen may be used for annual weeds in middles between rows of citrus crops, tree fruits, tree nuts and vine crops. This mixture is recommended when weeds are stressed or growing in dense populations. 12-24 fl oz (0.38-0.75 lb ae)/A of this product plus the labeled rate of oxyfluorfen will control annual weeds with a maximum height or diameter of 6 inches, including crabgrass, common groundsel, junglerice, common lambsquarters, redroot pigweed, London rocket, common ryegrass, shepherdspurse, annual sowthistle, common cheeseweed (malva), filaree (suppression), horseweed/marestail (*Conyza canadensis*), stinging nettle and common purslane (suppression). 12-24 fl oz (0.38-0.75 lb ae)/A of this product plus labeled rate of oxyfluorfen will control common cheeseweed (malva) or hairy fleabane (*Conyza bonariensis*), with a maximum height or diameter of 3 inches.

Strips (in rows of trees, vines, or bushes)

TANK MIXTURES: This product may be applied in rows of tree or vine crops and may also be tank mixed with the following products. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

bromacil + diuron	oryzalin
diuron	oxyfluorfen
napropamide	pendimethalin
norflurazon	simazine

Restriction: Not Registered for Use on above tank mixtures by Puerto Rico.

PERENNIAL GRASS SUPPRESSION

This product will suppress perennial grasses such as bahiagrass, bermudagrass, tall fescue, orchardgrass, Kentucky bluegrass, and quackgrass that are grown as ground covers in tree and vine crops.

For suppression of tall fescue, fine fescue, orchardgrass and quackgrass, apply 6 fl oz (0.19 lb ae) of this product in 10 to 20 gallons of water per acre.

For suppression of Kentucky bluegrass covers, apply 4.5 fl oz (0.14 lb ae) of this product per acre. Do not add ammonium sulfate.

For best results, mow cool season grass covers in the spring to even their height and apply this product 3 to 4 days after mowing.

For suppression of vegetative growth and seedhead inhibition of bahiagrass for approximately 45 days, apply 4.5 fl oz (0.14 lb ae) of this product in 10 to 25 gallons of water per acre. Apply 1 to 2 weeks after full green-up or after mowing to a uniform height of 3 to 4 inches. This application must be made prior to seedhead emergence.

For suppression up to 120 days, apply 3 fl oz (0.09 lb ae) of this product per acre, followed by an application of 1.5 – 3 fl oz (0.05-0.09 lb ae) per acre about 45 days later. Make no more than 2 applications per year.

For burndown of bermudagrass, apply 1.5 – 3 pints (0.75-1.5 lbs ae) of this product in 3 to 20 gallons of water per acre. Use this treatment only if reduction of the bermudagrass stand can be tolerated. When burndown is required prior to harvest, allow at least 21 days to ensure sufficient time for burndown to occur.

For suppression of bermudagrass, apply 4.5-12 fl oz (0.14-0.38 lb ae) of this product per acre east of the Rocky Mountains and 12 fl oz (0.38 lb ae) of this product per acre west of the Rocky Mountains. Apply in a total spray volume of 3 to 20 gallons per acre, no sooner than 1 to 2 weeks after full green-up. If the bermudagrass is mowed prior to application, maintain a minimum of 3 inches in height. Sequential applications up to the maximum labeled rate may be made when regrowth occurs and bermudagrass injury and stand reduction can be tolerated. East of the Rocky Mountains, rates of 4.5 – 7.5 fl oz (0.14-0.23 lb ae) per acre should be used in shaded conditions or where a lesser degree of suppression is desired.

CUT STUMPS (Tree crops)		
LABELED CROPS: <u>Citrus Trees:</u> Calamondin, Chironja, Citron, Citrus hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin (Tangerine), Orange (all), Pummelo, Tangelo, Tangor. <u>Fruit Trees:</u> Apply, Apricot, Cherry (sweet, sour), Crabapple, Loquat, Mayhaw, Nectarine, Olive, Peach, Pear, Plum/Prune (all), Quince. <u>Nut Trees:</u> Almond, Beechnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (hazelnut), Hickory Nut, Macadamia, Pecan, Pistachio, Walnut (black, English).		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
Suitable Hand-held Equipment	<p>Cut stump applications of this product may be made during site preparation or site renovation, prior to transplanting tree crops. This product will control regrowth of cut stumps and resprouts of many types of tree species, some of which are listed below.</p> <p>Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut trees or resprouts close to the soil surface. Apply a 50 to 100 percent solution of this product to the freshly cut surface immediately after cutting. Delays in application may result in reduced performance. For best results, applications should be made during periods of active growth and full leaf expansion.</p> <p>PRECAUTION: Some sprouts, stems, or trees may share the same root system. Adjacent trees having a similar age, height and spacing may signal shared roots. Whether grafted or shared, injury is likely to occur to non-treated stems/trees when one or more trees sharing common roots are treated.</p>	<p>See Section 10.0</p> <p>Do not make cut stump applications when the roots of adjacent desirable trees may be grafted to the roots of the cut stump. Injury resulting from root grating may occur in adjacent trees.</p>

10.1 - BERRY CROPS		
LABELED CROPS: Blackberry (including bingleberry, black satin berry, boysenberry, Cherokee blackberry, chesterberry, Cheyenne blackberry, coryberry, darrowberry, dewberry, Dirksen thronless berry, Himalayaberry, hullberry, juneberry, lavacaberry, lowberry, lucretiaberry, marionberry, nectarberry, olallie berry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, and youngberry), Blueberry, Cranberry, Currant, Elderberry, Gooseberry, Huckleberry, Loganberry, Raspberry (Black, Red), Salai		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
See Section 10.0	See Use Directions under Section 10.0	<p>See Section 10.0</p> <p>Do not permit herbicide solution to contact desirable vegetation, including green shoots, canes, or foliage.</p> <p>Pre-harvest Interval (PHI): Allow a minimum of 14 days between last application and harvest of labeled berry crops.</p> <p>Pre-harvest Interval (PHI): Allow a minimum of 30 days between last application and harvest of cranberries.</p> <p>Do not make directed sprays within the cranberry bush areas prior to berry harvest.</p>
Spot Treatment (Cranberry)	May be used to control weeds growing in dry ditches (interior and perimeter) of cranberry production areas. Handheld sprayers or other appropriate application	Pre-harvest Interval (PHI): Allow a minimum of 30 days between last application and harvest of cranberries.

10.1 - BERRY CROPS		
LABELED CROPS: Blackberry (including bingleberry, black satin berry, boysenberry, Cherokee blackberry, chesterberry, Cheyenne blackberry, coryberry, darrowberry, dewberry, Dirksen thronless berry, Himalayaberry, hullberry, juneberry, lavacaberry, lowberry, lucretiaberry, marionberry, nectarberry, olallie berry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, and youngberry), Blueberry, Cranberry, Currant, Elderberry, Gooseberry, Huckleberry, Loganberry, Rasperry (Black, Red), Salai		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
production)	<p>equipment listed under "Application Equipment and Techniques" in this label may be used. Drop water level to remove standing water in ditches prior to application. In hand-held sprayers, use 1 to 2 percent solution of this product. Spray to wet vegetation, not to run-off.</p> <p>For treatments after draw down of water in dry ditches, allow 2 or more days after treatment before reintroduction of water to achieve maximum weed control.</p> <p>Apply this product within 1 day after draw down to ensure application to actively growing weeds.</p> <p>Use nozzles that emit medium- to large-sized droplets to minimize drift in order to avoid crop injury.</p>	<p>Do not apply this material through irrigation system.</p> <p>Do not make applications by air.</p> <p>Do not apply directly to water.</p>
Post-harvest (Cranberry Production)	<p>Best results will be obtained if applications are made to vines that appear dormant (after they have turned red). Hand-held sprayers, wipers or other appropriate application equipment listed under "Application Equipment and Techniques" in this label may be used. If using hand-held sprayers, use a 0.5 to 1 percent solution of this product.</p> <p>Spray to wet vegetation, not to run off. If using hand-held boom sprayers, apply 1.5-3 quarts (1.5-3 lbs ae) of this product per acre.</p> <p>PRECAUTION: Even though vines appear dormant, contact of the herbicide solution with desirable vegetation may result in damage or severe plant injury. Cranberry plants that are directly sprayed may be killed.</p>	<p>Do not treat more than 10 percent of the total bog.</p> <p>Allow a minimum of 6 months after the last application and next harvest of cranberries.</p> <p>Do not apply this product through the irrigation system.</p> <p>Do not make applications by air.</p> <p>Do not apply directly to water.</p> <p>Make applications only after cranberries have been harvested to control weeds growing within the field.</p>

10.2 - CITRUS		
LABELED CROPS: Calamondin, Chironja, Citron, Citrus Hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Orange (All), Pummelo, Satsuma Mandarin, Tangelo (ugli), Tangor		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
See Section 10.0	<p>See Use Directions under Section 10.0</p> <p>Florida and Texas only: For burn down or control of the weeds listed below, apply the labeled rates of this product in 3 to 40 gallons of water per acre. Where weed foliage is dense, use 10 to 30 gallons of water per acre.</p> <p>For goatweed, apply 1.5-2.25 quarts (1.5-2.25 lbs ae) of this product per acre. Apply in 20 to 30 gallons of water per acre when plants are actively growing. Use 3 pints (1.5 lbs ae) per acre when plants are less than 8 inches tall and 2.25 pints (1.13 lbs ae) per acre when plants are greater than 8 inches tall. If goatweed is greater than 8 inches tall, the addition of bromacil + diuron or diuron may improve control.</p> <p>It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.</p>	<p>See Section 10.0</p> <p>Pre-harvest Interval (PHI): Allow a minimum of 1 day between last application and harvest.</p> <p>For citron groves apply as directed sprays only.</p>

10.2 - CITRUS				
LABELED CROPS: Calamondin, Chironja, Citron, Citrus Hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Orange (All), Pummelo, Satsuma Mandarin, Tangelo (ugli), Tangor				
TYPES OF APPLICATIONS	USE DIRECTIONS		RESTRICTIONS	
Perennial weeds:	S=Suppression	B=Burndown	PC=Partial Control	C=Control
	AgSaver™ Glyphosate 5.4 Plus Rate Per Acre			
Weed Species	1.5 pints (0.75 lb ae)	1.5 quarts (1.5 lbs ae)	2.25 quarts (2.25 lbs ae)	3.75 quarts (3.75 lbs ae)
Bermudagrass	B	-	PC	C
Guineagrass				
Texas and Florida Ridge	B	C	C	C
Florida Flatwoods	-	B	C	C
Paragrass	B	C	C	C
Torpedograss	S	-	PC	C

10.3 – MISCELLANEOUS TREE FOOD CROPS		
LABELED CROPS: Cactus (fruits & pads), Palm (heart, leaves), Palm (oil)		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
See Section 10.0	See Use Directions under Section 10.0	See Section 10.0

10.4 – NON-FOOD TREE CROPS		
LABELED CROPS: Pine, Poplar, Eucalyptus, Christmas Trees, Other Non-food Tree Crops.		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
See Section 10.0	See Use Directions under Section 10.0	See Section 10.0
Directed sprays, Spot treatments, Wiper applications	<p>This product may be used as a post-directed spray and spot treatment around established poplar, eucalyptus, Christmas Trees, and other non-food tree crops.</p> <p>PRECAUTION: Care must be exercised to avoid contact of spray drift or mist with foliage or green bark of established Christmas trees and other pine trees. Desirable plants may be protected from the spray solution by using shields or coverings made of cardboard or other impermeable material.</p>	Do not use this product as an over-the-top broadcast spray in Christmas trees and other pine tree.
Site Preparation	<p>This product may be used prior to planting non-food tree crops.</p> <p>Precautions must be taken to protect non-target plants during site preparations applications.</p>	
Directed Spray (Eucalyptus and Poplar Production)	<p>This product can be used around established eucalyptus and poplar trees to control undesirable vegetation.</p> <p>Use a 1 to 2 percent spray solution to control herbaceous weeds in eucalyptus farms. Use a 2 percent spray solution for control of undesirable woody brush and trees. For “hard-to-control” weeds, use a 5 to 10 percent spray solution. Avoid contact of spray, drift, or mist with foliage, green bark, or non-woody surface roots of plants.</p>	Do not allow herbicide spray to contact desirable vegetation.
Wiper Application (Eucalyptus and Poplar Production)	<p>This product may be used through wick or other suitable wiper applicators for control or partial control of grass and broadleaf weeds listed in the “WEEDS CONTROLLED” section of this label. For wick applicators, mix 3 quarts (3 lbs ae) of this product with 2 gallons water to make a 33% solution. For wiper systems that can handle thicker solutions, such as force-fed systems, a 33 to 100% solution may be used. For best results, ensure that the herbicide solution is allowed to contact the maximum amount of leaf surface. As weed densities increase, decrease equipment speed to allow sufficient herbicide flow to wet all weed surfaces contacted. Weeds not contacted will be unaffected.</p>	

10.5 – POME FRUIT		
LABELED CROPS: Apple, Crabapple, Loquat, Mayhaw, Pear (including oriental pear), Quince		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
See Section 10.0	See Use Directions under Section 10.0	See Section 10.0 Pre-harvest Interval (PHI): Allow a minimum of 1 day between last application and harvest in pome fruits.

10.6 – STONE FRUIT		
LABELED CROPS: Apricot, Cherry (Sweet, Tart), Nectarine, Olive, Peach, Pear, Plum/Prune (All types), Plumcot.		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
See Section 10.0	See Use Directions under Section 10.0 Avoid application near trees with recent pruning wounds or other mechanical injury.	See Section 10.0 Pre-harvest Interval (PHI): Allow a minimum of 17 days between last application and harvest in stone fruit crops. For olive groves, apply as directed sprays only.

RESTRICTIONS ON APPLICATION EQUIPMENT:

For cherries, any application equipment listed in Section 10.0 may be used in all states.

Any application equipment listed in Section 10.0 may be used in apricots, nectarines, peaches, and plums/prunes growing in Arizona, California, Colorado, Idaho, Kansas, Kentucky, New Jersey, North Dakota, Oklahoma, Oregon, Texas, Utah, and Washington, except for peaches grown in the states specified in the following paragraph. In AK, AL, AR, CT, DE, FL, GA, HI, IA, IL, IN, LA, MA, MD, ME, MI, MN, MO, MS, MT, NC, NE, NH, NM, NV, NY, OH, PA, RI, SC, SD, TN, VA, VT, and WI, use wiper equipment only.

For Peaches grown in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina, and Tennessee only, apply with a shielded boom spray or shielded wiper applicator, which prevents any contact of this product with the foliage or bark of trees. Apply no later than 90 days after first bloom. Applications made after this time may result in severe damage. Remove suckers and low hanging limbs at least 10 days prior to application. Apply only near trees that have been planted in the orchard for 2 or more years.

EXTREME CARE MUST BE TAKEN TO ENSURE NO PART OF THE PEACH TREE IS CONTACTED.

10.7 - TREE NUTS		
LABELED CROPS: Almond, Beechnut, Betelnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Coconut, Filbert (Hazelnut), Hickory nut, Macadamia, Pecan, Pine nut, Pistachio, Walnut (Black, English)		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
See Section 10.0	See Use Directions under Section 10.0	See Section 10.0 Pre-harvest Interval (PHI): Allow a minimum of 3 days between last application and harvest of tree nuts, except coconut. Allow 14 days between application and harvest in coconuts.

10.8 - TROPICAL CROPS & SUBTROPICAL TREES & FRUITS		
LABELED CROPS: Ambarella, Atemoya, Avocado, Banana, Barbados Cherry (acerola), Biriba, Blimbe, Breadfruit, Cacao (cocoa) bean, Canistel, Carambola (starfruit), Cherimoya, Coffee, Custard apple, Dates, Durian, Feijoa, Figs, Governor's plum, Guava, Ilama, Imbe, Imbu, Jaboticaba, Jackfruit, Longan, Lychee, Mamey apple, Mango, Mangosteen, Marmaladebox (genip), Mountain papaya, Papaya, Pawpaw, Plantain, Persimmon, Pomegranate, Pulasan, Rambutan, Rose apple, Sapodilla, Sapote (black, mamey, white), Spanish lime, Soursop, Star apple, Sugar apple, Surinam cherry, Tamarind, Tea, Ti (roots & leaves), Wax jambu.		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
See Section 10.0	See Use Directions under Section 10.0 This product may be applied for weed control or for site preparation prior to transplanting crops listed in this section.	See Section 10.0 Pre-harvest Interval (PHI): Allow a minimum of 1 day between last application and harvest of banana, guava, papaya, and plantain crops. Allow a minimum of 14 days between last application and harvest of any other tropical or subtropical tree fruit. Allow a minimum of 28 days between last

10.8 - TROPICAL CROPS & SUBTROPICAL TREES & FRUITS		
LABELED CROPS: Ambarella, Atemoya, Avocado, Banana, Barbados Cherry (acerola), Biriba, Blimbe, Breadfruit, Cacao (cocoa) bean, Canistel, Carambola (starfruit), Cherimoya, Coffee, Custard apple, Dates, Durian, Feijoa, Figs, Governor's plum, Guava, Ilama, Imbe, Imbu, Jaboticaba, Jackfruit, Longan, Lychee, Mamey apple, Mango, Mangosteen, Marmaladebox (genip), Mountain papaya, Papaya, Pawpaw, Plantain, Persimmon, Pomegranate, Pulasan, Rambutan, Rose apple, Sapodilla, Sapote (black, mamey, white), Spanish lime, Soursop, Star apple, Sugar apple, Surinam cherry, Tamarind, Tea, Ti (roots & leaves), Wax jambu.		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
		application and harvest in coffee crops. In coffee and banana, delay applications 3 months after transplanting to allow the new coffee or banana plant to become established.
Bananacide (Banana only)	This product may be used to destroy banana plants infected with the Banana Bunchy Top Virus as well as non-infected banana plants to establish a disease free buffers around plantations. Remove all fruit from the plants within the treatment area prior to treatment. Inject 1 mL of this product's concentrate per 2 to 3 inches of pseudostem diameter. Make the injection at least one foot above ground, except for very small plants, which should be injected vertically into the top. Any subsequent regrowth must also be destroyed. All plants and mats (or units) adjacent (within a 4-foot radius) to a treated mat shall be mechanically destroyed. For control of the Banana Bunchy Top Virus, it is critical that the grower follow a strict control program involving monitoring for diseased plants, spraying to control the aphid vector, and destruction of all infected mats (or units). An infected plant may not show symptoms of the disease for up to 125 days; therefore, it is critical that the entire mat (or unit) containing the diseased plant be destroyed immediately.	Do not apply more than 11.25 mL of this product's concentrate per mat (or units). Remove all fruit from plants and mats (or units) prior to treatment. Do not harvest any fruit or plant materials from treated mats (or units) following injection. Do not allow livestock to consume treated materials. Following transplant of new banana plants into treated areas, allow plants to become established for 3 months before applying this product for weed control.

10.9 - VINE CROPS		
LABELED CROPS: Grapes (raisin, table, wine), Hops, Kiwi, Passion fruit		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
See Section 10.0	See Use Directions under Section 10.0 In the northeast and Great Lakes regions, applications must be made prior to the end of bloom stage of grapes to avoid injury or make applications with shielded sprayers or wiper equipment.	See Section 10.0 Pre-harvest Interval (PHI): Allow a minimum of 14 days between last application and harvest. Do not use selective equipment in kiwi Applications must not be made when green shoots, canes or foliage are in the spray zone.

11.0 - PASTURE GRASSES, FORAGE LEGUMES & RANGELANDS

RESTRICTIONS:

- **DO NOT** apply more than 7.5 pints (3.75 lbs ae) per acre per application, unless otherwise specified for an individual use.
- **DO NOT** apply more than 4.5 quarts (4.5 lbs ae) per acre per year as a combined total of all treatments, except for applications in non-agricultural sites or in tree, vine, or shrub crops and unless otherwise specified for an individual use.
- **DO NOT** apply more than 6 quarts (6 lbs ae) per acre per year as a combined total of all treatments for applications in non-agricultural sites, unless otherwise specified for an individual use.

11.1 – ALFALFA, CLOVER, & OTHER FORAGE LEGUMES		
LABELED CROPS: Alfalfa, Clover, Kenaf, Kudzu, Lespedeza, Leucaena, Lupin, Sainfoin, Trefoil, Velvet bean, Vetch (all types)		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
		See Section 11.0
Pre-plant, Pre-emergence,	This product may be applied before, during or after planting crops listed.	If a single application is made at rates of 1.5 quarts (1.5 lbs ae) per acre or less, no waiting period between

11.1 – ALFALFA, CLOVER, & OTHER FORAGE LEGUMES		
LABELED CROPS: Alfalfa, Clover, Kenaf, Kudzu, Lespedeza, Leucaena, Lupin, Sainfoin, Trefoil, Velvet bean, Vetch (all types)		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
At-Planting	Make applications according to the rates listed in Annual Weeds, Perennial Weeds, and Woody Brush & Trees rate tables in this label.	treatment and feeding or grazing is required. If application rates greater than 1.5 quarts (1.5 lbs ae) per acre are made, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting.
Spot treatment, Over-the-Top Wiper applications (Alfalfa and Clover only)	This product may be applied as a spot treatment in alfalfa or clover. This product may be applied with wiper applicators to control or suppress the weeds listed under "WIPER APPLICATORS" in the "SELECTIVE EQUIPMENT" section of this label. Applications may be made in the same area at 30-day intervals.	For spot treatment and wiper applications, apply in areas where the movement of domestic livestock can be controlled. No more than one-tenth of any acre can be treated at one time. Pre-harvest Interval (PHI): Remove domestic livestock before application and wait 14 days after application before grazing livestock or harvesting.
Dormant (Alfalfa Only)	This product will control or suppress many weeds including quackgrass, downy brome and cheatgrass in dormant alfalfa. Apply 6-9 fl oz (0.19-0.28 lb ae) per acre of this product. Apply in the spring to alfalfa that is dormant. Applications should be made after spring temperatures have warmed enough to encourage resumption of weed growth, but prior to initiation of trifoliate leaf expansion of the alfalfa. Applications made after expansion of the first trifoliate leaf of the alfalfa will cause growth reduction and reduced crop yield. Slight discoloration of the alfalfa may occur, but the alfalfa will regreen and regrow under moist soil conditions as effects of this product wear off. PRECAUTION: Application of this product can cause crop injury.	Do not use ammonium sulfate when spraying dormant alfalfa with AgSaver™ Glyphosate 5.4 Plus. Do not use this product where a slight yield reduction in the first cutting of alfalfa cannot be tolerated. Do not make more than one application per year. Pre-harvest Interval (PHI): Allow 36 hours after application before grazing livestock or harvesting.
Pre-harvest (Alfalfa Only)	This product may be used in declining alfalfa stands or any stand of alfalfa where crop destruction is acceptable. This application will severely injure or destroy the stand of alfalfa. This product will control annual and perennial weeds including quackgrass, when applied prior to the harvest of alfalfa. Use up to 1.5 pints (0.75 lb ae) of this product per acre. Applications may be made at any time of the year. For control of quackgrass, apply in the spring, late summer or fall when quackgrass is actively growing. Treatments for quackgrass must be followed by deep tillage for complete control.	Make only one application to an existing stand of alfalfa per year. Do not apply more than 1.5 quarts (0.75 lb ae) per acre as a pre-harvest treatment. Do not apply more than 1.5 quarts (0.75 lb ae) per acre per year as a pre-harvest treatment. Do not use for alfalfa grown for seed. Pre-harvest Interval (PHI): Wait 36 hours before treated crop and weeds can be harvested and fed to livestock.
Renovation	This product may be applied as a broadcast spray to existing stands of alfalfa, clover, and other labeled forage legumes. Labeled crops may be planted into the treated area. Make applications according to the rates listed in Annual Weeds, Perennial Weeds and Wood Brush & Trees Rate Tables in this label.	Remove domestic livestock before application. If application rates of 1.5 quarts (1.5 lbs ae) per acre or less are used wait 36 hours after application before grazing or harvesting. If application rates greater than 1.5 quarts (1.5 lbs ae) per acre are used, wait 8 weeks after application before grazing or harvesting.

11.2 - CONSERVATION RESERVE PROGRAM (CRP)		
LABELED CROPS: Conservation Reserve Program (CRP) Acres		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
		See Section 11.0
Renovation (rotating out of CRP), Site preparation	This product may be used to prepare CRP land for crop production. Refer to Federal, state, or local use guides for CRP renovation recommendations. Make applications according to the rates listed in Annual Weeds, Perennial Weeds and Wood Brush & Trees Rate Tables in this label. PRECAUTION: Some stunting of CRP perennial grasses will occur if broadcast	Do not apply more than 2.25 quarts (2.25 lbs ae) per acre per year onto CRP grasses. For any crop not listed in the "CROPS" sections of this

	applications are made when plants are not dormant.	label applications must be made at least 30 days prior to planting.
Post-emergence Weed control in Dormant Acres, Over-the-Top Wiper Application	This product may be used to suppress competitive growth and seed production of undesirable vegetation in CRP acres. Such applications may be made with wiper application equipment or as a broadcast or spot treatment to dormant CRP grasses. For selective applications with broadcast spray equipment, apply 9 – 12 fl oz (0.28-0.38 lb ae) of this product per acre in early spring before desirable CRP grasses, such as crested and tall wheatgrass, break dormancy and initiate green growth. Late fall applications can be made after desirable perennial grasses have reached dormancy.	

11.3 - GRASS or TURFGRASS SEED PRODUCTION		
LABELED CROPS: Any grass (Gramineae family) except corn, sorghum, sugarcane, and those listed under "CEREAL CROPS"		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
		See Section 11.0
Pre-plant, Pre-emergence, Renovation, Site preparation	<p>This product may be applied before, during, or after planting or for renovation of turf or forage grass areas grown for seed production.</p> <p>Make applications according to the rates listed in Annual Weeds, Perennial Weeds and Wood Brush & Trees Rate Tables in this label.</p> <p>Applications must be made prior to the emergence of the crop to avoid injury.</p> <p>For maximum control of existing vegetation, delay planting to determine if any regrowth from escaped underground plant parts occurs. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm-season grasses, such as bermudagrass, summer or fall applications provide best control.</p>	<p>Do not disturb soil or underground plant parts before treatment. Tillage or renovation techniques such as vertical mowing, coring, or slicing must be delayed for 7 days after application to allow proper translocation into underground plant parts.</p> <p>If application rates total 2.25 quarts (2.25 lbs ae) per acre or less, no waiting period between treatment and feeding or livestock grazing is required.</p> <p>If the rate is greater 2.25 quarts (2.25 lbs ae). per acre, remove domestic livestock and wait 8 weeks following application before grazing or harvesting.</p>
Shielded Sprayer	<p>Apply 24-72 fl oz (0.75-2.25 lbs ae) of this product as a broadcast spray in 10 to 20 gallons of total spray volume per acre. Uniform planting in straight rows aid in shielded sprayer applications. Best results are obtained when the grass seed crop is small enough to easily pass by or through the protective shields.</p> <p>PRECAUTION: Contact of this product in any manner to any vegetation to which treatment is not intended may cause damage.</p> <p>To the extent consistent with applicable law, grower assumes all responsibility for crop losses from misapplication.</p>	
Over-the-Top Wiper Applications	<p>This product may be applied with wiper applicators to control or suppress the weeds listed under "WIPER APPLICATORS" in the "SELECTIVE EQUIPMENT" section of this label.</p> <p>Contact of the herbicide solution with desirable vegetation may result in damage or destruction. Applicators must be adjusted so that the wiper contact point is at least 2 inches above the desirable vegetation.</p> <p>Weeds should be a minimum of 6 inches above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations, or when weed height varies so that not all weeds are contacted. In these instances, repeat treatments up to the maximum application rate may be necessary. Better results may be obtained if 2 applications are made in opposite directions.</p>	

11.3 - GRASS or TURFGRASS SEED PRODUCTION		
LABELED CROPS: Any grass (Gramineae family) except corn, sorghum, sugarcane, and those listed under "CEREAL CROPS"		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
Spot treatments	Use a 1- to 1.5 percent solution. Apply this product prior to heading of grasses.	The crop receiving the spray in the treated area will be killed. Do not allow drift or spray outside of the target area for the same reason.
Creating Rows in Annual Ryegrass	Use 12-24 fl oz (0.38-0.75 lb ae) of this product per acre. Use the higher rate when the ryegrass is greater than 6 inches tall. Best results are obtained when applications are made before the ryegrass reaches 6 inches in height. PRECAUTION: Set nozzle height to allow the establishment of the desired row spacing while preventing spray droplets, spray fines, or drift to contact the ryegrass plants not treated. Use low-pressure nozzles, or drop nozzles designed to target the application over a narrow band. To the extent consistent with applicable law, grower assumes all responsibility for crop losses from misapplication.	

11.4 - PASTURES		
LABELED CROPS: Any grass (Gramineae family) except corn, sorghum, sugarcane, and those listed under "CEREAL CROPS." Including Bahiagrass, Bermudagrass, Bluegrass, Brome, Fescue, Guineagrass, Kikuygrass, Orchardgrass, Pangola grass, Ryegrass, Timothy, Wheatgrass.		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
		See Section 11.0
Spot treatment, Over-the Top Wiper Applications	This product may be applied as a spot treatment or with wiper applicators in pastures. Applications may be made in the same area at 30-day intervals. For spot treatments or wiper application methods using rates of 2.25 quarts (2.25 lbs ae). per acre or less, the entire field or any portion of it may be treated.	When spot treatment or wiper applications are made using rates above 2.25 quarts (2.25 lbs ae). per acre, no more the 10 percent of the total pasture may be treated at any one time. Pre-harvest Interval (PHI): Remove domestic livestock before application and wait 7 days after application before grazing livestock or harvesting.
Pre-plant, Pre-emergence, Pasture renovation, Stand Removal	This product may be applied prior to planting or emergence of forage grasses. In addition, this product may be used to control perennial pasture species listed on this label prior to re-planting. Make applications according to the rates listed in Annual Weeds, Perennial Weeds and Wood Brush & Trees Rate Tables in this label.	If application rates total 2.25 quarts (2.25 lbs ae) per acre or less, no waiting period between treatment and feeding or livestock grazing is required. If the rate is greater 2.25 quarts (2.25 lbs ae) per acre, remove domestic livestock and wait 8 weeks following application before grazing or harvesting.
Chemical Mowing (Bermudagrass Pastures Prior To Spring Growth Or Immediately After First Cutting)	This product may be applied at 12 fl oz (0.38 lb ae) per acre to control the weeds listed below and most other winter annual grass and broadleaf weeds in established coastal bermudagrass pastures. Annual bluegrass, Cheat, Crabgrass, Henbit, Johnsongrass seedling, Little barley, Oats, Ryegrass, Sandbur field, Wheat, Wild mustard <u>Applications prior to spring growth:</u> Apply this product in the late winter or early spring but before new coastal bermudagrass growth begins in the spring. Applications to new growth can damage the bermudagrass. <u>Applications following the first cutting:</u> Apply this product after the first bermudagrass cutting when the bermudagrass has not yet begun to regrow. Applications made after regrowth has begun can damage the bermudagrass.	Do not apply more than 7.5 pints (3.75 lbs ae) per acre per year. Labeled application rates totaling 2.25 quarts (2.25 lbs ae) per acre or less do not require a waiting period between treatment and feeding or livestock grazing. If the rate is greater than 2.25 quarts (2.25 lbs ae) per acre, remove domestic livestock and wait 8 weeks following application before grazing or harvesting. Only make one application per year to treated fields. Do not make a spring application prior to growth and an application following the first cutting on the field during the same year.

STATE SPECIFIC DIRECTIONS FOR PASTURES

Colorado, Idaho, Iowa, Kansas, Montana, Nebraska, North Dakota, Oregon, South Dakota, Utah, Washington, and Wyoming Only

Bromus Species: This product may be used to treat downy brome (*Bromus tectorum*), Japanese brome (*Bromus japonicus*), soft chess (*Bromus mollis*) and cheatgrass (*Bromus secalinus*) found in industrial, rangeland and pasture sites. Apply 6 to 12 fl oz (0.19-0.38 lb ae) of product per acre on a broadcast basis. For best results, treatment should coincide with early seedhead emergence of the most mature plants. Delaying the application until this growth stage will maximize the emergence of other weedy grass flushes. Applications should be made to the same site each year until seed banks are depleted and the desirable perennial grasses are able to become reestablished on the site.

Medusahead: To treat medusahead, apply 12 fl oz (0.38 lb ae) of this product per acre as soon as plants are actively growing, and prior to the 4 leaf stage. Applications may be made in the fall or spring.

Application Equipment and Techniques: Applications may be made using ground or aerial equipment. Aerial applications for these uses may be made using fixed wing or helicopter equipment. For aerial applications, apply in 2 to 10 gallons of water per acre. For applications using ground equipment, apply in 10 to 20 gallons of water per acre.

When applied as directed there are no grazing restrictions.

11.5 – RANGELANDS		
LABELED CROPS: Rangeland (Perennial cool and warm season grass rangelands)		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
		See Section 11.0
Post-emergence	<p>This product will control or suppress many annual weeds growing in perennial cool and warm-season grass rangelands.</p> <p>Preventing viable seed production is key to the successful control and invasion of annual grassy weeds in rangelands. Follow-up applications in sequential years should eliminate most of the viable seeds.</p> <p>Grazing of treated areas should be delayed to encourage growth of desirable perennials. Allowing desirable perennials to flower and reseed in the treated area will encourage successful transition.</p> <p>Apply 9 – 12 fl oz (0.28-0.38 lb ae) per acre to control or suppress many annual weeds growing in perennial cool and warm-season grass rangelands including downy brome, cheat grass, cereal rye and jointed goatgrass. Apply when most mature brome plants are in early flower and before the plants, including seedheads, turn color.</p> <p>Allowing for secondary weed flushes to occur in the spring following rain events further depletes the seed reserve and encourage perennial grass conversion on weedy sites. Fall applications are possible, and recommended, where spring moisture is usually limited and fall germination allows for good weed growth.</p> <p>For medusahead, apply 12 fl oz (0.38 lb ae) per acre at the 3-leaf stage. Delaying applications beyond this stage will result in reduced or unacceptable control. Fire may be useful in eliminating the thatch layer produced by slow decaying culms prior to application. Allow new growth to occur before spraying after a burn.</p> <p>Slight discoloration of the desirable grasses may occur, but they will regreen and regrow under moist soil conditions as effects of this product wear off.</p>	<p>Do not apply more than 2.25 quarts (2.25 lbs ae) per acre per year.</p> <p>Do not use ammonium sulfate when spraying rangeland grasses with this product.</p>

11.6 – TURF GRASS SOD PRODUCTION		
LABELED CROPS: Turfgrass for Sod		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
		See Section 11.0
Pre-plant, Pre-emergence, Renovation, Site Preparation	<p>This product controls most existing vegetation prior to renovating turf grass areas or establishing turf grass grown for sod. Broadcast of hand-held equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested. For maximum control of existing vegetation, delay planting or sodding to determine if any regrowth from escaped underground plant parts occurs. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm-season grasses such as bermudagrass, summer or fall applications provide the best control. Where existing vegetation is growing under mowed turfgrass management, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the spray.</p> <p>Make applications according to the rates listed in Annual Weeds, Perennial Weeds and Wood Brush & Trees Rate Tables in this label.</p> <p>Desirable turfgrasses may be planted following the above procedures.</p>	<p>If application rates total 2.25 quarts (2.25 lbs ae) per acre or less, no waiting period between treatment and feeding or livestock grazing is required.</p> <p>If the rate is greater than 2.25 quarts (2.25 lbs ae) per acre, remove domestic livestock and wait 8 weeks following application before grazing or harvesting.</p> <p>Do not disturb soil or underground plant before treatment.</p> <p>Tillage or renovation techniques such as vertical mowing, coring, or slicing must be delayed for 7 days after application to allow translocation into underground plant parts.</p>
Spot treatment	Hand-held equipment may be used for spot treatment of unwanted vegetation growing in existing turf grass	
Turfgrass Renovation for sod production	<p>This product controls most existing vegetation prior to renovating turfgrass areas or establishing turfgrass grown for seed or sod. For maximum control of existing vegetation, delay planting or sodding to determine if any regrowth from escaped underground plant parts occurs. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm-season grasses such as bermudagrass, summer or fall applications provide the best control. Where existing vegetation is growing under mowed turfgrass management, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the spray.</p> <p>Tillage or renovation techniques such as vertical mowing, coring, or slicing should be delayed for 7 days after application to allow translocation into underground plant parts.</p> <p>Desirable turfgrass may be planted following the above procedures.</p> <p>Hand-held equipment may be used for spot treatment of unwanted vegetation growing in existing turfgrass. Broadcast or hand-held equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested.</p>	<p>Do not feed or graze turfgrass grown for seed or sod production for 8 weeks following application.</p> <p>Do not disturb soil or underground plant parts before treatment.</p>

11.7 - RELEASE OF BERMUDAGRASS OR BAHIA GRASS

Dormant applications

This product may be used to control or partially control many winter annual weeds and tall fescue for effective release of dormant bermudagrass or bahiagrass. Treat only when turf is dormant and prior to spring greenup. This product may also be tank mixed with sulfometuron for residual control. Tank mixtures of this product with sulfometuron may delay greenup. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

For best results on winter annuals, treat when plants are in an early growth stage (below 6 inches in height) after most have germinated. For best results on tall fescue, treat when fescue is at or beyond the 4 to 6-leaf stage.

Apply 6 – 48 fl oz (0.19-1.5 lbs ae) of this product per acre alone or in a tank mixture with the labeled rate of sulfometuron. Apply the labeled rates in 10 to 40 gallons of water per acre. Use only in areas where bermudagrass or bahiagrass are desirable ground covers and where some temporary injury or discoloration can be tolerated.

Actively growing bermudagrass:

This product may be used to control or partially control many annual and perennial weeds for effective release of actively growing bermudagrass. Apply 12 – 36 fl oz (0.38-1.13 lbs ae) of this product in 10 to 40 gallons of spray solution per acre. Use the lower rate when treating annual weeds below 6 inches in height (or runner length). Use the higher rate as weeds increase in size or as they approach flower or seedhead formation. These rates will also provide partial control of the following perennial species:

Bahiagrass	Johnsongrass
Bluestem, silver	Trumpet creeper
Fescue, tall	Vaseygrass

This product may be tank mixed with sulfometuron. If tank-mixed, use no more than 12 – 24 fl oz (0.38-0.75 lb ae) of this product with the labeled rate of sulfometuron per acre. These rates will also provide partial control of the following perennial weeds:

Bahiagrass	Fescue, tall
Broomsedge	Johnsongrass
Bluestem, silver	Poorjoe
Dallisgrass	Trumpet creeper
Dock, curly	Vaseygrass
Dogfennel	Vervain, blue

For suppression of vegetative growth and seedhead inhibition of bahiagrass for approximately 45 days, apply 4.5 fl oz (0.14 lb ae) of this product in 10 to 40 gallons of water per acre. Apply 1 to 2 weeks after full green-up or after mowing to a uniform height of 3 to 4 inches. This application must be made prior to seedhead emergence.

For suppression up to 120 days, apply 3 fl oz (0.09 lb ae) of this product per acre, followed by an application of 1.5 – 3 fl oz (0.05-0.09 lb ae) per acre about 45 days later. Make no more than 2 applications per year.

A tank mixture of this product plus sulfometuron may be used. Apply 4.5 fl oz (0.14 lb ae) of this product plus the labeled rate of sulfometuron per acre 1 to 2 weeks following an initial spring mowing. Make only one application per year.

RESTRICTIONS

- For application to actively growing bermudagrass and tank mixture of this product plus sulfometuron, do not apply more than 24 fl oz (0.75 lb ae) per acre per application.
- For suppression of vegetative growth and seedhead inhibition of bahiagrass up to 120 days, do not apply more than 3 fl oz (0.09 lb ae) per acre per application.
- For suppression of vegetative growth and seedhead inhibition of bahiagrass up to 120 days, do not apply more than 6 fl oz (0.19 lb ae) per acre per application.
- For suppression of vegetative growth and seedhead inhibition of bahiagrass and tank mixture of this product plus sulfometuron, do not apply more than 4.5 fl oz (0.14 lb ae) per acre per application.
- For suppression of vegetative growth and seedhead inhibition of bahiagrass and tank mixture of this product plus sulfometuron, do not apply more than 4.5 fl oz (0.14 lb ae) per acre per year.
- Use only on well-established bermudagrass.
- Bermudagrass injury may result from the treatment, but regrowth will occur under moist conditions.
- Do not repeat applications of the tank mix in the same year since severe injury may occur. Actively growing bahiagrass.

12.0 - ROUNDUP READY CROPS

The following instructions include all applications which can be made onto the specified Roundup Ready crops during the complete cropping season. Do NOT combine these instructions with those listed for crop varieties that do not contain the Roundup Ready gene, in the “ANNUAL AND PERENNIAL CROPS (ALPHABETICAL)” section of this label.

THIS PRODUCT IS TO BE USED FOR POSTEMERGENCE APPLICATION ONLY ON CROP VARIETIES

DESIGNEATED AS CONTAINING A ROUNDUP READY GENE OR GLYPHOSATE TOLERANT GENE.

Applying this product to crop varieties that are not designated as glyphosate tolerant will result in severe crop injury and yield loss. Avoid contact with foliage, green stems, or fruit of crops, or any desirable plants that do not contain a Roundup Ready or glyphosate tolerant gene, since severe injury or destruction will result.

The Roundup Ready designation indicates that the crop variety contains a patented gene that provides tolerance to this product. Information on Roundup Ready crop varieties may be obtained from your seed supplier. Roundup Ready crop varieties must be purchased from an authorized licensed seed supplier.

NOTE: Roundup Ready seed, and the method of selectivity controlling weeds using glyphosate on a Roundup Ready crop, are protected under several U.S. Patents. A license to use Roundup Ready seed must be obtained prior to use.

For Ground Applications with broadcast equipment, apply this product in 5 to 20 gallons of spray solution per acre. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment use flat spray nozzles. Check for even distribution of spray droplets.

For Aerial Applications apply this product in 3 to 15 gallons of water per acre. See the "APPLICATION EQUIPMENT AND TECHNOLOGIES" section of this label for procedures to avoid spray drift that may cause injury to any vegetation not intended for treatment. Use of appropriate buffer zones will help prevent injury to adjacent vegetation.

ATTENTION: AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN A GLYPHOSATE TOLERANT GENE.

See the "MIXING and APPLICATION EQUIPMENT AND TECHNIQUES" sections of this label for additional directions and restrictions on the application of this product.

To prevent crop injury, tank mixtures with other herbicides, insecticides, fungicides, micronutrients, or fertilizers may result in reduced weed control or crop injury and are not recommended for over-the-top applications of this product unless otherwise specified in this product label.

Ammonium sulfate may be mixed with this product for applications to Roundup Ready crops. Refer to the "MIXING" section for USE DIRECTIONS for ammonium sulfate.

Sprayer Preparation: It is important that sprayer and mixing equipment be clean and free of pesticide residue before making applications of this product. Follow the cleaning procedures specified on the label of the product(s) previously used. THOROUGHLY CLEAN THE SPRAY TANK AND ALL LINES AND FILTERS TO ELIMINATE POTENTIAL CONTAMINATION FROM OTHER HERBICIDES PRIOR TO MIXING AND APPLYING THIS PRODUCT.

NOTE: The following use directions are based on a clean start at planting by using a burndown application or tillage to control existing weeds before crop emergence. In no-till and stale seedbed systems, use a preplant burn-down treatment of this product to control existing weeds prior to crop emergence. Some weeds, such as black nightshade, broadleaf signalgrass, sicklepod, Texas panicum, sandbur, annual morningglory, woolly cupgrass, shattercane, wild proso millet, burcucumber, and giant ragweed with multiple germination times or suppressed (stunted) weeds may require a second application of this product for complete control. The second application should be made after some regrowth has occurred and at least 10 days after a previous application of this product.

RESTRICTIONS

- **DO NOT** apply more than 7.5 pints (3.75 lbs ae) per acre per application, unless otherwise specified for an individual use.

12.1 - ROUNDUP READY CANOLA (Spring Varieties)		
<p>LABELED CROPS: Roundup Ready spring canola is defined as those Roundup Ready canola varieties that are seeded in the spring and harvested in the fall and do not enter a winter dormancy period.</p> <p>Do not use this product on spring canola with a Roundup Ready gene planted in the following states: Alabama, Delaware, Florida, Georgia, Kentucky, Maryland, New Jersey, North Carolina, South Carolina, Tennessee, Virginia, and West Virginia, except for uses in wildlife food plots that will not be for human or livestock food.</p>		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
		<p>See Section 12.0.</p> <p>Do not apply more than 4.5 pints (2.25 lbs ae) per acre per year total for all application types.</p>
Pre-plant, At-Planting, Pre-emergence	This product may be applied before, during or after planting Roundup Ready spring canola.	Do not apply more than 3 pints (1.5 lbs ae) per acre per year.
Post-emergence (In-crop)	<p>This product may be applied via post-emergence in-crop (over-the-top) broadcast applications to Roundup Ready spring canola from emergence through the 6-leaf stage of development. Applications made during bolting or flowering may result in crop injury and yield loss. To maximize yield potential, make applications early to eliminate competing weeds.</p> <p><u>Single Application</u> – 8.25 – 12 fl oz (0.26-0.38 lb ae) of this product per acre no later than the 6-leaf stage for the control of annual weeds. Avoid overlapping applications as this may result in temporary yellowing, delayed flowering, and or growth reduction. Similar crop injury may result when applications of more than 8.25 fl oz (0.26 lb ae) per acre are applied after the 4-leaf stage.</p> <p><u>Sequential Application</u> – Apply 8.25 fl oz (0.26 lb ae) of this product per acre to 1- to 3-leaf canola followed by a sequential application at a minimum interval of 10 days, but no later than the 6-leaf stage. Sequential applications can be made for early emerged annual weeds and perennial weeds such as Canada thistle and quackgrass, or when multiple applications are needed for adequate weed control.</p>	<p>Do not apply more than 1.5 pints (0.75 lb ae) per acre per year.</p> <p>No more than two in-crop (over-the-top) broadcast applications may be made from crop emergence through the 6-leaf stage of development and the total of all in-crop applications must not exceed 24 fl. oz.(0.75 lb ae) of this product per acre.</p> <p>Pre-harvest Interval (PHI): Allow a minimum of 60 days between last application and canola harvest.</p>

12.2 - ROUNDUP READY CANOLA (Fall & Winter Varieties)		
<p>LABELED CROPS: Roundup Ready winter canola is defined as those Roundup Ready canola varieties that are seeded in early fall and harvested the following spring or summer. Winter canola varieties are intended to enter a cold period dormancy in the winter.</p>		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
		<p>See Section 12.0.</p> <p>Do not apply more than 6 pints (3 lbs ae) per acre per year total for all application types.</p>
Pre-plant, At-Planting, Pre-emergence	This product may be applied before, during or after planting Roundup Ready winter canola.	Do not apply more than 3 pints (1.5 lbs ae) per acre per year.
Post-emergence (In-crop)	<p>This product may be applied in-crop (over-the-top) to Roundup Ready winter canola varieties from emergence to canopy closure in the fall and prior to bolting in the spring. Applications made during or after bolting may result in crop injury and yield loss. To maximize yield potential, make applications early to eliminate competing weeds.</p> <p>Some weeds with multiple germination times, or suppressed (stunted) weeds, or weeds that have overwintered may require sequential applications of this product for control. The second application should be made after some re-growth has occurred and at least 60 days after a previous application of this product.</p> <p><u>Single Application</u> – Apply 16.5-24 fl oz (0.52-0.75 lb ae) of this product per acre in the fall. Applications in the fall should be made when weeds are small and actively growing. Use the higher rate in the labeled range when weed densities are high, when weeds have</p>	<p>Do not apply more than 3 pints (1.5 lbs ae) per acre per year.</p> <p>Pre-harvest Interval (PHI): Allow a minimum of 60 days between last application and harvest of canola grain.</p> <p>No waiting period is required between application and open grazing of livestock.</p>

12.2 - ROUNDUP READY CANOLA (Fall & Winter Varieties)

LABELED CROPS: Roundup Ready winter canola is defined as those Roundup Ready canola varieties that are seeded in early fall and harvested the following spring or summer. Winter canola varieties are intended to enter a cold period dormancy in the winter.

TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
	<p>overwintered or when weeds become large and well established. Applications of greater than 12 fl oz (0.38 lb ae) per acre prior to the 6-leaf stage may result in reduced crop growth in the fall. Avoid overlaps. Spray overlaps may result in temporary yellowing and/or growth reduction.</p> <p><u>Sequential Applications</u> – Apply 12 – 24 fl oz (0.38-0.75 lb ae) of this product per acre to 2-leaf or larger canola in the fall, followed by a sequential application at the same rate and at a minimum interval of 60 days, but before bolting in the spring. Sequential applications can be made for early emerging annual weeds and winter emerging weeds such as downy brome, jointed goatgrass and ryegrass, and for weeds that have overwintered. This product will control or suppress most of perennial weeds. For some perennial weeds, sequential applications may be required to reduce competition with the crop.</p>	

12.3 - ROUNDUP READY CORN

TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
		<p>See Section 12.0.</p> <p>Do not apply more than 6 quarts (6 lbs ae) per acre per year total for all application types.</p>
Pre-plant, Pre-emergence, At-Planting	<p>This product may be applied alone or in a tank-mixture before, during or after planting corn.</p> <p>TANK MIXTURES: This product may be tank mixed with acetochlor or acetochlor + atrazine at 50 to 100 percent of labeled rate. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.</p> <p>If less than the maximum rates of the above listed residual products was used in pre-plant and pre-emergence treatment, then a post-emergence (in-crop) application of this product should be applied for maximum weed control.</p> <p>Avoid drift. Extreme care must be used when applying this product to prevent injury to desirable plants and crops which do not contain glyphosate tolerant gene.</p> <p>Make applications according to the rates listed in Annual Weeds, Perennial Weeds, and Woody Brush & Trees rate tables in this label.</p>	<p>Do not apply more than 3.75 quarts (3.75 lbs ae) per acre per year.</p> <p>Do not apply this product to crop varieties that are not designated as glyphosate tolerant.</p> <p>Do not allow contact with foliage, green stems, or fruit of crops, or any desirable plants that do not contain a Roundup Ready or glyphosate tolerant gene, since severe injury or destruction will result.</p> <p>Avoid drift. Extreme care must be used when applying this product to prevent injury to desirable plants and crops which do not contain a glyphosate tolerant gene.</p> <p>See the "Mixing and Application Equipment and Techniques" sections of this label for additional directions and restrictions on the application of this product.</p>
Post-emergence (in-crop)	<p>When applied as directed, this product controls labeled annual grass and broadleaf weeds in Roundup Ready corn. Many perennial grasses and broadleaf weeds will be controlled or suppressed with one or more application of this product. The post-emergent application of 18 – 36 fl oz (0.56-1.13 lbs ae) per acre of this product should be made before the weeds reach a height and/or density that the weeds become competitive with the crop, generally 4 inch tall weeds or less.</p> <p>This product may be applied over-the-top to Roundup Ready corn from emergence through the V8 stage (8 leaves with collars) or until corn height reaches 48 inches, whichever comes first.</p> <p>Use drop nozzles when corn height is 24 to 30 inches (free</p>	<p>Do not apply more than 2.25 pints (1.13 lbs ae) per acre per application.</p> <p>Do not apply more than 4.5 pints (2.25 lbs ae) per acre per year.</p> <p>Allow a minimum of 10 days between in-crop applications of this product.</p> <p>Pre-harvest Interval (PHI): Allow a minimum of 50 days between application of this product and harvest of corn forage.</p>

12.3 - ROUNDUP READY CORN		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
	<p>standing), for optimum spray coverage and weed control.</p> <p>For corn heights 30 to 48 inches (free standing), apply this product ONLY using ground application equipment with drop nozzles adjusted to avoid spraying into the whorls of the corn plants.</p> <p>If product is applied to whorls of corn, plant injury and yield reduction can occur.</p> <p>Maximum single in-crop application rate of this product up to 48-inch corn is 2.25 pints (1.13 lbs ae) per acre.</p> <p>See the "ROUNDUP READY CROPS" section of this label for precautionary instructions for use in Roundup Ready crops.</p> <p>TANK MIXTURES: This product may be applied in tank mixture with acetochlor and acetochlor + atrazine at 50 to 100 percent of labeled rate. This product may be applied in tank mixture with Permit and Atrazine at labeled rates. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.</p>	
Pre-Harvest	In Roundup Ready corn, up to 1.5 pints (0.75 lb ae) per acre of this product can be applied pre-harvest. Make applications at 35 percent grain moisture or less. Ensure that maximum kernel fill is complete, and the corn is physiologically mature (black layer formed).	<p>Do not apply more than 1.5 pints (0.75 lb ae) per acre per application.</p> <p>Do not apply more than 1.5 pints (0.75 lb ae) per acre per year.</p> <p>Pre-harvest Interval (PHI): Allow a minimum of 7 days between application and harvest.</p>
Post-Harvest	This product may be applied after harvest of corn. A 2.25 pints (1.13 lbs ae) per acre rate may be required for control of large weeds that were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.	Pre-harvest Interval (PHI): Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

12.4 - ROUNDUP READY COTTON		
Use of this product in accordance with label directions is expected to result in normal growth of Roundup Ready cotton. However, various environmental conditions, agronomic practices and other factors make it impossible to eliminate all risks associated with this product, even when application are made in conformance with the label specifications. In some cases, these factors can result in boll loss, delayed maturity and/or yield loss.		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
		<p>See Section 12.0.</p> <p>Do not apply more than 6 quarts (6 lbs ae) per acre per year total for all application types.</p>
Pre-plant, Pre-emergence, At-planting	<p>This product may be applied before, during or after planting cotton.</p> <p>Make applications according to the rates listed in Annual Weeds, Perennial Weeds, and Woody Brush & Trees rate tables in this label.</p> <p>See the "ROUNDUP READY CROPS" section of this label for precautionary instructions for use in Roundup Ready crops.</p>	Do not apply more than 3.75 quarts (3.75 lbs ae) per acre per year.
Post-emergence (Over-the-Top)	This product may be applied by aerial or ground application equipment at rates up to 1.5 pints (0.75 lb ae) per acre per application post-emergence to Roundup Ready cotton from the ground cracking stage until the 4-leaf (node) stage of development (until the fifth true leaf reaches the size of a quarter). Over-the-top applications made after the 4-leaf (node) stage of development may result in boll loss, delayed	<p>Do not apply more than 1.5 pints (0.75 lb ae) per acre per application.</p> <p>From ground cracking to layby, do not apply more than 3 quarts (3 lbs ae) per acre per year.</p> <p>Do not apply more than 4.5 quarts (4.5 lbs ae) per acre per year.</p>

12.4 - ROUNDUP READY COTTON

Use of this product in accordance with label directions is expected to result in normal growth of Roundup Ready cotton. However, various environmental conditions, agronomic practices and other factors make it impossible to eliminate all risks associated with this product, even when application are made in conformance with the label specifications. In some cases, these factors can result in boll loss, delayed maturity and/or yield loss.

TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
	<p>maturity and/or yield loss.</p> <p>Salvage Treatment. This treatment may be used after the 4-leaf stage of development and must only be used where weeds threaten to cause the loss of the crop. 1.5 pints (0.75 lb ae) per acre may be applied either as an over-the-top applications or as a post-directed treatments sprayed higher on the cotton plants and over the weeds.</p> <p>Salvage treatments will result in significant boll loss, delayed maturity and/or yield loss.</p> <p>See the "ROUNDUP READY CROPS" section of this label for precautionary instructions for use in Roundup Ready crops.</p>	<p>Do not apply more than 1 salvage treatment application per year.</p> <p>Sequential in-crop over-the-top or post-directed applications of this product must be at least 10 days apart and cotton must have at least two nodes of incremental growth between applications.</p> <p>Pre-harvest Interval (PHI): Allow a minimum of 7 days between application and harvest.</p>
Selective Equipment	<p>This product may be applied using precision post-directed or hooded sprayers at rates up to 1.5 pints (0.75 lb ae) per acre per application to Roundup Ready cotton through layby. At this stage, post-directed equipment must be used which directs the spray to the base of the cotton plants.</p> <p>Contact of the spray with cotton leaves should be avoided to the maximum extent possible. To minimize spray onto the leaves of the cotton plants, place nozzles in a low position directing a horizontal spray pattern under the cotton leaves to contact weeds in the row and maintain low spray pressure (less than 30 psi). For best results, make applications while weeds are small (less than 3 inches).</p>	<p>Do not apply more than 1.5 pints (0.75 lb ae) per acre per application.</p> <p>See the "SELECTIVE EQUIPMENT" part of the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for information on proper use and calibration of this equipment.</p>
Pre-harvest	<p>This product may be applied for pre-harvest annual and perennial weed control as a broadcast treatment to Roundup Ready cotton after 20 percent boll crack. Up to 3 pints (1.5 lbs ae) of this product may be applied using either aerial or ground spray equipment.</p> <p>TANK MIXTURES: This product may be tank mixed with tribufos, diuron + thidiazuron, or ethephon. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.</p> <p>Refer to manufacturers' labels for use of additives (such as surfactants, stickers and spreaders) for preharvest application to cotton.</p> <p>This product will not enhance the performance of these harvest aids when applied to Roundup Ready cotton.</p>	<p>Do not apply more than 3 pints (1.5 lbs ae) per acre per application.</p> <p>Do not apply more than 3 pints (1.5 lbs ae) per acre per year.</p> <p>Pre-harvest Interval (PHI): Allow a minimum of 7 days between application and harvest of cotton.</p> <p>Do not apply this product to cotton grown for seed, as a reduction in germination or vigor may occur.</p>

12.5 - ROUNDUP READY FLEX COTTON

ATTENTION: USE OF THIS PRODUCT IN ACCORDANCE WITH LABEL DIRECTIONS IS EXPECTED TO RESULT IN NORMAL GROWTH OF ROUNDUP READY COTTON, HOWEVER, VARIOUS ENVIRONMENTAL CONDITIONS, AGRONOMIC PRACTICES AND OTHER FACTORS MAKE IT IMPOSSIBLE TO ELIMINATE ALL RISKS ASSOCIATED WITH THIS PRODUCT, EVEN WHEN APPLICATIONS ARE MADE IN CONFORMANCE WITH THE LABEL SPECIFICATIONS. IN SOME CASES, THESE FACTORS CAN RESULT IN BOLL LOSS, DELAYED MATURITY AND/OR YIELD LOSS.

TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
		See Section 12.0. Do not apply more than 6 quarts (6 lbs ae) per acre per year total for all application types.
Pre-plant, Pre-emergence, At-planting	<p>This product may be applied before, during or after planting Roundup Ready Flex cotton. Always plant into a weed free seedbed. In no-till and stale seedbed systems, always burn down existing weeds before cotton emerges.</p> <p>Make applications according to the rates listed in Annual Weeds, Perennial Weeds, and Woody Brush & Trees rate tables in this label.</p> <p>See the "ROUNDUP READY CROPS" section of this label for precautionary instructions for use in Roundup Ready crops.</p>	Do not apply more than 3.75 quarts (3.75 lbs ae) per acre per year.
Post-emergence (Over-the-Top)	<p>When applied in accordance with this label, AgSaver™ Glyphosate 5.4 Plus herbicide will control labeled annual grasses and broadleaf weeds in Roundup Ready Flex cotton. To maximize yield potential spray cotton early to eliminate competing weeds. Many perennial weeds will be controlled or suppressed with one or more applications of this product.</p> <p>In-crop application rates above 1.5 pints (0.75 lb ae) per acre made alone or with the addition of other crop chemical products containing surfactant may cause a crop response including leaf speckling or leaf necrosis.</p> <p>Make an initial application of 1.5 pints (0.75 lb ae) per acre on 1 to 3 inch tall annual grass and broadleaf weeds. This product may be applied by ground application equipment at rates up to 2.25 pints (1.13 lbs ae) per acre per application post-emergence to Roundup Ready Flex cotton. In addition to broadcast applications, post-directed equipment may be used to achieve weed coverage.</p> <p>Application after 10th leaf or 10th node may result in plant injury and yield loss.</p> <p>NOTE: For specific rates of application and instructions, refer to the "ANNUAL WEEDS" and "PERENNIAL WEEDS RATE SECTION" in the label booklet for AgSaver™ Glyphosate 5.4 Plus herbicide.</p>	<p>Do not apply more than 2.25 pints (1.13 lbs ae) per acre per application by ground.</p> <p>Do not apply more than 1.5 pints (0.75 lb ae) per acre per application by air.</p> <p>Between layby and 60 percent open bolls, do not apply more than 3 pints (1.5 lbs ae) per acre per year.</p> <p>Do not apply more than 4.5 quarts (4.5 lbs ae) per acre per year.</p>
Pre-harvest	<p>This product may be applied for pre-harvest annual and perennial weed control as a broadcast treatment to Roundup Ready Flex cotton after 60 percent boll crack. Up to 3 pints (1.5 lbs ae) of this product may be applied using either aerial or ground spray equipment.</p> <p>NOTE: This product will not enhance the performance of harvest aids when applied to Roundup Ready Flex cotton.</p>	<p>Do not apply more than 3 pints (1.5 lbs ae) per acre per application.</p> <p>Do not apply more than 3 pints (1.5 lbs ae) per acre per year.</p> <p>Pre-harvest Interval (PHI): Allow a minimum of 7 days between application and harvest of cotton.</p> <p>Do not apply this product to cotton grown for seed, as a reduction in germination or vigor may occur.</p> <p>The use of additives, other than those listed on this label, for preharvest application to cotton is prohibited.</p>

12.6 - ROUNDUP READY SOYBEANS[*]		
[* Not Registered for Use by California].		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
		See Section 12.0.
Pre-plant, Pre-emergence, At-planting	<p>This product may be applied before, during or after planting soybeans. Make applications according to the rates listed in Annual Weeds, Perennial Weeds, and Woody Brush & Trees rate tables in this label.</p> <p>See the "ROUNDUP READY CROPS" section of this label for precautionary instructions for use in Roundup Ready crops.</p>	
Post-emergence (In-Crop)	<p>When applied as directed, this product will control labeled annual grasses and broadleaf weeds in Roundup Ready soybeans. Applications of this product can be made in Roundup Ready soybeans from emergence (cracking) throughout flowering. Refer to the "ANNUAL WEEDS RATE TABLE" in this label for specific rates on various annual weeds.</p> <p>Make an initial application of 1.5 pints (0.75 lb ae) per acre on 2- to 8-inch tall weeds for best results. Weeds will generally be 2 to 8 inches tall, 2 to 5 weeks after planting. If the initial application is delayed and weeds are larger, apply a higher rate of this product. This product may be used up to 3 pints (1.5 lbs ae) per acre in any single in-crop application for control of annual weeds and where heavy weed densities exist.</p> <p>1.5-3 pints (0.75-1.5 lbs ae) per acre per acre rate (single or multiple applications) of this product will control or suppress perennial weeds such as: bermudagrass, Canada thistle, common milkweed, field bindweed, hemp dogbane, horsenettle, marestail (horseweed), nutsedge, quackgrass, rhizome johnsongrass, redvine, trumpetcreeper, swamp smartweed and wirestem muhly. For best results, allow perennial weed species to achieve at least 6 inches of growth before spraying with this product.</p> <p>Under adverse growing conditions such as drought, hail, wind damage or a poor soybean stand that slows or delays canopy closure, a sequential application of this may be necessary to control late flushes of weeds. In the Southern States, a sequential application of this product will be required to control new flushes of weeds in the Roundup Ready Soybean crop. To control giant ragweed, apply up to 1.5 pints (0.75 lb ae) per acre of this product when the weed is 8 to 12 inches tall to increase control and possibly avoid the need for a sequential application.</p>	<p>Do not apply more than 3 pints (1.5 lbs ae) per acre per application.</p> <p>During flowering, do not apply more than 3 pints (1.5 lbs ae) per acre per year.</p> <p>Do not apply more than 4.5 pints (2.25 lbs ae) per acre per year.</p>

12.7 - ROUNDUP READY SUGAR BEETS		
<p>The Roundup Ready designation indicates that the sugar beet contains a patented gene, which provides tolerance to this product. Information on Roundup Ready sugarbeet may be obtained from your seed supplier or AgSaver representative. Roundup Ready crop varieties must be purchased from an authorized licensed seed supplier.</p> <p>Do NOT combine these instructions with those listed for crop varieties that do not contain a Roundup Ready gene listed in the "ANNUAL AND PERENNIAL CROPS (Alphabetical)" sections of the AgSaver™ Glyphosate 5.4 Plus herbicide label booklet.</p>		
TYPES OF APPLICATIONS	USE RESTRICTIONS	RESTRICTIONS
		<p>See Section 12.0.</p> <p>Do not apply more than 6 quarts (6 lbs ae) per acre per year total for all application types.</p>

12.7 - ROUNDUP READY SUGAR BEETS

The Roundup Ready designation indicates that the sugar beet contains a patented gene, which provides tolerance to this product. Information on Roundup Ready sugarbeet may be obtained from your seed supplier or AgSaver representative. Roundup Ready crop varieties must be purchased from an authorized licensed seed supplier.

Do NOT combine these instructions with those listed for crop varieties that do not contain a Roundup Ready gene listed in the "ANNUAL AND PERENNIAL CROPS (Alphabetical)" sections of the AgSaver™ Glyphosate 5.4 Plus herbicide label booklet.

TYPES OF APPLICATIONS	USE RESTRICTIONS	RESTRICTIONS
Pre-plant, At-Planting, Pre-emergence	This product may be applied before, during or after planting of Roundup Ready sugar beets. Make applications according to the rates listed in Annual Weeds, Perennial Weeds, and Woody Brush & Trees rate tables in this label.	Do not apply more than 7.5 pints (3.75 lbs ae) per acre per year.
Post-emergence (In-crop)	This product may be applied over the top of Roundup Ready sugar beets for control of annual grasses and broadleaf weeds from emergence to 30 days prior to harvest. To maximize yield potential, spray sugar beets early to eliminate competing weeds. Up to 4 sequential applications of this product may be made with at least 10 days between applications. This product will control or suppress most perennial weeds. For some perennial weeds, repeat applications may be required to eliminate crop competition throughout the growing season.	From emergence to the 8-leaf stage, do not apply more than 2.25 pints (1.13 lbs ae) per acre per application. From the 8-leaf stage to canopy closure, do not apply more than 1.5 pints (0.75 lb ae) per acre per application. From emergence to the 8-leaf stage, do not apply more than 3.75 pints (1.88 lbs ae) per acre per year. From the 8-leaf stage to canopy closure, do not apply more than 3 pints (1.5 lbs ae) per acre per year. Do not apply more than 6.75 pints (3.38 lbs ae) per acre per year. Pre-harvest Interval (PHI): Allow a minimum of 30 days between last application and sugar beet harvest.

12.8 - ROUNDUP READY ALFALFA

FOR POSTEMERGENCE APPLICATION ONLY ON ALFALFA VARIETIES DESIGNATED AS CONTAINING A ROUNDUP RADY GENE.

The Roundup Ready designation indicates that the alfalfa contains a patented gene, which provides tolerance to this product. Information on Roundup Ready alfalfa varieties may be obtained from your seed supplier or AgSaver representative. Roundup Ready crop varieties must be purchased from an authorized licensed seed supplier.

TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
		See Section 12.0. For year of establishment, do not apply more than 5.8 quarts (5.8 lbs ae) per acre per year total for all application types. For newly established and established stands, do not apply more than 4.5 quarts (4.5 lbs ae) per acre per year total for all application types.
Pre-plant, At-planting, Pre-emergence and Post-emergence	This product will control many troublesome emerged weeds with over-the-top applications in Roundup Ready alfalfa. For ground applications with broadcast equipment, apply this product in 3 to 40 gallons of spray solution per acre. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets. For aerial application: Use the labeled rates of this product in 3 to 15 gallons of spray solution per acre. New Stand Establishment (seeding year) Prior to First Cutting During New Stand Establishment: From emergence up to 4 trifoliate leaves 3 pints (1.5 lbs ae) per acre From 5 trifoliate leaves up to 5 days before first cutting 3 pints (1.5 lbs ae) per acre After First Cutting in Newly Established Stands: In-crop application, per cutting, up to 5 days before cutting 3 pints (1.5 lbs ae) per acre Established Stands (non-seeding year) In-Crop applications, per cutting, up to 5 days before cutting 3 pints (1.5 lbs ae) per acre	For pre-plant, at-planting, and pre-emergence applications, do not apply more than 3 pints (1.5 lbs ae) per acre per application. Do not apply more than 3 pints (1.5 lbs ae) per acre per over-the-top or aerial application. Do not apply more than 4.5 quarts (4.5 lbs ae) per acre per year. Sequential applications of this production must be at least 7 days apart. Remove domestic livestock before application and wait a minimum of 5 days after last application before grazing or cutting and feeding of Roundup Ready alfalfa forage and hay.

12.8 - ROUNDUP READY ALFALFA

FOR POSTEMERGENCE APPLICATION ONLY ON ALFALFA VARIETIES DESIGNATED AS CONTAINING A ROUNDUP RADY GENE.

The Roundup Ready designation indicates that the alfalfa contains a patented gene, which provides tolerance to this product. Information on Roundup Ready alfalfa varieties may be obtained from your seed supplier or AgSaver representative. Roundup Ready crop varieties must be purchased from an authorized licensed seed supplier.

TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
	<p>During stand establishment, due to the biology and breeding constraints of alfalfa, up to 10% of the seedlings may not contain the Roundup Ready gene and will not survive after the first application of this product. To eliminate the undesirable effects of stand gaps created by the loss of plants not containing a Roundup Ready gene, a single application of at least 1.5 pints (0.75 lb ae) per acre of this product should be applied at or before the 3 to 4 trifoliate growth stage.</p> <p>In both newly seeded and established stands, in order to maximize yield and quality potential of forage and hay, applications of this product should be made after weeds have emerged but before alfalfa growth or re-growth interferes with application spray coverage of the target weeds.</p> <p>In addition to those weeds listed in the AGSAVER™ GLYPHOSATE 5.4 PLUS herbicide label booklet, this product will suppress or control the parasitic weed, Dodder (<i>Cuscuta</i> spp.) in Roundup Ready alfalfa. Repeat applications may be necessary for complete control.</p>	
Over-the-top applications	<p>This product may be applied post-emergence to Roundup Ready alfalfa from emergence until 5 days prior to cutting. Any single over-the-top applications of this product must not exceed 3 pints (1.5 lbs ae) per acre.</p> <p>ATTENTION: Where Roundup Ready alfalfa is grown with a companion or cover crop, or is over seeded with a second species, over-the-top applications of this product will eliminate the non-Roundup Ready species.</p>	<p>Do not apply more than 3 pints (1.5 lbs ae) per acre per application.</p> <p>Sequential applications of this production must be at least 7 days apart.</p> <p>Tank mixtures with other herbicides, insecticides, or fungicides may result in crop injury or reduced weed control and must not be used for over-the-top applications of this product.</p>

13.0 - NON-CROP USES AROUND THE FARMSTEAD

RESTRICTIONS:

- **DO NOT** apply more than 7.5 pints (3.75 lbs ae) per acre per application, unless otherwise specified for an individual use.
- **DO NOT** apply more than 6 quarts (6 lbs ae) per acre per year as a combined total of all treatments for applications in non-agricultural sites, unless otherwise specified for an individual use.

13.1 - WEED CONTROL & TRIM-AND-EDGE

LABELED SITES: Non-crop Areas including building foundations, along and in fences, in dry ditches and canals, along ditchbanks, farm roads, shelterbelts, prior to landscape plantings and equipment storage areas.

TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
Any suitable application equipment described in Section 8.0 of this label	<p>This product may be used to control annual weeds, perennials weeds and woody brush which are found in any part of the farmstead.</p> <p>Make applications according to the rates listed in Annual Weeds, Perennial Weeds, and Woody Brush & Trees rate tables in this label.</p> <p>TANK MIXTURES: This product may be tank mixed with the following products. Refer to these product labels for approved farmstead sites and application rates. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.</p> <p>For annual weeds, use 1.5 pints (0.75 lb ae) per acre of this product when weeds are less than 6 inches tall and 2.25 pints (1.13 lbs ae) per acre when weeds are greater than 6 inches tall. For perennial weeds, apply 1.5-3.75 quarts (1.5-3.75 lbs ae) per acre in these tank mixes.</p>	Not Registered for Use by California for dicamba tank mixtures applied by air.

13.1 - WEED CONTROL & TRIM-AND-EDGE				
LABELED SITES: Non-crop Areas including building foundations, along and in fences, in dry ditches and canals, along ditchbanks, farm roads, shelterbelts, prior to landscape plantings and equipment storage areas.				
TYPES OF APPLICATIONS	USE DIRECTIONS			RESTRICTIONS
	For tank mixtures with these products through backpack sprayers, handguns or other high-volume spray-to-wet applications, see the "HAND-HELD AND HIGH VOLUME EQUIPMENT" section of this label for allowable application rates.			
	2,4-D bromacil + diuron chlorsulfuron dicamba, diglycolamine salt diuron	diuron + imazapyr imazapic imazapyr metsulfuron oryzalin oxadiazon	pendimethalin prodiamine simazine sulfometuron	
	For control or partial control of the following perennial weeds, apply 1.5-3 pints (0.75-1.5 lbs ae) of this product plus the labeled rate of sulfometuron per acre. Bahiagrass Bermudagrass Broomsedge Dallisgrass Dock, curly Dogfennel			
			Fescue, tall Johnsongrass Poorjoe Quackgrass Vaseygrass Vervain, blue	

13.2 - GREENHOUSE/SHADEHOUSE		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
Spot Spray, Directed Spray	<p>This product may be used to control weeds in and around greenhouses and shadehouses.</p> <p>Make applications according to the rates listed in Annual Weeds, Perennial Weeds, and Woody Brush & Trees rate tables in this label.</p>	<p>Air circulation fans must be turned off during application.</p> <p>Desirable vegetation must not be present during application.</p>

13.3 – CHEMICAL MOWING		
LABELED USES: Farm Ditches and Other Parts of Farmsteads		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
Any suitable application equipment described in Section 8.0 of this label	<p>This product will suppress perennial grasses listed in this section to serve as a substitute for mowing. Use 6 fl oz (0.19 lb ae) of this product per acre when treating tall fescue, fine fescue, orchardgrass or quackgrass covers. Use 4.5 fl oz (0.14 lb ae) of this product per acre when treating Kentucky bluegrass. Use 12 fl oz (0.38 lb ae) of this product when treating bermudagrass. Use 3 pints (1.5 lbs ae) of this product when treating torpedograss or paragrass. Apply treatments in 10 to 20 gallons of spray solution per acre.</p>	Use only in areas where some temporary injury or discoloration of perennial grasses can be tolerated.

13.4 – CUT STUMPS		
LABELED USES: Cut Stumps (on any non-crop site listed on this label)		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
Suitable Hand-Held Equipment	<p>This product will control regrowth of cut stumps and resprouts of many types of woody brush and tree species, some of which are listed below. Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut trees or resprouts close to the soil surface. Apply a 50 to 100 percent solution of this product to the freshly cut surface immediately after cutting.</p> <p>Delays in application may result in reduced performance. For best results, applications should be made during periods of active growth and full leaf expansion.</p> <p>Some sprouts, stems, or trees may share the same root system. Adjacent trees having a similar age, height and spacing may signal shared roots. Whether grafted or shared, injury is likely to occur to non-treated stems/trees when one or more trees sharing common roots are treated.</p>	

13.4 – CUT STUMPS				
LABELED USES: Cut Stumps (on any non-crop site listed on this label)				
TYPES OF APPLICATIONS	USE DIRECTIONS			RESTRICTIONS
	Alder Eucalyptus Madrone Oak	Pepper, brazilian Pine, Austrian Reed, giant Salt cedar	Sweetgum Tan oak Willow	

13.5 – HABITAT MANAGEMENT		
LABELED USES: Habitat Restoration & Maintenance, Wildlife Food Plots		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
Any suitable application equipment described in Section 8.0 of this label	<p>This product may be used to control exotic and other undesirable vegetation in habitat management and natural areas including rangeland and wildlife refuges. Applications can be made to allow recovery of native plant species, prior to planting desirable native species, and for similar broad spectrum vegetation control requirements in habitat management areas.</p> <p>Make applications according to the rates listed in Annual Weeds, Perennial Weeds, and Woody Brush & Trees rate tables in this label.</p> <p>Spot treatments can be made to selectively remove unwanted plants for habitat maintenance and enhancement.</p> <p>This product may be used as a site preparation treatment to control annual and perennial weeds prior to planting wildlife food plots. Any wildlife food species may be planted after applying this product, or native species may be allowed to repopulate the area.</p>	If tillage is needed to prepare a seedbed, wait 7 days after application before tillage to allow translocation into underground plant parts.

14.0 – FORESTRY, INDUSTRIAL, TURF & ORNAMENTAL

14.1 – FORESTRY SITE PREPARATION										
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS								
Boom Sprayers, Shielded Boom Sprayers, High-Volume Off-Center Nozzles, Hand-Held Equipment, And Similar Equipment.	<p>This product can be used for the control or partial control of woody brush, trees, and herbaceous weeds in forestry, as well as for use in preparing or establishing wildlife openings with these sites and maintaining logging roads.</p> <p>Make applications according to the rates listed in Annual Weeds, Perennial Weeds, and Woody Brush & Trees rate tables in this label.</p> <p>This product can be used in site preparation prior to planting any tree species, including Christmas trees, eucalyptus, hybrid tree cultivars and silvicultural nursery sites.</p> <p>Use higher rates of this product within the labeled range for control or partial control of woody brush, trees, and hard-to-control perennial herbaceous weeds. For best results, apply to actively growing woody brush and trees after full leaf expansion and before fall color and leaf drop. Increase rates within the labeled range for control of perennial herbaceous weeds any time after emergence and before seedheads, flowers or berries appear.</p> <p>Use the lower rates of this product within the labeled range for control of annual herbaceous weeds and actively growing perennial herbaceous weeds after seedheads, flowers or berries appear. Apply to the foliage of actively growing annual herbaceous weeds any time after emergence.</p> <p>TANK MIXTURES: Tank mixtures of this product may be used to increase the spectrum of vegetation controlled. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.</p> <p>Any labeled rate of this product may be used in a tank mix with the following products for forestry site preparation.</p> <table><tr><td>chlorsulfuron + sulfometuron</td><td>sulfometuron</td></tr><tr><td>hexazinone + sulfometuron</td><td>triclopyr, butoxyethyl ester</td></tr><tr><td>imazapyr</td><td>triclopyr, triethylamine salt</td></tr><tr><td>metsulfuron</td><td></td></tr></table> <p>For control of herbaceous weeds, use the lower labeled tank mixture rates. For control of dense stands or tough-to-control woody brush and trees, use the higher labeled rates.</p>	chlorsulfuron + sulfometuron	sulfometuron	hexazinone + sulfometuron	triclopyr, butoxyethyl ester	imazapyr	triclopyr, triethylamine salt	metsulfuron		<p>Do not apply this product as an over-the-top broadcast spray for forestry conifer or hardwood release.</p>
chlorsulfuron + sulfometuron	sulfometuron									
hexazinone + sulfometuron	triclopyr, butoxyethyl ester									
imazapyr	triclopyr, triethylamine salt									
metsulfuron										

14.2 – NONCROP AREAS & INDUSTRIAL SITES

Labeled Uses: Non-crop areas including airports, apartment complexes, Christmas tree farms, commercial sites, Conservation Reserve Program (CRP) areas, ditch banks, dry ditches, dry canals, fencerows, gold courses, greenhouses, industrial sites, landscape areas, lumber yards, manufacturing sites, municipal sites, natural areas, office complexes, ornamentals, parks, parking areas, pastures, petroleum tank farms and pumping installations, plant nurseries, public areas, railroads, rangeland, recreational areas, residential areas, rights-of-way, roadsides, schools, sod or turf seed farms, sports complexes, storage areas, substations, turfgrass areas, utility sites, warehouse areas, and wildlife management areas.																																									
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS																																							
This product may be applied with any suitable application equipment described in Section 8.0 of this label.	<p>This product may be used to trim-and-edge around objects in non-crop sites, for spot treatment of unwanted vegetation and to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. This product may be used prior to planting an areas to ornamentals, flowers, turfgrass (sod or seed), or prior to laying asphalt or beginning construction projects.</p> <p>Make applications according to the rates listed in Annual Weeds, Perennial Weeds, and Woody Brush & Trees rate tables in this label.</p> <p>Repeated applications of this product may be used, as weeds emerge, to maintain bare ground.</p> <p>TANK MIXTURES: This product may be tank mixed with the following products provided that the specific product is registered for use on the target site. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.</p> <table> <tr> <td>2,4-D</td><td>dicamba</td><td>oxyfluorfen</td></tr> <tr> <td>2,4-D + triclopyr</td><td>diuron</td><td>pendimethalin</td></tr> <tr> <td>butoxyethyl ester</td><td>hexazinone</td><td>proflaminate</td></tr> <tr> <td>atrazine</td><td>imazapic</td><td>sethoxydim</td></tr> <tr> <td>bromacil + diuron</td><td>imazapyr</td><td>simazine</td></tr> <tr> <td>chlorsulfuron</td><td>isoxaben</td><td>sulfometuron</td></tr> <tr> <td>chlorsulfuron + sulfometuron</td><td>metsulfuron</td><td>sulfosulfuron</td></tr> <tr> <td>clopyralid</td><td>oryzalin</td><td>triclopyr, butoxyethyl ester</td></tr> <tr> <td></td><td>oxadiazon</td><td>triclopyr, triethylamine salt</td></tr> </table> <p>When applied as a tank mixture for bare ground, this product provides control of the emerged annual weeds and control of partial control of emerged perennial weeds, woody brush, and trees.</p> <p>For control or partial control of the following perennial weeds, apply 1.5 – 3 pints (0.75-1.5 lbs ae) of this product plus the labeled rate of sulfometuron per acre.</p> <table> <tr> <td>Bahiagrass</td><td>Dock, curly</td><td>Poorjoe</td></tr> <tr> <td>Bermudagrass</td><td>Dogfennel</td><td>Quackgrass</td></tr> <tr> <td>Broomsedge</td><td>Fescue, tall</td><td>Vaseygrass</td></tr> <tr> <td>Dallisgrass</td><td>Johnsongrass</td><td>Vervain, blue</td></tr> </table>	2,4-D	dicamba	oxyfluorfen	2,4-D + triclopyr	diuron	pendimethalin	butoxyethyl ester	hexazinone	proflaminate	atrazine	imazapic	sethoxydim	bromacil + diuron	imazapyr	simazine	chlorsulfuron	isoxaben	sulfometuron	chlorsulfuron + sulfometuron	metsulfuron	sulfosulfuron	clopyralid	oryzalin	triclopyr, butoxyethyl ester		oxadiazon	triclopyr, triethylamine salt	Bahiagrass	Dock, curly	Poorjoe	Bermudagrass	Dogfennel	Quackgrass	Broomsedge	Fescue, tall	Vaseygrass	Dallisgrass	Johnsongrass	Vervain, blue	Not Registered for Use by California for dicamba tank mixtures applied by air.
2,4-D	dicamba	oxyfluorfen																																							
2,4-D + triclopyr	diuron	pendimethalin																																							
butoxyethyl ester	hexazinone	proflaminate																																							
atrazine	imazapic	sethoxydim																																							
bromacil + diuron	imazapyr	simazine																																							
chlorsulfuron	isoxaben	sulfometuron																																							
chlorsulfuron + sulfometuron	metsulfuron	sulfosulfuron																																							
clopyralid	oryzalin	triclopyr, butoxyethyl ester																																							
	oxadiazon	triclopyr, triethylamine salt																																							
Bahiagrass	Dock, curly	Poorjoe																																							
Bermudagrass	Dogfennel	Quackgrass																																							
Broomsedge	Fescue, tall	Vaseygrass																																							
Dallisgrass	Johnsongrass	Vervain, blue																																							

14.3 – INJECTION & FRILL (Woody Brush & Trees)

Labeled Sites: Woody brush & Trees in non-crop areas												
Types of Applications	USE DIRECTIONS	RESTRICTIONS										
Injection or Frill Applications	<p>Apply this product using suitable equipment which must penetrate into the living tissue. Apply the equivalent of 0.75 mL of this product per each 2 to 3 inches of trunk diameter at breast height (DBH). This is best achieved by applying a 50 to 100 percent concentration of this product either to a continuous frill around the tree or as cuts evenly spaced around the tree below all branches. As tree diameter increases in size, better results are achieved by applying diluted material to a continuous frill or more closely spaced cuttings.</p> <p>For best results, application should be made during periods of active growth and after full leaf expansion. This product will control many species, some of which are listed below:</p> <table><tr><td><u>Control</u></td><td><u>Partial Control</u></td></tr><tr><td>Oak</td><td>Black gum</td></tr><tr><td>Poplar</td><td>Dogwood</td></tr><tr><td>Sweetgum</td><td>Hickory</td></tr><tr><td>Sycamore</td><td>Maple, red</td></tr></table>	<u>Control</u>	<u>Partial Control</u>	Oak	Black gum	Poplar	Dogwood	Sweetgum	Hickory	Sycamore	Maple, red	<p>Do not use application techniques that allow runoff to occur from frilled or cut areas in species that exude sap freely.</p> <p>In species such as this, make the frill or cuts at an oblique angle to produce a cupping effect and use a 100 percent concentration of this product.</p>
<u>Control</u>	<u>Partial Control</u>											
Oak	Black gum											
Poplar	Dogwood											
Sweetgum	Hickory											
Sycamore	Maple, red											

14.4 – HOLLOW STEM INJECTION

LABELED SITES: Hollow-stem plants growing in any non-crop site specified on this label.		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
Hand-Held Injection Devices That Deliver Labeled Amounts Of This Product	<p>For control of the following hollow-stem plants, use the application rates below:</p> <p>Japanese Knotweed, <i>Polygonum cuspidatum</i> Inject 3.75 mL per stem of this product between second and third internode.</p> <p>Bohemian Knotweed, <i>Polygonum bohemicum</i> Inject 3.75 mL per stem of this product between the second and third internode.</p> <p>Giant Hogweed, <i>Hercleum mantegazzianum</i> Inject one leaf cane per plant 12 inches above the root brown with 3.75 mL of a 5% v/v solution of this product.</p> <p>Poison Hemlock, <i>Conium maculatum</i> Inject one leaf cane per plant 10 to 12 inches above the root crown with 3.75 mL a 5% v/v solution of this product.</p> <p>Field horsetail, <i>Equisetum arvense</i> Inject one segment above the root crown with 3.75 mL per stem of this product. Use a small syringe that calibrates to this rate.</p> <p>Canada Thistle, <i>Cirsium arvense</i> Cut 8 to 9 of the tallest plants at bud stage in a clump with clippers. Use a cavity needle that is pushed into the stem center and then slowed removed as 3.75 mL per stem of this product is injected into the stem.</p>	<p>Do not apply more than 5.25 quarts (5.25 lbs ae) per acre per year.</p> <p>At 3.75 mL per stem, 5.25 quarts (5.25 lbs ae) should treat approximately 1300 stems per acre.</p>

14.5 – ORNAMENTALS, PLANT NURSERIES & CHRISTMAS TREES

LABELED SITES: Plant Nurseries, Christmas Tree farms & other non-food tree production sites		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
Post-Directed, Trim-and-Edge	<p>This product may be used as a post-directed spray around established woody ornamental species (including arborvitae, azalea, boxwood, crabapple, eucalyptus, euonymus, fir, Douglas fir, jojoba, hollies, lilac, magnolia, maple, oak, poplar, privet, pine, spruce and yew, growing in plant nurseries, on Christmas tree farms, or on other non-food tree production sites), or to trim and edge around trees, buildings, sidewalks, roads, potted plants and other objects in a production setting.</p> <p>Apply at a concentration labeled by Section 15.0 or Section 16.0 or Section 16.1 or Section 17.0 appropriate to the species of weed to be controlled.</p> <p>Avoid contact of spray, drift or mist with foliage or green bark of desirable ornamental species. Desirable plants may be protected from the spray solution by using shields or coverings made of cardboard or other impermeable material.</p>	This product is not allowed for use as an over-the-top broadcast spray in ornamentals and Christmas trees.
Site Preparation	This product may be used prior to planting any tree, shrub, or vine, including Christmas tree species, in a nursery or production setting.	
Wiper Application	This product may be used through wick or other suitable wiper applicators to control or partially control undesirable vegetation around established trees, shrubs, or vines. See the "SELECTIVE EQUIPMENT" section of this label for further information about the proper use of wiper applicators.	

14.6 – PARKS, RECREATIONAL & RESIDENTIAL AREAS

LABELED SITES: Around Trees, Fences, Paths, Driveways, Around Buildings, Patios, Sidewalks, Flower Beds, Around Shrubs and other Ornamental Plants		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
Trim-and-Edge, Spot Treatment	<p>This product may be used to eliminate unwanted weeds growing in areas listed above.</p> <p>Use suitable handheld equipment for directed spraying according to instructions in Section 7.3 "MIXING FOR HAND-HELD SPRAYERS."</p> <p>If necessary, use cardboard or plastic to shield desirable plants.</p>	<p>Do not allow spray, drift, or mist to contact foliage or green bark of desirable ornamental species.</p> <p>Do not use for spot weed control in lawns since desirable lawn grass will also be killed.</p>
Site Preparation, Lawn Renovation	<p>This product may be used prior to planting an area to ornamentals, flowers, turfgrass (sod or seed), lawn renovation or prior to laying asphalt or beginning construction projects.</p> <p>Make applications according to the rates listed in Annual Weeds, Perennial Weeds,</p>	<p>Do not mow for 7 days before or after treatment.</p>

14.6 – PARKS, RECREATIONAL & RESIDENTIAL AREAS		
LABELED SITES: Around Trees, Fences, Paths, Driveways, Around Buildings, Patios, Sidewalks, Flower Beds, Around Shrubs and other Ornamental Plants		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
	<p>and Woody Brush & Trees rate tables in this label.</p> <p>Apply using suitable broadcast or directed spray equipment.</p> <p>For lawn renovation, thorough coverage is necessary to kill all weeds and old lawn.</p> <p>For best results, apply when daytime temperatures are at least 60°F.</p> <p>7 days after application, soil may be tilled, fertilized, and seeded.</p>	

14.7 – RAILROADS																													
LABELED SITES: Railroad Rights-of-Way, Railroad Ballast areas																													
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS																											
<p>Boom Sprayers, Shielded Boom Sprayers, High-Volume Off-Center Nozzles, Hand-Held Equipment</p>	<p>Observe application precautions in Section 8.0.</p> <p>All of the instructions in the "NONCROP AREAS AND INDUSTRIAL SITES" section apply to railroads.</p> <p>Make applications according to the rates listed in Annual Weeds, Perennial Weeds, and Woody Brush & Trees rate tables in this label.</p> <p>This product may be used to maintain bare ground on railroad ballast and shoulders. Repeat applications of this product may be used, as weeds emerge, to maintain bare ground. This product may be used to control tall-growing weeds to improve line-of-sight at railroad crossings and reduce the need for mowing along rights-of-way. For crossing applications, up to 80 gallons of spray solution per acre may be used.</p> <p>This product may be used to maintain bare ground on railroad ballast and shoulders. Repeat applications of this product may be used, as weeds emerge, to maintain bare ground. This product may be used to control tall-growing weeds to improve line-of-sight at railroad crossings and reduce the need for mowing along rights-of-way. For crossing applications, up to 80 gallons of spray solution per acre may be used.</p> <p>Avoid application to non-target plants due to drift, overspray, or runoff.</p> <p>TANK MIXTURES: This product may be tank mixed with the following products for ballast, shoulder, spot, bare ground and crossing treatments provided that the specific product is registered for use on such sites. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.</p> <table border="0"> <tr> <td>2,4-D</td><td>diuron</td><td>sulfometuron</td></tr> <tr> <td>bromacil</td><td>diuron + imazapyr</td><td>tebuthiuron</td></tr> <tr> <td>bromacil + diuron</td><td>hexazinone</td><td>triclopyr, butoxyethyl ester</td></tr> <tr> <td>chlorsulfuron</td><td>imazapyr</td><td>triclopyr, triethylamine salt</td></tr> <tr> <td>dicamba</td><td>metsulfuron</td><td></td></tr> </table> <p>Brush control</p> <p>This product may be used to control woody brush and trees on railroad rights-of-way. Apply 3-7.5 quarts (3-7.5 lbs ae) of this product per acre as a broadcast spray, using boom-type or boomless nozzles. Up to 80 gallons of spray solution per acre may be used. Apply a 3/4 to 2 percent solution of this product when using high-volume spray-to-wet applications. Apply a 5 to 10 percent solution of this product when using low volume directed sprays for spot treatment. This product may be mixed with the following products for enhanced control of woody brush and trees:</p> <table border="0"> <tr> <td>chlorsulfuron</td><td>fosamine</td><td>picloram</td></tr> <tr> <td>clopyralid</td><td>hexazinone</td><td>triclopyr, butoxyethyl ester</td></tr> <tr> <td>dicamba, diglycolamine salt</td><td>imazapyr</td><td>triclopyr, triethylamine salt</td></tr> <tr> <td></td><td>metsulfuron</td><td></td></tr> </table>	2,4-D	diuron	sulfometuron	bromacil	diuron + imazapyr	tebuthiuron	bromacil + diuron	hexazinone	triclopyr, butoxyethyl ester	chlorsulfuron	imazapyr	triclopyr, triethylamine salt	dicamba	metsulfuron		chlorsulfuron	fosamine	picloram	clopyralid	hexazinone	triclopyr, butoxyethyl ester	dicamba, diglycolamine salt	imazapyr	triclopyr, triethylamine salt		metsulfuron		
2,4-D	diuron	sulfometuron																											
bromacil	diuron + imazapyr	tebuthiuron																											
bromacil + diuron	hexazinone	triclopyr, butoxyethyl ester																											
chlorsulfuron	imazapyr	triclopyr, triethylamine salt																											
dicamba	metsulfuron																												
chlorsulfuron	fosamine	picloram																											
clopyralid	hexazinone	triclopyr, butoxyethyl ester																											
dicamba, diglycolamine salt	imazapyr	triclopyr, triethylamine salt																											
	metsulfuron																												

14.8 – ROADSIDES																	
LABELED SITES: Roadside Rights of Way areas (including Shoulders, Guardrails and Signposts)																	
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS															
Boom Sprayers, Shielded Boom Sprayers, High-Volume Off-Center Nozzles, Hand-Held Equipment, And Similar Equipment.	<p>Observe application precautions in Section 8.0.</p> <p>All the instructions in the “NONCROP AREAS AND INDUSTRIAL SITES” section apply to roadsides.</p> <p>Make applications according to the rates listed in Annual Weeds, Perennial Weeds, and Woody Brush & Trees rate tables in this label.</p> <p>This product may be used on road shoulders, under guardrails and around signposts and other objects along roadsides that may be obstacles to mowing.</p> <p>Avoid application to non-target plants due to drift, overspray, or runoff.</p> <p>TANK MIXTURES: This product may be tank-mixed with the following products for shoulder, guardrail, spot and bare ground treatments:</p> <table> <tr> <td>2,4-D</td><td>diuron</td><td>pendimethalin</td></tr> <tr> <td>bromacil + diuron</td><td>diuron + imazapyr</td><td>proflaminate</td></tr> <tr> <td>chlorsulfuron</td><td>metsulfuron</td><td>simazine</td></tr> <tr> <td>dicamba,</td><td>oryzalin</td><td>sulfometuron</td></tr> <tr> <td>diglycolamine salt</td><td>oxadiazon</td><td></td></tr> </table> <p>It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.</p> <p>See the “NONCROP AREAS AND INDUSTRIAL SITES” section of this label for instructions for tank mixing.</p>	2,4-D	diuron	pendimethalin	bromacil + diuron	diuron + imazapyr	proflaminate	chlorsulfuron	metsulfuron	simazine	dicamba,	oryzalin	sulfometuron	diglycolamine salt	oxadiazon		
2,4-D	diuron	pendimethalin															
bromacil + diuron	diuron + imazapyr	proflaminate															
chlorsulfuron	metsulfuron	simazine															
dicamba,	oryzalin	sulfometuron															
diglycolamine salt	oxadiazon																
Spot treatment	This product may be used as a spot treatment to control unwanted vegetation growing along roadsides.																

14.9 – UTILITY SITES		
LABELED SITES: Electrical Power, Pipeline And Telephone Rights-Of-Way, And In Other Sites Associated With These Rights-Of-Way, Including Substations, Roadsides, Railroads Or Similar Rights-Of-Way That Run In Conjunction With Utilities.		
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS
Boom Sprayers, Shielded Boom Sprayers, High-Volume Off-Center Nozzles, Hand-Held Equipment, And Similar Equipment.	<p>Observe application precautions in Section 8.0.</p> <p>This product may be used in utility sites and substations to control unwanted vegetation and to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. This product may be used prior to planting a utility site to ornamentals, flowers, turfgrass (sod or seed), or beginning construction projects.</p> <p>Make applications according to the rates listed in Annual Weeds, Perennial Weeds, and Woody Brush & Trees rate tables in this label.</p> <p>Avoid application to non-target plants due to drift, overspray, or runoff.</p> <p>A follow up application of this product may be used, up to the maximum labeled rate, as weeds emerge, to maintain bare ground.</p> <p>This product can also be used when preparing or establishing wildlife openings within these sites, maintaining access roads and for side trimming along utility rights-of-way.</p> <p>For control of herbaceous weeds, use the lower labeled tank mixture rates. For control of dense stands of tough-to-control woody brush and trees, use the higher labeled rates.</p> <p>TANK MIXTURES: Tank mixtures of this product may be used to increase the spectrum of control for herbaceous weeds, woody brush, and trees. This product may be tank mixed with the following products. Refer to these products' labels for approved non-crop sites and application rates. It is the pesticide user's responsibility to ensure that all products are registered for the intended use.</p>	

14.9 – UTILITY SITES																													
LABELED SITES: Electrical Power, Pipeline And Telephone Rights-Of-Way, And In Other Sites Associated With These Rights-Of-Way, Including Substations, Roadsides, Railroads Or Similar Rights-Of-Way That Run In Conjunction With Utilities.																													
TYPES OF APPLICATIONS	USE DIRECTIONS	RESTRICTIONS																											
	<p>Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.</p> <table> <tr> <td>2,4-D</td><td>fosamine</td><td>simazine</td></tr> <tr> <td>atrazine</td><td>hexazinone</td><td>sulfometuron</td></tr> <tr> <td>bromacil + diuron</td><td>imazapic</td><td>sulfosulfuron</td></tr> <tr> <td>chlorsulfuron</td><td>imazapyr</td><td>triclopyr, butoxyethyl ester</td></tr> <tr> <td>clopyralid</td><td>metsulfuron</td><td>triclopyr,</td></tr> <tr> <td>dicamba,</td><td>oryzalin</td><td>triethylamine salt</td></tr> <tr> <td>diglycolamine salt</td><td>oxadiazon</td><td></td></tr> <tr> <td>diuron</td><td>pendimethalin</td><td></td></tr> <tr> <td>diuron + imazapyr</td><td>proflaminate</td><td></td></tr> </table>	2,4-D	fosamine	simazine	atrazine	hexazinone	sulfometuron	bromacil + diuron	imazapic	sulfosulfuron	chlorsulfuron	imazapyr	triclopyr, butoxyethyl ester	clopyralid	metsulfuron	triclopyr,	dicamba,	oryzalin	triethylamine salt	diglycolamine salt	oxadiazon		diuron	pendimethalin		diuron + imazapyr	proflaminate		
2,4-D	fosamine	simazine																											
atrazine	hexazinone	sulfometuron																											
bromacil + diuron	imazapic	sulfosulfuron																											
chlorsulfuron	imazapyr	triclopyr, butoxyethyl ester																											
clopyralid	metsulfuron	triclopyr,																											
dicamba,	oryzalin	triethylamine salt																											
diglycolamine salt	oxadiazon																												
diuron	pendimethalin																												
diuron + imazapyr	proflaminate																												

15. ANNUAL WEEDS RATE TABLES ALPHABETICALLY BY SPECIES

Water carrier volumes of 3 to 10 gallons per acre for ground applications and 3 to 5 gallons per acre for aerial applications are required.

This product may be used up to 2.25 pints (1.13 lbs ae) per acre where heavy weed densities exist. For weeds that have been mowed, grazed, or cut, allow regrowth to occur prior to treatment.

RESTRICTIONS:

Do not tank mix with soil residual herbicides when using these rates unless otherwise specified.

West Region	Alaska, Arizona, California, Colorado, Idaho, Kansas (west of Highway 83), Montana, Nebraska (west of Highway 83), Nevada, New Mexico, North Dakota (west of Highway 83), Oregon, South Dakota (west of Highway 83), Texas, Utah, Washington, Wyoming
North Region	Connecticut, Indiana (north of I-70), Iowa, Kansas (east of Highway 83 & north of I-35), Maine, Massachusetts, Minnesota, Missouri (north of I-44), Nebraska (east of Highway 83), New Hampshire, New Jersey, New York, North Dakota (east of Highway 83), Ohio, Pennsylvania, Rhode Island, South Dakota (east of Highway 83), Vermont, Wisconsin
South Region	Alabama, Arkansas, Delaware, Florida, Georgia, Hawaii, Illinois(south of I-70), Indiana (south of I-70), Kansas (east of Highway 83 & south of I-35), Kentucky, Louisiana, Maryland, Mississippi, Missouri (south of I-44), North Carolina, Oklahoma (east of I-35), South Carolina, Tennessee, Texas (east of I-35), Virginia, Washington DC, West Virginia

ANNUAL WEEDS RATE TABLE, NORTH AND SOUTH REGIONS

WEED SPECIES	REGION	RATE					
		(FLUID OUNCES PER ACRE)					
		9	12	18	24	30	36
		(0.28 lb ae)	(0.38 lb ae)	(0.56 lb ae)	(0.75 lb ae)	(0.94 lb ae)	(1.13 lbs ae)
MAXIMUM HEIGHT/LENGTH							
Annoda, spurred		-	1"	2"	3"	5"	8"
Barley		-	18"	18" +	-	-	-
Barnyardgrass	South	-	3"	5"	7"	9"	12"
	North	-	-	6"	12"	-	-
Bassia, fivehook		-	-	-	6"	-	-
Bittercress		-	12"	20"	-	-	-
Bluegrass, annual		-	10"	-	-	-	-
Brome, downy		6"	-	-	-	-	-
Brome. Japanese		-	6"	-	24"	-	-

WEED SPECIES	REGION	RATE					
		(FLUID OUNCES PER ACRE)					
		9	12	18	24	30	36
		(0.28 lb ae)	(0.38 lb ae)	(0.56 lb ae)	(0.75 lb ae)	(0.94 lb ae)	(1.13 lbs ae)
MAXIMUM HEIGHT/LENGTH							
Browntop panicum		-	6"	8"	12"	-	24"
Burcucumber		-	-	6"	12"	-	-
Buttercup		-	12"	20"	-	-	-
Carolina foxtail		-	20"	-	-	-	-
Carolina geranium		-	-	-	4"	-	9"
Carpetweed		-	-	6"	12"	-	-
Cheat		-	6"	20"	-	-	-
Chervil		-	20"	-	-	-	-
Chickweed		-	12"	18"	-	-	-
Cocklebur		-	12"	18"	24"	-	-
Copperleaf, hophornbeam		-	1"	2"	3"	4"	6"
Copperleaf, Virginia		-	1"	2"	3"	4"	6"
Corn		-	12"	20"	-	-	-
Corn speedwell		-	12"	-	-	-	-
Crabgrass		-	12"	18"	-	-	-
Cutleaf evening primrose		-	-	-	3"	3"	6"
Dwarf dandelion		-	20"	-	-	-	-
Eastern mannagrass		-	8"	12"	-	-	-
Eclipta		-	4"	8"	12"	-	-
Fall panicum	South	-	4"	6"	8"	12"	24"
	North	-	6"	12"	18"	-	-
Falsedandelion		-	20"	-	-	-	-
Falseflax, smallseed		-	12"	-	-	-	-
Fiddleneck		-	-	-	6"	6"	12"
Field pennycress		-	6"	12"	-	-	-
Filaree		-	-	-	-	-	12"
Fleabane, annual		-	6"	20"	-	-	-
Fleabane, hairy (<i>Conyza bonariensis</i>)		-	6"	-	-	-	-
Fleabane, rough		-	3"	6"	12"	-	-
Florida pusley		-	-	-	4"	4"	6"
Foxtail	South	-	8"	12"	20"	-	-
	North	18"	18" +	-	-	-	-
Goatgrass, jointed		-	6"	-	-	-	-
Goosegrass		-	3"	5"	8"	-	18"
Grain sorghum (milo)		-	6"	12"	20"	-	-
Groundsel, common		-	6"	-	-	-	-
Hemp sesbania		-	-	2"	4"	6"	8"
Henbit		-	-	-	6"	-	20"
Horseweed/Marestail (<i>Conyza canadensis</i>)	South	-	-	12"	30"	-	-
	North	-	6"	12"	18"	-	-
Itchgrass		-	6"	12"	18"	-	-
Johnsongrass, seedling	South	-	-	18"	-	-	-
	North	-	12"	18"	-	-	-

WEED SPECIES	REGION	RATE					
		(FLUID OUNCES PER ACRE)					
		9	12	18	24	30	36
		(0.28 lb ae)	(0.38 lb ae)	(0.56 lb ae)	(0.75 lb ae)	(0.94 lb ae)	(1.13 lbs ae)
MAXIMUM HEIGHT/LENGTH							
Junglerice		-	3"	5"	7"	9"	12"
Knotweed		-	3"	8"	12"	-	20"
Kochia ¹		-	3 to 6"	12"	-	-	-
Lambsquarters		-	6"	8"	12"	-	20"
Little barley		-	20"	-	-	-	-
London rocket		-	6"	-	-	-	-
Mayweed		-	-	2"	6"	12"	18"
Morningglory (<i>Ipomoea</i> spp.)		-	-	2"	4"	-	6"
Mustard, blue		6"	-	-	-	-	-
Mustard, tansy		6"	12"	20"	-	-	-
Mustard, tumble		6"	-	-	-	-	-
Mustard, wild		6"	12"	18"	-	-	-
Nightshade, black		-	6"	12"	-	-	-
Nightshade, hairy		-	6"	12"	-	-	-
Oats		-	-	6"	20"	-	-
Pigweed		-	12"	18"	24"	-	-
Prickly lettuce		-	6"	12"	20"	-	-
Purslane		-	-	-	6"	6"	12"
Ragweed, common	South	-	4"	6"	8"	-	11"
	North	-	6"	12"	18"	-	-
Ragweed, giant		-	-	4"	6"	-	11"
Red rice		-	-	-	4"	-	-
Russian thistle		-	-	-	6"	-	-
Rye	South	-	6"	20"	60"	-	-
	North	-	18"	18" +	-	-	-
Ryegrass		-	-	-	6"	-	7" +
Sandbur, field		12"	-	-	-	-	-
Shattercane		-	12"	18"	-	-	-
Sheperd's purse		-	6"	12"	-	-	-
Sicklepod		-	-	2"	4"	-	8"
Signalgrass, broadleaf		-	3"	5"	7"	9"	12"
Smartweed, ladysthumb		-	4"	6"	8"	-	12"
Smartweed, Pennsylvania		-	4"	6"	8"	-	12"
Sowthistle, annual		-	-	-	6"	-	12"
Spanishneedles		-	-	-	8"	-	18"
Speedwell, purslane		-	12"	-	-	-	-
Sprangletop		-	6"	12"	20"	-	-
Spurge, prostrate		-	6"	12"	20"	-	-
Spurge, spotted		-	6"	12"	20"	-	-
Spurry, umbrella		6"	-	-	-	-	-
Stinkgrass		12"	-	-	-	-	-
Sunflower		-	12"	18"	-	-	-
Teaweed/Prickly sida		-	1"	2"	3"	4"	6"

WEED SPECIES	REGION	RATE					
		(FLUID OUNCES PER ACRE)					
		9	12	18	24	30	36
		(0.28 lb ae)	(0.38 lb ae)	(0.56 lb ae)	(0.75 lb ae)	(0.94 lb ae)	(1.13 lbs ae)
MAXIMUM HEIGHT/LENGTH							
Texas panicum		-	6"	8"	12"	-	24"
Velvetleaf	South	-	2"	3"	4"	5"	8"
	North	-	3"	6"	12"	-	-
Virginia pepperweed		-	18"	-	-	-	-
Waterhemp		-	-	6"	12"	-	-
Wheat	South	-	6"	30"	-	-	-
	North	-	18"	18" +	-	-	-
Wheat (overwintered)		-	6"	18"	-	-	-
Wild oats		-	12"	-	-	-	-
Wild Proso Millet		-	-	6"	12"	12"	18"
Witchgrass		-	12"	-	-	-	-
Woolly cupgrass		-	6"	12"	-	-	-
Yellow rocket		-	-	12"	20"	-	-

¹Do not treat kochia in the button stage.

ANNUAL WEEDS RATE TABLE, WEST REGION

WEED SPECIES	RATE				
	(FLUID OUNCES PER ACRE)				
	9	12	18	24	36
	(0.28 lb ae)	(0.38 lb ae)	(0.56 lb ae)	(0.75 lb ae)	(1.13 lbs ae)
MAXIMUM HEIGHT/LENGTH					
Barley	12"	-	-	-	-
Barnyardgrass	6"	-	-	-	-
Bluegrass, annual	6"	-	-	-	-
Bluegrass, bulbous	-	6"	-	-	-
Brome, downy ¹	6"	-	-	-	-
Buttercup	-	12"	-	-	-
Cheat	-	6"	-	-	-
Chickweed	-	6"	-	-	-
Cocklebur	-	12"	-	-	-
Corn	-	6"	-	-	-
Crabgrass	-	12"	-	-	-
Dwarf dandelion	-	12"	-	-	-
Fall panicum	-	12"	-	-	-
False flax, smallseed	-	12"	-	-	-
Field pennycress	-	6"	-	-	-
Filaree	-	-	-	-	12"
Fleabane, hairy (<i>Conyza bonariensis</i>)	-	6"	-	-	-
Florida pusley	-	-	-	12"	-
Foxtail	6 fl oz (0.19 lb ae) for up to 12"				
Goatgrass, jointed	-	6"	-	-	-
Groundsel, common	-	6"	-	-	-

WEED SPECIES	RATE				
	(FLUID OUNCES PER ACRE)				
	9 (0.28 lb ae)	12 (0.38 lb ae)	18 (0.56 lb ae)	24 (0.75 lb ae)	36 (1.13 lbs ae)
	MAXIMUM HEIGHT/LENGTH				
Henbit	-	6"	-	-	-
Horseweed/Marestail (<i>Conyza canadensis</i>)	-	6"	-	-	-
Johnsongrass, seedling	-	12"	-	-	-
Lambsquarters	-	6"	-	-	-
London rocket	-	6"	-	-	-
Morningglory (<i>Ipomoea</i> spp.)	-	2"	-	-	-
Mustard, blue	6"	-	-	-	-
Mustard, tansy	6"	-	-	-	-
Mustard, tumble	6"	-	-	-	-
Mustard, wild	6"	-	-	-	-
Pigweed	-	12"	-	-	-
Rye	12"	-	-	-	-
Ryegrass, Italian	-	6"	-	-	-
Sandbur, field	12"	-	-	-	-
Shattercane	12"	-	-	-	-
Sheperd's purse	-	6"	-	-	-
Sowthistle, annual	-	6"	-	-	-
Spurge, annual	-	6"	-	-	-
Stinkgrass	12"	-	-	-	-
Texas panicum	-	12"	-	-	-
Wheat	18"	-	-	-	-
Wild oats	-	12"	-	-	-
Witchgrass	-	12"	-	-	-

¹For control of Downy brome in no-till systems, use 12 fl oz (0.38 lb ae) per acre.

15.1 Annual Weeds – 10 to 40 Gallons Per Acre in Water

Apply 1.5 pints to 2.25 pints (0.75-1.13 lbs ae) of this product per acre. Use 1.5 pints (0.75 lb ae) per acre if weeds are less than 6 inches tall and 2.25 pints (1.13 lbs ae) per acre if weeds are over 6 inches tall.

These rates will provide control of weeds listed in the annual weed control tables when water carrier volumes are 10 to 40 gallons per acre for ground applications.

15.2 Annual Weeds – Tank Mixtures with 2,4-D or Dicamba

9 to 12 fl oz (0.28-0.38 lb ae) of this product plus the labeled rate of dicamba or the labeled rate of 2,4-D per acre will control the following weeds with the maximum height or length indicated: prickly lettuce, marestail/horseweed (*Conyza canadensis*), morningglory (*Ipomoea* spp.), kochia (dicamba only); cocklebur, lambsquarters, pigweed, Russian thistle.

12 fl oz (0.38 lb ae) of this product plus the labeled rate of 2,4-D per acre will control the following weeds when they are a maximum height or length of 6 inches: common ragweed, giant ragweed, Pennsylvania smartweed, and velvetleaf.

9 fl oz (0.28 lb ae) of the product plus the labeled rate of dicamba or the labeled rate of 2,4-D per acre will control foxtail.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

RESTRICTION:

Not Registered for Use by California for dicamba tank mixtures applied by air.

16.0 PERENNIAL WEEDS RATE TABLE ALPHABETICALLY BY SPECIES

Apply to actively growing perennial weeds.

A second treatment may be necessary to control weeds regenerating from underground parts or seed. The second treatment must be made prior to crop emergence.

Unless otherwise stated, allow 7 or more days after application before tillage.

RESTRICTION:

If weeds have been mowed or tilled, do not treat until plants have resumed active growth and have reached the labeled stages.

Best results are obtained when soil moisture is adequate for active weed growth.

Weed Species	Rate (PT/A)	Water Volume	Hand-Held % Solution	Directions
Alfalfa	1.5-3 (0.75-1.5 lbs ae)	3-10	1.5%	Make applications after the last hay cutting in the fall. Allow alfalfa to regrow to a height of 6 to 8 inches or more prior to treatment. Applications should be followed with deep tillage at least 7 days after treatment, but before soil freeze-up.
Alligatorweed	6 (3 lbs ae)	3-20	1.25%	Partial control. Apply when most of the plants are in bloom. Repeat applications will be required to maintain control.
Anise (fennel)	-	-	0.75-1.5%	Apply as a spray-to-wet treatment. Optimum results are obtained when plants are treated at the bud to full-bloom stage of growth.
Bahiagrass	4.5-7.5 (2.25-3.75 lbs ae)	3-20	1.5%	Apply when most plants have reached the early head stage.
Bentgrass	2.25 (1.13 lbs ae)	10-20	1.5%	For suppression in grass seed production areas. For ground applications only. Ensure entire crown area has resumed growth prior to a fall application. Bentgrass should have at least 3 inches of growth. Tillage prior to treatment should be avoided. Till 7 to 10 days after application for best results.
Bermudagrass	4.5-7.5 (2.25-3.75 lbs ae)	3-20	1.5%	For control, apply 7.5 pints (3.75 lbs ae) of this product per acre. For partial control, apply 4.5 pints (2.25 lbs ae) per acre. Treat when bermudagrass is actively growing and seedheads are present. A second treatment may be necessary to maintain control.

Weed Species	Rate (PT/A)	Water Volume	Hand-Held % Solution	Directions
Bermudagrass, water (knotgrass)	1.5-2.25 (0.75-1.13 lbs ae)	5-10	1.5%	<p>Apply 2.25 pints (1.13 lbs ae) of this product in 5 to 10 gallons of water per acre. Apply when water bermudagrass is 12 to 18 inches in length. Allow 7 or more days before tilling, flushing, or flooding the field.</p> <p>Fall applications only: Apply 1.5 pints (0.75 lb ae) of this product in 5 to 10 gallons of water per acre.</p> <p>Fallow fields should be tilled prior to application.</p> <p>Apply prior to frost on water bermudagrass that is 12 to 18 inches in length.</p> <p>RESTRICTION: Not Registered for Use on water bermudagrass by California.</p>
Bindweed, field	0.75-7.5 (0.38-3.75 lbs ae)	3-20	1.5%	<p>Do not treat when weeds are under drought stress as good soil moisture is necessary for active growth.</p> <p>For control, apply 6 to 7.5 pints (3-3.75 lbs ae) of this product per acre west of the Mississippi River and 4.5 to 6 pints (2.25-3 lbs ae) east of the Mississippi River.</p> <p>Apply when the weeds are at or beyond full bloom. For best results, apply in late summer or fall. Fall treatments must be applied before a killing frost.</p> <p>Also, for control, apply 3 pints (1.5 lbs ae) of this product plus the labeled rate of dicamba in 10 to 20 gallons of water per acre. Do not apply by air.</p> <p>For suppression on irrigated agricultural land, apply 1.5 to 3 pints (0.75-1.5 lbs ae) of this product plus the specified label rate of a 2,4-D product in 10 to 20 gallons of water per acre with ground equipment only.</p> <p>It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.</p> <p>Applications should be made following harvest or in fall fallow ground when the bindweed is actively growing, and the majority of runners are 12 inches or more in length. The use of at least one irrigation will promote active bindweed growth.</p> <p>For suppression, apply 12 fl oz (0.38 lb ae) of this product plus the specified label rate of a 2,4-D product in 3 to 10 gallons of water per acre for ground applications and 3 to 5 gallons of water per acre for aerial applications.</p> <p>Apply by air in fallow and reduced tillage systems only. Delay applications until maximum emergence has occurred and when vines are between 6 to 18 inches in length.</p> <p>In California only, apply 1.5 to 7.5 pints (0.75-3.75 lbs ae) of this product per acre.</p> <p>Actual rate needed for suppression or control will vary within this range depending on local conditions. For suppression on irrigated land where annual tillage is performed, apply 1.5 pints (0.75 lb ae) of this product in 3 to 10 gallons of water per acre.</p> <p>Apply to bindweed that has reached a length of 12 inches or greater.</p> <p>Allow maximum weed emergence and runner growth. Allow 3 or more days after application before tillage.</p>

Weed Species	Rate (PT/A)	Water Volume	Hand-Held % Solution	Directions
Bluegrass, Kentucky	1.5-3 (0.75-1.5 lbs ae)	3-40	1.5%	Apply 3 pints (1.5 lbs ae) of this product in 10 to 40 gallons of water per acre when most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 1.5 to 2.25 pints (0.75-1.13 lbs ae) of this product in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height.
Blueweed, Texas	4.5-7.5 (2.25-3.75 lbs ae)	3-40	1.5%	Apply 6 to 7.5 pints (3-3.75 lbs ae) of this product per acre west of the Mississippi River and 4.5 to 6 pints (2.25-3 lbs ae) per acre east of the Mississippi River. Apply when plants are at or beyond full bloom. New leaf development indicates active growth. For best results, apply in late summer or fall. Fall treatments must be applied before a killing frost.
Brackenfern	4.5-6 (2.25-3 lbs ae)	3-40	0.75-1.5%	Apply to fully expanded fronds which are at least 18 inches long.
Bromegrass, smooth	1.5-3 (0.75-1.5 lbs ae)	3-40	1.5%	Apply 3 pints (1.5 lbs ae) of this product in 10 to 40 gallons of water per acre when most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 1.5 to 2.25 pints (0.75-1.13 lbs ae) of this product in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height.
Bursage, woolly-leaf	-	3-20	1.5%	For control, apply 3 pints (1.5 lbs ae) of this product plus the labeled rate of dicamba per acre. For partial control, apply 1.5 pints (0.75 lb ae) of this product plus the labeled rate of dicamba per acre. Apply when plants are producing new active growth which has been initiated by moisture for at least 2 weeks and when plants are at or beyond flowering.
Canarygrass, reed	3-4.5 (1.5-2.25 lbs ae)	3-40	1.5%	For best results, apply when most plants have reached the boot-to-head stage of growth.
Cattail	4.5-7.5 (2.25-3.75 lbs ae)	3-40	1.5%	Apply when most plants have reached the early head stage.
Clover; red, white	4.5-7.5 (2.25-3.75 lbs ae)	3-20	1.5%	Apply when most plants have reached the early bud stage.
Cogongrass	4.5-7.5 (2.25-3.75 lbs ae)	10-40	1.5%	Apply when cogongrass is at least 18 inches tall in late summer or fall. Due to uneven stages of growth and the dense nature of vegetation preventing good spray coverage, repeat treatments may be necessary to maintain control.
Dallisgrass	4.5-7.5 (2.25-3.75 lbs ae)	3-20	1.5%	Apply when most plants have reached the early head stage.
Dandelion	4.5-7.5 (2.25-3.75 lbs ae)	3-40	1.5%	Apply when most plants have reached the early bud stage of growth. Also, for control, apply 12 fl oz (0.38 lb ae) of this product plus the specified label rate of a 2,4-D product in 3 to 10 gallons of water per acre.

Weed Species	Rate (PT/A)	Water Volume	Hand-Held % Solution	Directions
				It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.
Dock, curly	4.5-7.5 (2.25-3.75 lbs ae)	3-40	1.5%	Apply when most plants have reached the early bud stage of growth. Also, for control, apply 12 fl oz (0.38 lb ae) of this product plus the specified label rate of a 2,4-D product in 3 to 10 gallons of water per acre.
Dogbane, hemp	6 (3 lbs ae)	3-40	1.5%	Apply when most plants have reached the late bud to flower stage of growth. Following crop harvest or mowing, allow weeds to regrow to a mature stage prior to treatment. For best results, apply in late summer or fall. For suppression, apply 12 fl oz (0.38 lb ae) of this product plus the labeled rate of 2,4-D in 3 to 10 gallons of water per acre for ground applications and 3 to 5 gallons of water per acre for aerial applications. Delay applications until maximum emergence of dogbane has occurred.
Fescue (except tall)	4.5-7.5 (2.25-3.75 lbs ae)	3-20	1.5%	Apply when most plants have reached the early head stage.
Fescue, tall	1.5-4.5 (0.75-2.25 lbs ae)	3-40	1.5%	Apply 4.5 pints (2.25 lbs ae) of this product per acre when most plants have reached boot-to-early seedhead stage of development. Fall applications only: Apply 1.5 pints (0.75 lb ae) of this product in 3 to 10 gallons of water per acre. Apply to fescue in the fall when plants have 6 to 12 inches of new growth. A sequential application of 12 fl oz (0.38 lb ae) per acre of this product will improve long-term control and control seedlings germinating after fall treatments or the following spring.
Guineagrass	4.5 (2.25 lbs ae)	3-40	0.75%	Apply when most plants have reached at least the 7-leaf stage of growth. Ensure thorough coverage when using hand-held equipment.
Horsenettle	4.5-7.5 (2.25-3.75 lbs ae)	3-20	1.5%	Apply when most plants have reached the early bud stage.
Horseradish	6 (3 lbs ae)	3-40	1.5%	Apply when most plants have reached the late bud to flower stage of growth. For best results, apply in late summer or fall.
Iceplant	-	-	1.5%	Iceplant should be at or beyond the early bud stage of growth. Thorough coverage is necessary for best control.
Jerusalem artichoke	4.5-7.5 (2.25-3.75 lbs ae)	3-20	1.5%	Apply when most plants are in the early bud stage.
Johnsongrass	0.75-4.5 (0.38-2.25 lbs ae)	3-40	0.75%	In annual cropping systems apply 1.5 to 3 pints (0.75-1.5 lbs ae) of this product per acre. Apply 1.5 pints (0.75 lb ae) of this product in 3 to 10 gallons of water per acre. Use 3 pints (1.5 lbs ae) of this product when applying 10 to 40 gallons of water per acre. In noncrop, or areas where annual tillage (no-till) is not practiced, apply 3 to 4.5 pints (1.5-2.25 lbs ae) of this product in 10 to 40 gallons of water per acre.

Weed Species	Rate (PT/A)	Water Volume	Hand-Held % Solution	Directions
				<p>For best results, apply when most plants have reached the boot-to-head stage of growth or in the fall prior to frost.</p> <p>Allow 7 or more days after application before tillage. Do not tank-mix with residual herbicides when using the 1 quart (1 lb ae) per acre rate.</p> <p>For burndown of Johnsongrass, apply 12 fl oz (0.38 lb ae) of this product in 3 to 10 gallons of water per acre before the plants reach a height of 12 inches.</p> <p>For this use, allow at least 3 days after treatment before tillage.</p> <p>Spot treatment (partial control or suppression) – Apply a 3/4 percent solution of this product when Johnsongrass is 12 to 18 inches in height.</p> <p>Coverage must be uniform and complete.</p>
Kikuyugrass	3-4.5 (1.5-2.25 lbs ae)	3-40	1.5%	Spray when most kikuyugrass is at least 8 inches in height (3 or 4-leaf stage of growth). Allow 3 or more days after application before tillage.
Knapweed	6 (3 lbs ae)	3-40	1.5%	<p>Apply when most plants have reached the late bud to flower stage of growth.</p> <p>For best results, apply in late summer or fall.</p>
Lantana	-	-	0.75-1.0%	<p>Apply at or beyond the bloom stage of growth.</p> <p>Use the higher application rate for plants that have reached the woody stage of growth.</p>
Lespedeza	4.5-7.5 (2.25-3.75 lbs ae)	3-20	1.5%	Apply when most plants have reached the early bud stage.
Milkweed, common	4.5 (2.25 lbs ae)	3-40	1.5%	Apply when most plants have reached the late bud to flower stage of growth.
Muhly, wirestem	1.5-3 (0.75-1.5 lbs ae)	3-40	1.5%	<p>Use 1.5 pints (0.75 lb ae) of this product in 3 to 10 gallons of water per acre. Use 3 pints (1.5 lbs ae) of this product when applying 10 to 40 gallons of water per acre or in pasture, sod, or noncrop areas.</p> <p>Spray when the wirestem muhly is 8 inches or more in height.</p> <p>Do not till between harvest and fall applications or in the fall or spring prior to spring applications.</p> <p>Allow 3 or more days after application before tillage.</p>
Mullein, common	4.5-7.5 (2.25-3.75 lbs ae)	3-20	1.5%	Apply when most plants are in the early bud stage.
Napiergrass	4.5-7.5 (2.25-3.75 lbs ae)	3-20	1.5%	Apply when most plants are in the early head stage.
Nightshade, silverleaf	3 (1.5 lbs ae)	3-10	1.5%	Applications should be made when at least 60 percent of the plants have berries. Fall treatments must be applied before a killing frost.
Nutsedge; purple, yellow	0.75-4.5 (0.38-2.25 lbs ae)	3-40	0.75-1.5%	<p>Apply 4.5 pints (2.25 lbs ae) of this product per acre or apply a 3/4 to 1 1/2 percent solution for control of nutsedge plants and immature nutlets attached to treated plants.</p> <p>Treat when plants are in flower or when new nutlets can be found at rhizome tips. Nutlets which have not germinated will not be controlled and may germinate following treatment.</p> <p>Repeat treatments will be required for long-term control of ungerminated tubers.</p> <p>Sequential applications: 1.5 to 3 pints (0.75-1.5 lbs ae) of this product in 3 to 10 gallons of water per acre will also provide control. Make applications when a majority of the plants are</p>

Weed Species	Rate (PT/A)	Water Volume	Hand-Held % Solution	Directions
				<p>in the 3 to 5-leaf stage (less than 6 inches tall). Repeat this application, as necessary, when newly emerging plants reach the 3 to 5-leaf stage.</p> <p>Subsequent applications will be necessary for long-term control.</p> <p>For partial control of existing plants, apply 12 fl oz to 3 pints (0.38-1.5 lbs ae) of this product in 3 to 40 gallons of water per acre.</p> <p>Treat when plants have 3 to 5 leaves, and most are less than 6 inches tall.</p> <p>Repeat treatments will be required to control subsequent emerging plants or regrowth of existing plants.</p>
Orchardgrass	1.5-3 (0.75-1.5 lbs ae)	3-40	1.5%	<p>Apply 3 pints (1.5 lbs ae) of this product in 10 to 40 gallons of water per acre when most plants have reached boot-to-early seedhead stage of development.</p> <p>For partial control in pasture or hay crop renovation, apply 1.5 to 2.25 pints (0.75-1.13 lbs ae) of this product in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height.</p> <p>Orchardgrass sods going to no-till corn: Apply 1.5 to 2.25 pints (0.75-1.13 lbs ae) of this product in 3 to 10 gallons of water per acre.</p> <p>Apply to orchardgrass that is a minimum of 12 inches tall for spring applications and 6 inches tall for fall applications.</p> <p>Allow at least 3 days following application before planting. A sequential application of atrazine will be necessary for optimum results.</p>
Pampasgrass	-	-	1.5%	Pampasgrass should be at or beyond the boot stage of growth. Thorough coverage is necessary for best control.
Paragrass	4.5-7.5 (2.25-3.75 lbs ae)	3-20	1.5%	Apply when most plants are in the early head stage.
Phragmites	4.5-7.5 (2.25-3.75 lbs ae)	10-40	0.75-1.5%	<p>For partial control. For best results, treat during late summer or fall months or when plants are actively growing and in full bloom.</p> <p>Treatment before or after this stage may lead to reduced control.</p> <p>Due to the dense nature of the vegetation, which may prevent good spray coverage or uneven stages of growth, repeat treatments may be necessary to maintain control.</p> <p>Visual control symptoms will be slow to develop.</p>
Poison hemlock	-	-	0.75-1.5%	<p>Apply as a spray-to-wet treatment.</p> <p>Optimum results are obtained when plants are treated at the bud to full-bloom stage of growth.</p>
Pokeweed, common	1.5 (0.75 lb ae)	3-40	1.5%	Apply to actively growing plants up to 24 inches tall.
Quackgrass	1.5-4.5 (0.75-2.25 lbs ae)	3-40	1.5%	<p>In annual cropping systems, or in pastures and sods followed by deep tillage:</p> <p>Apply 1.5 pints (0.75 lb ae) of this product in 3 to 10 gallons of water per acre. For 10 to 40 gallons of water per acre, apply 3 pints (1.5 lbs ae) of this product. Do not tank mix with residual herbicides when using the 1.5 pint (0.75 lb ae) rate.</p>

Weed Species	Rate (PT/A)	Water Volume	Hand-Held % Solution	Directions
				<p>Spray when quackgrass is 6 to 8 inches in height. Do not till between harvest and fall applications or in fall or spring prior to spring application.</p> <p>Allow 3 or more days after application before tillage.</p> <p>In pastures or sods, use a moldboard plow for best results.</p> <p>In pastures, sods or noncrop areas where deep tillage does not follow application: Apply 3 to 4.5 pints (1.5-2.25 lbs ae) of this product in 10 to 40 gallons of water per acre when the quackgrass is greater than 8 inches tall.</p>
Redvine	1.25-3 (0.63-1.5 lbs ae)	5-10	1.5%	<p>For suppression, apply 18 fl oz (0.56 lb ae) of this product per acre at each of two applications 7 to 14 days apart or a single application of 3 pints (1.5 lbs ae) per acre.</p> <p>Apply labeled rates in 5 to 10 gallons of water per acre.</p> <p>Apply in late September or early October to plants which are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least 1 week before a killing frost.</p>
Reed, giant	-	-	1.5%	<p>Best results are obtained when applications are made in late summer to fall.</p>
Ryegrass, perennial	1.5-4.5 (0.75-2.25 lbs ae)	3-40	0.75%	<p>In annual cropping systems apply 1.5 to 3 pints (0.75-1.5 lbs ae) of this product per acre. Apply 1.5 pints (0.75 lb ae) of this product in 3 to 10 gallons of water per acre. Use 3 pints (1.5 lbs ae) of this product when applying 10 to 40 gallons of water per acre. In noncrop, or areas where annual tillage (no-till) is not practiced, apply 3 to 4.5 pints (1.5-2.25 lbs ae) of this product in 10 to 40 gallons water per acre.</p> <p>For best results, apply when most plants have reached the boot-to-head stage of growth or in the fall prior to frost. Do not tank-mix with residual herbicides when using the 1.5 pint (0.75 lb ae) per acre rate.</p>
Smartweed, swamp	4.5-7.5 (2.25-3.75 lbs ae)	3-40	1.5%	<p>Apply when most plants have reached the early bud stage of growth.</p> <p>Also, for control, apply 12 fl oz (0.38 lb ae) of this product plus the specified label rate of a 2,4-D product in 3 to 10 gallons of water per acre in the late summer or fall.</p> <p>It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.</p>
Sowthistle, perennial	3-4.5 (1.5-2.25 lbs ae)	3-40	1.5%	<p>Apply when most plants are at or beyond the bud stage of growth. After harvest, mowing or tillage in the late summer or fall, allow at least 4 weeks for initiation of active growth and rosette development prior to the application of this product.</p> <p>Fall treatments must be applied before a killing frost. Allow 3 or more days after application before tillage.</p>
Spurge, leafy	-	3-10	1.5%	<p>For suppression, apply 12 fl oz (0.38 lb ae) of this product plus the specified label rate of a 2,4-D product in 3 to 10 gallons of water per acre in the late summer or fall.</p> <p>If mowing has occurred prior to treatment, apply when most of the plants are 12 inches tall.</p>
Starthistle, yellow	3 (1.5 lbs ae)	10-40	1.5%	<p>Best results are obtained when applications are made during the rosette, bolting and early flower stages.</p>

Weed Species	Rate (PT/A)	Water Volume	Hand-Held % Solution	Directions
Sweet potato, wild	-	-	1.5%	Partial control. Apply to plants that are at or beyond the bloom stage of growth. Repeat applications may be required.
Thistle, artichoke	-	-	1.5%	Partial control. Apply to plants that are at or beyond the bloom stage of growth. Repeat applications may be required.
Thistle, Canada	3-4.5 (1.5-2.25 lbs ae)	3-40	1.5%	Apply when most plants are at or beyond the bud stage of growth. After harvest, mowing or tillage in the late summer or fall, allow at least 4 weeks for initiation of active growth and rosette development prior to the application of this product. Fall treatments must be applied before a killing frost. Allow 3 or more days after application before tillage. For suppression, apply 1.5 pints (0.75 lb ae) of this product, or 12 fl oz (0.38 lb ae) of this product plus the specified label rate of a 2,4-D product in 3 to 10 gallons of water per acre in the late summer or fall after harvest, mowing or tillage. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Allow rosette regrowth to a minimum of 6 inches in diameter before treating. Applications can be made as long as leaves are still green, and plants are actively growing at the time of application. Allow 3 or more days after application before tillage. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.
Timothy	3-4.5 (1.5-2.25 lbs ae)	3-40	1.5%	For best results, apply when most plants have reached the boot-to-head stage of growth.
Torpedograss	6-7.5 (3-3.75 lbs ae)	3-40	1.5%	For partial control. Apply when most plants are at or beyond the seedhead stage of growth. Repeat applications will be required to maintain control. Fall treatments must be applied before frost.
Trumpet creeper	3 (1.5 lbs ae)	5-10	1.5%	Partial control. Apply in late September or October, to plants which are at least 18 inches tall and have been growing 45-60 days since the last tillage operation. Make applications at least 1 week before a killing frost.
Vaseygrass	4.5-7.5 (2.25-3.75 lbs ae)	3-20	1.5%	Apply when most plants are in the early head stage.
Velvetgrass	4.5-7.5 (2.25-3.75 lbs ae)	3-20	1.5%	Apply when most plants are in the early head stage.
Wheatgrass, western	3-4.5 (1.5-2.25 lbs ae)	3-40	1.5%	For best results, apply when most plants have reached the boot-to-head stage of growth.

17.0 WOODY BRUSH AND TREES RATE TABLE ALPHABETICALLY BY SPECIES

Apply this product after full leaf expansion, unless otherwise directed. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation.

In arid areas, best results are obtained when applications are made in the spring to early summer when brush species are at high moisture content and are flowering.

Ensure thorough coverage when using hand-held equipment. Symptoms may not appear prior to frost or senescence with fall treatments.

Allow 7 or more days after application before tillage, mowing or removal. A second treatment may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatments are made following a frost.

Weed Species	Rate (PT/A)	Water Volume	Hand-Held % Solution	Directions
Alder	4.5-6 (2.25-3 lbs ae)	3-40	0.75-1.5%	For control
Ash	3-7.5 (1.5-3.75 lbs ae)	3-40	0.75-1.5%	Partial control
Aspen, quaking	3-4.5 (1.5-2.25 lbs ae)	3-40	0.75-1.5%	For control
Bearmat (Bearclover)	3-7.5 (1.5-3.75 lbs ae)	3-40	0.75-1.5%	Partial control
Beech	3-7.5 (1.5-3.75 lbs ae)	3-40	0.75-1.5%	Partial control
Birch	3 (1.5 lbs ae)	3-40	0.75%	For control
Blackberry	4.5-6 (2.25-3 lbs ae)	10-40	0.75-1.5%	For control. Make applications after plants have reached full leaf maturity. Best results are obtained when applications are made in late summer or fall. Applications may also be made after leaf drop and until a killing frost or as long as stems are green. After berries have set or dropped in late fall, blackberry can be controlled by applying a ¾ percent solution of this product. For control of blackberries after leaf drop and until a killing frost or as long as stems are green, apply 4.5 to 6 pints (2.25-3 lbs ae) of this product in 10 to 40 gallons of water per acre.
Blackgum	3-7.5 (1.5-3.75 lbs ae)	3-40	0.75-1.5%	For control
Bracken	3-7.5 (1.5-3.75 lbs ae)	3-40	0.75-1.5%	For control
Broom; French, Scotch	-	-	1.5%	For control
Buckwheat, California	-	-	0.75-1.5%	For partial control. Thorough coverage of foliage is necessary for best results.
Cascara	3-7.5 (1.5-3.75 lbs ae)	3-40	0.75-1.5%	Partial control
Catsclaw	-	-	0.75-1.5%	Partial control
Ceanothus	3-7.5 (1.5-3.75 lbs ae)	3-40	0.75-1.5%	Partial control
Chamise	-	-	0.75%	For control. Thorough coverage of foliage is necessary for best results.
Cherry; bitter, black, pin	3-4.5 (1.5-2.25 lbs ae)	3-40	0.75-1.5%	For control
Coyote brush	-	-	1.5%	For control. Apply when at least 50 percent of the new leaves are fully developed.
Dogwood	3-7.5 (1.5-3.75 lbs ae)	3-40	0.75-1.5%	Partial control
Elderberry	3 (1.5 lbs ae)	3-40	0.75%	For control

Weed Species	Rate (PT/A)	Water Volume	Hand-Held % Solution	Directions
Elm	3-7.5 (1.5-3.75 lbs ae)	3-40	0.75-1.5%	Partial control
Eucalyptus	-	-	1.5%	For control of eucalyptus resprouts, apply when resprouts are 6 to 12 feet tall. Ensure complete coverage. Avoid application to drought-stressed plants.
Florida holly (Brazilian Peppertree)	3-7.5 (1.5-3.75 lbs ae)	3-40	0.75-1.5%	Partial control
Gorse	3-7.5 (1.5-3.75 lbs ae)	3-40	0.75-1.5%	Partial control
Hasardia	-	-	0.75-1.5%	Partial control. Thorough coverage of foliage is necessary for best results.
Hawthorn	3-4.5 (1.5-2.25 lbs ae)	3-40	0.75-1.5%	For control
Hazel	3 (1.5 lbs ae)	3-40	0.75%	For control
Hickory	3-7.5 (1.5-3.75 lbs ae)	3-40	0.75-1.5%	Partial control
Honeysuckle	3-6 (1.5-3 lbs ae)	3-40	0.75-1.5%	For control
Hornbeam, American	3-7.5 (1.5-3.75 lbs ae)	3-40	0.75-1.5%	Partial control
Kudzu	6 (3 lbs ae)	3-40	1.5%	For control. Repeat applications may be required to maintain control.
Locust, black	3-6 (1.5-3 lbs ae)	3-40	0.75-1.5%	Partial control
Madrone resprouts	-	-	1.5%	Partial control. Apply to resprouts that are 3 to 6 feet tall. Best results are obtained with spring/early summer treatments.
Manzanita	3-7.5 (1.5-3.75 lbs ae)	3-40	0.75-1.5%	Partial control
Maple, red	3-6 (1.5-3 lbs ae)	3-40	0.75-1.5%	For control, apply a 0.75 to 1.5 percent solution when at least 50 percent of the new leaves are fully developed. For partial control, apply 3 to 6 pints (1.5-3 lbs ae) of this product per acre.
Maple, sugar	-	-	0.75-1.5%	For control. Apply when at least 50 percent of the new leaves are fully developed.
Monkey flower	-	-	0.75-1.5%	Partial control. Thorough coverage of foliage is necessary for best results.
Oak; black, white	3-6 (1.5-3 lbs ae)	3-40	0.75-1.5%	Partial control
Oak, post	4.5-6 (2.25-3 lbs ae)	3-40	0.75-1.5%	For control
Oak; northern, pin	-	-	0.75-1.5%	For control. Apply when at least 50 percent of the new leaves are fully developed.
Oak, southern, red	3-4.5 (1.5-2.25 lbs ae)	3-40	0.75-1.5%	For control
Persimmon	3-7.5 (1.5-3.75 lbs ae)	3-40	0.75-1.5%	Partial control
Pine	3-7.5 (1.5-3.75 lbs ae)	3-40	0.75-1.5%	For control
Poison ivy/Poison oak	6-7.5 (3-3.75 lbs ae)	3-40	1.5%	For control. Repeat applications may be required to maintain control. Fall treatments must be applied before leaves lose green color.
Poplar, yellow	3-7.5 (1.5-3.75 lbs ae)	3-40	0.75-1.5%	Partial control

Weed Species	Rate (PT/A)	Water Volume	Hand-Held % Solution	Directions
Redbud, eastern	3-7.5 (1.5-3.75 lbs ae)	3-40	0.75-1.5%	For control
Rose, multiflora	3 (1.5 lbs ae)	3-40	0.75%	For control. Treatments should be made prior to leaf deterioration by leaf-eating insects.
Russian olive	3-7.5 (1.5-3.75 lbs ae)	3-40	0.75-1.5%	Partial control
Sage, black	-	-	0.75%	For control. Thorough coverage of foliage is necessary for best results.
Sage, white	3-7.5 (1.5-3.75 lbs ae)	3-40	0.75-1.5%	Partial control
Sage brush, California	-	-	0.75%	For control. Thorough coverage of foliage is necessary for best results.
Salmonberry	3 (1.5 lbs ae)	3-40	0.75%	For control
Salt-cedar	3-7.5 (1.5-3.75 lbs ae)	3-40	0.75-1.5%	For control
Sassafras	3-7.5 (1.5-3.75 lbs ae)	3-40	0.75-1.5%	Partial control
Sourwood	3-7.5 (1.5-3.75 lbs ae)	3-40	0.75-1.5%	Partial control
Sumac; poison, smooth, winged	3-6 (1.5-3 lbs ae)	3-40	0.75-1.5%	Partial control
Sweetgum	3-4.5 (1.5-2.25 lbs ae)	3-40	0.75-1.5%	For control
Swordfern	3-7.5 (1.5-3.75 lbs ae)	3-40	0.75-1.5%	Partial control
Tallowtree, Chinese	-	-	0.75%	For control. Thorough coverage of foliage is necessary for best results.
Tan oak resprouts	-	-	1.5%	For partial control. Apply to resprouts that are less than 3 to 6 feet tall. Best results are obtained with fall applications.
Thimbleberry	3 (1.5 lbs ae)	3-40	0.75%	For control
Tobacco, tree	-	-	0.75-1.5%	Partial control
Trumpet creeper	3-4.5 (1.5-2.25 lbs ae)	3-40	0.75-1.5%	For control
Vine maple	3-7.5 (1.5-3.75 lbs ae)	3-40	0.75-1.5%	Partial control
Virginia creeper	3-7.5 (1.5-3.75 lbs ae)	3-40	0.75-1.5%	For control
Waxmyrtle, southern	3-7.5 (1.5-3.75 lbs ae)	3-40	0.75-1.5%	Partial control
Willow	4.5 (2.25 lbs ae)	3-40	0.75%	For control

18.0 WARRANTY DISCLAIMER, INHERENT RISKS OF USE, LIMITATION OF REMEDIES

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Remedies before using this product.

If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following warranty disclaimer, inherent risks of use and limitation of remedies.

WARRANTY DISCLAIMER

AgSaver, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, AGSAVER, LLC MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label directions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of AgSaver, LLC or the seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

LIMITATION OF REMEDIES

To the extent consistent with applicable law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at AgSaver, LLC's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent consistent with applicable law, AgSaver, LLC shall not be liable for losses or damages resulting from handling or use of this product unless AgSaver, LLC is promptly notified of such loss or damage in writing. In no case shall AgSaver, LLC be liable for consequential or incidental damages or losses. The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of AgSaver, LLC or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

AgSaver is a trademark of AgSaver, LLC

[EPA APPROVAL DATE]

AgSaver™ Glyphosate 5.4 Plus

{ABN:} [CropSmart Glyphosate 5.4 Plus]

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

	% BY WT.
ACTIVE INGREDIENT: Glyphosate*, N-(phosphonomethyl)glycine, in the form of its isopropylamine salt...	53.8%
OTHER INGREDIENTS:	46.2%
	TOTAL..... 100.0%

*Contains 648 grams per liter or 5.4 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its isopropylamine salt. Equivalent to 480 grams per liter or 4 pounds per U.S. gallon of the acid, glyphosate.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DOMESTIC ANIMALS: This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE CALL CHEMTREC (800) 424-9300

EPA Reg. No. 83772-[xx]

EPA Est. No.

[[[(When more than one EPA Est. No. is used this may appear:)] Last letter in lot number corresponds to the EPA Est. No. used.]

Manufactured by:

AgSaver, LLC 203 East Ash Street McGehee, AR 71654

[Net Contents:]

[Batch Code:]

{Net Contents: May appear on label and/or container.} {Batch Code: May appear on label or container.}

[AgSaver is a trademark of AgSaver, LLC]

PHYSICAL OR CHEMICAL HAZARDS

Spray solutions of this product must be mixed, stored, and applied using only stainless steel, aluminum, fiberglass, plastic, or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette, or other ignition source.

ENVIRONMENTAL HAZARDS

FOR AQUATIC USES ONLY: Do not contaminate water when disposing of equipment waste waters. Treatment of aquatic weeds can result in oxygen depletion or loss due to decomposition of dead plants. This oxygen loss can cause fish suffocation.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

STORAGE: STORE ABOVE 10° F (-12° C) TO KEEP FROM CRYSTALIZING. Crystals will settle to the bottom. If crystals form, allow product to warm above 50° F (10° C) and mix well or recirculate to redissolve.

DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed must be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state, or local procedures.

Emptied container retains vapor and product residue. Observe all labeled safeguards until container is cleaned, reconditioned, or destroyed.

To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing process two more times.

CONTAINER HANDLING: (See the Net Contents section on the container to determine if it non-refillable or refillable.) APPROPRIATE BOX MUST BE CHECKED.

Non-refillable containers (1 and 2.5 gallon): Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Non-refillable containers (>5 gallon): Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Refillable containers: Refillable container. Refill this container with glyphosate only. Do not reuse this container for any other purpose.

When this container is empty, replace the cap and seal all openings that have been made during usage and return the container to the point of purchase, or to an alternate location designated by the manufacturer at the time of purchase of this product. If not returned, clean container the empty container and offer for recycling, if available.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the re-filler.

To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or re-circulate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing process two more times.

If the container cannot be refilled, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Read the entire label before using this product. Use only according to label instructions.

Not all products labeled on this product are registered for use in California. Check the registration status of each product in California before using.

Read the "CONDITIONS OF SALE AND WARRANTY" statement at the end of the label before buying or using. If terms are not acceptable, return at once unopened.

USE INFORMATION

This product, a water-soluble liquid, mixes readily with water and nonionic surfactant to be applied as a foliar spray for the control or destruction of many herbaceous and woody plants.

This product moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days but on most perennial brush species may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow the activity of this product and delay visual effects of control. Visible effects are a gradual wilting and yellowing of the plant which advances to complete browning of above-ground growth and deterioration of underground plant parts.

Unless otherwise directed on this label, delay application until vegetation has emerged and reached the stages described for control of such vegetation under the "Weeds Controlled" section of this label. Unemerged plants arising from unattached underground rhizomes or root stocks of perennials or brush will not be affected by the spray and will continue to grow. For this reason, best control of most perennial weeds or brush is obtained when treatment is made at late growth stages approaching maturity.

Always use the higher rate of this product per acre within the specific range when vegetation is heavy or dense.

Do not treat weeds or brush under poor growing conditions such as drought stress, disease or insect damage, as reduced control may result. Reduced results may also occur when treating weeds or brush heavily covered with dust.

Reduced control may result when applications are made to any weed or brush species that have been mowed, grazed, or cut, and have not been allowed to regrow to the specific stage for treatment.

Rainfall or irrigation occurring within 6 hours after application may reduce effectiveness. Heavy rainfall or irrigation within 2 hours after application may wash the product off the foliage and a repeat treatment may be required.

When this product comes in contact with soil (on the soil surface or as suspended soil or sediment in water) it is bound to soil particles. Under use situations, once this product is bound to soil particles, it is not available for plant uptake and will not harm off-site vegetation where roots grow into the treatment area or if the soil is transported off-site. Under use conditions, the strong affinity of this product to soil particles prevents this product from leaching out of the soil profile and entering ground water. The affinity between this product and soil particles remains until this product is degraded, which is primarily a biological degradation process carried out under both aerobic and anaerobic conditions by soil microflora.

This product does not provide residual weed control. For subsequent residual weed control, follow a label-approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

To the extent consistent with applicable law, the buyer and all users are responsible for all loss or damage in connection with the use or handling or mixtures of this product or other materials that are not expressly in this label. Mixing this product with herbicides or other materials not in this label may result in reduced performance.

USE RESTRICTIONS:

- Except otherwise specified, for noncrop uses, the combined total of all treatments must not exceed 6.0 quarts (6 lbs ae) of this product per acre per year.
- The maximum use rates stated throughout this product's labeling apply to this product combined with the use of all other herbicides containing glyphosate or sulfosate as the active ingredient, whether applied as mixtures or separately. Calculate application rates and ensure that the total use of this and other glyphosate or sulfosate containing products does not exceed stated maximum use rate.

ATTENTION

Avoid drift. Extreme care must be used when applying this product to prevent injury to desirable plants and crops.

Do not allow the herbicide solution to mist, drip, drift, or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of injury occurring from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing or when there are other meteorological conditions that favor spray drift. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) which are likely to drift.

Keep container closed to prevent spills and contamination.

MIXING AND APPLICATION INSTRUCTIONS

Apply these spray solutions in properly maintained and calibrated equipment capable of delivering desired volumes. Hand-gun applications should be properly directed to avoid spraying desirable plants. Reduced results may occur if water containing soil is used, such as water from ponds and unlined ditches.

Mixing

This product mixes readily with water. Mix spray solutions of this product as follows: fill the mixing or spray tank with the required amount of water while adding the required amount of this product (see the “Directions for Use” and “Weeds Controlled” sections of this label). Near the end of the filling process, add the required surfactant and mix well. Remove hose from tank immediately after filling to avoid siphoning back into the water source. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, avoid the use of mechanical agitators, place the filling hose below the surface of the spray solution, terminate by-pass and return lines at the bottom of the tank and if needed use an approved anti-foam or defoaming agent.

Keep by-pass line on or near bottom of tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50 mesh. Carefully select correct nozzle to avoid spraying a fine mist. For best results with conventional ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

When using this product, mix 2 or more quarts of a nonionic surfactant per 100 gallons of spray solution. Use a nonionic surfactant labeled for use with herbicides. The surfactant must contain 50 percent or more active ingredient.

Always read and follow the manufacturer’s surfactant label directions for best results. Carefully observe all cautionary statements and other information appearing in the surfactant label.

These surfactants should not be used in excess of 1 quart per acre when making broadcast applications.

Colorants or marking dyes approved for use with herbicides may be added to spray mixtures of this product. Colorants or dyes used in spray solutions of this product may reduce performance, especially at lower rates or dilutions. Use colorants or dyes according to the manufacturer’s label directions. Clean sprayer and parts immediately after using this product by thoroughly flushing with water and dispose of rinsate according to labeled use or disposal instructions.

Observe all cautionary statements and other information appearing in the surfactant label.

APPLICATION EQUIPMENT AND TECHNIQUES

Aerial Equipment

Use the specified rates of this product and surfactant in 3 to 20 gallons of water per acre as a broadcast spray, unless otherwise specified. See the “Weeds Controlled” section of this label for specific rates. Aerial applications of this product may only be made as specified in this label.

Avoid drift – do not apply during inversion conditions, when winds are gusty or under any other condition which will allow drift. Drift may cause damage to any vegetation contacted to which treatment is not intended. To prevent injury to adjacent desirable vegetation, appropriate buffer zones must be maintained.

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine spray droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure.

Drift control additives may be used. When a drift control additive is used, read, and carefully observe the cautionary statements and all other information appearing in the additive label.

Ensure uniform application – To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR ARE MOST SUSCEPTIBLE. The maintenance of an organic coating (paint) which meets aerospace specification MIL-C 38413 may prevent corrosion.

AERIAL SPRAY DRIFT MANAGEMENT

SPRAY DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator.

The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed $\frac{3}{4}$ the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.
3. Where states have more stringent regulations, they must be observed.

The applicator must be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory.

Aerial Drift Reduction Advisory

This section is advisory in nature and does not supersede the mandatory label requirements.

INFORMATION ON DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (See Wind, Temperature and Humidity, and Temperature Inversions).

CONTROLLING DROPLET SIZE

- Volume – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure – Do not exceed the nozzle manufacturer's specified pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of nozzles – Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation – Orient nozzles so that the spray is released parallel to the airstream which produces larger droplets than other orientations. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- Nozzle Type – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

BOOM LENGTH

For some use patterns, reducing the effective boom length to less than $\frac{3}{4}$ of the wingspan or rotor length may further reduce drift without reducing swath width.

APPLICATION HEIGHT

Applications should not be made at a height greater than 10 feet above the top of the target plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

SWATH ADJUSTMENT

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.)

WIND

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator must be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

TEMPERATURE INVERSIONS

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small, suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SENSITIVE AREAS

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

For Aerial Application in California Only

Aquatic and Other Noncrop Sites:

When applied as directed and under the conditions described in the "Weeds Controlled" section of this label booklet, this product will control or partially control the labeled weeds growing in the following industrial, recreational, and public areas or other similar sites.

Aquatic Sites – Including all bodies of fresh and brackish water which may be flowing, nonflowing, or transient. This includes lakes, rivers, streams, ponds, seeps, irrigation and drainage ditches, canals, reservoirs, estuaries, and similar sites.

If aquatic sites are present in the noncrop area and are part of the intended treatment, read and observe the following directions:

There is no restriction on the use of treated water for irrigation, recreation, or domestic purposes. Consult local state fish and game agency and water control authorities before applying this product to public water. Permit may be required to treat such water.

RESTRICTIONS:

Do not apply this product within ½ mile upstream of an active potable water intake in flowing water (i.e., river, stream, etc.) or within ½ mile of an active potable water intake in a standing body of water such as lake, pond, or reservoir. To make aquatic applications around and within ½ mile of active potable water intakes, the water intake must be turned off for a minimum period of 48 hours after the application. The water intake may be turned on prior to 48 hours if the glyphosate level in the intake water is below 0.7 part per million as determined by laboratory analysis. These aquatic applications may be made ONLY in those cases where there are alternative water sources or holding ponds which would permit the turning off of an active potable water intake for a minimum period of 48 hours after the application.

This product does not control plants which are completely submerged or have a majority of their foliage under water.

Aerial Applications:

Only make aerial applications with helicopters.

Use the following guidelines when aerial applications are to be made near perennial crops after bud break and before total leaf drop and/or near emerged annual crops.

1. Do not apply within a minimum of 100 feet of all crops.
2. If wind up to 5 miles per hour is blowing toward the crop(s), do not apply within a minimum of 500 feet of the crop(s).
3. Winds blowing from 5 to 10 miles per hour toward the crops(s) may require buffer zones in excess of the 500 feet minimum.
4. Do not apply when winds are in excess of 10 miles per hour or when inversion conditions exist. For Aerial Application in Fresno County, California Only from February 15 through March 31 Only

Applicable Area:

The area contained inside the following boundaries within Fresno County, California.

North:	Fresno County line
South:	Fresno County line
East:	State Highway 99
West:	Fresno County line

Use Information:

Always read and follow the label directions, restrictions and precautionary statements for all products used in the aerial application.

Observe the following directions to minimize off-site movement during aerial application of this product.

Minimization of off-site movement is the responsibility of the grower, Pest Control Advisor, and aerial applicator.

Written Recommendations:

A written recommendation **MUST** be submitted by or on behalf of the applicator to the Fresno County Agricultural Commissioner 24 hours prior to the application. This written recommendation **MUST** state the proximity of surrounding crops, and that conditions of each manufacturer's applicable product label and this label have been satisfied.

Aerial Applicator Training and Equipment:

Aerial application of this product is limited to pilots who have successfully completed a Fresno County Agricultural Commissioner and California Department of Pesticide Regulation approved training program for aerial application of herbicides. All aircraft must be inspected, critiqued in flight, and certified at a Fresno County Agricultural Commissioner approved fly-in. Test and calibrate spray equipment at intervals sufficient to ensure that proper rates of herbicides and adjuvants are being applied during commercial use. Applicator must document such calibrations and testing. Demonstration of performance at Fresno County Agricultural Commissioner approved "fly-ins" constitutes such documentation, or other written records showing calculations and measurements of flight and spray parameters acceptable to the Fresno County Agricultural Commissioner.

Applications at Night:

Do not apply this product by air earlier than 30 minutes prior to sunrise and/or later than 30 minutes after sunset without prior permission from the Fresno County Agricultural Commissioner.

Note: For aerial application from April 1 through February 14, refer to the "For Aerial Application in California Only" section of this label.

Boom Equipment

For control of weed or brush species listed in this section using conventional boom equipment – Use the specified rates of this product and surfactant in 3 to 30 gallons of water per acre as a broadcast spray, unless otherwise specified. See the "Weeds Controlled" section of this label for specific rates. As density of vegetation increases, spray volume should be increased within this range to ensure complete coverage. Carefully select correct nozzle to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

Hand-Held and High-Volume Equipment Use Coarse Sprays Only

For control of weeds listed in this section using knapsack sprayers or high-volume spraying equipment utilizing handguns or other suitable nozzle arrangements – Prepare a $\frac{3}{4}$ to 2 percent solution of this product in water, add a nonionic surfactant and apply to foliage of vegetation to be controlled. For specific rates of application and instructions for control of various annual and perennial weeds, see the "Weeds Controlled" section of this label.

Applications should be made on a spray-to-wet basis. Spray coverage should be uniform and complete. Do not spray to point of runoff.

This product may be used as a 5 to 8 percent solution for low volume directed sprays for spot treatment of trees and brush. It is most effective in areas where there is a low density of undesirable trees or brush.

If a straight stream nozzle is used, start the application at the top of the targeted vegetation and spray from top to bottom in a lateral zigzag motion. Ensure that at least 50 percent of the leaves are contacted by the spray solution. For flat fan and cone nozzles and with hand-directed mist blowers, mist the application over the foliage of the targeted vegetation. Small, open-branched trees need only be treated from one side. If the foliage is thick or there are multiple root sprouts, applications must be made from several sides to ensure adequate spray coverage.

Prepare the desired volume of spray solution by mixing the amount of this product in water, shown in the following table:

Spray Solution

AMOUNT OF PRODUCT

Desired Volume	3/4%	1%	1 1/4%	1 1/2%	5%	10%
1 Gal	1 fl oz (0.03 lb. ae)	1 1/3 fl oz (0.04 lb. ae)	1 2/3 fl oz (0.052 lb ae)	2 fl oz (0.06 lb. ae)	6 1/2 fl oz (0.2 lb. ae)	10 1/4 fl oz (0.32 lb ae)
25 Gal	1 1/2 pt (0.75 lb. ae)	1 qt (1 lb. ae)	1 1/4 qt (1.25 lb ae)	1 1/2 qt (1.5 lbs. ae)	5 qt (5 lbs. ae)	2 gal (8 lb ae)
100 Gal	3 qt (3 lbs. ae)	1 gal (4 lbs. ae)	1 1/4 gal (5 lb ae)	1 1/2 gal (6 lbs. ae)	5 gal (20 lbs. ae)	8 gal (32 lb ae)

2 tablespoons = 1 fluid ounce

For use in knapsack sprayers, it is suggested that the labeled amount of this product be mixed with water in a larger container. Fill sprayer with the mixed solution and add the correct amount of surfactant.

WEEDS CONTROLLED

Annual Weeds

Apply to actively growing annual grasses and broadleaf weeds.

Allow at least 3 days after application before disturbing treated vegetation. After this period, the weeds may be mowed, tilled, or burned. See "Directions for Use," "Use Information" and "Mixing and Application Instructions" for labeled uses and specific application instructions.

Broadcast Application – Use 1.5 pints (0.75 lb ae) of this product per acre plus 2 or more quarts of a nonionic surfactant per 100 gallons of spray solution if weeds are less than 6 inches tall. If weeds are greater than 6 inches tall, use 2.5 pints (1.25 lbs ae) of this product per acre plus 2 or more quarts of an approved nonionic surfactant per 100 gallons of spray solution.

Hand-Held, High-Volume Application – Use a ¾ percent solution of this product in water plus 2 or more quarts of a nonionic surfactant per 100 gallons of spray solution and apply to foliage of vegetation to be controlled.

When applied as directed under the conditions described in this label, this product plus nonionic surfactant will control the following annual weeds:

Balsam apple** <i>Momordica charantia</i>	Foxtail, Carolina <i>Alopecurus carolinianus</i>	Rye <i>Secale cereale</i>
Barley <i>Hordeum vulgare</i>	Groundsel, common <i>Senecio vulgaris</i>	Ryegrass, Italian* <i>Lolium multiflorum</i>
Barnyardgrass <i>Echinochola crus-galli</i>	Horseweed/Marestail <i>Conyza canadensis</i>	Sandbur, field <i>Cenchrus</i> spp.
Bassia, fivehook <i>Bassia hyssopifolia</i>	Kochia <i>Kochia scoparia</i>	Shattercane <i>Sorghum bicolor</i>
Bluegrass, annual <i>Poa annua</i>	Lambsquarters, common <i>Chenopodium album</i>	Shepherdspurse <i>Capsella bursa-pastoris</i>
Bluegrass, bulbous <i>Poa bulbosa</i>	Lettuce, prickly <i>Lactuca serriola</i>	Signalgrass, broadleaf <i>Brachiaria platyphylla</i>
Brome <i>Bromus</i> spp.	Morningglory <i>Ipomoea</i> spp.	Smartweed, Pennsylvania <i>Polygonum pennsylvanicum</i>
Buttercup <i>Ranunculus</i> spp.	Mustard, blue <i>Chorispora tenella</i>	Sowthistle, annual <i>Sonchus oleraceus</i>
Cheat <i>Bromus secalinus</i>	Mustard, tansy <i>Descurainia pinnata</i>	Spanishneedles* <i>Bidens bipinnata</i>
Chickweed, mouseear <i>Cerastium vulgatum</i>	Mustard, tumble <i>Sisymbrium altissimum</i>	Stinkgrass <i>Eragrostis cilianensis</i>
Cocklebur	Mustard, wild	Sunflower

<i>Xanthium strumarium</i>	<i>Sinapis arvensis</i>	<i>Helianthus annuus</i>
Corn, volunteer	Oats, wild	Thistle, Russian
<i>Zea mays</i>	<i>Avena fatua</i>	<i>Salsola kali</i>
Crabgrass	Panicum	Spurry, umbrella
<i>Digitaria</i> spp.	<i>Panicum</i> spp.	<i>Holosteum umbellatum</i>
Dwarf dandelion	Pennycress, field	Velvetleaf
<i>Krigia cespitosa</i>	<i>Thlaspi arvense</i>	<i>Abutilon theophrasti</i>
False flax, smallseed	Pigweed, redroot	Wheat
<i>Camelina microcarpa</i>	<i>Amaranthus retroflexus</i>	<i>Triticum aestivum</i>
Fiddleneck	Pigweed, smooth	Witchgrass
<i>Amsinkia</i> spp.	<i>Amaranthus hybridus</i>	<i>Panicum capillare</i>
Flaxleaf fleabane	Ragweed, common	
<i>Conyza bonariensis</i>	<i>Ambrosia artemisiifolia</i>	
Fleabane	Ragweed, giant	
<i>Erigeron</i> spp.	<i>Ambrosia trifida</i>	
Foxtail	Rocket, London	
<i>Setaria</i> spp.	<i>Sisymbrium irio</i>	

*Apply 3 pints (1.5 lbs ae) of this product per acre.

**Apply with hand-held equipment only.

Annual weeds will generally continue to germinate from seed throughout the growing season. Repeat treatments will be necessary to control later germinating weeds.

Perennial Weeds

Apply this product as follows to control or destroy most vigorously growing perennial weeds. Unless otherwise directed, allow at least 7 days after application before disturbing vegetation.

Add 2 or more quarts of a nonionic surfactant per 100 gallons of spray solution to the rates of this product given in this list. See the "Directions for Use" and "Mixing and Application" sections in this label for specific uses and application instructions.

NOTE: If weeds have been mowed or tilled, do not treat until regrowth has reached the recommended stages.

RESTRICTION: Fall treatments must be applied before a killing frost.

A second application may be necessary to control weeds regenerating from underground parts or seed. When applied as directed under the conditions described, this product plus surfactant will control the following perennial weeds:

Alfalfa
Medicago sativa
 Alligatorweed*
Alternanthera philoxeroides
 Anise/Fennel
Foeniculum vulgare
 Artichoke, Jerusalem
Helianthus tuberosus
 Bahiagrass
Paspalum notatum
 Beachgrass, European***
Ammophila arenaria
 Bermudagrass
Cynodon dactylon
 Bindweed, field
Convolvulus arvensis
 Bluegrass, Kentucky
Poa pratensis
 Blueweed, Texas
Helianthus ciliaris
 Brackenfern
Pteridium spp.
 Bromegrass, smooth
Bromus inermis
 Canarygrass, reed
Phalaris arundinacea
 Cattail
Typha spp.
 Clover, red
Trifolium pratense
 Clover, white
Trifolium repens
 Cogongrass
Imperata cylindrica
 Quackgrass
Agropyron repens
 Reed, giant
Arundo donax
 Ryegrass, perennial
Lolium perenne
 Smartweed, swamp
Polygonum coccineum
 Spatterdock
Nuphar luteum
 Starthistle, yellow
Centaurea solstitialis
 Sweet potato, wild*
Ipomoea pandurata
 Thistle, artichoke
Cynara cardunculus
 Thistle, Canada
Cirsium arvense

Cordgrass
Spartina spp.
 Cutgrass, giant*
Zizaniopsis miliacea
 Dallisgrass
Paspalum dilatatum
 Dandelion
Taraxacum officinale
 Dock, curly
Rumex crispus
 Dogbane, hemp
Apocynum cannabinum
 Fescue
Festuca spp.
 Fescue, tall
Festuca arundinacea
 Guineagrass
Panicum maximum
 Hemlock, poison
Conium maculatum
 Horsenettle
Solanum carolinense
 Horseradish
Armoracia rusticana
 Ice Plant
Mesembryanthemum crystallinum
 Johnsongrass
Sorghum halepense
 Kikuyugrass
Pennisetum clandestinum
 Knapweed
Centaurea repens
 Lantana
Lantana camara
 Timothy
Phleum pratense
 Torpedograss*
Panicum repens
 Tules, common
Scirpus acutus
 Vaseygrass
Paspalum urvillei
 Velvetgrass
Holcus spp.
 Waterhyacinth
Eichornia crassipes
 Waterlettuce
Pistia stratiotes
 Waterprimrose
Ludwigia spp.
 Wheatgrass, western
Agropyron smithii

Lespedeza: common, serices
Lespedeza striata
Lespedeza cuneata
 Loosestrife, purple
Lythrum salicaria
 Lotus, American
Nelumbo lutea
 Maidencane
Panicum hematomon
 Milkweed
Asclepias spp.
 Muhly, wirestem
Muhlenbergia frondosa
 Mullein, common
Verbascum thapsus
 Napiergrass
Pennisetum purpureum
 Nightshade, silverleaf
Solanum elaeagnifolium
 Nutsedge: purple, yellow
Cyperus rotundus
Cyperus esculentus
 Orchardgrass
Dactylis glomerata
 Pampasgrass
Cortaderia jubata
 Paragrass
Brachiaria mutica
 Phragmites**
Phragmites spp.

*Partial control.

**Partial control in southeastern states. See specific instructions below.

***Washington and Oregon only.

Alligatorweed – Apply 6 pints of this product per acre as a broadcast spray or as a 1¼ percent solution with hand-held equipment to provide partial control of alligatorweed. Apply when most of the target plants are in bloom. Repeat applications will be required to maintain such control.

Beachgrass, European (Washington and Oregon only) – Best results are obtained when applications are made when European beachgrass is actively growing through the boot to the full heading stages of growth. Applications should be made prior to the loss of more than 50% green leaf color in the fall.

Applications made during any period of plant (drought) stress, or beyond the recommended active growth period in the fall, will likely result in reduced performance.

A second application of AgSaver™ Glyphosate 5.4 Plus may be necessary to treat skips. Monitor treated acres prior to reseeding of desirable vegetation.

Spray-to-Wet Applications:

Apply an 8 percent solution of this product plus 0.5 to 1.5 percent nonionic surfactant on a spray-to-wet basis for control of European beachgrass.

Spray coverage should be uniform and complete but not to the point of runoff.

Wiper Applications:

For selective control of European beachgrass, apply a 33 1/3 percent solution of this product plus 1 to 2.5 percent nonionic surfactant during active growth. Avoid contact of herbicide solution with desirable vegetation. Wiping the plants in opposite directions may improve performance. Maximizing the amount of individual leaf tissue contacted with the wiping equipment will result in optimal performance.

Bermudagrass – Apply 7.5 pints (3.75 lbs ae) of this product per acre as a broadcast spray or as a 1½ percent solution with hand-held equipment. Apply when target plants are actively growing and when seed heads appear.

Bindweed, field/Silverleaf Nightshade/Texas Blueweed – Apply 6 to 7.5 pints (3 to 3.75 lbs ae) of this product per acre as a broadcast spray west of the Mississippi River and 4.5 to 6 pints (2.25 to 3 lbs ae) of this product per acre east of the Mississippi River. With hand-held equipment, use a 1 ½ percent solution. Apply when target plants are actively growing and are at or beyond full bloom. For silverleaf nightshade, best results can be obtained when application is made after berries are formed. Do not treat when weeds are under drought stress. New leaf development indicates active growth. For best results apply in late summer or fall.

Brackenfern – Apply 4.5 to 6 pints (2.25 to 3 lbs ae) of this product per acre as a broadcast spray or as a ¾ to 1 percent solution with hand-held equipment. Apply to fully expanded fronds which are at least 18 inches long.

Cattail – Apply 4.5 to 6 pints (2.25 to 3 lbs ae) of this product per acre as a broadcast spray or as a ¾ percent solution with hand-held equipment. Apply when target plants are actively growing and are at or beyond the early-to-full bloom stage of growth. Best results are achieved when application is made during the summer or fall months.

Cogongrass – Apply 4.5 to 7.5 pints (2.25 to 3.75 lbs ae) of this product per acre as a broadcast spray. Apply when cogongrass is at least 18 inches tall and actively growing in late summer or fall. Allow 7 or more days after application before tillage or mowing. Due to uneven stages of growth and the dense nature of vegetation preventing good spray coverage, repeat treatments may be necessary to maintain control. Cordgrass – Broadcast Applications (Air) – Apply 4 to 7.5 pints (2 to 3.75 lbs ae) of this product in 5-20 gallons of spray solution per acre. Add 1 to 2 quarts of nonionic surfactant per 100 gallons of spray solution.

Broadcast Applications (Ground) – Apply 4 to 7.5 pints (2 to 3.75 lbs ae) of this product in 10 to 60 gallons of spray solution per acre. For best results, ensure that complete coverage of cordgrass clumps is achieved. Add 1 to 2 quarts of a nonionic surfactant per 100 gallons of spray solution.

Hand-Held and High Volume Equipment - Apply a 2 to 8 percent solution of this product. Ensure that complete coverage of cordgrass clumps is achieved. Do not spray to the point of run-off. Add 1 to 2 quarts of a nonionic surfactant per 100 gallons of spray solution.

Wiper Applications - For wick or wiper applications, mix 1 gallon of this product with 2 gallons of clean water to make a 33 percent solution. Include a nonionic surfactant at a rate of 10 percent by volume of the total herbicide solution.

In heavy stands, a double application in opposite directions may improve results.

Application Conditions - Schedule applications in order to allow 6 hours before treated plants are covered by tidewater. Rainfall or immersion within 6 hours after application may reduce effectiveness.

The presence of debris and silt on the cordgrass plants will reduce performance of this product. It may be necessary to wash targeted plants prior to application to improve uptake of this product into the plant. Where cordgrass has been cut or mowed prior to application with AgSaver Glyphosate 5.4 Plus, ensure adequate regrowth of cordgrass occurs to allow for interception or absorption of the herbicide solution.

Cutgrass, giant – Apply 6 pints (3 lbs ae) of this product per acre as a broadcast spray or as a 1 percent solution with hand-held equipment to provide partial control of giant cutgrass. Repeat applications will be required to maintain such control, especially where vegetation is partially submerged in water. Allow for substantial regrowth to the 7 to 10-leaf stage prior to retreatment.

Dogbane, hemp/Knapweed/Horseradish – Apply 6 pints (3 lbs ae) of this product per acre as a broadcast spray or as a 1-½ percent solution with hand-held equipment. Apply when target plants are actively growing, and most have reached the late bud-to-flower stage of growth. For best results, apply in late summer or fall. Fescue, tall – Apply 4.5 pints (2.25 lbs ae) of this product per acre as a broadcast spray or as a 1 percent solution with hand-held equipment. Apply when target plants are actively growing, and most have reached the boot-to-head stage of growth. When applied prior to the boot stage, less desirable control may be obtained.

Guineagrass – Apply 4.5 pints (2.25 lbs ae) of this product per acre as a broadcast spray or as a ¾ percent solution with hand-held equipment. Apply when target plants are actively growing and when most have reached at least the 7-leaf stage of growth.

Johnsongrass/Bluegrass, Kentucky/Bromegrass, smooth/Canarygrass, reed/Orchardgrass/Ryegrass, perennial/Timothy/Wheatgrass, western – Apply 3 to 4.5 pints (1.5 to 2.25 lbs ae) of this product per acre as a broadcast spray or as a ¾ percent solution with hand-held equipment. Apply when target plants are actively growing, and most have reached the boot-to-head stage of growth. When applied prior to the boot stage, less desirable control may be obtained. In the fall, apply before plants have turned brown.

Lantana – Apply this product as a ¾ to 1 percent solution with hand-held equipment. Apply to actively growing lantana at or beyond the bloom stage of growth. Use the higher application rate for plants that have reached the woody stage of growth.

Loosestrife, purple – Apply 4 pints (2 lbs ae) of this product per acre as a broadcast spray or as a 1 to 1-½ percent solution using hand-held equipment. Treat when plants are actively growing at or beyond the bloom stage of growth. Best results are achieved when application is made during summer or fall months. Fall treatments must be applied before a killing frost.

Lotus, American – Apply 4 pints (2 lbs ae) of this product per acre as a broadcast spray or as a ¾ percent solution with hand-held equipment. Treat when plants are actively growing at or beyond the bloom stage of growth. Best results are achieved when application is made during summer or fall months. Fall treatments must be applied before a killing frost. Repeat treatment may be necessary to control regrowth from underground parts and seeds.

Maidencane/Paragrass – Apply 6 pints (3 lbs ae) of this product per acre as a broadcast spray or as a $\frac{3}{4}$ percent solution with hand-held equipment. Repeat treatments will be required, especially to vegetation partially submerged in water. Under these conditions, allow for regrowth to the 7 to 10-leaf stage prior to retreatment.

Milkweed, common – Apply 4.5 pints (2.25 lbs ae) of this product per acre as a broadcast spray or as a $1\frac{1}{2}$ percent solution with hand-held equipment. Apply when target plants are actively growing, and most have reached the late bud-to-flower stage of growth.

Nutsedge: purple, yellow – Apply 4.5 pints (2.25 lbs ae) of this product per acre as a broadcast spray, or as a $\frac{3}{4}$ percent solution with hand-held equipment to control existing nutsedge plants and immature nutlets attached to treated plants. Apply when target plants are in flower or when new nutlets can be found at rhizome tips. Nutlets which have not germinated will not be controlled and may germinate following treatment. Repeat treatments will be required for long-term control.

Pampasgrass – Apply a $1\frac{1}{2}$ percent solution of this product with hand-held equipment when plants are actively growing.

Phragmites – For partial control of phragmites in Florida and the counties of other states bordering the Gulf of Mexico, apply 7.5 pints per acre as a broadcast spray or apply a $1\frac{1}{2}$ percent solution with hand-held equipment. In other areas of the U.S., apply 4 to 6 pints per acre as a broadcast spray or apply a $\frac{3}{4}$ percent solution with hand-held equipment for partial control. For best results, treat during late summer or fall months when plants are actively growing and in full bloom. Due to the dense nature of the vegetation, which may prevent good spray coverage and uneven stages of growth, repeat treatments may be necessary to maintain control. Visual control symptoms will be slow to develop.

Quackgrass/Kikuyugrass/Muhly, wirestem – Apply 3 to 4.5 pints (1.5 to 2.25 lbs ae) of this product per acre as a broadcast spray or as a $\frac{3}{4}$ percent solution with hand-held equipment when most quackgrass or wirestem muhly is at least 8 inches in height (3 to 4-leaf stage of growth) and actively growing. Allow 3 or more days after application before tillage.

Reed, giant/ice plant – For control of giant reed and ice plant, apply a $1\frac{1}{2}$ percent solution of this product with hand-held equipment when plants are actively growing. For giant reed, best results are obtained when applications are made in late summer or fall.

Spatterdock – Apply 6 pints (3 lbs ae) of this product per acre as a broadcast spray or as a $\frac{3}{4}$ percent solution with hand-held equipment. Apply when most plants are in full bloom. For best results, apply during the summer or fall months.

Sweet potato, wild – Apply this product as a $1\frac{1}{2}$ percent solution using hand-held equipment. Apply to actively growing weeds that are at or beyond the bloom stage of growth. Repeat applications will be required. Allow the plant to reach the recommended stage of growth before retreatment.

Thistle: Canada, artichoke – Apply 3 to 4.5 pints (1.5 to 2.25 lbs ae) of this product per acre as a broadcast spray or as a $1\frac{1}{2}$ percent solution with hand-held equipment for Canada thistle. To control artichoke thistle, apply a 2 percent solution as a spray-to-wet application. Apply when target plants are actively growing and at or beyond the bud stage of growth.

Torpedograss – Apply 6 to 7.5 pints (3 to 3.75 lbs ae) of this product per acre as a broadcast spray or as a $\frac{3}{4}$ to $1\frac{1}{2}$ percent solution with hand-held equipment to provide partial control of torpedograss. Use the lower rates under terrestrial conditions, and the higher rates under partially submerged or a floating mat condition. Repeat treatments will be required to maintain such control.

Tules, common – Apply this product as a $1\frac{1}{2}$ percent solution with hand-held equipment. Apply to actively growing plants at or beyond the seedhead stage of growth. After application, visual symptoms will be slow to appear and may not occur for 3 or more weeks.

Water hyacinth – Apply 5 to 6 pints (2.5 to 3 lbs ae) of this product per acre as a broadcast spray or apply a ¾ to 1 percent solution with hand-held equipment. Apply when target plants are actively growing and at or beyond the early bloom stage of growth. After application, visual symptoms may require 3 or more weeks to appear with complete necrosis and decomposition usually occurring within 60 to 90 days. Use the higher rates when more rapid visual effects are desired.

Waterlettuce – For control, apply a ¾ to 1 percent solution of this product with hand-held equipment to actively growing plants. Use higher rates where infestations are heavy. Best results are obtained from mid-summer through winter applications. Spring applications may require retreatment.

Waterprimrose – Apply this product as a ¾ percent solution using hand-held equipment. Apply to plants that are actively growing at or beyond the bloom stage of growth, but before fall color changes occur. Thorough coverage is necessary for best control.

Other perennials listed on this label – Apply 4.5 to 7.5 pints (2.25 to 3.75 lbs ae) of this product per acre as a broadcast spray or as a ¾ to 1 ½ percent solution with hand-held equipment. Apply when target plants are actively growing, and most have reached early head or early bud stage of growth.

WOODY BRUSH AND TREES

When applied as directed under the conditions described, this product plus surfactant CONTROLS or PARTIALLY CONTROLS the following woody brush plants and trees:

Alder	Broom:
<i>Alnus</i> spp.	Scotch
Ash*	<i>Cytisus scoparius</i>
<i>Fraxinus</i> spp.	Buckwheat, California*
Aspen, quaking	<i>Eriogonum fasciculatum</i>
<i>Populus tremuloides</i>	Cascara*
Bearclover, Bearmat	<i>Rhamnus purshiana</i>
<i>Chamaebatia foliolosa</i>	Catsclaw*
Birch	<i>Acacia greggi</i>
<i>Betula</i> spp.	Ceanothus
Blackberry	<i>Ceanothus</i> spp.
<i>Rubus</i> spp.	Chamise
Broom:	<i>Adenostoma fasciculatum</i>
French	Maple:
<i>Cytisus monspessulanus</i>	Red**
Cherry:	<i>Acer rubrum</i>
Bitter	Sugar
<i>Prunus emarginata</i>	<i>Acer saccharum</i>
Black	Vine*
<i>Prunus serotina</i>	<i>Acer circinatum</i>
Pin	Monkey Flower*
<i>Prunus pensylvanica</i>	<i>Mimulus guttatus</i>
Coyote brush	Oak:
<i>Baccharis consanguinea</i>	Black*
Creeper, Virginia*	<i>Quercus velutina</i>
<i>Parthenocissus quinquefolia</i>	Northern pine
Dewberry	<i>Quercus palustris</i>
<i>Rubus trivialis</i>	Post
Dogwood	<i>Quercus stellata</i>
<i>Cornus</i> spp.	Red
Elderberry	<i>Quercus rubra</i>
<i>Sambucus</i> spp.	Southern red

Elm*	<i>Quercus falcata</i>
<i>Ulmus</i> spp.	White*
Eucalyptus, bluegum	<i>Quercus alba</i>
<i>Eucalyptus globules</i>	Persimmon*
Hasardia*	<i>Diospyros</i> spp.
<i>Haplopappus squamosus</i>	Poison Ivy
Hawthorn	<i>Rhus radicans</i>
<i>Crataegus</i> spp.	Poison Oak
Hazel	<i>Rhus toxicodendron</i>
<i>Corylus</i> spp.	Poplar, yellow*
Hickory	<i>Liriodendron tulipifera</i>
<i>Carya</i> spp.	Prunus
Holly, Florida; Brazilian Peppertree	<i>Prunus</i> spp.
<i>Schinus terebinthifolius</i>	Raspberry
Honeysuckle	<i>Rubus</i> spp.
<i>Lonicera</i> spp.	Redbud, eastern
Hornbeam, American	<i>Cercis canadensis</i>
<i>Carpinus caroliniana</i>	Rose, multiflora
Kudzu	<i>Rosa multiflora</i>
<i>Pueraria lobata</i>	Russian-olive
Locust, black*	<i>Elaeagnus angustifolia</i>
<i>Robinia pseudoacacia</i>	Sweet gum
Manzanita	<i>Liquidambar styraciflua</i>
<i>Arctostaphylos</i> spp.	Swordfern*
Sage: black, white	<i>Polystichum munitum</i>
<i>Salvia</i> spp.	Tallowtree, Chinese
Sagebrush, California	<i>Sapium sebiferum</i>
<i>Artemisia californica</i>	Thimbleberry
Salmonberry	<i>Rubus parviflorus</i>
<i>Rubus spectabilis</i>	Tobacco, tree*
Salt cedar*	<i>Nicotiana glauca</i>
<i>Tamarix</i> spp.	Trumpetcreeper
Saltbush, Sea myrtle	<i>Campsis radicans</i>
<i>Baccharis halimifolia</i>	Waxmyrtle, southern*
Sassafras	<i>Myrica cerifera</i>
<i>Sassafras aibidum</i>	Willow
Sourwood*	<i>Salix</i> spp.
<i>Oxydendrum arboreum</i>	
Sumac:	
Poison*	
<i>Rhus vernix</i>	
Smooth*	
<i>Rhus glabra</i>	
Winged*	
<i>Rhus copallina</i>	

*Partial Control

**See below for control or partial control instruction.

NOTE: If brush has been mowed or tilled or trees have been cut, do not treat until regrowth has reached the recommended stage of growth.

Apply the directed rate of this product plus 2 or more quarts of a nonionic surfactant per 100 gallons of spray solution when plants are actively growing and, unless otherwise directed, after full-leaf expansion. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation.

In arid areas, best results are obtained when application is made in the spring or early summer when brush species are at high moisture content and are flowering. Ensure thorough coverage when using hand-held equipment. Symptoms may not appear prior to frost or senescence with fall treatment.

Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatments are made following a frost.

See the "Directions for Use" and "Mixing and Application Instructions" sections in this label for labeled use and specific application instructions.

Applied as a 5 to 8 percent solution as a directed application as described in the "Hand-Held and High-Volume Equipment" section, this product will control or partially control all species listed in this section of this label. Use the higher rate of application for dense stands and larger woody brush and trees.

Apply the product as follows to control or partially control the following woody brush and trees.

Alder/Blackberry/Dewberry/Honeysuckle/Oak, Post/Raspberry – For control, apply 4.5 to 6 pints (2.25 to 3 lbs ae) of this product per acre as a broadcast spray or as a $\frac{3}{4}$ to 1 $\frac{1}{4}$ percent solution with hand-held equipment.

Aspen, Quaking/Hawthorn/Trumpet creeper – For control, apply 3 to 4.25 pints (1.5 to 3.75 lbs ae) of this product per acre as a broadcast spray or as a $\frac{3}{4}$ to 1 $\frac{1}{4}$ percent solution with hand-held equipment.

Birch/Elderberry/Hazel/Salmonberry/Thimbleberry – For control, apply 3 pints (1.5 lbs ae) per acre of this product as a broadcast spray or as a $\frac{3}{4}$ percent solution with hand-held equipment.

Broom: French, Scotch – For control, apply a 1 $\frac{1}{4}$ to 1 $\frac{1}{2}$ percent solution with hand-held equipment.

Buckwheat, California/Hasardia/Monkey Flower/Tobacco, Tree – For partial control of these species, apply a $\frac{3}{4}$ to 1 $\frac{1}{2}$ percent solution of this product as a foliar spray with hand-held equipment. Thorough coverage of foliage is necessary for best results.

Catsclaw – For partial control, apply a 1 $\frac{1}{4}$ to 1 $\frac{1}{2}$ percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed.

Cherry: Bitter, Black, Pin/Oak, Southern Red/Sweet Gum/Prunus – For control, apply 3 to 7.5 pints (1.5 to 3.75 lbs ae) of this product per acre as a broadcast spray or as a 1 to 1 $\frac{1}{2}$ percent solution with hand-held equipment.

Coyote brush – For control, apply a 1 $\frac{1}{4}$ to 1 $\frac{1}{2}$ percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed.

Dogwood/Hickory/Salt cedar – For partial control, apply a 1 to 2 percent solution of this product with hand-held equipment or 6 to 7.5 pints (3 to 3.75 lbs ae) per acre as a broadcast spray.

Eucalyptus, bluegum – For control of eucalyptus resprouts, apply a 1- $\frac{1}{2}$ percent solution of this product with hand-held equipment when resprouts are 6 to 12-feet tall. Ensure complete coverage. Apply when plants are actively growing. Avoid application to drought-stressed plants.

Holly, Florida (Brazilian peppertree (*Schinus terebinthifolius*)) – For partial control, apply this product as a 1- $\frac{1}{2}$ percent solution with hand-held equipment.

Alternatively, when applied as directed, this product with QuikSorb™ Penetrant will control or partially control Brazilian peppertree in areas such as dry drainage ditches and canals, wildlife habitat restoration and management areas, roadsides, railroads, fence rows, and similar non-crop areas.

The recommended application technique is directed spot treatment of Brazilian peppertree using hand-held equipment only. Apply this product using backpack, hand-held, handgun or similar equipment. Use flat fan, cone, or similar nozzles that will provide effective spray coverage of target vegetation. Do not apply to Brazilian peppertree growing in water. Do not use aerial, boom-type, or other broadcast spray equipment. These applications are more effective on small brush less than 15 feet in height or 3-inch stem diameter.

Basal and Selective Stem Application:

Apply a solution consisting of 25% v/v of this product and 75% v/v of QuikSorb™ penetrant. Completely cover the lower 18-24 inches of the brush stems or trunks. For larger stems over 3 inches in diameter, treat up to 48 inches or higher from the ground level. For better control of large trees, apply spray solution directly to upper foliage of plant canopy. Spray coverage should be uniform, covering at least 40 to 60% of the upper foliage and stems. Application is best when made to young, actively growing stems, branches, and foliage. Spray-to-wet but not to the point of run-off.

Read and carefully observe the label claims, cautionary statements, and all information on the labels of all products used in this tank mixture.

Kudzu – For control, apply 6 pints (3 lbs ae) of this product per acre as a broadcast spray or as a 1-½ percent solution with hand-held equipment. Repeat applications will be required to maintain control.

Maple, Red – For control, apply as a ¾ to 1 ¼ percent solution with hand-held equipment when leaves are fully developed. For partial control, apply 2 to 7.5 pints (1 to 3.75 lbs ae) of this product per acre as a broadcast spray.

Maple, Sugar/Oak: Northern Pin, Red – For control, apply as a ¾ to 1 ¼ percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed.

Poison Ivy/Poison Oak – For control, apply 6 to 7.5 pints (3 to 3.75 lbs ae) of this product per acre as a broadcast spray or as a 1-½ percent solution with hand-held equipment. Repeat applications may be required to maintain control. Fall treatments must be applied before leaves lose green color.

Rose, multiflora – For control, apply 3 pints (1.5 lbs ae) of this product per acre as a broadcast spray or as a ¾ percent solution of this product as a foliar spray with hand-held equipment. Thorough coverage of foliage is necessary for best results.

Sage, black/Sagebrush, California/Chamise/Tallowtree, Chinese – For control of these species apply as a ¾ percent solution of this product as a foliar spray with hand-held equipment. Thorough coverage of foliage is necessary for best results.

Saltbush, Sea myrtle – For control, apply this product as a 1 percent solution with hand-held equipment.

Waxmyrtle, southern – For partial control, apply this product as a 1-½ percent solution with hand-held equipment.

Willow – For control, apply 4.5 pints (2.25 lbs ae) of this product per acre as a broadcast spray or as a ¾ percent solution with hand-held equipment.

Other woody brush and trees listed in this label – For partial control, apply 3 to 7.5 pints (1.5 to 3.75 lbs ae) of this product per acre as a broadcast spray or as a ¾ to 1 ½ percent solution with hand-held equipment.

AQUATIC SITES

When applied as directed and under the conditions described in the “Weeds Controlled” section in this label, this product will control or partially control the labeled weeds growing in the following industrial, recreational, and public areas or other listed aquatic and terrestrial sites.

RESTRICTION: When applying AgSaver™ Glyphosate 5.4 Plus directly to water the use of surfactants must only be used if the surfactant is known to be non-toxic to aquatic species.

Aquatic Sites – This product may be applied to emerged weeds in all bodies of fresh and brackish water which may be flowing, nonflowing or transient. This includes lakes, rivers, streams, ponds, estuaries, rice levees, seeps, irrigation and drainage ditches, canals, reservoirs, wastewater treatment facilities, wildlife habitat restoration and management areas, and similar sites.

If aquatic sites are present in the noncrop area and are part of the intended treatment, read and observe the following directions:

This product does not control plants which are completely submerged or have a majority of their foliage under water.

There is no restriction on the use of treated water for irrigation, recreation, or domestic purposes.

Consult local state fish and game agency and water control authorities before applying this product to public water. Permits may be required to treat such water.

For treatments after drawdown of water or in dry ditches, allow 7 or more days after treatment before reintroduction of water to achieve maximum weed control. Apply this product within 1 day after drawdown to ensure application to actively growing weeds.

Floating mats of vegetation may require retreatment. Avoid wash-off of sprayed foliage by spray boat or recreational boat backwash or by rainfall within 6 hours of application. Do not re-treat within 24 hours following the initial treatment.

Applications made to moving bodies of water must be made while traveling upstream to prevent concentration of this herbicide in water. When making any bankside applications, do not overlap more than 1 foot into open water. Do not spray in bodies of water where weeds do not exist. The maximum application rate of 7.5 pints (3.75 lbs ae) per acre must not be exceeded in any single broadcast application that is being made over water.

When emerged infestations require treatment of the total surface area of impounded water, treating the area in strips may avoid oxygen depletion due to decaying vegetation. Oxygen depletion may result in fish kill.

RESTRICTIONS:

- Do not apply this product directly to water within ½ mile up-stream of an active potable water intake in flowing water (i.e., river stream, etc.) or within ½ mile of an active potable water intake in a standing body of water such as lake, pond, or reservoir.
- To make aquatic applications around and within ½ mile of active potable water intakes, the water intake must be turned off for a minimum period of 48 hours after application.
- The water intake may be turned on prior to 48 hours if the glyphosate level in the intake water is below 0.7 parts per million as determined by laboratory analysis.
- These aquatic applications may be made ONLY in those cases where there are alternative water sources or holding ponds which would permit the turning off of an active potable water intake for a minimum period of 48 hours after the applications. This restriction does not apply to intermittent inadvertent overspray of water in terrestrial use sites.

OTHER NONCROP TYPE SITES

This product may be used to control the listed weeds in terrestrial noncrop sites and/or in aquatic sites within these areas.

Airports
Golf Courses
Habitat Restoration & Management Areas
Highways & Roadsides

Industrial Plant Sites
Lumberyards
Parking Areas
Parks
Petroleum Tank Farms
Pipeline, Power, Telephone & Utility Rights-of-Way
Pumping Installations
Railroads
Schools
Storage Areas
Listed Similar Sites

TANK MIXTURES

NOTE: Read and carefully observe the label directions, cautionary statements, and all information on the labels of products used in these tank mixtures before proceeding with these directions.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

AGSAVER™ GLYPHOSATE 5.4 PLUS Triclopyr, butoxyethyl ester

For burndown and partial control or suppression of woody brush and weeds in industrial sites:

When applied as directed for "Noncrop Uses" under the conditions described, this product, and an approved surfactant plus triclopyr, butoxyethyl ester, provides burndown and partial control or suppression of woody brush and vegetation labeled for this product. This tank mixture is recommended for use on rights-of-way (utility, railroad, highway, pipeline), fencerows, roadsides, nonirrigation ditchbanks, wasteland and similar noncrop or industrial sites. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Hand-Held and High-Volume Applications:

Use 3 to 6 pints (1.5 to 3 lbs ae) of AgSaver™ Glyphosate 5.4 Plus herbicide and 2 or more quarts of an approved surfactant, plus the labeled rate of triclopyr, butoxyethyl ester per 100 gallons of spray solution and apply to foliage of actively growing woody brush and weeds. Applications should be made on a spray to wet basis. Spray coverage should be uniform and complete. Do not spray to point of runoff.

Broadcast Applications with Ground Equipment:

Use 3 to 6 pints (1.5 to 3 lbs ae) of AgSaver™ Glyphosate 5.4 Plus the labeled rate of triclopyr, butoxyethyl ester in sufficient water and make 20 to 100 gallons of total spray per acre. Use 2 to 4 quarts of an approved surfactant per 100 gallons of spray solution with this product.

Aerial Application (Helicopter Only):

Use 3 to 6 pints (1.5 to 3 lbs ae) of AgSaver™ Glyphosate 5.4 Plus surfactant plus the labeled rate of triclopyr, butoxyethyl ester and apply in a total spray volume of 10 to 20 gallons per acre. Aerial sprays should be applied using suitable drift control. Use 2 to 4 quarts of an approved surfactant per 100 gallons of spray solution with this product.

Apply when plants are actively growing and after full leaf expansion of woody brush. Use the higher rates of these products where vegetation is heavy or dense, or where hard-to-control brush species are prevalent. Repeat applications may be necessary to maintain control and to suppress areas where canopied vegetation prevents good spray coverage and penetrations.

Drift control additives may be used. When a drift control additive is used, read, and carefully observe the cautionary statements and all other information appearing on the additive label.

AGSAVER™ GLYPHOSATE 5.4 PLUS IMAZAPYR

When applied as directed, this tank mixture will control or partially control labeled woody brush, trees, and herbaceous weeds in noncrop areas. In addition to the weeds listed on this label, this tank mixture will control arrowweed, salt cedar and yaupon. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Hand-Held and High-Volume Applications:

Use 6 to 12 pints (3 to 6 lbs ae) of AgSaver™ Glyphosate 5.4 Plus the labeled rate of imazapyr per 100 gallons of spray solution. Add 2 to 4 quarts of nonionic surfactant per 100 gallons of spray solution.

Apply to foliage of actively growing vegetation. Applications should be made on a spray-to-wet basis. Spray coverage should be uniform and complete. Do not spray to the point of runoff.

Broadcast Applications with Ground Equipment:

Use 3 to 7.5 pints (1.5 to 3.75 lbs ae) of AgSaver™ Glyphosate 5.4 Plus the labeled rate of imazapyr in sufficient water to apply in a total spray volume of 10 to 20 gallons per acre. Add 2 to 4 quarts of nonionic surfactant per 100 gallons of spray solution. Apply to foliage of actively growing vegetation.

Apply to woody brush and trees after full leaf expansion until initiation of fall color. Avoid direct applications to any body of water.

RESTRICTION:

Do not apply on ditches used to transport irrigation water.

AGSAVER™ GLYPHOSATE 5.4 PLUS 2,4-D AMINE

When applied as a tank mixture, this product will control the annual weeds listed in this label booklet. This tank mixture will control or partially control the listed perennial weeds, woody brush, and trees. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Use 1.5 to 2.5 pints (0.75 to 1.25 lbs ae) of this product plus the labeled rate of 2,4-D amine (labeled for aquatic sites) for control of annual weeds.

Use 3 to 7.5 pints (1.5 to 3.75 lbs ae) of this product plus the labeled rate of 2,4-D amine (labeled for aquatic sites) for control or partial control of perennial weeds, woody brush, and trees. The tank mixture may be used on alligatorweed, smartweed, waterprimrose, waxmyrtle and other labeled weeds.

When using this product, mix 2 or more quarts of a nonionic surfactant per 100 gallons of spray solution. Always read and follow the surfactant manufacturer's label directions.

Always predetermine the compatibility of the tank mixtures of this herbicide and 2,4-D amine by mixing small proportional quantities in advance.

Mix in the following sequence: Fill sprayer tank one-half full with water, add AgSaver Glyphosate 5.4 Plus, then 2,4-D amine and finally surfactant. Fill sprayer tank to final volume with water.

RESTRICTIONS:

- Do not mix AgSaver™ Glyphosate 5.4 Plus amine concentrates without water carrier.
- Do not mix AgSaver™ Glyphosate 5.4 Plus and 2,4-D amine in bypass injector-type spray equipment.

WETLAND SITES

This product may be used in and around water (aquatic areas) and wetlands found in forestry and in power, telephone, and pipeline rights-of-way sites, including where these sites are adjacent to or surrounding domestic water supply reservoirs, supply streams, lakes, and ponds. Read and observe the following before making applications in and around water.

Consult local public water control authorities before applying this product in and around public water. Permits may be required to treat in such areas.

There is no restriction on the use of treated water for irrigation, recreation, or domestic purposes.

RESTRICTIONS:

- Do not apply this product directly to water within ½ mile up-stream of an active potable water intake in flowing water (i.e., river, stream, etc.) or within ½ mile of an active potable water intake in a standing body of water such as lake, pond, or reservoir. To make aquatic applications around and within ½ mile of active potable water intakes, the water intake must be turned off for a minimum period of 48 hours after the application. These aquatic applications may be made ONLY in those cases where there are alternative water sources or holding ponds which would permit the turning off of an active potable water intake for a minimum period of 48 hours after the applications. This restriction does not apply to intermittent inadvertent overspray of water in terrestrial use sites.
- Do not spray open bodies of water where woody brush, trees and herbaceous weeds do not exist.
- The maximum application rate of 3.75 quarts per acre must not be exceeded in a single over-water broadcast application except as follows, where any labeled rate may be applied:
 - Stream crossings in utility rights-of-way.
 - Where applications will result in less than 20 percent of the total water area being treated.

WILDLIFE HABITAT RESTORATION AND MANAGEMENT AREAS

This product can be used for the restoration and/or maintenance of native habitat and in wildlife management areas.

Habitat Restoration and Maintenance – When applied as directed, exotic and other undesirable vegetation may be controlled in habitat management areas. Applications may be made to allow recovery of native plant species, to open up water to attract waterfowl, and for similar broad-spectrum vegetation control requirements in habitat management areas. Spot treatments may be made to selectively remove unwanted plants for habitat enhancement. For spot treatments, care should be exercised to keep spray off of desirable plants.

Wildlife Food Plots – This product may be used as a site preparation treatment prior to planting wildlife food plots. Apply as directed to control vegetation in the plot area. Any wildlife food species may be planted after applying this product, or native species may be allowed to reinfest the area. If tillage is needed to prepare a seedbed, wait 7 days after applying this product before tilling to allow for maximum effectiveness.

WIPER APPLICATIONS

For wick or wiper applications, mix 1 gallon of this product with 2 gallons of clean water to make a 33 percent solution. Include a nonionic surfactant at a rate of 10 percent by volume of total herbicide solution.

Wiper applications can be used to control or suppress annual and perennial weeds listed on this label. In heavy weed stands, a double application in opposite directions may improve results. See the "Weeds Controlled" section in this label for timing, growth stage and other instructions for achieving optimum results.

Bromegrass (smooth), Canarygrass (reed), Dock (curly), Mullein (common), Quackgrass and Canada thistle: This product may be applied through a wiper applicator after dilution with water and thorough mixing to these weeds growing in or along aquatic sites.

Wiper applicators, including wick devices, apply the herbicide solution by rubbing the weed with an absorbent material containing the herbicide solution.

Contact of the herbicide solution with desirable vegetation may result in damage or destruction. Applicators used above desired vegetation should be adjusted so that the lowest wiper contact point is at least two (2) inches above this vegetation. Application made above desirable vegetation should be made when the weeds are a minimum of six (6) inches above this vegetation.

Best results may be attained when more of the weed is exposed to the herbicide solution. Weeds not contacted (wiped) with the herbicide solution will not be affected. This may occur in dense clumps, severe infestations, or when the height of the weed varies so that not all weeds are contacted.

In severe infestations, reduce equipment ground speed to ensure that adequate amounts of this herbicide solution are wiped onto the weeds. When wiping moderate weed infestations an adequate flow rate should be 3 to 4 quarts of herbicide solution per mile of canal (wiping 4 foot band). For best results, do not allow wiper applicator to contact water.

Note:

- Maintain wiper equipment in good operating condition.
- Adjust height of wiper applicator to ensure adequate contact with weeds.
- Keep wiping surfaces clean.
- Keep wiper material at proper degree of saturation with herbicide solution.
- DO NOT use wiper equipment when weeds are wet or under conditions where wave action or other water immersions will wash the solution off the weed.
- DO NOT operate equipment at ground speeds of greater than 5 MPH. As weed density increases, reduce equipment ground speed to ensure good coverage of weeds.
- Be aware that on sloping ground, the herbicide solution may migrate, causing dripping on the lower end and drying on the upper end of the wiper applicator.
- Variation in equipment design may affect weed control. With wiper applicators, the wiping material and its orientation must allow delivery of sufficient quantities of the herbicide solution directly to the weeds.
- Mix only the amount of solution to be used during a one day period as reduced activity may result from use of leftover solutions.

Mixing Instructions:

Mix 2.5 gallons (10 lbs ae) of AgSaver™ Glyphosate 5.4 Plus herbicide with 7.5 gallons of water to prepare a 25 percent solution. Add 1 quart of an approved surfactant per 10 gallons of herbicide solution (2 ½ percent surfactant by total volume). Apply this solution to weeds listed above.

CUT STUMP APPLICATION

Woody vegetation may be controlled by treating freshly cut stumps of trees and resprouts with this product. Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut vegetation close to the soil surface. Apply a 50 to 100 percent solution of this product to freshly cut surface immediately after cutting. Delay in applying this product may result in reduced performance. For best results, trees should be cut during periods of active growth and full leaf expansion.

When used according to directions for cut stump application, this product will CONTROL, PARTIALLY CONTROL, or SUPPRESS most woody brush and tree species, some of which are listed below:

Alder	Poplar*
<i>Alnus</i> spp.	<i>Populus</i> spp.
Coyote brush*	Reed, giant
<i>Baccharis consanguinea</i>	<i>Arundo donax</i>
Dogwood*	Salt cedar
<i>Cornus</i> spp.	<i>Tamarix</i> spp.
Eucalyptus	Sweet gum*
<i>Eucalyptus</i> spp.	<i>Liquidambar styraciflua</i>
Hickory*	Sycamore*
<i>Carya</i> spp.	<i>Platanus occidentalis</i>
Madrone	Tan oak
<i>Arbutus menziesii</i>	<i>Lithocarpus densiflorus</i>
Maple*	Willow
<i>Acer</i> spp.	<i>Salix</i> spp.
Oak	
<i>Quercus</i> spp.	

* Not Registered for Use in California.

INJECTION AND FRILL APPLICATIONS

Woody vegetation may be controlled by injection or frill application of this product. Apply this product using suitable equipment which must penetrate into living tissue. Apply the equivalent of 1 mL of this product per 2 to 3 inches of trunk diameter. This is best achieved by applying 25 to 100 percent concentration of this product either to a continuous frill around the tree or as cuts evenly spaced around the tree below all branches. As tree diameter increases in size, better results are achieved by applying dilute material to a continuous frill or more closely spaced cuttings. Avoid application techniques that allow runoff to occur from frill or cut areas in species that exude sap freely after frills or cutting. In species such as these, make frill or cut at an oblique angle so as to produce a cupping effect and use undiluted material. For best results, applications should be made during periods of active growth and full leaf expansion.

This treatment WILL CONTROL the following woody species:

Oak
Quercus spp.
Poplar
Populus spp.
Sweet gum
Liquidambar styraciflua
Sycamore
Platanus occidentalis

This treatment WILL SUPPRESS the following woody species:

Black gum*

Nyssa sylvatica

Dogwood

Cornus spp.

Hickory

Carya spp.

Maple, red

Acer rubrum

* Not Registered for Use in California.

INDUSTRIAL TURF

Apply 3 to 5 fl oz (0.09 to 0.16 lb ae) of this product per acre alone or as directed for a tank mixture, at spray volumes of 10 to 40 gallons per acre.

When using this product, mix 2 quarts of a nonionic surfactant per 100 gallons of spray solution. This product can be used for growth and seedhead suppression of:

Tall Fescue Smooth Brome

For best results, apply this product in a tank mixture to actively growing turfgrasses after greenup in the spring of the year. For suppression of seedheads, applications must be made before boot-to-seedhead stage of development. Applications made from seedhead emergence until maturity may result in turf discoloration or injury.

After mowing or removal of seedheads, this product in a tank mixture may also be used to suppress the growth of certain turfgrasses. Allow turf to recover from stress caused by heat, drought or mowing before making applications. Applications made to turf under stress may increase the potential for discoloration or injury.

Annual Grasses

For growth suppression of some annual grasses such as annual ryegrass, wild barley, and wild oats, apply 3 to 4 fl oz (0.09 to 0.13 lb ae) of this product in 10 to 40 gallons of spray solution per acre. Applications should be made when annual grasses are actively growing and before the seedheads are in the boot stage of development. Treatments made after seedhead emergence may cause injury to the desired grasses.

TANK MIXTURES FOR INDUSTRIAL TURFGRASSES

For the following tank mixtures, consult each product label for weeds controlled and the proper stage of application. Do not treat turf under stress. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Tank Mixtures plus 2,4-D Amine

For additional weed control benefits, the labeled rate of 2,4-D amine may be added to the following tank mixtures.

TALL FESCUE

AgSaver™ Glyphosate 5.4 Plus Chlorsulfuron

For suppression of tall fescue growth and seedheads, and control or partial control of some annual weeds, apply this tank mixture after greenup and prior to boot-to-seedhead stage of development. Use up to ½ ounce of Chlorsulfuron per acre.

This tank mixture can also be applied after mowing or removal of tall fescue seedheads for turf growth suppression and control or partial control of some annual weeds. Make only one of the above applications per year.

AgSaver™ Glyphosate 5.4 Plus Sulfometuron

For suppression of tall fescue growth and seedheads, and control or partial control of some annual weeds, apply this tank mixture after greenup and prior to boot-to-seedhead stage of development. Use the labeled rate of Sulfometuron.

AgSaver™ Glyphosate 5.4 Plus Metsulfuron

This tank mixture can be applied after mowing or removal of tall fescue seedheads for turf growth suppression and control or partial control of some annual weeds. Use the labeled rate of Metsulfuron.

SMOOTH BROME AgSaver™ Glyphosate 5.4 Plus Sulfometuron

For suppression of smooth brome growth and seedheads and control or partial control of some annual weeds, apply this tank mixture after greenup and prior to boot-to-seedhead stage of development. Use the labeled rate of Sulfometuron.

RELEASE OF BERMUDAGRASS OR BAHIAGRASS ON NONCROP SITES

RELEASE OF DORMANT BERMUDAGRASS AND BAHIAGRASS

When applied as directed, this product will provide control or suppression of many winter annual weeds and tall fescue for effective release of dormant bermudagrass or bahiagrass. Make applications to dormant bermudagrass or bahiagrass.

For best results on winter annuals, treat when weeds are in an early growth stage (below 6 inches in height) after most have germinated. For best results on tall fescue, treat when fescue is in or beyond the 4 to 6-leaf stage.

WEEDS CONTROLLED

Rate specifications for control or suppression of winter annuals and tall fescue are listed below.

Apply the specified rates of this product in 10 to 25 gallons of water per acre plus 2 quarts nonionic surfactant per 100 gallons of total spray volume.

WEEDS CONTROLLED OR SUPPRESSED

NOTE: C = Control
S = Suppression

WEED SPECIES	AgSaver™ Glyphosate 5.4 Plus fl oz/acre					
	6 (0.19 lb ae)	9 (0.28 lb ae)	12 (0.38 lb ae)	18 (0.56 lb ae)	24 (0.75 lb ae)	48 (1.5 lbs ae)
Barley, little <i>Hordeum pusillum</i>	S	C	C	C	C	C
Bedstraw, catchweed <i>Galium aparine</i>	S	C	C	C	C	C
Bluegrass, annual <i>Poa annua</i>	S	C	C	C	C	C
Chervil <i>Chaerophyllum tainturieri</i>	S	C	C	C	C	C

WEED SPECIES	AgSaver™ Glyphosate 5.4 Plus fl oz/acre					
	6 (0.19 lb ae)	9 (0.28 lb ae)	12 (0.38 lb ae)	18 (0.56 lb ae)	24 (0.75 lb ae)	48 (1.5 lbs ae)
Chickweed, common <i>Stellaria media</i>	S	C	C	C	C	C
Clover, crimson <i>Trifolium incarnatum</i>	•	S	S	C	C	C
Clover, large hop <i>Trifolium campestre</i>	•	S	S	C	C	C
Speedwell, corn <i>Veronica arvensis</i>	S	C	C	C	C	C
Fescue, tall <i>Festuca arundinacea</i>	•	•	•	•	S	S
Geranium, Carolina <i>Geranium carolinianum</i>	•	•	S	S	C	C
Henbit <i>Lamium amplexicaule</i>	•	S	C	C	C	C
Ryegrass, Italian <i>Lolium multiflorum</i>	•	•	S	C	C	C
Vetch, common <i>Vicia sativa</i>	•	•	S	C	C	C

*These rates apply only to sites where an established competitive turf is present.

RELEASE OF ACTIVELY GROWING BERMUDAGRASS

Use only on sites where bahiagrass or bermudagrass are desired for ground cover and some temporary injury or yellowing of the grasses can be tolerated.

When applied as directed, this product will aid in the release of bermudagrass by providing control of annual species listed in the "Weeds Controlled" section in this label, and suppression or partial control of certain perennial weeds.

For control or suppression of those annual species listed in this label, use $\frac{3}{4}$ to 2 $\frac{1}{4}$ pints (0.38 to 1.13 lbs ae) of this product as a broadcast spray in 10 to 25 gallons of spray solution per acre, plus 2 quarts of a nonionic surfactant per 100 gallons of total spray volume. Use the lower rate when treating annual weeds below 6 inches in height (or length of runner in annual vines). Use the higher rate as size of plants increases or as they approach flower or seedhead formation.

Use the higher rate for partial control or longer-term suppression of the following perennial species. Use lower rates for shorter-term suppression of growth.

Bahagrass
Dallisgrass
Fescue (tall)
Johnsongrass**
Trumpet creeper*
Vaseygrass

*Suppression at the higher rate only.

**Johnsongrass is controlled at the higher rate.

Use only on well-established bermudagrass. Bermudagrass injury may result from the treatment, but regrowth will occur under moist conditions. Do not repeat applications in the same year since severe injury may result.

BAHIAGRASS SEEDHEAD AND VEGETATIVE SUPPRESSION

When applied as directed in the “Noncrop Sites” section in this label, this product will provide significant inhibition of seedhead emergence and will suppress vegetative growth for a period of approximately 45 days with single applications and approximately 120 days with sequential applications.

Apply this product 1 to 2 weeks after full green-up of bahiagrass or after the bahiagrass has been mowed to a uniform height of 3 to 4 inches. Applications must be made prior to seedhead emergence. Apply 5 fl oz (0.16 lb ae) per acre of this product, plus 2 quarts of an approved nonionic surfactant per 100 gallons of total spray volume in 10 to 25 gallons of water per acre.

Sequential applications of this product plus nonionic surfactant may be made at approximately 45-day intervals to extend the period of seedhead and vegetative growth suppression. For continued vegetative growth suppression, sequential applications must be made prior to seedhead emergence.

Apply no more than 2 sequential applications per year. As a first sequential application, apply 3 fl oz (0.09 lb ae) of this product per acre plus nonionic surfactant. A second sequential application of 2 to 3 fl oz per acre plus nonionic surfactant may be made approximately 45 days after the last application.

ANNUAL GRASS GROWTH SUPPRESSION

For growth suppression of some annual grasses, such as annual ryegrass, wild barley and wild oats growing in coarse turf on roadsides or other industrial areas, apply 3 to 4 fl oz (0.09 to 0.13 lb ae) of this product in 10 to 40 gallons of spray solution per acre. Mix 2 quarts of a nonionic surfactant per 100 gallons of spray solution. Applications should be made when annual grasses are actively growing and before the seedheads are in the boot stage of development. Treatments made after seedhead emergence may cause injury to the desired grasses.

18.0 WARRANTY DISCLAIMER, INHERENT RISKS OF USE, LIMITATION OF REMEDIES

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Remedies before using this product.

If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following warranty disclaimer, inherent risks of use and limitation of remedies.

WARRANTY DISCLAIMER

AgSaver, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, AGSAVER, LLC MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label directions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of AgSaver, LLC or the seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

LIMITATION OF REMEDIES

To the extent consistent with applicable law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at AgSaver, LLC's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent consistent with applicable law, AgSaver, LLC shall not be liable for losses or damages resulting from handling or use of this product unless AgSaver, LLC is promptly notified of such loss or damage in writing. In no case shall AgSaver, LLC be liable for consequential or incidental damages or losses. The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of AgSaver, LLC or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

USES WITH OTHER PRODUCTS (TANK MIXES)

When used in combination as directed by AgSaver, to the extent consistent with applicable law, the liability of AgSaver, shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the AgSaver product in such combination use.

AgSaver is a trademark of AgSaver, LLC

[EPA APPROVAL DATE]