



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

AmTide LLC
c/o Mr. Michael Kellogg
Pyxis Regulatory Consulting, Inc.
4110 136th St. NW
Gig Harbor, WA 98332

JAN - 6 2012

Subject: AmTide TEBU 3.6F Foliar Fungicide
EPA Reg. No. 83851-9
Your amendment dated October 27, 2011
Decision Number 445331

Dear Mr. Kellogg:

The label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act as amended is acceptable provided the following conditions are met:

- 1) On page 21 under "Ornamental Use Restrictions and Precautions", add a bullet stating "Do not apply more than 10 fl oz per acre in a single application".

One copy of the label stamped "Accepted, with comments" is enclosed for your records. Please submit one copy of the final printed label before the product is released for shipment. If you have any questions, please contact Tracy Keigwin of my team at (703) 305-6605 and/or via email at keigwin.tracy@epa.gov.

Sincerely,

A handwritten signature in cursive script that reads "Mary L. Waller".

Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7505P)

Enclosure: Label stamped "Accepted, with comments"

AmTide TEBU 3.6F Foliar Fungicide

ACTIVE INGREDIENT:

Tebuconazole,

Alpha-[2-(4-chlorophenyl)ethyl]alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol.....40.53%

OTHER INGREDIENTS..... 59.47%**TOTAL**..... 100.00%

Contains 3.6 pounds Tebuconazole per gallon.

STOP-Read the label before use
KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail)

FIRST AID	
If swallowed:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For MEDICAL Emergencies call Chemtrec at 1-800-424-9300. For PRODUCT USE information call 949-753-4723.	
Note to Physician: No specific antidote. Treat symptomatically. Symptoms of Poisoning: The compound does not cause any definite symptoms that would be diagnostic. Contact with eyes may cause irritation.	

EPA Reg. No.: 83851-9

EPA Est. No.:

Manufactured for:

AmTide, LLC

21Hubble

Irvine, CA 92618, USA

ACCEPTED
with **COMMENTS**
In EPA Letter Dated:

1/6/2012

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticides
registered under EPA Reg. No.

NET CONTENTS:

83851-9

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately, if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory: Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams and springs will reduce the potential for contamination of water from rainfall- runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact worker or other persons, either directly or through drift. Only protected handlers or protected supervisors may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirement in this box only applies to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) listed in the application directions for the crop being treated.

PPE required for early to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate, or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box only apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Turf and Landscape Uses: Keep children and pets out of treated areas until sprays have dried.

GENERAL INFORMATION FOR AGRICULTURAL USES
SHAKE WELL BEFORE USING

Spray Volume: Apply TEBU 3.6F Foliar Fungicide with ground or aerial equipment using sufficient volume of spray to provide thorough coverage. TEBU 3.6F Foliar Fungicide may be applied in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment. Check equipment calibration frequently. Continuous agitation is required to keep the material in suspension.

Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied. If necessary, increase the spray volume per acre for complete crop coverage. Use the higher rate under conditions of severe disease pressure. Also, see local State Extension Service recommendations for application schedules.

Chemigation: Do not apply this product through any type of irrigation system.

Mixing: Add specified amount of TEBU 3.6F Foliar Fungicide into the spray tank while filling with water to the desired level. Operate the agitator while mixing. If other materials are added to the spray tank, the TEBU 3.6F Foliar Fungicide should be thoroughly dispersed prior to the addition of other materials.

NOTE: Do not tank mix with products containing a prohibition against tank mixing. Follow the most restrictive labeling requirements of any tank mix product.

Compatibility: To determine the compatibility of TEBU 3.6F Foliar Fungicide with other products, the following procedure should be followed: Pour the recommended proportions of the product into a suitable container of water, mix thoroughly and allow to stand at least five minutes. If the combination remains mixed or can be re-mixed readily, the mixture is considered physically compatible.

DISEASE CONTROL IN CROPS

Crop	Diseases	Rate in fl. oz. of Tebu 3.6F Foliar Fungicide/ Acre	Remarks
Asparagus	Rust (<i>Puccinia</i> spp.)	4 – 6	Apply as a foliar spray to the developing ferns after harvest of spears is completed. Apply at the earliest sign of rust pustules or when weather conditions are conducive for rust development. Apply the specified amount (0.11 to 0.17 lb. of active ingredient per acre) in alternation with another effective fungicide. Under conditions of severe rust pressure, use the higher rate. Repeat applications on a 14-day interval as necessary to maintain control of rust. Do not apply to harvestable spears. Do not make more than three foliar applications per season (18 fl. oz./acre or 0.51 lb of active ingredient per acre).

General Comments: Applications may be made using ground or aerial application equipment. A 50 foot spray drift buffer zone is required for all aerial applications. For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with TEBU 3.6F Foliar Fungicide. TEBU 3.6F Foliar Fungicide is a sterol demethylation inhibitor (DMI) fungicide (Group 3). Alternating TEBU 3.6F Foliar Fungicide with other DMI fungicides may lead to resistance.

Restricted-entry interval (REI) = 12 hours

Pre-harvest interval (PHI) = 100 days (California); 180 days (all other states)

Crop	Diseases	Rate in fl. oz. of TEBU 3.6F Foliar Fungicide/ Acre	Remarks
Barley*	Rusts <i>(Puccinia spp)</i>	4	Apply TEBU 3.6F Foliar Fungicide in a minimum of 10 gallons of spray solution per acre by ground or in a minimum of 5 gallons of spray solution per acre by air. A maximum of 4 fl. oz. may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw cut after harvest may be fed or used for bedding. Grazing livestock or feeding of green forage is permitted 6 or more days after the last application of TEBU 3.6F Foliar Fungicide. Barley fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development.
*Not registered for this use in California.	Head Blight <i>(Fusarium spp.)</i>		
Application Timing of TEBU 3.6F Foliar Fungicide for Optimum Control: Rusts: Apply TEBU 3.6F Foliar Fungicide at the earliest sign of rust pustules on foliage. Fusarium head blight: Optimal timing of TEBU 3.6F Foliar Fungicide for Fusarium head blight suppression is when stem heads have fully emerged (Feekes 10.5) on 50% of the plants.			
General Comments: For optimum disease control, the lowest recommended rate of a spray surfactant should be tank-mixed with TEBU 3.6F Foliar Fungicide. TEBU 3.6F Foliar Fungicide must have two to			

6/24

four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBU 3.6F Foliar Fungicide will be resistant to weathering. TEBU 3.6F Foliar Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours

Crop	Diseases	Rate in fl. oz. of TEBU 3.6F Foliar Fungicide/ Acre	Remarks
Beans (fresh & dry, except succulent shelled)	Rust (<i>Uromyces appendiculatus</i>)	4 – 6	Apply TEBU 3.6F Foliar Fungicide in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 14-day intervals, or as necessary to maintain control. Fresh beans: Do not apply more than 24 fl. oz. of TEBU 3.6F Foliar Fungicide per acre per crop season. Dry beans: Do not apply more than 12 fl. oz. of TEBU 3.6F Foliar Fungicide per acre per crop season.

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with TEBU 3.6F Foliar Fungicide. TEBU 3.6F Foliar Fungicide must have two to four hours of drying time on bean foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBU 3.6F Foliar Fungicide will be resistant to weathering. TEBU 3.6F Foliar Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours

Pre-harvest interval (PHI) = 7 days (fresh beans); 14 days (dry beans)

Crop	Diseases	Rate in fl. oz. of TEBU 3.6F Foliar Fungicide/ Acre	Remarks
Corn (sweet corn, field corn, field corn grown for seed, and popcorn)	Rust (<i>Puccinia spp.</i>) Northern Leaf Blight (<i>Helminthosporium turcicum</i>) Southern Leaf Blight (<i>Helminthosporium maydis</i>) Northern Leaf Spot (<i>Helminthosporium carbonum</i>) Gray Leaf Spot (<i>Cercospora zeae-maydis</i>)	4 - 6	Apply TEBU 3.6F Foliar Fungicide in a protective spray schedule or when weather conditions are favorable for disease development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. A maximum of 24 fl. oz. (1.5 pint) of TEBU 3.6F Foliar Fungicide may be applied per acre per crop season.

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with TEBU 3.6F Foliar Fungicide. TEBU 3.6F Foliar Fungicide must have two to four hours of drying time on corn foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBU 3.6F Foliar Fungicide will be resistant to weathering. TEBU 3.6F Foliar Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) for sweet corn = 19 days

Pre-harvest interval (PHI) for sweet corn = 7 days (ears or forage); 49 days (fodder)

Restricted-entry interval (REI) for all corn except sweet corn = 12 hours

Pre-harvest interval (PHI) for field, seed or popcorn = 21 days (forage); 36 days (grain or fodder)

7/24

Crop	Diseases	Rate in fl. oz. of TEBU 3.6F Foliar Fungicide/ CWT	Remarks
Corn (seed treatment*) (Sweet, Field, Popcorn) *Not registered for this use in California.	Soilborne and Seedborne Fusarium	0.071	Apply as a seed treatment using standard slurry or mist-type seed treatment equipment. Uniform application of seed is necessary to ensure seed safety and best disease protection. Seed should be sound and well cured prior to treatment. Product should be diluted with sufficient water to ensure complete seed coverage. Consult a seed treatment specialist regarding slurry rates recommended for the crop to be treated with TEBU 3.6F Foliar Fungicide. The length of control will vary depending on the rate used.
	Soilborne and Seedborne Head Smut	0.27 – 0.54	

General Comments: To meet U.S. Federal Seed Act requirements, all seed treated with TEBU 3.6F Foliar Fungicide must be labeled:
TREATED SEED. DO NOT USE FOR FOOD, FEED, OR OIL PURPOSES. Treated with Tebuconazole.

When using formulations that do not contain dye, to comply with 40 CFR 153.155, all seed treated with an economic poison must be colored with an EPA approved dye such as one of the dyes listed in 40 CFR Section 180.910 or 180.950 to distinguish and prevent subsequent inadvertent use as a food for man or feed for animals.

Crop	Diseases	Rate in fl. oz. of TEBU 3.6F Foliar Fungicide/ Acre	Remarks
Cotton	Southwestern Cotton Rust (<i>Puccinia cacabata</i>)	6 - 8	Apply TEBU 3.6F Foliar Fungicide in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. Do not apply more than 24 fl. oz. of TEBU 3.6F Foliar Fungicide per acre per crop season.

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with TEBU 3.6F Foliar Fungicide. TEBU 3.6F Foliar Fungicide must have two to four hours of drying time on cotton foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBU 3.6F Foliar Fungicide will be resistant to weathering. TEBU 3.6F Foliar Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) =12 hours
Pre-harvest interval (PHI) = 30 days

8/29

Crop	Diseases	Rate in fl. oz. of TEBU 3.6F Foliar Fungicide/ Acre	Remarks
Cucurbit Vegetables Group Chayote Chinese Waxgourd Citron Melon Cucumber Gherkin Edible Gourd (includes Hyotan, Cucuzza, Hechima, and Chinese Okra) <i>Momordica</i> spp. (includes Balsam Apple, Balsam Pear, Bitter Melon and Chinese Cucumber) Muskmelon (includes Cantaloupe, Casaba, Crenshaw Melon, Golden Pershaw Melon, Honeydew Melon, Honey Balls, Mango Melon, Persian Melon, Pineapple Melon, Santa Clause Melon and Snake Melon) Pumpkin Summer Squash (includes Crookneck Squash, Scallop Squash, Straightneck Squash, Vegetable Marrow and Zucchini) Winter Squash (includes Butternut Squash, Calabaza, Hubbard Squash, Acorn Squash and Spaghetti Squash) Watermelon	Powdery Mildew (<i>Sphaerotheca fuliginea</i> / <i>Podosphaera xanthii</i>) (<i>Erysiphe cichoracearum</i>)	4 - 6	Apply specified dosage in a protective spray schedule to foliage and fruit. Repeat applications at 10- to 14-day intervals. Do not apply more than 24 fl. oz. of TEBU 3.6F Foliar Fungicide per acre per crop season.
	Gummy Stem Blight - suppression (<i>Didymella bryonae</i>) (watermelon, squash, pumpkin, and melons only)	8	
<p>General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with TEBU 3.6F Foliar Fungicide. TEBU 3.6F Foliar Fungicide must have two to four hours of drying time for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBU 3.6F Foliar Fungicide will be resistant to weathering. TEBU 3.6F Foliar Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restricted-entry interval (REI) = 12 hours Pre-harvest interval (PHI) = 7 days</p>			

9/24

Crop	Diseases	Rate in fl. oz. of TEBU 3.6F Foliar Fungicide/ Acre	Remarks
Dry Bulb Onion Garlic Great-headed (Elephant) Garlic Welch Onion Shallot	White Rot (<i>Sclerotium cepivorum</i>)	White rot: 20.5 fl. oz. per acre applied in a 4 to 6 inch band over/into each furrow	White Rot: For the control of white rot, make one application in the furrow at the time of planting. The in-furrow application should be made at the rate of 20.5 fl. oz. TEBU 3.6F Foliar Fungicide per acre. Apply the entire per acre rate in a 4 to 6 inch band over/into each furrow. Additional control may be obtained by including two foliar applications at 4 - 6 fl. oz./acre. Rust: For the control of rust, make foliar applications at the rate of 4 - 6 fl. oz./acre per application. Repeat at an interval of 10 - 14 days. Apply TEBU 3.6F Foliar Fungicide in a protective spray schedule or when weather conditions are favorable for rust development. Do not apply more than 32.5 fl. oz. TEBU 3.6F Foliar Fungicide per acre per season if an in-furrow treatment is made. If TEBU 3.6F Foliar Fungicide is not applied as an in-furrow treatment, then do not apply more than 12 fl. oz./acre per season as a foliar spray.
	Rust (<i>Puccinia allii</i> , <i>Puccinia porri</i>)		
	Purple Blotch* (<i>Alternaria porri</i>)	4 - 6	

*Not registered for this use on Garlic in California.

General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest labeled rate of a spray surfactant may be tank-mixed with TEBU 3.6F Foliar Fungicide. TEBU 3.6F Foliar Fungicide must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBU 3.6F Foliar Fungicide will be resistant to weathering. TEBU 3.6F Foliar Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) =12 hours
Pre-harvest interval (PHI) = 7 days

10/24

Crop	Diseases	Rate in fl. oz. of TEBU 3.6F Foliar Fungicide/ Acre	Remarks
Garden Beet roots and tops (leaves)	Cercospora Leaf Spot (<i>Cercospora beticola</i>)	3 - 7.2	Make applications on 14-day intervals. Do not apply more than 28.8 fl. oz./acre per season.
<p>General Comments: For optimum results, use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest labeled rate of a spray surfactant may be tank-mixed with TEBU 3.6F Foliar Fungicide. TEBU 3.6F Foliar Fungicide must have two to four hours of drying time on beet foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBU 3.6F Foliar Fungicide will be resistant to weathering. TEBU 3.6F Foliar Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restricted-entry interval (REI) = 12 hours Pre-harvest interval (PHI) = 7 days</p>			

Crop	Diseases	Rate in fl. oz. of TEBU 3.6 F Foliar Fungicide/ Acre	Remarks
Grasses Grown for seed	Rusts (<i>Puccinia</i> spp.)	4 to 8 fl. oz. per acre	Apply the specified rate of TEBU 3.6F Foliar Fungicide as soon as weather conditions are favorable for rust development or when first rust pustules are present. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl. oz./A and apply at shorter spray intervals.
	Powdery mildew	4 to 8 fl. oz. per acre	Apply the specified rate of TEBU 3.6F Foliar Fungicide when powdery mildew first appears on the leaves. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl. oz./A and apply at shorter spray intervals.
<p>General Information: Apply the specified rate of TEBU 3.6F Foliar Fungicide in a minimum of 20 gallons of water per acre with aircraft. Thorough coverage is important for optimum disease control. For optimum benefit, the lowest recommended rate of a spray surfactant should be more than 16 fl. oz. (0.45 lb a.i.) per acre per crop season. Chaff, screenings and straw from treated areas may be used for feed purposes; however, do not forage, cut green crop, or use seed for feed purposes. Regrowth may be grazed starting 17 days after last application.</p> <p>Preharvest interval: 4 days. Restricted-entry interval (REI) = 12 hours</p>			

Crop	Diseases	Rate in fl. oz. of TEBU 3.6F Foliar Fungicide/ Acre	Remarks
Green Onion Green Eschalots Green Shallots Japanese Bunching	White Rot (<i>Sclerotium cepivorum</i>) (suppression only) Rust	4 - 6	For the control of diseases, make foliar applications using an interval of 10 - 14 days. Apply TEBU 3.6F Foliar Fungicide in a protective spray schedule or when weather conditions are favorable for rust development. Do not apply more

1/24

Crop	Diseases	Rate in fl. oz. of TEBU 3.6F Foliar Fungicide/ Acre	Remarks
Onion Leek Scallion Spring Onion	(<i>Puccinia allii</i> , <i>Puccinia porri</i>) Purple Blotch (<i>Alternaria porri</i>)		than 24 fl. oz./acre per season.

General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with TEBU 3.6F Foliar Fungicide. TEBU 3.6F Foliar Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBU 3.6F Foliar Fungicide will be resistant to weathering. TEBU 3.6F Foliar Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours

Pre-harvest interval (PHI) = 7 days

Crop	Diseases	Rate in fl. oz. of TEBU 3.6F Foliar Fungicide/ Acre	Remarks
Hops	Powdery Mildew (<i>Sphaerotheca humuli</i> / <i>Sphaerotheca maculans</i>)	4 - 8	Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 10- to 14-day intervals. Do not apply more than 32 fl. oz. of TEBU 3.6F Foliar Fungicide per acre per crop season. Increase the spray volume and the application rate as vine growth increases during the season.

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with TEBU 3.6F Foliar Fungicide. TEBU 3.6F Foliar Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBU 3.6F Foliar Fungicide will be resistant to weathering. TEBU 3.6F Foliar Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours

Pre-harvest interval (PHI) = 14 days

12/28

Crop	Diseases	Rate in fl. oz. of TEBU 3.6F Foliar Fungicide/ Acre	Remarks
Leafy Brassica Greens Broccoli Raab Chinese Cabbage (Bok Choy) Collards Kale Mizuna Mustard Greens Mustard Spinach Rape Greens Turnip Greens	Cercospora Leaf Spot <i>(Cercospora brassicicola)</i> Powdery Mildew <i>(Erysiphe cruciferarum)</i> Alternaria Leaf Spot <i>(Alternaria brassicicola)</i>	3 - 4	Do not apply more than 16 fl. oz./acre per season. Do not apply more often than once every 10 days.
<p>General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with TEBU 3.6F Foliar Fungicide. TEBU 3.6F Foliar Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBU 3.6F Foliar Fungicide will be resistant to weathering. TEBU 3.6F Foliar Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restriction: Application to turnip greens is limited to east of the Rockies.</p> <p>Restricted-entry interval (REI) = 12 hours</p> <p>Pre-harvest interval (PHI) = 7 days</p>			

Crop	Diseases	Rate in fl. oz. of TEBU 3.6F Foliar Fungicide/ Acre	Remarks
Lychee	Anthracnose <i>(Colletotrichum gloeosporioides)</i>	4 - 6	Begin first application of TEBU 3.6F Foliar Fungicide as panicle emerges. Spray up to 6 fl. oz./acre every 10 days thereafter for a total of 8 sprayings. Apply specified dosage in a minimum of 50 gallons of spray solution per acre by ground only. Do not apply more than 48 fl. oz./acre per season.
<p>General Comments: For optimum disease control, the lowest labeled rate of a non-ionic spray surfactant should be tank-mixed with TEBU 3.6F Foliar Fungicide. TEBU 3.6F Foliar Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBU 3.6F Foliar Fungicide will be resistant to weathering. TEBU 3.6F Foliar Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restricted-entry interval (REI) = 2 days</p> <p>Pre-harvest interval (PHI) = 0 (zero) days</p>			

Crop	Diseases	Rate in fl. oz. of TEBU 3.6F Foliar Fungicide/ Acre	Remarks
Okra	Cercospora Leaf Spot (<i>Cercospora</i> spp.)	4 - 6	Apply specified dosage of TEBU 3.6F Foliar Fungicide in a preventative spray program. Use the highest rate when disease conditions are favorable and in areas where high disease pressure is expected. Applications may be repeated at 14-day intervals in order to maintain control of the disease. Apply specified dosage as a foliar spray in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. Do not apply more than 24 fl. oz./acre per season.
<p>General Comments: For optimum disease control, the lowest labeled rate of spray surfactant should be tank-mixed with TEBU 3.6F Foliar Fungicide. TEBU 3.6F Foliar Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBU 3.6F Foliar Fungicide will be resistant to weathering. TEBU 3.6F Foliar Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restricted-entry interval (REI) = 12 hours Pre-harvest interval (PHI) = 3 days</p>			

14/24

Crop	Diseases	Rate in fl. oz. of TEBU 3.6F Foliar Fungicide/ Acre	Remarks
Peanut	SOILBORNE: Cyindrocladium Black Rot (suppression) Rhizoctonia Limb Rot Rhizoctonia Pod Rot (Virginia and North Carolina only) Sclerotium Stem and Pod Rot (White Mold, Southern Blight, Southern Stem Rot) FOLIAR: Early Leaf Spot Late Leaf Spot Leaf Rust Pepper Spot (<i>Leptosphaerulina</i>) Web Blotch (<i>Phoma</i>)	7.2 fl. oz. per acre	FOUR-APPLICATION SPRAY PROGRAM: Apply the specified rate in a preventive spray schedule. See table below for proper timing of applications. Applications of chlorothalonil should be made prior to and following applications of TEBU 3.6F Foliar Fungicide to discourage development of resistant strains of fungi. For optimum control of foliar diseases such as leaf rust, web blotch, and pepper spot, the lowest label recommended rate of a spray surfactant should be tank-mixed with TEBU 3.6F Foliar Fungicide. LEAF SPOT ADVISORY SCHEDULE: For control of soilborne diseases in an advisory schedule, apply TEBU 3.6F Foliar Fungicide in the first advisory spray in July and continue TEBU 3.6F Foliar Fungicide applications at 14-day intervals. When applying TEBU 3.6F Foliar Fungicide after August 15, tank mix with chlorothalonil for resistance management purposes. General Information: For optimum control of the specified soilborne diseases, four consecutive applications of TEBU 3.6F Foliar Fungicide must be made at 14-day intervals. A maximum of 28.8 fl. oz. (0.81 lb a.i.) of TEBU 3.6F Foliar Fungicide may be applied per crop season. TEBU 3.6F Foliar Fungicide may be applied up to 14 days before harvest. Do not feed hay or threshings or allow livestock to graze in treated areas. TEBU 3.6F Foliar Fungicide is a sterol demethylation inhibitor (DMI) fungicide. Chlorothalonil may be tank mixed at the rate of 12 ounces of active ingredient with TEBU 3.6F Foliar Fungicide as a leaf spot resistance management strategy. A spray surfactant is not necessary when TEBU 3.6F Foliar Fungicide is tank mixed with chlorothalonil. Mixing or alternating TEBU 3.6F Foliar Fungicide with other DMI fungicides may lead to resistance. TEBU 3.6F Foliar Fungicide must be carried by rainfall or irrigation into the root and pod zone for control of root and pod rots caused by <i>Sclerotium rolfsii</i> and <i>Rhizoctonia solani</i> . Drought conditions will decrease the effectiveness of TEBU 3.6F Foliar Fungicide against the root and pod rots. Use TEBU 3.6F Foliar Fungicide in conjunction with cultural practices that are known to reduce the severity of soilborne diseases, such as proper crop rotation practices. Restricted-entry interval (REI) = 12 hours
Application Timing of TEBU 3.6F Foliar Fungicide for Optimum Control of White Mold and Rhizoctonia Limb and Pod Rot			
Spraying Program		TEBU 3.6F Foliar Fungicide Application No.	Chlorothalonil Application No.
7 applications		3,4,5 and 6	1,2, and 7

Crop	Diseases	Rate in fl. oz. of TEBU 3.6F Foliar Fungicide/ Acre	Remarks
Pecan	Brown Leaf Spot (<i>Sirosporium diffusum</i>) Downy Spot (<i>Mycosphaerella caryigena</i>) Liver Spot (<i>Gnomonia caryae</i>) Scab (<i>Cladosporium caryigenum</i>) Vein Spot (<i>Gnomonia nerviseda</i>) Zonate Leaf Spot (<i>Grovesinia pyramidalis</i>)	4 - 8	<p>Apply TEBU 3.6F Foliar Fungicide in a preventative spray schedule beginning at early bud break (young leaves unfolding), and continue applications at 10- to 14-day intervals through the pollination period. TEBU 3.6F Foliar Fungicide should be applied at 4 fl. oz./acre in a tank-mix with the recommended rate of Super-Tin® in cover sprays. Follow label directions for the use of Super-Tin®. Do not add a surfactant to the spray solution when tank-mixing TEBU 3.6F Foliar Fungicide with Super-Tin®.</p> <p>Apply TEBU 3.6F Foliar Fungicide in a spray volume of 15 or more gallons per acre by air or 50 or more gallons per acre by ground. Apply 7 - 8 fl. oz./acre of TEBU 3.6F Foliar Fungicide to full-size mature trees, and 4 - 6 fl. oz./acre to smaller trees. Apply the high rate to varieties that are highly susceptible to the indicated diseases, or when severe disease conditions exist. The lowest labeled rate of a surfactant may be added to the spray solution for optimum control of the indicated diseases. Do not apply after shucks begin to split. Do not apply more than 32 fl. oz./acre per crop season Do not cut cover crops in treated areas for feed or allow livestock to graze treated areas.</p>

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with TEBU 3.6F Foliar Fungicide. TEBU 3.6F Foliar Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBU 3.6F Foliar Fungicide will be resistant to weathering. TEBU 3.6F Foliar Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). It may be applied in a tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy.

Restricted-entry interval (REI) = 12 hours

Pre-harvest interval (PHI) = do not apply after shucks begin to split

10/24

Crop	Diseases	Rate in fl. oz. of TEBU 3.6F Foliar Fungicide/ Acre	Remarks
Soybean	Rust (<i>Phakopsora pachyrhizi</i>) Powdery Mildew (<i>Microsphaera diffusa</i>)	3 -4	Apply TEBU 3.6F Foliar Fungicide as a broadcast foliar spray as a preventative spray or at first visible symptoms of disease. Repeat applications on a 10- to 14-day spray interval if environmental conditions are favorable for continued disease development. Use of the higher rates and shorter spray intervals are recommended when disease pressure is severe. The lowest label recommended rate of a spray surfactant may be tank-mixed with TEBU 3.6F Foliar Fungicide. TEBU 3.6F Foliar Fungicide should be applied in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons per acre by aircraft spray equipment. Do not apply more than 12 fl. oz./acre per use season. Do not make more than three applications per season.
Restricted-entry interval (REI) = 12 hours Pre-harvest interval (PHI) = 21 days			

Crop	Diseases	Rate in fl. oz. of TEBU 3.6F Foliar Fungicide/ Acre	Remarks
Sunflower	Rust (<i>Puccinia helianthi</i>)	4 - 6	Apply specified dosage of TEBU 3.6F Foliar Fungicide at the earliest sign of infection (rust pustules developing) or when weather conditions are favorable for rust development. Apply higher rate to highly susceptible varieties and/or under severe disease conditions. Application may be repeated at 14 days if necessary to maintain control of the disease. Apply specified dosage in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. Do not apply more than 16 fl. oz./acre per season.
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with TEBU 3.6F Foliar Fungicide. Contact your state extension service for a list of approved surfactants. TEBU 3.6F Foliar Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBU 3.6F Foliar Fungicide will be resistant to weathering. TEBU 3.6F Foliar Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).			
Restricted-entry interval (REI) = 12 hours Pre-harvest interval (PHI) = 50 days			

Crop	Diseases	Rate in fl. oz. of TEBU 3.6F Foliar Fungicide/ Acre	Remarks
Turnip	Cercospora Leaf Spot (<i>Cercospora brassicicola</i>)	4 – 7.2	<p>Application is limited to east of the Rockies.</p> <p>Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 12- to 14-day intervals. Do not apply more than 28.8 fl. oz./acre per crop season.</p>
<p>General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with TEBU 3.6F Foliar Fungicide. TEBU 3.6F Foliar Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBU 3.6F Foliar Fungicide will be resistant to weathering. TEBU 3.6F Foliar Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restricted-entry interval (REI) = 12 hours Pre-harvest interval (PHI) = 7 days</p>			

Crop	Diseases	Rate in fl. oz. of TEBU 3.6F Foliar Fungicide/ Acre	Remarks
Wheat* *Not registered for this use in California.	Rusts leaf, stem, and stripe (<i>Puccinia</i> spp.) Head blight or scab (<i>Fusarium</i> spp.) — Suppression	4.0	<p>Notes: Wheat fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. A maximum of 4 fl. oz. of TEBU 3.6F Foliar Fungicide may be applied per acre per crop season. Straw may be fed or used for bedding. Do not allow livestock to graze or feed green forage to livestock prior to 6 days after treatment with TEBU 3.6F Foliar Fungicide. Apply TEBU 3.6F Foliar Fungicide in a minimum of 10 gallons of spray solution per acre by ground, or in a minimum of 5 gallons of spray solution per acre by air.</p> <p>Application timing: Rusts: Apply TEBU 3.6F Foliar Fungicide at the earliest sign of rust pustules on foliage. Fusarium head blight: Optimal timing of TEBU 3.6F Foliar Fungicide for Fusarium head blight suppression is the beginning of flowering on main stem heads (Feekes 10.51).</p>
<p>General Comments: For optimum disease control, the lowest recommended rate of a spray surfactant should be tank-mixed with TEBU 3.6F Foliar Fungicide. TEBU 3.6F Foliar Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBU 3.6F Foliar Fungicide will be resistant to weathering. TEBU 3.6F Foliar Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restricted-entry interval (REI) = 12 hours Pre-harvest interval (PHI) = 30 days</p>			

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.

Apply only during alternate years in fields adjacent to aquatic areas listed above.

Do not apply by ground or air within 100 feet of aquatic areas listed above.

Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip.

Spray Drift Management: For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wingspan or rotor diameter.

Use the largest droplet size consistent with pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure. Apply in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment.

Spray should be released at the lowest possible height consistent with good pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

Make aerial or ground applications when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Low humidity and high temperatures increase the evaporation rate spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.

Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

ROTATIONAL CROPS

Treated areas may be replanted with any crop specified on this label as soon as practical after last application. Any crop not specified on this label may be planted into treated areas 120 days after last applications.

DISEASE CONTROL IN GOLF COURSE TURF, FIELD, NURSERY AND CONTAINER ORNAMENTALS AND COMMERCIAL AND RESIDENTIAL LANDSCAPES

Chemigation: Do not apply this product through any type of irrigation system.

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.

- Do not apply within 100 feet of aquatic areas listed above.
- Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetation filter strip.
- See Spray Drift Management section for further information.

Spray Drift Management

Make ground application when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

Risks of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperatures.

Do not make ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near

the ground surface.

Spray Volume: For best results, apply this product in 66-132 gallons of water per acre for turf using ground based equipment. For ornamentals, 50-300 gallons of finished spray per acre are recommended depending upon equipment, plant species and plant growth stage at time of application. For the most effective results, equipment calibration should be checked regularly. When using lower spray volumes, be sure to maintain uniform application and full crop coverage so as to ensure effective control. Increase spray volume to ensure proper application, if required.

Compatibility Test for Mix Components:

Before mixing components, always perform a compatibility jar test. For 66 gallons per acre spray volume, use 5 cups of water in a clear, clean mixing jar. For other spray volumes adjust accordingly. Only use water from the intended source at the source temperature. Add components in the sequence indicated below in Mixing Order using 3 teaspoons for each pound of dry product or 1 1/2 teaspoons for each pint of liquid product of recommended label rate per acre. Always cap the jar and invert 10 cycles between component additions. When the components have all been added to the jar and fully mixed, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent and use the compatibility agent as directed on its label.

Mixing:

Continuous agitation is required during mixing. When mixing this product and water, use the specified application rates as listed for each crop on this label. Before combining any other substances with the mixture, ensure that this product is completely dispersed in the mixture.

Recommended Mixing Procedure:

1. Water. Add three-quarters of the required volume to a thoroughly clean sprayer tank
2. Agitation. Start agitation and maintain constant agitation throughout mixing and application.
3. Inductor. If an inductor is used, rinse it thoroughly after each component has been added.
4. Products in PVA Bags. Place any product contained in water soluble PVA bags into the mixing tank. Wait until all water soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5. Water Dispersible Products. Including dry flowables (DF), wettable powders (WP), suspension concentrates (SC) or suspo-emulsions (SE).
6. Water-soluble products.
7. Emulsifiable concentrates (such as oil concentrates when applicable).
8. Water soluble additives (such as AMS or UAN when applicable).
9. Remaining quantity of water.

Resistance Management Information

The active ingredient in this product is a member of the DMI (Demethylation Inhibitor) fungicide group (FRAC grouping 3) and exhibits no known cross-resistance to products with the same mode of action when used repeatedly in the same location or in successive years as the primary method of control for targeted diseases. Because the speed and scope of resistant population development cannot be predicted, the use of this product should conform to resistance management strategies established for the crop and use area. Such strategies may include the rotation and/or tank mixing with products utilizing different modes of action or limiting the number of applications per season. Contact your local university or extension specialist and/or manufacturer for fungicide resistance management recommendations.

DISEASE CONTROL IN GOLF COURSE TURF

TURF USE RESTRICTIONS AND PRECAUTIONS

- For use on golf course turf only.

20/24

- Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (i.e., elementary, middle and high schools), campgrounds, churches, and theme parks.
- Not for homeowner use.
- Not for use on turf being grown for sale or commercial use as sod.
- Do not use clippings for animal feed.
- Do not exceed 3.6 fl. oz of this product per 1000 sq. ft. per year.
- Do not apply more than 6 applications per year.

Product Information

For use on all Golf turf applications of cool season and warm season grasses (such as Bentgrasses, Bluegrasses, Fescues, Ryegrasses, St. Augustine grasses, and Zoysia) or their mixtures. This product is not phytotoxic to any of the above mentioned grasses when used in accordance with the label.

Note: Bermudagrass can be sensitive to this product under certain conditions. Do not apply consecutive applications during or just after dormancy break. Avoid applications when the temperatures are expected to exceed 85 degrees F.

Use this product for the prevention and control of the diseases mentioned in the table below. Begin applications when conditions favor disease development and repeat applications as long as these conditions persist. Preventative treatments can be applied using 28 day intervals as indicated. When treating golf greens, always treat aprons and approaches. Spray uniformly over the area to be treated with properly calibrated equipment.

Apply the specified amount of this product in sufficient water for thorough coverage. A volume of 66 – 132 gallons per acre (1.5-3.0 gallons per 1,000 sq ft) is recommended. Apply using properly calibrated low volume, hand held, mechanical or motorized ground broadcast equipment. Application to small areas may be made with low-pressure hand wand or backpack equipment. Maintain constant agitation during application.

Depending on the disease, water this product into the crown and active root zone for best results. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. For best results, use spray mixture the same day it is prepared.

Golf Course Turf Disease Control

DISEASE	RATE of AmTide TEBU 3.6F Foliar Fungicide (Fl. Oz./1000 Sq Ft)	Notes
Dollar Spot (<i>Sclerotinia homoeocarpa</i>) Copper Spot (<i>Gloeocercospora sorghi</i>) Powdery Mildew (<i>Erysiphe graminis</i>) Corticium Red Thread (<i>Laetisaria fuciformis</i>) Rusts (<i>Puccinia</i> spp.)	0.6	For prevention, begin applications when conditions are favorable for disease development. Do not make two consecutive applications of this product. Alternate with another fungicide with a different mode of action. A second application may be made after 28 days.
Brown Patch/Rhizoctonia Blight, Large Patch (<i>Rhizoctonia solani</i>) Brown Ring Patch (<i>R. circinata</i>)	0.6	For prevention, begin applications when conditions are favorable for disease development. Do not make two consecutive applications of this product.
Anthracnose-Basal and Foliar (<i>Colletotrichum cereale</i>) Red Thread (<i>Laetisaria fuciformis</i>) Pink Patch (<i>Limonomyces rosipellis</i>)	0.6	Alternate with another fungicide with a different mode of action. A second application may be made after 28 days.
Bermuda Grass decline	0.6	Immediately after fungicide is applied,

21/24

DISEASE	RATE of AmTide TEBU 3.6F Foliar Fungicide (Fl. Oz./1000 Sq Ft)	Notes
(<i>Gaeumannomyces graminis</i> var. <i>graminis</i>)		irrigate the area with sufficient water to move the active ingredient down into the crown and root zone of the turf. The amount of water is dependent on the depth of the root zone. For prevention, begin applications two or four weeks prior to the historical appearance of disease symptoms. Initiate cultural control practices at the same time the fungicide is applied. Refer to your local County Extension Service for this information. Apply subsequent applications at 28 day intervals.
Take All Patch (<i>Gaeumannomyces graminis</i>)	0.6	For prevention, apply in the fall when soil temperature reaches 55-65° F and again in the spring under similar soil temperature conditions. Applications in both fall and spring may be necessary. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
Gray Leaf Spot (<i>Pyricularia grisea</i>)	0.6	Apply when conditions are favorable for disease development at 28 day intervals. This product can be tank mixed with a registered contact fungicide at the label rate.
Stipe Smut (<i>Ustilago striiformis</i>)	0.6	Make a single application to historical disease areas in spring as grass growth begins.
Spring Dead Spot (<i>Leptosphaeria korrea</i> , <i>L. narmari</i> , <i>Ophiosphaerella herpotricha</i> , <i>Gaeumannomyces graminis</i>) Necrotic Ring Spot (<i>Leptosphaeria korrea</i>)	0.6	For prevention, apply in fall when soil temperature reaches 65° F and again in spring under similar soil temp conditions or after dormancy break. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
Fusarium Patch (<i>Fusarium roseum</i>)	0.6	Apply first application in mid-June or 28 days prior to time this blight normally becomes evident. Make applications at no less than 28 day intervals.
Summer Patch (<i>Magnaporthe poae</i>)	0.6	Apply beginning in the spring. Do not make two consecutive applications of this product. Alternate with another fungicide with a different mode of action. Second and third applications may be made at 28 day intervals. See local university recommendations for suggested timing. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
Zoysia Patch, Large Patch of zoysia (<i>Rhizoctonia solani</i>)	0.6	Make first application in early fall (mid-September to mid-October) prior to development of disease symptoms. A

DISEASE	RATE of AmTide TEBU 3.6F Foliar Fungicide (Fl. Oz./1000 Sq Ft)	Notes
		second application in early spring may be necessary in areas where disease pressure is known to be heavy.
Gray Snow Mold/Typhula Blight (<i>Typhula incarnate</i>) Pink Snow Mold/Microdochium Patch (<i>Microdochium nivalis</i>)	0.6	Apply in the fall, before anticipated turf dormancy and before the first snow cover. If turf breaks dormancy during winter months, a second application may be made. Do not apply over snow cover, or when turf is dormant. It is recommended that this product be tank-mixed with other registered snow mold products for best season long results.
NOTE: Apply the specified amount of this product in 1.5 to 3.0 gallons of water per 1000 sq. ft. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. Do not use clippings for animal feed. Do not exceed 3.6 fl. oz of this product per 1000 sq. ft. per year. Do not exceed 6 applications per year.		

DISEASE CONTROL IN FIELD, NURSERY AND CONTAINER ORNAMENTALS AND COMMERCIAL and RESIDENTIAL LANDSCAPES

ORNAMENTAL USE RESTRICTIONS AND PRECAUTIONS

- For use on ornamental plants only, not for woodlands or forest management.
- Not for homeowner use.
- Do not apply more than 0.94 gallons (120 fl oz) of this product (equal to 3.38 lbs of tebuconazole) per acre per year.
- Do not make more than 4 applications per year at highest rate.
- Do not apply to bearing fruit trees or vegetables.

Use this product in a preventative and curative disease control program for the listed plant types and disease in the table below. Optimum disease management is obtained when this product is used in conjunction with sound disease management practices.

Apply material with properly calibrated hand held, mechanical or motorized spray equipment. Begin applications when disease first appears and repeat at 14-21 day intervals during the growing season. Use the shortest interval when conditions are unusually favorable for the development of disease. For hand held, mechanical, or motorized applications, mix as directed below and apply as a full coverage spray to drip for the prevention and control of the diseases listed below. Choose a finished spray volume appropriate for the size of the plants and amount of foliage, which will provide thorough coverage throughout the canopy. Allow sprays to dry before overhead irrigation is applied.

Apply this product at rates of 4 – 10 fl oz per acre in 100 gallons of water. Spray volume may range from 50 up to 300 gallons of finished spray per acre depending upon equipment, plant species and plant growth stage at time of application.

Note: The "Directions for Use" of this product reflect the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied to any plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results. This product is not recommended for use on African Violets, Begonias, Boston Fern, and Geraniums.

23/24

ORNAMENTALS DISEASE CONTROL

PLANTS	DISEASE	APPLICATION	
		To Prevent Disease	To Treat Existing Disease
Roses	Black Spot Powdery Mildew Rust	Apply every 14-21 days during the growing season, starting when leaves first appear.	Apply every 14 days for a total of 3 applications beginning at the first sign of disease.
Flowers	Leaf Spot Powdery Mildew Rust Southern Blight	Apply at least 3 times per year, 14-21 days apart, beginning with Spring bud break.	
Crabapples (Ornamental), Dogwoods and Other Landscape (Ornamental) Trees	Anthrachnose Leaf Spot Powdery Mildew Rust Scab	Rotation or Tank mixing with barrier protectant fungicides is recommended for resistance management.	
Azaleas, Camellias, Rhododendrons and Other Landscape (Ornamental) Shrubs	Anthrachnose Black Spot Leaf Spot Petal Blight	Petal Blight – Apply 2-3 times per week into the flowers as they open and develop color.	
Ground Covers and Vines	Powdery Mildew Rust Southern Blight		
HOW MUCH TO USE FOR SMALL PLANTINGS: ADD 1 TEASPOON TO 2.5 GALLONS OF WATER.			

Pump Style Sprayers

1. Add the appropriate amounts of concentrate and water to the sprayer tank.
2. Close the sprayer, shake well and pressurize.
3. Adjust nozzle to a coarse spray pattern and apply.
4. Occasionally re-pressurize the sprayer if needed to maintain a good spray pattern.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, secure, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If container is leaking invert to prevent leakage. If the container is leaking or material is spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticide below. In spill or leak incidents, keep unauthorized people away.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND DISCLAIMER

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of AmTide, LLC. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: AMTIDE, LLC MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. NO AGENT OF AMTIDE, LLC IS AUTHORIZED TO MAKE ANY WARRANTIES BEYOND THOSE CONTAINED HEREIN OR TO MODIFY THE WARRANTIES CONTAINED HEREIN. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, AMTIDE, LLC DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT AMTIDE, LLC'S ELECTION, THE REPLACEMENT OF PRODUCT.

[EPA approval date]