



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

May 10, 2021

Michael Kellogg
Agent for Ragan and Massey, Inc
c/o Pyxis Regulatory Consulting Inc.
4110 136th Street Ct NW
Gig Harbor, WA 98332

Subject: Label Amendment – Clarify use directions and add optional marketing claims
Product Name: RM43 Total Vegetation Control
EPA Registration Number: 84009-3
Application Date: August 20, 2018
Decision Number: 546883

Dear Michael Kellogg:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.


Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2
EPA Reg. No. 84009-3
Decision No. 546883

with FIFRA section 6. If you have any questions, please contact Lydia Crawford by phone at 703-347-0622, or via email at Crawford.Lydia@epa.gov.

Sincerely,



Emily Schmid, Product Manager 25
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

Note to Reviewer - Brackets [] indicate optional text

GLYPHOSATE	GROUP	9	HERBICIDE
IMAZAPYR	GROUP	2	HERBICIDE

RM43™ Total Vegetation Control

[Alternate Brand Name: HDX Vegetation Killer and Preventer]

[Contains 43.68% Glyphosate]

[With Weed Preventer]

ACTIVE INGREDIENTS:

Glyphosate, isopropylamine salt* 43.68%

Imazapyr, isopropylamine salt** 0.78%

OTHER INGREDIENTS: 55.54%

TOTAL: 100.00%

ACID EQUIVALENT:

*Glyphosate 3.158 lbs./gal

**Imazapyr 0.062 lbs./gal

KEEP OUT OF REACH OF CHILDREN

DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 seven days a week, 6:30 am to 4:30 pm Pacific Time or your poison control center at 1-800-222-1222.	
NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.	

[See Inside Booklet for Complete Precautionary Statements, First Aid, Directions for Use & Storage and Disposal.]

[See back booklet for complete Precautionary Statements, Directions for Use and Storage and Disposal.]


EPA Reg. No. 84009-3

EPA Est. No.

Manufactured For:

Ragan & Massey, Inc.
101 Ponchatoula Parkway
Ponchatoula, LA 70454

Net Contents:



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER - PELIGRO**

Corrosive. Causes irreversible eye damage. Harmful if swallowed. Harmful if absorbed through skin. Do not get in eyes or on clothing. Avoid contact with skin.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, loaders, applicators, and other handlers must wear:

- Protective eyewear (goggles, face shield, or safety glasses)
- Long-sleeved shirt and long pants,
- Shoes plus socks,
- Waterproof gloves

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands with plenty of soap and water before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product.
- Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to plants. Drift and run-off may be hazardous to plants in water adjacent to treated areas. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. See Directions for Use for additional precautions and requirements.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read the entire label before using this product. Use strictly in accordance with label precautionary statements and directions.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not enter or allow others to enter treated areas until sprays have dried.

Do not use on feed or food crops.

Keep people and pets out of the area during application.

Applicators are required to use a nozzle height below 4 feet above the ground or plant canopy and coarse or coarser droplet size (ASABE S572) or, if specifically using a spinning atomizer nozzle, applicators are required to use a volume mean diameter (VMD) of 385 microns or greater. Applications with wind speeds greater than 10 mph are prohibited. Applications into temperature inversions are prohibited.

To delay herbicide resistance, take one or more of the following steps:

- Avoid the consecutive use of this product or other target site of action Group 9/Group 2 herbicides that might have a similar target site of action, on the same weed species.
- Use tank mixtures or premixes with herbicides from different target site of action Groups as long as the involved products are all registered for the same use, have different sites of action and are both effective at the tank mix or prepack rate on the weed(s) of concern.
- Base herbicide use on a comprehensive Integrated Pest Management (IPM) program.
- Scout fields prior to application to identify the weed species present and their growth state to determine if the intended application will be effective.
- Scout fields after application to verify that the treatment was effective and to monitor weed populations for early signs of resistance development.
- Contact your local extension specialist, certified crop advisors and/or manufacturer for herbicide resistance management and/or integrated weed management for specific crops and resistant weed biotypes.

Suspected herbicide-resistant weeds may be identified by these indicators:

- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
- A spreading patch of non-controlled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.

Report any incidence of non-performance of this product against a particular weed species to your Ragan and Massey, Inc. retailer, representative or call 800-264-5281. If resistance is suspected, treat weed escapes with an herbicide having a different mechanism of action and/or use non-chemical means to remove escapes, as practical, with the goal of preventing further seed production.

WHERE TO USE

Use on non-crops areas: Industrial sites, Petroleum tank farms, Lumberyards, Manufacturing Sites, Storage Areas, Airports, Roadsides, Curbs, Sidewalks, Driveways, Parking areas, Brick walks, Gravel Paths, Patios, Vacant lots, Paved areas (including crack and crevice), Warehouse areas, Urban areas, Recreational areas, Play areas, Tennis courts, Fencerows, Along fences of paddocks, corrals, feedlots, and livestock holding areas, Hedgerows, Dry Ditchbanks, Around domestic dwellings, Around farm buildings and barns, Agricultural uncultivated areas, Shelter beds, Farm yards, Farm premises, Outdoor buildings, structures, and shelters including beef barns, hog barns, dairy barns, stables, dog kennels and poultry houses, and for establishment and maintenance of wildlife openings

Only apply to locations where no vegetation growth is desired for up to one year.

WHEN TO USE

Apply to bare ground or when weeds are present. If weeds are taller than 6 inches, mow or cut down weeds and remove debris before treatment, and allow for 2 weeks of new growth. If woody vines or brush is mowed or cut, wait until full leaf formation before treatment.

To prevent growth of weeds under asphalt driveways, play areas, tennis courts and parking areas, apply this product before laying surfacing materials.

Treat area uniformly to maximize results.

Re-treatment every 6 to 8 months may be necessary in areas of heavy rainfall.

RESTRICTIONS

- Do not apply to or allow product to drift onto desirable plants.
- Only apply to areas where no vegetation is desired for one year.
- Do not apply near bodies of water including, fish pools, ponds, lakes or streams.
- Do not apply over the root systems of desirable plants. For trees and shrubs do not apply closer than twice the distance from the trunk to the drip line as roots may be within this area.
- Do not apply to slopes in landscape as movement on soil surfaces may damage desirable plants down slope.
- Do not apply next to a fence if desirable vegetation is growing on the other side, or if future planting is intended.
- Do not use before planting lawns, fruits, vegetables, flowers, or other plants because this product remains active in soil for up to one year.

- Do not use on desirable lawns or turf.
- Do not apply when wind speeds are greater than 10 mph.
- Do not apply to powdery, dry or light sandy soils when the soil may be blown by wind. If necessary, soil surface must be settled by rain or irrigation.

HOW TO APPLY

Apply on a sunny day with temperatures above 60°F. Avoid product runoff to nearby desirable plants. In soils with heavy clay, weed control may exceed one year.

Total Vegetation Control (Bareground)

Use hand operated spray equipment such as backpack sprayers, pump-up sprayers or sprinkling cans, or ATV/Tractor mounted sprayers.

Mix 7.4 fl. oz. (0.46 pints) of this product with water to produce 1 to 10 gallons of spray solution. Uniformly apply spray mixture to 1,000 square feet where total vegetation control and bareground is desired. Optimum performance will be achieved from applications to actively growing weeds in early summer.

Maximum application rate of this product is 2.5 gallons per acre per year (7.4 fl. oz. per 1,000 sq. ft. per year).

Spot Control of Brush, Vines, and Weeds

Use hand operated spray equipment such as backpack sprayers, pump-up sprayers or sprinkling cans, or ATV/Tractor mounted sprayers. Mix 6 fl. oz. of this product per 1 gallon of spray solution. Thoroughly and uniformly spray foliage of woody plants to wet. For optimum performance:

- Apply to actively growing brush, vines and weeds.
- Woody brush and vines are controlled better in late summer and fall after fruit formation.
- If woody vines and brush are mowed or cut, wait until full leaf formation before treatment.
- Do not cut or mow brush, vines or perennial weeds for 2 months following treatment.

Maximum application rate of this product is 2.5 gallons per acre per year (7.4 fl. oz. per 1,000 sq. ft. per year).

Weeds Controlled

GRASS/NUTSEDGE WEED SPECIES CONTROLLED			
Bahiagrass <i>Paspalum notatum</i>	Cogongrass <i>Imperata cylindrical</i>	Muhly, wirestem <i>Muhlenbergia frondosa</i>	Quackgrass <i>Agropyron repens</i>
Bentgrass <i>Agrostis</i> spp.	Dallisgrass <i>Paspalum dilatatum</i>	Napier grass <i>Pennisetum purpureum</i>	Reed, giant <i>Arundo donax</i>
Bermudagrass <i>Cynodon dactylon</i>	Fescues <i>Festuca</i> spp.	Nutsedge; purple, yellow <i>Cyperus rotundus</i> <i>Cyperus esculentus</i>	Ryegrass, perennial <i>Lolium perenne</i>
Bluegrass, Kentucky <i>Poa</i> spp.	Fescue, tall <i>Festuca arundinacea</i>	Orchardgrass <i>Dactylis glomerata</i>	Timothy <i>Phleum pratense</i>
Brome <i>Bromus</i> spp.	Guineagrass <i>Panicum maximum</i>	Pampas grass <i>Cortaderia</i> spp.	Torpedograss* <i>Panicum repens</i>
Bromegrass, smooth <i>Bromus inermis</i>	Johnsongrass <i>Sorghum halepense</i>	Panicum <i>Panicum</i> spp.	Vaseygrass <i>Paspalum urvillei</i>
Canarygrass, reed <i>Phalaris arundinacea</i>	Kikuyugrass <i>Pennisetum clandestinum</i>	Para grass <i>Brachiaria mutica</i>	Velvetgrass <i>Holcus</i> spp.
Cattail <i>Typha</i> spp.	Knotgrass <i>Paspalum distichum</i>	Phragmites* <i>Phragmites</i> spp.	Wheatgrass, western <i>Agropyron smithii</i>

*Partial Control

BROADLEAF WEED SPECIES CONTROLLED		
Alfalfa <i>Medicago sativa</i>	Fiddleneck <i>Amsinckia</i> spp.	Poison hemlock <i>Conium maculatum</i>
Alligatorweed* <i>Alternanthera philoxeroides</i>	Fleabane <i>Erigeron</i> spp.	Ragweed, common <i>Ambrosia artemisiifolia</i>
Anise (fennel) <i>Foeniculum vulgare</i>	Fleabane, hairy <i>Conyza bonariensis</i>	Ragweed, giant <i>Ambrosia trifida</i>

Artichoke, Jerusalem <i>Helianthus tuberosus</i>	Horsenettle <i>Solanum carolinense</i>	Smartweed, Pennsylvania <i>Polygonum pensylvanicum</i>
Bassia, fivehook <i>Bassia hyssopifolia</i>	Horseradish <i>Armoracia rusticana</i>	Smartweed, swamp <i>Polygonum coccineum</i>
Bindweed, field <i>Convolvulus arvensis</i>	Ice plant <i>Mesembryanthemum crystallinum</i>	Sowthistle, annual <i>Sonchus oleraceus</i>
Bluweed, Texas <i>Helianthus ciliaris</i>	Knapweed <i>Centaurea repens</i>	Spurge, leafy* <i>Euphorbia esula</i>
Bracken fern <i>Pteridium aquilinum</i>	Kochia <i>Kochia scoparia</i>	Starthistle, yellow <i>Centaurea solstitialis</i>
Bursage, woollyleaf <i>Franseria tomentosa</i>	Lantana <i>Lantana camara</i>	Sunflower <i>Helianthus annuus</i>
Clover, red <i>Trifolium pretense</i>	Lespedeza <i>Lespedeza spp.</i>	Thistle, artichoke <i>Cynara cardunculus</i>
Clover, white <i>Trifolium repens</i>	Lettuce, prickly <i>Lactuca serriola</i>	Thistle, Canada <i>Cirsium arvense</i>
Dandelion <i>Taraxacum officinale</i>	Milkweed <i>Asclepias spp.</i>	Thistle, Russian <i>Salsola kali</i>
Dock, curly <i>Rumex crispus</i>	Mullein, common <i>Verbascum thapsus</i>	Velvetleaf <i>Abutilon theophrasti</i>
Dogbane, hemp <i>Apocynum cannabinum</i>	Nightshade, silverleaf <i>Solanum elaeagnifolium</i>	

*Partial Control

TREE, BRUSH AND VINE SPECIES CONTROLLED			
Alder <i>Alnus spp.</i>	Coyote brush <i>Baccharis consanguinea</i>	Maple: Red* <i>Acer rubrum</i>	Russian Olive** <i>Elaeagnus angustifolia</i>
Ash <i>Fraxinum spp.</i>	Creper, Virginia* <i>Parthenocissus quinquefolia</i>	Sugar <i>Acer saccharum</i>	Sage: black, white <i>Salvia spp.</i>
Aspen, quaking <i>Populus tremuloides</i>	Dewberry <i>Rubus trivialis</i>	Vine <i>Acer circinatum</i>	Sagebrush, California <i>Artemisia californica</i>
Balsam apple* <i>Momordica charantia</i>	Dogwood* <i>Cornus spp.</i>	Monkey Flower* <i>Mimulus guttatus</i>	Salmonberry <i>Rubus spectabilis</i>
Bearmat (Bearclover) <i>Chamaebatia foliolosa</i>	Elderberry <i>Sambucus spp.</i>	Oak: Black* <i>Quercus velutina</i>	Salt Cedar <i>Tamarix spp.</i>
Beech <i>Fagus grandifolia</i>	Elm* <i>Ulmus spp.</i>	Northern Pin <i>Quercus palustris</i>	Sassafras <i>Sassafras albidum</i>
Birch <i>Betula spp.</i>	Eucalyptus <i>Eucalyptus spp.</i>	Post <i>Quercus stellata</i>	Sourwood <i>Oxydendrum arboreum</i>
Blackberry <i>Rubus spp.</i>	Gorse <i>Ulex europaeus</i>	Red <i>Quercus rubra</i>	Sweet potato, wild* <i>Ipomoea pandurata</i>
Blackgum <i>Nyssa spp.</i>	Hasardia* <i>Haplopappus squamosus</i>	Southern Red <i>Quercus falcate</i>	Sumac: Poison* <i>Rhus vernix</i>
Bracken <i>Peridium spp.</i>	Hawthorn <i>Crataegus spp.</i>	White* <i>Quercus alba</i>	Smooth* <i>Rhus glabra</i>
Broom: French <i>Cytisus monspessulanus</i>	Hazel <i>Corylus spp.</i>	Persimmon* <i>Diospyros spp.</i>	Winged* <i>Rhus copallina</i>
Scotch <i>Cytisus scoparius</i>	Hickory* <i>Carya spp.</i>	Pine <i>Pinus spp.</i>	Sweetgum <i>Liquidambar styraciflua</i>
Buckwheat, California* <i>Eriogonum fasciculatum</i>	Holly, Florida/Brazilian Peppertree* <i>Schinus terebinthifolius</i>	Poison Ivy <i>Rhus radicans</i>	Swordfern* <i>Polystichum munitum</i>
Cascara*	Honeysuckle	Poison Oak	Tallowtree, Chinese

TREE, BRUSH AND VINE SPECIES CONTROLLED			
<i>Rhamnus purshiana</i>	<i>Lonicera</i> spp.	<i>Rhus toxicodendron</i>	<i>Sapium sebiferum</i>
Catsclaw* <i>Acacia greggi</i>	Hornbean, American* <i>Carpinus caroliniana</i>	Poplar, Yellow* <i>Liriodendron tulipifera</i>	Tan Oak <i>Lithocarpus densiflorus</i>
Ceanothus* <i>Ceanothus</i> spp.	Kudzu <i>Pueraria lobata</i>	Raspberry <i>Rubus</i> spp.	Thimbleberry <i>Rubus parviflorus</i>
Chamise <i>Adenostoma fasciculatum</i>	Locust, black* <i>Robinia pseudoacacia</i>	Redbud, eastern <i>Cercis canadensis</i>	Tobacco, tree* <i>Nicotiana glauca</i>
Cherry: Bitter <i>Prunus emarginata</i>	Madrone <i>Arutus menziesii</i>	Redvine* <i>Brunnichia ovate</i>	Trumpetcreeper* <i>Campsis radicans</i>
Black <i>Prunus serotina</i>	Manzanita <i>Arctostaphylos</i> spp.	Rose, multiflora <i>Rosa multiflora</i>	Waxmyrtle, southern* <i>Myrica cerifera</i>
Pin <i>Prunus pensylvanica</i>			Willow <i>Salix</i> spp.

*Partial Control

**This product is not registered in California for use on Russian olive

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

PESTICIDE STORAGE: Store product in original container in a safe place and protect from freezing.

PESTICIDE DISPOSAL: Pesticide wastes may be acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

Nonrefillable containers smaller than 5 gallons:

Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable containers larger than 5 gallons:

Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. If recycling or reconditioning is not available, puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable containers larger than 5 gallons:

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10%

full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Ragan and Massey, Inc. or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Ragan and Massey, Inc. and Seller harmless for any claims relating to such factors.

Ragan and Massey, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Ragan and Massey, Inc., and Buyer and User assume the risk of any such use. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, RAGAN AND MASSEY, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

To the extent consistent with applicable law, neither Ragan and Massey, Inc. nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF RAGAN AND MASSEY, INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF RAGAN AND MASSEY, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Ragan and Massey, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Ragan and Massey, Inc.

RM43™ is a trademark of Tangi·Pac, LLC

[Optional Product Claims:

Concentrate
Outdoor Use Only
Fast-acting
Kills weeds and their germinating seeds
Kills weeds and prevents weeds up to 1 year
Two way killer
Prevents new weeds for up to 1 year
No weeds for up to one year
Works two ways
Kills weeds and grasses, roots and all
One application controls weeds for up to 1 year
To keep treated ground clear of weeds for up to 1 year
Not for use on desirable vegetation
Prevents weeds from emerging for up to 1 year
Rainfast within 1-hour
Season long control
Total Vegetation Control
Do not apply in root zone of desirable trees, plants & shrubs
Treats [X] square feet

Use on Fencerows and around Farm Buildings and Barns
The red jug with the white cap is trade dress of TangiPac, LLC
Marketed Through: [Company Name & Address]

[EPA approval date]