



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs  
Registration Division (7505P)  
1200 Pennsylvania Ave., N.W.  
Washington, D.C. 20460

EPA Reg. Number:

84009-37

Date of Issuance:

3/24/21

NOTICE OF PESTICIDE:

Registration  
 Reregistration  
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

RM3

Name and Address of Registrant (include ZIP Code):

Michael Kellogg  
Regulatory Consultant  
Ragan and Massey, Inc.  
c/o Pyxis Regulatory Consulting, Inc.  
4110 136th St. Ct. NW  
Gig Harbor, WA 98332

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Erik Kraft, Product Manager 24  
Fungicide Herbicide Branch, Registration Division (7505P)

Date:

3/24/21

2. You are required to comply with the data requirements described in the DCI identified below:
  - a. Diquat GDCI- EPA-HQ-OPP-2009-0846
  - b. Imazapyr GDCI- EPA-HQ-OPP-2014-0200

You must comply with all of the data requirements within the established deadlines. If you have questions about the DCI listed above, you may contact the Chemical Review Manager in the Pesticide Reevaluation Division: <http://iaspub.epa.gov/apex/pesticides/f?p=chemicalsearch:1>

3. Make the following label changes before you release the product for shipment:
  - Revise the EPA Registration Number to read, “EPA Reg. No. 84009-37.”
4. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.


If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 02/17/2021

If you have any questions, please contact Manjula Unnikrishnan by phone at 703-347-8520, or via email at [unnikrishnan.manjula@epa.gov](mailto:unnikrishnan.manjula@epa.gov) .

Enclosure

[Note to reviewer: [Text] in brackets denotes optional text].  
[Note to reviewer: {Text} in braces denotes where in the final label text will appear.]



{[BOOKLET FRONT PANEL LANGUAGE]}

## RM3

**[Alternate Brand Names: RM100 Total Vegetation Control; Compare-N-Save® Extended Weed Control; Compare-N-Save® Total Vegetation Control; FarmWorks® Extended Weed Control; Ground Shield Total Vegetation Control; RM-X1 Total Vegetation Control]**

**FOR [QUICK ACTING] NONSELECTIVE WEED CONTROL OF WEEDS IN NONCROP AREAS**

GLUFOSINATE	GROUP	10	HERBICIDE
IMAZAPYR	GROUP	2	HERBICIDE
DIQUAT DIBROMIDE	GROUP	22	HERBICIDE

TO PREVENT ACCIDENTAL POISONING, NEVER STORE THIS PRODUCT IN FOOD, DRINK, OR UNLABELED CONTAINERS.

### ACTIVE INGREDIENTS:

Glufosinate ammonium..... 11.33%  
Imazapyr, isopropylamine salt ..... 1.10%  
Diquat dibromide..... 1.09%

**OTHER INGREDIENTS:** ..... 86.48%

**TOTAL:** ..... 100.00%

Equivalent to 1 lb. of glufosinate active ingredient; 0.1 lb. of imazapyr acid active ingredient; and 0.1 lb. of diquat dibromide active ingredient per U.S. gallon.

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION - PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

[See] [inside] [attached] [label] [booklet] [for] [First Aid][,] [additional] [Precautionary Statements][,] [and] [Directions for Use] [including] [and] [Storage and Disposal] [instructions].

**EPA Reg. No.** 84009-xx  
**EPA Est. No.**

**Manufactured For:**  
Ragan & Massey, Inc.  
101 Ponchatoula Parkway  
Ponchatoula, LA 70454

[Batch/Lot No.]

**Net Contents:**

## {LANGUAGE INSIDE BOOKLET}

<b>FIRST AID</b>	
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• <b>DO NOT</b> induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• <b>DO NOT</b> give anything by mouth to an unconscious person.</li> </ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• <b>IMMEDIATELY</b> hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>HOT LINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product call your poison control center at 1-800-222-1222.	
<b>NOTE TO PHYSICIAN:</b> If in eyes, treat symptomatically. Symptoms may develop gradually. Severe damage may be caused by apparently trivial contact and healing may be delayed. Medical supervision should be continued until healing is complete.	

### **PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION**

Harmful if swallowed or inhaled. Causes moderate eye irritation. Avoid breathing spray mist. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Wear long-sleeved shirt and long pants; chemical-resistant gloves such as barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, polyvinyl chloride (PVC) ≥14 mils, or Viton® ≥14 mils; and shoes plus socks. Protective eyewear is strongly recommended when mixing and applying this product.

### **ENVIRONMENTAL HAZARDS**

This product is toxic to plants and aquatic invertebrates. Drift and run-off may be hazardous to plants in water adjacent to treated areas. **DO NOT** apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** clean equipment or dispose of equipment washwaters in a manner that will contaminate water resources or arable land. Glufosinate-ammonium and its degradates have those properties normally associated with pesticides that have been detected in groundwater. Use of this product in areas with coarse soils and high water tables may result in groundwater contamination.

**NON-TARGET ORGANISM ADVISORY:** This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

### **PHYSICAL OR CHEMICAL HAZARDS**

**DO NOT** mix or allow contact with oxidizing agents. Hazardous chemical reaction may occur.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**DO NOT** use this product until you have read the entire label.

**Restrictions:**

- In the State of New York only: Not for use in Nassau and Suffolk Counties.
- Keep people and pets out of the area during application.
- **DO NOT** use on feed or food crops.
- **DO NOT** apply this product via backpack sprayer.
- **DO NOT** apply by air.
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply near bodies of water including fish pools, ponds, lakes or streams.
- **DO NOT** apply over the root systems of desirable plants. For trees and shrubs do not apply closer than twice the distance from the trunk to the drip line as roots may be within this area.
- **DO NOT** apply to slopes in landscape as movement on soil surfaces may damage desirable plants down slope.
- **DO NOT** apply next to a fence if desirable vegetation is growing on the other side, or if future planting is intended.
- **DO NOT** use before planting lawns, fruits, vegetables, flowers, or other plants because this product remains active in soil for up to one year.
- **DO NOT** use on desirable lawns or turf.
- **DO NOT** apply to powdery, dry or light sandy soils when the soil may be blown by wind. If necessary, soil surface can be settled by rain or irrigation.
- **DO NOT** apply directly to or allow drift to contact desirable green tissue or green, thin, or uncalloused bark of desirable vegetation.
- **DO NOT** apply or allow product to drift onto desirable plants.
- **DO NOT** allow animals intended for slaughter to graze on treated vegetation.

**PRODUCT INFORMATION**

RM3 is a nonselective water-soluble herbicide for application as a foliar spray for the control of a broad spectrum of emerged annual and perennial grass and broadleaf weeds. RM3 will also control certain woody species. THOROUGH SPRAY COVERAGE IS IMPORTANT. Visual effects and control from application of RM3 occur within 2 to 4 days after application under good growing conditions.

This product is nonselective and will injure or kill all green vegetation contacted by the spray. Avoid all contact with foliage or green tissue of desirable vegetation. Avoid direct spray contact with green, thin, or uncalloused bark of desirable vegetation or plant injury may result. If desirable vegetation is contacted, rinse the sprayed portion with water immediately.

Weed control may be reduced when applications are made to weeds under stress due to drought or cool temperatures. Weeds under stress or in dense populations will require application at the highest rate. Refer to the How to Apply section of this label.

**WHERE TO USE**

Curbs, Sidewalks, Driveways, Parking areas, Brick walks, Gravel Paths, Patios, Vacant lots, Paved areas (including crack and crevice), Play areas, Tennis courts, Fencerows, Along fences of paddocks, corrals, Hedgerows, Dry Ditchbanks, Around domestic dwellings, Around farm buildings and barns, Shelter beds, Farm yards, Farm premises, Outdoor buildings, structures, and shelters including stables, and dog kennels.

Only apply to locations where no vegetation growth is desired for up to one year.

**WHEN TO APPLY**

Thorough spray coverage is important. Apply to bare ground or when weeds are present. If weeds are taller than 6 inches, mow or cut down weeds and remove debris before treatment and allow for 2 weeks of new growth. Contact with soil is important for extended control. If woody vines or brush is mowed or cut, wait until full leaf formation before treatment.

To prevent growth of weeds under asphalt driveways, play areas, tennis courts and parking areas, apply this product before laying surfacing materials.

Treat area uniformly to maximize results.

Re-treatment every 6 to 8 months may be necessary in areas of heavy rainfall.

### HOW TO MIX

RM3 must be mixed with water to make a finished spray solution as follows:

1. Fill the spray tank with the required amount of water.
2. Add the proper amount of this product, then mix thoroughly.

### HOW TO APPLY

Apply on a sunny day with temperatures above 60°F. Avoid product runoff to nearby desirable plants. In soils with heavy clay, weed control may exceed one year.

### TOTAL VEGETATION CONTROL (BAREGROUND)

Use hand operated spray equipment including pump-up sprayers or sprinkling cans. Mix 4 fl. oz. RM3 (equivalent to 0.03 lb. ai glufosinate/0.003 lb. ai imazapyr/0.003 lb. ai diquat) with water to produce 1 to 10 gallons of spray solution and apply to 1,000 sq. ft. Uniformly spray the mix over area where total vegetation control and bareground is desired. Optimum performance will be achieved from applications to actively growing weeds in early summer. Use 1 to 10 gallons of water depending on spray speed and equipment used to deliver 4 fl. oz. RM3 per 1,000 sq. ft.

#### Restrictions:

- **DO NOT** apply more than 170 fl. oz. of product (1.33 lbs. ai glufosinate/0.13 lb. ai imazapyr/0.13 lb. ai diquat) per acre per application (4 fl. oz. per 1,000 sq. ft.).
- **DO NOT** apply more than 170 fl. oz. of product (1.33 lbs. ai glufosinate/0.13 lb. ai imazapyr/0.13 lb. ai diquat) per acre per year (4 fl. oz. per 1,000 sq. ft.).
- **DO NOT** make more than 1 application per year.

### SPOT OR DIRECTED APPLICATIONS

Use this product as a spot or directed spray application using 2 to 4 fluid ounces of product (0.016 lb. ai glufosinate/0.0016 lb. ai imazapyr/0.0016 lb. ai diquat to 0.03 lb. ai glufosinate/0.003 lb. ai imazapyr/0.003 lb. ai diquat) per gallon of water. Mix 2 to 4 fluid ounces of product per gallon of water depending upon the weed and stage of growth as shown in the following sections. Spray undesirable vegetation foliage on a spray-to-wet basis. Ensure uniform and complete coverage. Use a coarse spray. Thoroughly clean the sprayer following use.

#### Restrictions:

- **DO NOT** apply more than 4 fl. oz. of product (0.03 lb. ai glufosinate/0.003 lb. ai imazapyr/0.003 lb. ai diquat) per acre per application (0.09 fl. oz. per 1,000 sq. ft.).
- **DO NOT** apply more than 104 fl. oz. of product (0.78 lb. glufosinate/0.078 lb. ai imazapyr/0.078 lb. ai diquat) per acre per year (2.38 fl. oz. per 1,000 sq. ft.).
- **DO NOT** make more than 26 applications per year.
- Applications must be made at least 14 days apart.
- **DO NOT** apply beyond runoff
- **DO NOT** spray during windy conditions

#### Weeds Controlled by RM3 For spot application:

Apply 2 fl. oz. of product (0.016 lb. ai glufosinate/0.0016 lb. ai imazapyr/0.0016 lb. ai diquat) per gallon of water when the weed height or diameter is less than 6 inches.

Apply 3 fl. oz. of product (0.023 lb. ai glufosinate/0.0023 lb. ai imazapyr/0.0023 lb. ai diquat) per gallon of water when the weed height or diameter is 6 inches or greater.

#### Broadleaf Weeds

chickweed  
clover  
common cocklebur  
filaree  
jimsonweed  
kochia

#### Grasses and Sedges

barnyardgrass  
cupgrass  
fall panicum  
giant foxtail  
goosegrass  
green foxtail

London rocket  
malva (little mallow)  
marestail  
purslane  
shepherdspurge  
smartweed

Johnsongrass (rhizome)  
lovegrass  
shattercane  
smallflower Alexandergrass (signal grass)  
stinkgrass  
windgrass  
yellow foxtail

#### For spot application:

Apply 3 fl. oz. of product (0.023 lb. ai glufosinate/0.0023 lb. ai imazapyr/0.0023 lb. ai diquat) per gallon of water when the weed height or diameter is less than 6 inches.

Apply 4 fl. oz. of product (0.03 lb. ai glufosinate/0.003 lb. ai imazapyr/0.003 lb. ai diquat) per gallon of water when the weed height or diameter is 6 inches or greater.

#### Broadleaf Weeds

annual sowthistle  
bindweed  
buffalobur  
burdock  
Canada thistle  
curly dock  
dandelion  
dogbane (hemp)  
field gromwell  
fleabane  
goldenrod  
horsetail  
lambquarters  
leafy spurge  
mugwort  
musk thistle  
nettle  
nightshade  
pennycress  
pigweed, red root  
plantain

prickly lettuce  
ragweed  
Russian thistle  
tansy mustard  
velvetleaf  
vervain  
Virginia copperleaf  
white heath aster  
wild buckwheat  
wild mustard  
wild onion  
wild rose  
wild turnip  
woodsorrel  
yellow rocket

#### Grasses and Sedges

annual bluegrass  
bahiagrass  
barley  
Bermudagrass  
carpetgrass  
crabgrass  
dallisgrass  
downy brome  
fescue  
guineagrass  
Kentucky bluegrass  
nutsedge  
paragrass  
quackgrass  
ryegrass  
sandbur  
smooth brome  
torpedograss  
vaseygrass  
wheat  
wild oat

#### Use Notes

1. Use higher rates within the specified rate range for plant sizes listed when vegetation cover is dense or when weeds are growing under stressed conditions including drought or when average temperatures are below 50°F.

#### Use on Woody Species

When applied as directed, RM3 will provide control, partial control, or suppression of certain perennial woody weed species. Apply 64 to 192 fl. oz. of product (0.5 lb. ai glufosinate/0.05 lb. ai imazapyr/0.05 lb. ai diquat to 1.5 lbs. ai glufosinate/0.15 lb. ai imazapyr/0.15 lb. ai diquat) per acre (1.46 to 4.4 fl. oz. per 1,000 sq. ft.). Use the higher listed rates of this product when conditions are not optimum for spray penetration, including when vegetation growth is heavy or dense. Lower listed rates may be used when the target species is a conifer and when vegetation growth conditions allow for uniform spray coverage.

blackberry	<i>Rubus</i> spp.
deer brush	<i>Ceanothus integerrimus</i>
Douglas fir	<i>Pseudotsuga menziesii</i>
gallberry	<i>Ilex</i> spp.
hazel	<i>Corylus</i> spp.
honeysuckle	<i>Lonicera</i> spp.
huckleberry	<i>Gaylussacia</i> spp.

maple	<i>Acer</i> spp.
multiflora rose	<i>Rosa multiflora</i>
oak	<i>Quercus</i> spp.
pine	<i>Pinus</i> spp.
poison ivy	<i>Toxicodendron radicans</i>
poison oak	<i>Toxicodendron toxicarium</i>
roundleaf greenbriar	<i>Smilax rotundifolia</i>
salmonberry	<i>Rubus spectabilis</i>
sweet gum	<i>Liquidambar styraciflua</i>
sumac	<i>Rhus</i> spp.
thimbleberry	<i>Rubus parviflorus</i>
trumpetcreeper	<i>Campsis radicans</i>
vine maple	<i>Acer circinatum</i>
Western red cedar	<i>Thuja plicata</i>

### TRIMMING AND EDGING

RM3 may be used for trimming and edging landscape areas including: around individual trees and shrubs, landscape beds, foundations, fences, driveways, paths, and parking areas. For control of weeds emerging from seed, use RM3 in a tank mix with preemergence herbicides. If spraying in areas adjacent to desirable plants, use a shield made of cardboard, plywood, or sheet metal while spraying to help prevent spray from contacting foliage of desirable plants. Refer to the How to Apply section of this labeling for appropriate application rates to control specific weeds.

#### Restrictions:

- **DO NOT** apply more than 4 fl. oz. of product (0.03 lb. ai glufosinate/0.003 lb. ai imazapyr/0.003 lb. ai diquat) per acre per application (0.09 fl. oz. per 1,000 sq. ft.).
- **DO NOT** apply more than 104 fl. oz. of product (0.78 lb. glufosinate/0.078 lb. ai imazapyr/0.078 lb. ai diquat) per acre per year (2.38 fl. oz. per 1,000 sq. ft.).
- **DO NOT** make more than 26 applications per year.
- Applications must be made at least 14 days apart.

### FARMSTEADS

When applied as directed, this product controls undesirable plant vegetation in non-crop areas around farmstead building foundations, shelter belts, along fences, and nonselective farmstead weed control. Refer to the How to Apply section of this labeling for appropriate application rates to control specific weeds.

#### Restrictions for spot or directed applications:

- **DO NOT** allow animals intended for slaughter to graze on treated vegetation.
- **DO NOT** apply more than 4 fl. oz. of product (0.03 lb. ai glufosinate/0.003 lb. ai imazapyr/0.003 lb. ai diquat) per acre per application (0.09 fl. oz. per 1,000 sq. ft.).
- **DO NOT** apply more than 104 fl. oz. of product (0.78 lb. glufosinate/0.078 lb. ai imazapyr/0.078 lb. ai diquat) per acre per year (2.38 fl. oz. per 1,000 sq. ft.).
- **DO NOT** make more than 26 applications per year.
- Applications must be made at least 14 days apart.

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store product in original container only. Store in a cool, dry place.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

#### CONTAINER HANDLING:

**[Nonrefillable Container (five gallons or less):** Nonrefillable container. **DO NOT** reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later



use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration.]

**[Nonrefillable Container (larger than 5 gallons):** Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration.]

#### **CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Ragan and Massey, Inc. or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Ragan and Massey, Inc. and Seller harmless for any claims relating to such factors.

Ragan and Massey, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Ragan and Massey, Inc., and Buyer and User assume the risk of any such use. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, RAGAN AND MASSEY, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

To the extent consistent with applicable law, neither Ragan and Massey, Inc. nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF RAGAN AND MASSEY, INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF RAGAN AND MASSEY, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Ragan and Massey, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Ragan and Massey, Inc.

[Compare-N-Save®] [FarmWorks®] is a trademark of Tangi-Pac, LLC

[EPA approval date]

# {[LANGUAGE ON LABEL AFFIXED TO CONTAINER]}

## RM3

GLUFOSINATE	GROUP	10	HERBICIDE
IMAZAPYR	GROUP	2	HERBICIDE
DIQUAT DIBROMIDE	GROUP	22	HERBICIDE

TO PREVENT ACCIDENTAL POISONING, NEVER STORE THIS PRODUCT IN FOOD, DRINK, OR UNLABELED CONTAINERS.

### ACTIVE INGREDIENTS:

Glufosinate ammonium.....	11.33%
Imazapyr, isopropylamine salt .....	1.10%
Diquat dibromide .....	1.09%
<b>OTHER INGREDIENTS:</b> .....	<b>86.48%</b>
<b>TOTAL:</b> .....	<b>100.00%</b>

Equivalent to 1 lb. of glufosinate active ingredient; 0.1 lb. of imazapyr acid active ingredient; and 0.1 lb. of diquat dibromide active ingredient per U.S. gallon.

## KEEP OUT OF REACH OF CHILDREN CAUTION - PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li><b>DO NOT</b> induce vomiting unless told to do so by a poison control center or doctor.</li> <li><b>DO NOT</b> give anything by mouth to an unconscious person.</li> </ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product call your poison control center at 1-800-222-1222.	
<b>PHYSICIAN INSTRUCTIONS:</b> If in eyes, treat symptomatically. Symptoms may develop gradually. Severe damage may be caused by apparently trivial contact and healing may be delayed. Medical supervision should be continued until healing is complete.	

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed or inhaled. Causes moderate eye irritation. Avoid breathing spray mist. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear long-sleeved shirt and long pants; chemical-resistant gloves such as barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, polyvinyl chloride (PVC) ≥14 mils, or Viton® ≥14 mils; and shoes plus socks. Protective eyewear is strongly recommended when mixing and applying this product.

### ENVIRONMENTAL HAZARDS

This product is toxic to plants and aquatic invertebrates. Drift and run-off may be hazardous to plants in water adjacent to treated areas. **DO NOT** apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** clean equipment or dispose of equipment washwaters in a manner that will contaminate water resources or arable land. Glufosinate-ammonium and its degradates have those properties normally associated with pesticides that have been detected in groundwater. Use of this product in areas with coarse soils and high water tables may result in groundwater contamination.

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store product in original container only. Store in a cool, dry place.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

### CONTAINER HANDLING:

**[Nonrefillable Container (five gallons or less):** Nonrefillable container. **DO NOT** reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration.]

**[Nonrefillable Container (larger than 5 gallons):** Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration.]

[See] [inside] [attached] [label] [booklet] [for] [First Aid][,] [additional] [Precautionary Statements][,] [and] [Directions for Use] [including] [and] [Storage and Disposal] [instructions].

**EPA Reg. No.** 84009-xx  
**EPA Est. No.**

**Manufactured For:**  
Ragan & Massey, Inc.  
101 Ponchatoula Parkway  
Ponchatoula, LA 70454

**Net Contents:**