

84059-3

12/04/2012

1/65



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON D C 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

John Fournier
Regulatory Manager
Marriott Bio Innovations
2121 Second Street Suite B 107
Davis CA 95618

DEC 04 2012

Subject Label Amendment to Remove California Restrictions Language
Product Name Regalia Bioprotectant Concentrate
EPA Reg No 84059 3
Your Submission Dated September 7 2012

Dear Mr Fournier

The amendment referred to above submitted in connection with registration under FIFRA section 3(c)(5) is acceptable provided that you

- 1 Submit and/or cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data
- 2 Submit five (5) copies of your final printed labeling before you release the product for shipment Final printed labeling means the label or labeling of the product when distributed or sold Clearly legible reproductions or photo reductions will be accepted for unusual labels such as those silk screened directly onto glass or metal containers or large bags or drum labels

If these conditions are not complied with the registration will be subject to cancellation in accordance with FIFRA section 6(e) Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions Should you have any questions regarding this action you may contact Gina Burnett at (703) 605 0513 or via email at burnett_gina@epa.gov A stamped copy of the label is enclosed for your records

Sincerely

Linda A Hollis Chief
Biochemical Pesticides Branch
Biopesticides and Pollution
Prevention Division (7511P)

CONCURRENCE							
SYMBOL	▶ 7511P						
SURNAME	▶ Burnett						
DATE	▶ 12/3/12						
EPA	m						

MASTER LABEL

Regalia® Bioprotectant Concentrate

Alternate Brand Names Regalia® Biofungicide Concentrate

Sublabel A Agricultural Crops
Sublabel B Turf & Professional Landscape Use
Sublabel C Home & Garden Use

EPA Registration No 84059 3

ACCEPTED

12/4/2012

Under the Federal Insecticide Fungicide and Rodenticide Act as amended for the pesticide registered under EPA Reg No

84059-3

Sublabel A Agricultural Crops

Regalia® Bioprotectant Concentrate

A plant extract to boost the plants defense mechanisms to protect against certain fungal and bacterial diseases and to improve plant health

Active ingredient Extract of *Reynoutria sachalinensis* 5 %
Other ingredients 95 %
Total 100 %

EPA Reg No 84059 3

EPA Est No 085970 FL 001
EPA Est No 047857 CA 001
EPA Est No 61842 CA 001

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID	
IF SWALLOWED	Call poison control center or doctor immediately for treatment advice Have person sip a glass of water if able to swallow Do not induce vomiting unless told to do so by the poison control center or doctor Do not give anything by mouth to an unconscious person
IF ON SKIN OR CLOTHING	Take off contaminated clothing Rinse skin immediately with plenty of water for 15 20 minutes Call a poison control center or doctor for treatment advice
IF INHALED	Move person to fresh air If person is not breathing call 911 or an ambulance then give artificial respiration preferably by mouth to mouth if possible Call a poison control center or doctor for further treatment advice
IF IN EYES	Hold eye open and rinse slowly and gently with water for 15 20 minutes Remove contact lenses if present after the first 5 minutes then continue rinsing eye Call a poison control center or doctor for treatment advice
Have the product container or label with you when calling a poison control center or doctor or if going for treatment	



LOT # _____

Net Contents 1 pint, 1 quart 1 gallon 2 5 gallon 5 gallon, 55 gallon drum
Marrone Bio Innovations, Inc 2121 Second St Suite B 107, Davis CA 95618

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION Causes moderate eye irritation Avoid contact with eyes or clothing Wear goggles or safety glasses Wash thoroughly with soap and water after handling and before eating drinking chewing gum using tobacco or using the toilet

PERSONAL PROTECTIVE EQUIPMENT (PPE)

- Applicators and other handlers must wear
 - Long sleeved shirt and long pants
 - Shoes plus socks
 - Waterproof gloves
 - Protective eyewear

Follow manufacturer s instructions for cleaning and maintaining PPE If no such instructions for washables use detergent and hot water Keep and wash PPE separately from other laundry

ENVIRONMENTAL HAZARDS

For terrestrial uses Do not apply directly to water to areas where surface water is present or to intertidal areas below the mean high water mark Do not contaminate water when disposing of equipment washwater or rinsate

(Use the following additional statement for containers that hold 5 gallons or more Do not discharge effluent containing this product into lakes streams ponds estuaries oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority For guidance contact your State Water Board or Regional Office of the EPA)

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling Do not apply this product in a way that will contact workers or other persons either directly or through drift Only protected handlers may be in the area during application For any requirements specific to your State or Tribe consult the State or Tribal agency responsible for pesticide regulation

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170 This standard contains requirements for the protection of agricultural workers on farms forests nurseries and greenhouses and handlers of agricultural pesticides It contains requirements for training decontamination notification and emergency assistance It also contains specific instructions and exemptions pertaining to the statements on this label about personal protective equipment (PPE) and the restricted entry interval (REI) The requirements in this box onl apply to uses of this product that are covered by the Worker Protection Standard

Do not enter or allow worker entry into treated areas during the restricted entry interv2l (REI) of 4 hours

PPE required for early entry to treated areas that is permitted under the Worker Protectio 1 Standard and that involves contact with anything that has been treated such as plants soil or wa cr 1

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

GENERAL INFORMATION

Regalia® Bioprotectant Concentrate is an extract from the plant *Reynoutria* spp for use on ornamental plants turf and edible crops Regalia® Bioprotectant Concentrate applied to actively growing plants (see DIRECTIONS FOR USE) will improve plant health and will help make the treated portions resistant to certain plant diseases Plant health benefits often result in greater yields at harvest especially when crops are stressed by pathogens or environmental conditions Use Regalia® Bioprotectant Concentrate as a preventative rather than a curative application Apply prior to disease infestation to protect the growing leaf tissue See specific information for diseases controlled and use rates on ornamental plants turf and edible crops

Regalia® Bioprotectant Concentrate can be used as a seed treatment plant dip soil drench in furrow spray or applied through drip irrigation to control or suppress certain soil borne diseases and to promote healthy root growth See specific information for diseases controlled and use rates on treating seeds with Regalia® Bioprotectant Concentrate

MODE OF ACTION

The extract obtained from *Reynoutria* spp plant material contains active chemical compounds The extract when applied to the host plant increases the plant's defense system due to a five fold increase in phenolics and antioxidants and strengthens cell walls This induced resistance against important diseases is not systemic but provides some translaminar protection Repeat foliar applications at 7-14 day intervals to maintain induction and to protect new plant growth The resistance induction takes place within one to two days

Use Regalia® Bioprotectant Concentrate therefore as a preventative treatment

**MIXING AND APPLICATION INSTRUCTIONS
- SHAKE WELL PRIOR TO USE -**

Regalia® Bioprotectant Concentrate is a micro emulsion concentrate consisting of certain ingredients extracted from *Reynoutria* spp Use 50 mesh nozzle screens or larger

- See **AERIAL APPLICATION** section for aerial application use directions
- See **CHEMIGATION** section for chemigation use directions
- See **PRE PLANT DIP** section for pre plant dip use directions
- See **SEED TREATMENT** section for seed treatment use directions
- See **SOIL TREATMENT** section for soil application use directions

Use higher water volumes with larger sized crops and extensive foliage to secure thorough coverage

Regalia® Bioprotectant Concentrate alone Add 1/2 of the required amount of water to the mix tank With the agitator running add the Regalia® Bioprotectant Concentrate to the mix tank Continue agitation while adding the remainder of the water Begin application of the solution after the Regalia® Bioprotectant Concentrate has completely dispersed into the mix water Maintain agitation until all the mixture has been applied

Regalia® Bioprotectant Concentrate + tank mixtures Add 1/2-3/4 of the required amount of water to the mix tank Start the agitation before adding any tank mix partners In general tank mix partners should be added in the order wettable powders dry flowable formulations liquid flowable formulations and emulsifiable formulations such as Regalia® Bioprotectant Concentrate Always allow each tank mix partner to become completely dispersed before adding the next component Maintain continuous agitation until all components have been dispersed and throughout the application process After all components are completely dispersed add the remainder of the water Regalia® Bioprotectant Concentrate cannot be mixed with another product with a prohibition against mixing Use of the tank mix must be in accordance with the more restrictive label limitations and precautions **Do not pre mix Regalia® Bioprotectant Concentrate with any other tank mix component prior to adding to the spray tank**

Compatibility Do not combine Regalia® Bioprotectant Concentrate in the spray tank with pesticides adjuvants or fertilizers if there has been no previous experience or use of the combination to show it is physically compatible effective and non injurious under your use conditions

Regalia® Bioprotectant Concentrate is compatible with many commonly used pesticides fertilizers adjuvants and surfactants but has not been evaluated with all potential combinations To ensure compatibility of the tank mix

combinations evaluate prior to use as follows Using a suitable container add the proportional amounts of product to water Add wettable powders first then water dispersible granules then liquid flowables and lastly emulsifiable concentrates Mix thoroughly and let stand for at least five minutes If the combination stays mixed or can be remixed it is physically compatible Test the mix on a small portion of the crop to be treated to ensure that a phytotoxic response will not occur as a result of the application

AERIAL APPLICATION INSTRUCTIONS

Apply Regalia® Bioprotectant Concentrate by aerial application to the Edible Crops listed at the rate of 0.5 – 1 quart per acre in a minimum of 5 gallons of water per acre unless specified differently in the SELECTED CROPS section Increasing the amount of water applied per acre may improve product performance Follow all instructions to reduce aerial drift

AERIAL DRIFT REDUCTION ADVISORY INFORMATION

GENERAL Avoiding spray drift at the application site is the responsibility of the applicator The interaction of many equipment and weather related factors determine the potential for spray drift The applicator and the grower are responsible for considering all these factors when making decisions Where states have more stringent regulations they should be observed Note This section is advisory in nature and does not supersede the mandatory label requirements

INFORMATION ON DROPLET SIZE The most effective way to reduce drift potential is to apply large droplets The best drift management strategy is to apply droplets large enough to provide sufficient coverage and control Applying larger droplets reduces drift potential but will not prevent drift if applications are made improperly or under unfavorable environmental conditions (see Wind Temperature and Humidity and Temperature Inversions)

CONTROLLING DROPLET SIZE Use high flow rate nozzles to apply the highest practical spray volume Nozzles with higher rated flows produce larger droplets Pressure – Do not exceed the nozzle manufacturer s recommended pressures For many nozzle types lower pressure produces larger droplets When high flow rates are needed use higher flow rate nozzles instead of increasing pressure Number of Nozzles – Use the minimum number of nozzles that provide uniform coverage Nozzle Orientation – Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice Significant deflection from horizontal will reduce droplet size and increase drift potential Nozzle Type – Use a nozzle type that is designed for the intended application With most nozzle types narrower spray angles produce larger droplets Consider using low drift nozzles Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift

BOOM WIDTH For aerial applications the boom width must not exceed 75% of the wingspan or 90% of the rotary blade Use upwind swath displacement and apply only when wind speed is 3-10 mph as measured by an anemometer Use medium or coarser spray according to ASAE 572 definition for standard nozzles or VMD for spinning atomizer nozzles If application includes a no spray zone do not release spray at a height greater than 10 feet above the ground or crop canopy

APPLICATION HEIGHT Do not make application at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind

SWATH ADJUSTMENT When applications are made with a crosswind the swath will be displaced downward Therefore on the up and downwind edges of the field the applicator must compensate for this displacement by adjusting the path of the aircraft upwind Swath adjustment distance should increase with increasing drift potential (higher wind smaller drops etc)

WIND Drift potential is lowest between wind speeds of 2 10 mph However many factors including droplet size and equipment type determine drift potential at any given speed Application should be avoided below 2 mph due to variable wind direction and high inversion potential NOTE Local terrain can influence wind patterns Every applicator should be familiar with local wind patterns and how they affect spray drift

TEMPERATURE AND HUMIDITY When making applications in low relative humidity set up equipment to produce larger droplets to compensate for evaporation Droplet evaporation is most severe when conditions are both hot and dry

TEMPERATURE INVERSIONS Do not apply during a temperature inversion because drift potential is high Temperature inversions restrict vertical air mixing which causes small suspended droplets to remain in a concentrated cloud This cloud can move in unpredictable directions due to the light variable winds common during inversions Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind They begin to form as the sun sets and often continue into the morning Their presence can be indicated by ground fog however if fog is not present inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion while smoke that moves upward and rapidly dissipates indicates good vertical air mixing

SENSITIVE AREAS The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e g residential areas bodies of water known habitat for threatened or endangered species non target crops) is minimal (e g when wind is blowing away from the sensitive areas) Do not allow spray to drift from the application site and contact people structures people occupy at any time and the associated property parks and recreation areas non target crops aquatic and wetland areas woodlands pastures rangelands or animals

CHEMIGATION USE DIRECTIONS

Apply Regalia® Bioprotectant Concentrate at 1 4 quarts per acre according to the instructions below unless specified differently in the **SELECTED CROPS** section

CHEMIGATION

General Requirements

- 1) Apply this product only through a drip system or sprinkler including center pivot lateral move end tow side (wheel) roll traveler big gun solid set hand move flood (basin) furrow border or drip (trickle) irrigation systems Do not apply this product through any other type of irrigation system
- 2) Crop injury lack of effectiveness or illegal pesticide residues in the crop can result from non uniform distribution of treated water
- 3) If you have questions about calibration you should contact State Extension Service specialists equipment manufacturers or other experts
- 4) Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place
- 5) A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise

Specific Requirements for Chemigation Systems Connected to Public Water Systems

- 1) Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year

- 2) Chemigation systems connected to public water systems must contain a functional reduced pressure zone backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- 3) The pesticide injection pipeline must contain a functional automatic quick closing check valve to prevent the flow of fluid back toward the injection pump.
- 4) The pesticide injection pipeline must contain a functional normally closed solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- 5) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or in cases where there is no water pump when the water pressure decreases to the point where pesticide distribution is adversely affected.
- 6) Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- 7) Do not apply when wind speed favors drift beyond the area intended for treatment.

Specific Requirements for Sprinkler Chemigation

- 1) The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- 2) The pesticide injection pipeline must contain a functional automatic quick closing check valve to prevent the flow of fluid back toward the injection pump.
- 3) The pesticide injection pipeline must also contain a functional normally closed solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- 4) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- 5) The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- 6) Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being filled with a system interlock.
- 7) Do not apply when wind speed favors drift beyond the area intended for treatment.

Specific Requirements for Flood (Basin) Furrow and Border Chemigation

- 1) Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as a drop structure or weir box to decrease potential for water source contamination from backflow if water flow stops.
- 2) The systems utilizing a pressurized water and pesticide injection system must meet the following requirements:
 - a) The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
 - b) The pesticide injection pipeline must contain a functional automatic quick closing check valve to prevent the flow of fluid back toward the injection pump.
 - c) The pesticide injection pipeline must also contain a functional normally closed solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
 - d) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
 - e) The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

- f Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock

Specific Requirements for Drip (Trickle) Chemigation

- 1) The system must contain a functional check valve vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow
- 2) The pesticide injection pipeline must contain a functional automatic quick closing check valve to prevent the flow of fluid back toward the injection pump
- 3) The pesticide injection pipeline must also contain a functional normally closed solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down
- 4) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops
- 5) The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected
- 6) Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock

Application Instructions for All Types of Chemigation

- 1) Remove scale pesticide residues and other foreign matter from the chemical supply tank and entire injector system Flush with clean water Failure to provide a clean tank void of scale or residues may cause product to lose effectiveness or strength
- 2) Determine the treatment rates as indicated in the directions for use and make proper dilutions Product can be applied continuously or at any time during the water application
- 3) Prepare a solution in the chemical tank by filling the tank with the required water and then adding product as required

PRE PLANT DIP USE DIRECTIONS

Regalia® Bioprotectant Concentrate can be applied as a pre plant dip for improved plant health and suppression of certain soil borne diseases Apply Regalia® Bioprotectant Concentrate in 1-4 quarts product per 100 gallons of water as a pre plant dip immediately prior to transplanting unless specified differently in the SELECTED CROPS section

SEED TREATMENT USE DIRECTIONS

Regalia® Bioprotectant Concentrate can be applied as a seed dressing for suppression of soil borne diseases to improve early season root growth Regalia® Bioprotectant Concentrate may be applied as a water based slurry with other registered seed treatment insecticides and fungicides through standard slurry or mist type commercial seed treatment equipment

Mixing instructions Prepare no more mixture than is required for the immediate operation Agitate the solution continuously during mixing and application Mechanical mixing is recommended for proper mixing of Regalia® Bioprotectant Concentrate mixtures

Regalia® Bioprotectant Concentrate alone Add ½ of the required amount of water to the mix tank With the agitator running add the Regalia® Bioprotectant Concentrate to the mix tank Continue agitation while adding the remainder of the water Begin application of the solution after the Regalia® Bioprotectant Concentrate has completely dispersed into the mix water Maintain agitation until all the mixture has been applied

Regalia® Bioprotectant Concentrate+ tank mixtures Add ½ of the required amount of water to the mix tank Start the agitation before adding any tank mix partners In general tank mix partners should be added in this order wettable powders dry flowable formulations liquid flowable formulations and emulsifiable formulations such as Regalia® Bioprotectant Concentrate Always allow each tank mix partner to become completely dispersed before adding the next component Maintain continuous agitation until all components have been dispersed and throughout the application process

Note When using Regalia® Bioprotectant Concentrate in tank mixtures all products in water soluble packaging should be added to the tank before any other tank mix partner including Regalia® Bioprotectant Concentrate Allow the water soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank mix partner to the tank

If using Regalia® Bioprotectant Concentrate in a tank mixture with other seed treatment products observe all directions for use crops/sites use rates dilution ratios precautions and limitations which appear on the tank mix partner label No label dosage may be exceeded and the most restrictive label precautions and limitations must be followed This product should not be mixed with any product which prohibits such mixing

Note Federal law requires that bags containing treated seeds shall be labeled with the following information This seed has been treated with Regalia® Bioprotectant Concentrate fungicide Do not use for food feed or oil purposes Store away from feed and foodstuffs **Treated seed bagged for later use must contain an EPA approved dye or colorant that imparts an unnatural color to the seed**

SOIL TREATMENT USE DIRECTIONS

Regalia® Bioprotectant Concentrate can be applied by soil drench in furrow spray or soil injection to improve plant health and to protect against certain soil borne diseases

In general Regalia® Bioprotectant Concentrate can be applied by the following methods unless specified differently in the SELECTED CROPS section

Soil Drench Applications

Apply Regalia® Bioprotectant Concentrate at a concentration of 1 3 quarts per 100 gallons of water and at a sufficient rate to thoroughly soak the growing media and root zone Make an initial application of Regalia® Bioprotectant Concentrate during or shortly after transplant to reduce transplant shock suppress soil borne diseases and improve root growth Multiple drench applications can be made on a 10 14 day interval

Shanked In and Injected Applications

Regalia® Bioprotectant Concentrate can be shanked in or injected into the soil alone or with most types of liquid nutrients

In Furrow Applications

At planting apply Regalia® Bioprotectant Concentrate as an in furrow spray at the rate of 1 4 quarts per acre or 2 2 8 8 fluid ounces per 1000 feet of row according to the chart below Apply Regalia® Bioprotectant Concentrate in 5 15 gallons of water so as the spray is directed into the seed furrow just before the seeds are covered

Rate	In Furrow Application Rates Product per Acre (fl oz)					
	30 Rows	32 Rows	34 Rows	36 Rows	38 Rows	40 Rows
2 2 fl oz per 1000 ft row	38 3	36 0	33 8	32 0	30 3	28 7
8 8 fl oz per 1000 ft row	153 2	144 0	135 2	128 0	121 2	114 8

30 = 17 424 row ft /acre 32 = 16 315 row ft /acre 34 = 15 374 row ft /acre 36 = 14 520 row ft /acre 38 = 13 754 row ft /acre 40 = 13 068 row ft /acre

APPLICATION RATES FOR SELECTED CROPS

Regalia® Bioprotectant Concentrate used as specified will improve plant health and induce the defense system of the treated plants listed below towards the diseases specified below

The general recommended use rate for Regalia® Bioprotectant Concentrate applied alone or as an alternate spray is 2 4 quarts per 100 gallons of water (0.5 1.0% v/v dilution of Regalia® Bioprotectant Concentrate concentrate) applied at 50 100 gallons of water per acre. When tank mixed with another fungicide the use rate for Regalia is 1 4 quarts in 100 gallons of water applied at 50 100 gallons of water per acre. Use higher water volumes with larger sized crops and extensive foliage in order to secure thorough coverage. See specific application recommendations pertaining to each crop for additional details.

For greenhouse application on the crops and diseases listed the recommended use rate for Regalia® Bioprotectant Concentrate is 2 4 quarts in 100 gallons of water (0.5 1.0% v/v dilution of Regalia® Bioprotectant Concentrate) sprayed until just before point of runoff. When tank mixed with another fungicide the use rate for Regalia® Bioprotectant Concentrate is 1 4 quarts in 100 gallons of water. Repeat at 7 14 day intervals as needed. See specific application recommendations for each crop for additional details.

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
Artichoke	Powdery Mildew (<i>Erysiphe cichoracearum</i>) (<i>Leveillula taurica</i>)	Foliar (Ground)	1 – 4 quarts per acre	For ground applications apply this product in 50 100 gallons of water per acre. Apply this product preventatively or when the first disease symptoms are visible and reapply every 7 14 days.
		Foliar (Aerial)	0.5 1 quart per acre	For aerial applications apply this product in a minimum of 5 gallons of water per acre. Apply this product preventatively or when the first disease symptoms are visible and reapply every 7 14 days.
	Ramularia Leaf Spot (<i>Ramularia cynarae</i>)	Chemigation	1 – 4 quarts per acre	For chemigation applications for improved plant growth and suppression of soil borne diseases apply this product through drip irrigation immediately after transplant and at 14 day intervals or begin 14 days after transplant when soil drench applications are used.

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI)** of **0 days**
Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI)** of **4 hours**

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
------	----------------	--------------------	----------------------------------	--------------------------

Asparagus	Botrytis Blight (<i>Botrytis cinerea</i>)	Foliar (Ground)	1 – 4 quarts per acre	For ground applications apply this product in 50 100 gallons of water per acre Apply this product preventatively or when the first disease symptoms are visible and reapply every 7 14 days
	Rust (<i>Puccinia asparagi</i>)	Foliar (Aerial)	0.5 – 1 quart per acre	For aerial applications apply this product in a minimum of 5 gallons of water per acre Apply this product preventatively or when the first disease symptoms are visible and reapply every 7 14 days

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI)** of 0 days
Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI)** of 4 hours

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
Bushberries and Caneberries Blueberry Blackberry (all varieties) Cranberry Currant Elderberry Gooseberry Huckleberry Juneberry Ligonberry Loganberry Raspberry (red and black) Salal and other berry crops	Mummy Berry (<i>Monilinia vaccinii-corymbosi</i>) Alternaria Fruit Rot (<i>Alternaria</i> spp) Anthracnose Fruit Rot (<i>Colletotrichum acutatum</i>) Bacterial Canker (<i>Pseudomonas syringae</i>) Botrytis Blight (<i>Botrytis cinerea</i>) Leaf Rust (<i>Pucciniastrum vaccinii</i>) Leaf Spot and Blotch	Foliar (Ground)	1 – 4 quarts per acre	For ground applications apply this product in 50 100 gallons of water per acre Mummy Berry – Initiate application at bud break stage of development Apply this product preventatively and repeat on a 7 10 day interval or as needed For best performance tank mix this product with other registered fungicides for Mummy Berry control Botrytis Blight – Apply this product preventatively when the first disease symptoms are visible and reapply every 7 14 days Bacterial Canker – Apply this product prior to Fall rains and repeat applications during dormancy before Spring growth This product can be tank mixed with another registered fungicide for improved control of bacterial canker Anthracnose Fruit Rot and Alternaria Fruit Rot on blueberries – Initiate application at green tip and continue applications on a 7 10 day

	<p><i>(Mycosphaerella spp)</i> <i>(Septoria spp)</i></p> <p>Phomopsis Leaf Spot Twig Blight and Fruit Rot <i>(Phomopsis spp)</i></p> <p>Powdery Mildew <i>(Microsphaera alni)</i></p> <p>Spur Blight <i>(Didymella spp)</i> <i>(Phoma spp)</i></p>	<p>Foliar (Aerial)</p>	<p>0.5 - 1 quart per acre</p>	<p>For aerial applications apply this product in a minimum of 5 gallons of water per acre</p>
--	--	----------------------------	-------------------------------	---

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI) of 0 days**
Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI) of 4 hours**

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
<p>Bulb Vegetables</p> <p>Onion (Bulb and Green) Garlic Leek Shallot and other bulb vegetable crops</p>	<p>Botrytis Leaf Blight <i>(Botrytis squamosa)</i></p> <p>Botrytis Neck Rot <i>(Botrytis spp)</i></p> <p>Downy Mildew <i>(Peronospora spp)</i></p> <p>Onion Downy Mildew <i>(Peronospora destructor)</i></p> <p>Onion Purple Blotch <i>(Alternaria porri)</i></p> <p>Powdery Mildew <i>(Erysiphe spp)</i></p> <p>Rust</p>	<p>Foliar</p>	<p>1 - 4 quarts per acre</p>	<p>For foliar applications apply this product preventatively in 50 - 100 gallons of water per acre</p> <p>Repeat applications at 7 - 14 day intervals</p> <p>Under moderate to heavy disease pressure tank mix this product with another fungicide</p>

	<p><i>(Puccinia porri)</i></p> <p>Stemphyllium Leaf Blight <i>(Stemphyllium vesicarium)</i></p>			
	<p><i>Fusarium</i> spp</p> <p><i>Pythium</i> spp</p> <p><i>Rhizoctonia</i> spp</p>	Soil Drench	1 – 3 quarts per 100 gallons	For soil drench applications apply this product at a concentration of 1 – 3 quarts per 100 gallons of water and at a sufficient rate to thoroughly soak the growing media and root zone. Make an initial application of this product during or shortly after transplant to reduce transplant shock suppress soil borne diseases and improve root growth. Multiple drench applications can be made on a 10 – 14 day interval.
		In Furrow	1 – 4 quarts per acre 2.2 – 8.8 fl oz per 1000 ft row	For in furrow applications at planting apply this product as an in furrow spray at the rate of 1 – 4 quarts per acre or 2.2 – 8.8 fluid ounces per 1000 feet of row according to the chart in the SOIL TREATMENT USE DIRECTIONS section. Apply this product in 5 – 15 gallons of water so as the spray is directed into the seed furrow just before the seeds are covered.
		Chemigation	1 – 4 quarts per acre	For chemigation applications apply this product through irrigation at the rate of 1 – 4 quarts per acre immediately after transplant and at 14 day intervals or begin 14 days after transplant when plant dip or soil drench applications are used.
		Plant Dip	1 – 4 quarts per 100 gallons	For plant dip applications for improved plant growth and suppression of soil borne diseases apply this product in a 0.25 – 1% v/v suspension (1 – 4 quarts this product per 100 gallons water) as a pre plant dip immediately prior to transplanting.
		Seed Treatment	1.5 – 2.5 fl oz per 100 lbs seed	For suppression of soil borne diseases apply this product as a seed treatment at the rate of 1.5 – 2.5 fl oz per 100 lbs seed.

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI)** of **0 days**
Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI)** of **4 hours**

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
<p>Cereal Grains Barley Buckwheat Grain Amaranth Milo Oat Millets Rice Rye Sorghum Triticale Wheat and other cereal grain crops</p>	<p>Powdery Mildew (<i>Erysiphe graminis</i>) Bacterial Blight and Streak (<i>Xanthomonas</i> spp) Brown Rot Leaf Spots & Smuts (<i>Ceratobasidium</i> spp) (<i>Cercospora</i> spp) (<i>Cochliobolus</i> spp) (<i>Drechslera</i> spp)</p>	<p>Foliar (Ground)</p>	<p>1 – 2 quarts per acre</p>	<p>For ground applications to optimize disease control and to maximize yields apply this product in 15 – 40 gallons of water per acre It is important to apply this product at the flag leaf stage to maximize yield Apply this product preventatively or when the first disease symptoms appear Repeat applications in 7 – 14 day intervals depending upon crop growth and disease pressure When the plants are under high disease pressure tank mix this product with another fungicide for more effective control</p>
	<p>Rice Blast (<i>Pyricularia grisea</i>) Rust (<i>Puccinia</i> spp) Septoria Leaf Spot (<i>Septoria</i> spp) Sheath Spot and Blight (<i>Rhizoctonia oryzae</i>) (<i>Thanatephorus cucumeris</i>) Stem Rot (<i>Sclerotium oryzae</i>) Smut (<i>Tilletia barclayana</i>)</p>	<p>Foliar (Aerial)</p>	<p>0.5 – 1 quart per acre</p>	<p>For aerial applications apply this product in a minimum of 5 gallons water per acre It is important to apply this product at the flag leaf stage to maximize yield Apply this product preventatively or when the first disease symptoms appear Repeat applications in 7 – 14 day intervals depending upon crop growth and disease pressure When the plants are under high disease pressure tank mix this product with another registered fungicide for more effective control</p>

	<i>Fusarium</i> spp <i>Phytophthora</i> spp <i>Pythium</i> spp <i>Rhizoctonia</i> spp <i>Verticillium</i> spp	Seed Treatment	1.5 – 2.5 fl oz per 100 lbs seed	For suppression of soil borne diseases apply this product as a seed treatment at the rate of 1.5 – 2.5 fl oz per 100 lbs seed
--	---	----------------	----------------------------------	---

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI) of 0 days**
 Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI) of 4 hours**

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
Citrus Crops Orange Grapefruit Lemon Tangelo Tangerine Pummelo and other citrus crops	Bacterial Canker (<i>Xanthomonas</i> spp)	Foliar (Ground)	1 – 4 quarts per acre	For ground applications apply this product preventatively in 50-100 gallons of water per acre
	Alternaria Brown Spot (<i>Alternaria alternata</i>) Bacterial Blast (<i>Pseudomonas syringae</i>) Black Spot (<i>Guignardia citricarpa</i>) (<i>Phyllosticta citricarpa</i>) Greasy Spot (<i>Mycosphaerella citri</i>)			For improved performance use this product in a tank mix or rotational program with other registered fungicides Repeat applications at 7-14 day intervals Dilute applications this product can be applied by ground equipment to tree crops in dilute applications of 100-400 gallons of water. Apply this product at a rate of 2-4 quarts per acre when applied alone or at 1-4 quarts per acre when tank mixed with another fungicide. Avoid excessive amounts of water that result in the runoff of spray material.
	Melanose (<i>Diaporthe citri</i>) Postbloom Fruit Drop (<i>Colletotrichum acutatum</i>) Scab (<i>Elsinoe</i>)	Foliar (Aerial)	0.5 – 1 quart per acre	For aerial applications apply this product in a minimum of 5 gallons water per acre For improved performance use this product in a tank mix or rotational program with other registered fungicides Repeat applications at 7-14 day intervals

	<i>australis</i> (<i>Elsinoe fawcetti</i>)			
	<i>Fusarium</i> spp <i>Phytophthora</i> spp <i>Pythium</i> spp <i>Rhizoctonia</i> spp	Plant Dip	1 4 quarts per 100 gallons	For plant dip applications for improved plant growth and suppression of soil borne diseases apply this product in a 0.25 – 1 % v/v suspension (1 4 quarts this product per 100 gallons water) as a pre plant dip immediately prior to transplanting

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI)** of **0 days**
Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI)** of **4 hours**

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
Cole Crops (Brassicacae) Broccoli Broccoli Rabe Brussels Sprouts Cabbage Chinese Broccoli Chinese Cabbage (Bok Choy) Chinese Mustard Cabbage (Gai Choy) Cauliflower Cavalo Collards Kale Kohlrabi Mizuna Mustard Greens	Powdery Mildew (<i>Erysiphe cruciferarum</i>) (<i>Erysiphe polygona</i>) Alternaria Leaf Spot (<i>Alternaria</i> spp) Downy Mildew (<i>Peronospora parasitica</i>) Pin Rot Complex (<i>Alternaria/Xanthomonas</i>) Xanthomonas Leaf Spot (<i>Xanthomonas campestris</i>)	Foliar (Ground)	0.5 – 4 quarts per acre	For ground applications apply this product at 1 4 quarts per 50 gallons of water For concentrated ground applications apply this product at 0.5 – 1.5 quarts per acre in 10 25 gallons of water per acre Repeat applications at 7 14 day intervals Under moderate to heavy disease pressure tank mix this product with another fungicide
			Foliar (Aerial)	0.5 – 1.5 quarts per acre

Mustard Spinach Rape Greens Turnip and other cole crops	<i>Fusarium</i> spp	Seed Treatment	1.5 – 2.5 fl oz per 100 lbs seed	For suppression of soil borne diseases apply this product as a seed treatment at the rate of 1.5 – 2.5 fl oz per 100 lbs seed
	<i>Phytophthora</i> spp			
	<i>Pythium</i> spp			
	<i>Rhizoctonia</i> spp			
	<i>Verticillium</i> spp			

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI)** of 0 days
Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI)** of 4 hours

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
Corn Sweet Corn Field Corn Popcorn Silage Corn Seed Corn and other corn crops (includes crops grown for seed)	Anthracnose Leaf Blight (<i>Colletotrichum graminicola</i>)	Foliar (Ground)	0.5 – 2 quarts per acre	For ground applications to optimize disease control and to maximize yields apply 1 – 2 quarts of this product preventatively in 15 – 40 gallons of water per acre prior to disease development using sufficient volume for thorough coverage For improved performance apply 0.5 – 2 quarts this product in a tank mix with another registered fungicide. Consult your local Extension Specialist or Crop Consultant regarding the optimum timing of fungicide applications
	Eye Spot (<i>Aureobasidium zeae</i>)			
	Gray leafspot (<i>Cercospora zeae maydis</i>)			
	Rusts (<i>Puccinia</i> spp)	Foliar (Aerial)	0.5 – 1 quart per acre	For aerial applications apply this product in a minimum of 3 gallons of water per acre
	Northern Leaf Blight (<i>Exserohilum turcicum</i>)			
	Northern Leaf Spot (<i>Cochliobus carbonum</i>)			
Southern Leaf Blight (<i>Cochliobolus heterostrophus</i>)				

19/65

<i>Fusarium</i> spp <i>Phytophthora</i> spp <i>Pythium</i> spp <i>Rhizoctonia</i> spp <i>Verticillium</i> spp	Seed Treatment	1.5 – 2.5 fl oz per 100 lbs seed	For suppression of soil borne diseases apply this product as a seed treatment at the rate of 1.5 – 2.5 fl oz per 100 lbs seed
---	----------------	----------------------------------	---

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI)** of **0 days**
 Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI)** of **4 hours**

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
Cotton	Alternaria Leaf Spot Boll Rot (<i>Alternaria</i> spp) Anthracnose Boll Rot (<i>Glomeria</i> spp)	Foliar (Ground)	1 – 2 quarts per acre	For ground applications for foliar and Boll Rot disease control apply this product preventatively in 15 – 40 gallons of water per acre prior to disease development using sufficient volume for thorough coverage Repeat applications at 7 – 14 day intervals
	Ascochyta Blight Boll Rot (<i>Ascochyta</i> spp) Cercospora Blight and Leaf Spot (<i>Cercospora</i> spp) Diplodia Boll Rot (<i>Diplodia</i> spp) Hard Lock Boll Rot (<i>Fusarium</i> spp) Leaf Spot (<i>Corynespora cassicola</i>) Phoma Blight	Foliar (Aerial)	0.5 – 1 quart per acre	For aerial applications apply this product in a minimum of 3 gallons of water per acre

	Boll Rot (<i>Phoma</i> spp)			
	Rust (<i>Puccinia</i> spp) (<i>Phykopsora</i> spp)			
	Stemphyllium Leaf Spot (<i>Stemphyllium</i> spp)			
	<i>Fusarium</i> spp <i>Phytophthora</i> spp <i>Pythium</i> spp <i>Rhizoctonia</i> spp <i>Verticillium</i> spp	In Furrow	1 – 4 quarts per acre 2.2 8 8 fl oz per 1000 ft row	For in furrow applications at planting apply this product as an in furrow spray at the rate of 1 4 quarts per acre or 2.2 8 8 fluid ounces per 1000 feet of row according to the chart in the SOIL TREATMENT USE DIRECTIONS section Apply this product in 5 15 gallons of water so as the spray is directed into the seed furrow just before the seeds are covered
		Seed Treatment	7.5 – 15 fl oz per 100 lbs seed	For suppression of soil borne diseases apply this product as a seed treatment at the rate of 7.5 – 15 fl oz per 100 lbs seed

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI)** of 0 days
Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI)** of 4 hours

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
------	----------------	--------------------	----------------------------------	--------------------------

<p>Cucurbits Includes all types and hybrids of Chayote Chinese waxgourd Cucumber Citron melon Gherkin Pumpkin Watermelon</p> <p>Edible Gourd Chinese okra Cucuzza Hyotan</p> <p>Mormordica spp Balsam apple Balsam pear Bitter melon Chinese cucumber</p>	<p>Powdery Mildew (<i>Erysiphe cichoracearum</i>) (<i>Sphaerotheca fuliginea</i>)</p> <p>Anthracnose (<i>Colletotrichum lagenarium</i>)</p> <p>Alternaria Blight (<i>Alternaria cucumerina</i>)</p> <p>Cercospora Leaf Spot (<i>Cercospora citrulina</i>)</p> <p>Damping off (<i>Fusarium spp</i>) (<i>Pythium spp</i>) (<i>Phytophthora sp</i>) (<i>Rhizoctonia solani</i>)</p> <p>Downy Mildew (<i>Pseudoperonospora cubensis</i>)</p>	<p>Foliar (Ground)</p>	<p>1 – 4 quarts per acre</p>	<p>For ground applications apply this product preventatively in 25 100 gallons of water per acre or when the first symptoms of disease are visible Increase water volume as plant size increases</p> <p>Repeat applications in 7 14 day intervals depending upon crop growth and disease pressure</p> <p>When greenhouse cucurbits are under high disease conditions use the shorter spray interval</p> <p>Downy Mildew Tank mix this product with another fungicide labeled for Downy Mildew control and re apply at a 7 day interval or according to the label directions of the tank mix partner</p> <p>Phytophthora Blight Apply this product in combination with labeled rates of a copper fungicide or with another fungicide labeled for Phytophthora Blight control</p>
<p>Muskmelon Cantaloupe Casaba Crenshaw melon Golden pershaw melon Honeydew melon Honey balls Mango melon Persian melon Pineapple melon Santa Claus melon Snake melon</p> <p>Summer Squash Crookneck squash Scallop squash Straightneck squash</p>	<p>Gummy Stem Blight (<i>Didymella bryoniae</i>)</p> <p>Phytophthora Blight (<i>Phytophthora capsici</i>)</p>	<p>Foliar (Aerial)</p>	<p>0.5 – 1 quart per acre</p>	<p>For aerial applications apply this product in a minimum of 5 gallons of water per acre</p> <p>Repeat applications in 7 14 day intervals depending upon crop growth and disease pressure</p> <p>Downy Mildew Tank mix this product with another fungicide labeled for Downy Mildew control and re apply at a 7 day interval or according to the label directions of the tank mix partner</p> <p>Phytophthora Blight Apply this product in combination with labeled rates of a copper fungicide or with another fungicide labeled for Phytophthora Blight control</p>
<p>Vegetable marrow Zucchini</p> <p>Winter</p>	<p><i>Fusarium spp</i> <i>Phytophthora spp</i> <i>Pythium spp</i></p>	<p>Soil Drench</p>	<p>1 – 3 quarts per 100 gallons</p>	<p>For soil drench applications apply this product at a concentration of 1 3 quarts per 100 gallons of water and at a sufficient rate to thoroughly soak the growing media and root zone Make an</p>

<p>Squash Acorn squash Butternut squash Calabaza Hubbard squash Spaghetti squash and other cucurbit crops</p>	<p><i>Rhizoctonia</i> spp <i>Verticillium</i> spp</p>			<p>initial application of this product during or shortly after transplant to reduce transplant shock suppress soil borne diseases and improve root growth Multiple drench applications can be made on a 10 - 14 day interval</p>
		In Furrow	<p>1 - 4 quarts per acre 2.2 - 8.8 per 1000 ft row</p>	<p>For in furrow applications at planting apply this product as an in furrow spray at the rate of 1 - 4 quarts per acre or 2.2 - 8.8 fluid ounces per 1000 feet of row according to the chart in the SOIL TREATMENT USE DIRECTIONS section Apply this product in 5 - 15 gallons of water so as the spray is directed into the seed furrow just before the seeds are covered</p>
		Plant Dip	<p>1 - 4 quarts per 100 gallons</p>	<p>For plant dip applications for improved plant growth and suppression of soil borne diseases apply this product in a 0.25 - 1% v/v suspension (1 - 4 quarts this product per 100 gallons water) as a pre plant dip immediately prior to transplanting</p>
		Chemigation	<p>1 - 4 quarts per acre</p>	<p>For chemigation applications for improved plant growth and suppression of soil borne diseases apply this product through drip irrigation at the rate of 1 - 4 quarts per acre immediately after transplant and at 14 day intervals or begin 14 days after transplant when plant dip or soil drench applications are used</p>
		Seed Treatment	<p>1.5 - 2.5 fl oz per 100 lbs seed</p>	<p>For suppression of soil borne diseases apply this product as a seed treatment at the rate of 1.5 - 2.5 fl oz per 100 lbs seed</p>

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI)** of 0 days
Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI)** of 4 hours

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
------	----------------	--------------------	----------------------------------	--------------------------

Fruiting Vegetables Tomato Pepper Eggplant Ground Cherry Okra Tomatillo and other fruiting vegetable crops	Bacterial Blight (<i>Xanthomonas</i> spp)	Foliar (Ground)	1 – 3 quarts per acre	For ground applications apply this product preventatively in 25 – 100 gallons of water per acre Increase water volume as plant size increases Repeat applications at 7 – 10 day intervals Tank mix this product with other registered fungicides for improved disease control under heavy pressure Phytophthora Blight Apply this product in combination with labeled rates of a copper fungicide or with another fungicide labeled for Phytophthora Blight control
	Bacterial Spot (<i>Xanthomonas</i> spp)			
	Bacterial Speck (<i>Pseudomonas syringae</i>) Black Mold (<i>Alternaria alternata</i>) Damping off (<i>Fusarium</i> spp) (<i>Pythium</i> spp) (<i>Rhizoctonia solani</i>)			
	Early Blight (<i>Alternaria solani</i>) Gray Mold (<i>Botrytis cinerea</i>) Late Blight(<i>Phytophthora infestans</i>) Phytophthora Blight (<i>Phytophthora capsici</i>) Powdery Mildew (<i>Erysiphe</i> spp) (<i>Leveillula taurica</i>) (<i>Oidopsis taurica</i>) (<i>Sphaerotheca</i> spp) Target Spot (<i>Corynespora cassicola</i>)	Foliar (Aerial)	0.5 – 1 quart per acre	For aerial applications apply this product in a minimum of 10 gallons of water per acre Repeat applications at 7 – 10 day intervals Tank mix this product with other registered fungicides for improved disease control under heavy pressure Phytophthora Blight Apply this product in combination with labeled rates of a copper fungicide
	<i>Fusarium</i> spp <i>Phytophthora</i> spp <i>Pythium</i> spp <i>Rhizoctonia</i> spp <i>Verticillium</i> spp	Soil Drench	1 – 3 quarts per 100 gallons	For soil drench applications apply this product at a concentration of 1 – 3 quarts per 100 gallons of water and at a sufficient rate to thoroughly soak the growing media and root zone Make an initial application of this product during or shortly after transplant to reduce transplant shock suppress soil borne diseases and improve root growth Multiple drench applications can be made on a 10 – 14 day interval

		In Furrow	1 – 4 quarts per acre 2.2 – 8.8 per 1000 ft row	For in furrow applications at planting apply this product as an in furrow spray at the rate of 1 – 4 quarts per acre or 2.2 – 8.8 fluid ounces per 1000 feet of row according to the chart in the SOIL TREATMENT USE DIRECTIONS section. Apply this product in 5 – 15 gallons of water so as the spray is directed into the seed furrow just before the seeds are covered.
		Plant Dip	1 – 4 quarts per 100 gallons	For plant dip applications for improved plant growth and suppression of soil borne diseases apply this product in a 0.25 – 1% v/v suspension (1 – 4 quarts this product per 100 gallons water) as a pre plant dip immediately prior to transplanting.
		Chemigation	1 – 4 quarts per acre	For chemigation applications for improved plant growth and suppression of soil borne diseases apply this product through drip irrigation at the rate of 1 – 4 quarts per acre immediately after transplant and at 14 day intervals or begin 14 days after transplant when plant dip or soil drench applications are used.
		Seed Treatment	1.5 – 2.5 fl oz per 100 lbs seed	For suppression of soil borne diseases apply this product as a seed treatment at the rate of 1.5 – 2.5 fl oz per 100 lbs seed.

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI)** of 0 days
Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI)** of 4 hours

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
------	----------------	--------------------	----------------------------------	--------------------------

<p>Grape</p>	<p>Powdery Mildew (<i>Uncinula necator</i>)</p> <p>Angular Leaf Spot (<i>Mycosphaerella angulata</i>)</p> <p>Anthracnose (<i>Elsinoe ampelina</i>)</p> <p>Botrytis Bunch Rot (<i>Botrytis cinerea</i>)</p> <p>Black Rot (<i>Guignardia bidwellii</i>)</p> <p>Downy Mildew (<i>Plasmopara viticola</i>)</p> <p>Eutypa (<i>Eutypa lata</i>)</p> <p>Leaf Blight (<i>Pseudocercospora vitis</i>)</p> <p>Phomopsis Fruit Rot (<i>Phomopsis viticola</i>)</p> <p>Ripe Rot (<i>Colletotrichum gloeosporioides</i>)</p> <p>Sour Rot (<i>Alternaria tenuis</i>) (<i>Aspergillus spp</i>) (<i>Botrytis cinerea</i>) (<i>Cladosporium herbarum</i>) (<i>Penicillium spp</i>) (<i>Rhizopus arrhizus</i>)</p>	<p>Foliar</p>	<p>1 – 4 quarts per acre</p>	<p>For ground applications apply this product preventatively in 50 100 gallons of water per acre or when the first disease symptoms are visible</p> <p>Under high disease pressure use in a tank mix with another registered fungicide for more effective control</p> <p>Repeat applications in 7 14 day intervals depending upon crop growth and disease pressure</p> <p>Dilute applications this product can be applied by ground equipment to vine and tree crops in dilute applications of 100 400 gallons of water Apply this product at a rate of 2 4 quarts per acre when applied alone or at 1 4 quarts per acre when tank mixed with another fungicide Avoid excessive amounts of water that result in the runoff of spray material</p>
	<p><i>Phytophthora</i> spp</p> <p><i>Verticillium</i> spp</p>	<p>Plant Dip</p>	<p>1 4 quarts per 100 gallons</p>	<p>For plant dip applications for improved plant growth and suppression of soil borne diseases apply this product in a 0.25 – 1% v/v suspension (1 4 quarts this product per 100 gallons water) as a pre plant dip immediately prior to transplanting</p>

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI)** of 0 days
 Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI)** of 4 hours

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
Grass Seed	Powdery Mildew (<i>Erysiphe graminis</i>) (<i>Oidium</i> spp) (<i>Podosphaera</i> spp) (<i>Sphaerotheca</i> spp)	Foliar (Ground)	1 – 4 quarts per acre	For ground applications apply this product preventatively in 25 – 100 gallons of water per acre when disease symptoms are first visible or when environmental conditions are conducive to rapid disease development Continue sprays at 7 day intervals or as needed
	Rust (<i>Puccinia</i> spp)	Foliar (Aerial)	0.5 – 1 quart per acre	For aerial applications apply this product in a minimum of 5 gallons of water per acre
	<i>Fusarium</i> spp <i>Phytophthora</i> spp <i>Pythium</i> spp <i>Rhizoctonia</i> spp <i>Verticillium</i> spp	Seed Treatment	1.5 – 2.5 fl oz per 100 lbs seed	For suppression of soil borne diseases apply this product as a seed treatment at the rate of 1.5 – 2.5 fl oz per 100 lbs seed

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI)** of 0 days
 Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI)** of 4 hours

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
Hops	Downy Mildew (<i>Pseudoperonospora humuli</i>) Powdery Mildew (<i>Sphaerotheca macularis</i>)	Foliar	1 – 4 quarts per acre	Apply this product preventatively when disease symptoms are first visible or when environmental conditions are conducive to rapid disease development Continue sprays at 7 day intervals or as needed Minimum spray volumes for hop growth stages are as follows

				<p>Emergence to Training Apply 1 2 quarts this product per acre using a minimum spray volume of 20 gallons per acre Coverage will vary with the size of the vines and the type of spray equipment Apply adequate spray volume to achieve complete spray coverage</p> <p>Training to Wire Touch Apply 1 2 quarts this product per acre using a minimum spray volume of 50 gallons per acre Coverage will vary with the size of the vines and the type of spray equipment Apply adequate spray volume to achieve complete spray coverage</p> <p>Wire Touch through Harvest Apply 2 4 quarts of this product using a minimum of 100 gallons of water per acre Higher water volumes may be necessary to achieve thorough coverage after side arms develop Do not apply more than 4 quarts of product per acre per application Apply adequate spray volume to achieve complete spray coverage Use the higher rates when moderate to high disease pressure is present or expected</p> <p>For control of downy mildew tank mix this product with another fungicide labeled for Downy Mildew control and re apply at a 7 day interval or according to the label directions of the tank mix partner</p>
--	--	--	--	--

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI) of 0 days**
 Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI) of 4 hours**

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
Leafy Vegetable Crops Arugula Beet Celery Chervil Cilantro	Downy Mildew (<i>Bremia lactuca</i>) (<i>Peronospora</i> spp) Bacterial Blight/Rot (<i>Xanthomonas</i> spp)	Foliar (Ground)	0.5 – 4 quarts per acre	For ground applications apply this product at 0.5 – 4 quarts in 50 – 100 gallons of water per acre For concentrated ground applications apply this product at 0.5 – 1.5 quarts per acre in a minimum of 10 gallons of water per acre

Corn Salad Cress Dandelion Dock Edible Chrysanthemum Endive Fennel Garden Peas Head Lettuce Leaf Lettuce Parsley Purslane Radicchio Rhubarb Spinach Swiss Chard Watercress and other leafy vegetable crops	Cercospora leafspot (<i>Cercospora</i> spp) Late Blight (<i>Septoria apicola</i>)			Repeat applications at 7 - 14 day intervals
	Pink Rot (<i>Sclerotinia sclerotiorum</i>) Powdery Mildew (<i>Erysiphe cichoracearum</i>) Sclerotinia Head and Leaf Drop (<i>Sclerotinia minor</i>) (<i>Sclerotinia sclerotiorum</i>) White Rust (<i>Albugo occidentalis</i>)	Foliar (Aerial)	0.5 - 1.5 quarts per acre	West of the Rocky Mountains For aerial applications apply this product at 0.5 - 1.5 quarts per acre in a minimum of 10 gallons of water per acre East of the Rocky Mountains - For aerial applications apply this product at 0.5 - 1 quarts per acre in a minimum of 5 gallons of water per acre For California For aerial application apply Regalia at 1.3 pints per acre in 10-20 gallons of water per acre Repeat applications at 7 - 14 day intervals
	<i>Fusarium</i> spp <i>Phytophthora</i> spp <i>Pythium</i> spp <i>Rhizoctonia</i> spp <i>Verticillium</i> spp	Seed Treatment	1.5 - 2.5 fl oz per 100 lbs seed	For suppression of soil borne diseases apply this product as a seed treatment at the rate of 1.5 - 2.5 fl oz per 100 lbs seed

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI) of 0 days**
Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI) of 4 hours**

Restrictions

Regalia® Biofungicide Concentrate should be applied to healthy actively growing plants. Do not apply Regalia® Biofungicide Concentrate to plants that are stressed due to cold weather, drought, excessive moisture, etc. Do not apply when extended cold or cold and cloudy conditions are expected.

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
------	----------------	--------------------	----------------------------------	--------------------------

Legumes/Vegetables (not including soybeans and peanuts) Chick Peas Dry Beans Garbanzo Beans Green Beans Lentils Lima Beans Peas Shell Beans Snap Beans Split Peas and other legume crops (including those grown for seed or oil production)	Bacterial Blight <i>(Xanthomonas campestris)</i> Gray Mold <i>(Botrytis cinerea)</i> Pythium (aerial blight phase) <i>(Pythium spp)</i> Powdery Mildew <i>(Erysiphe spp)</i> Rust <i>(Puccinia spp)</i> <i>(Uromyces appendiculatus)</i> White Mold <i>(Sclerotinia sclerotiorum)</i>	Foliar	1 – 4 quarts per acre	For foliar applications apply this product preventatively in 20 100 gallons of water per acre For improved performance use this product in a tank mix or rotational program with another registered fungicide Repeat applications at 7 14 day intervals
	<i>Fusarium spp</i> <i>Phytophthora spp</i> <i>Pythium spp</i> <i>Rhizoctonia spp</i>	In Furrow	1 – 4 quarts per acre 2 2 8 8 per 1000 ft row	For in furrow applications at planting apply this product as an in furrow spray at the rate of 1 4 quarts per acre or 2 2 8 8 fluid ounces per 1000 feet of row according to the chart in the SOIL TREATMENT USE DIRECTIONS section Apply this product in 5 15 gallons of water so as the spray is directed into the seed furrow just before the seeds are covered
		Seed Treatment	1 5 – 2 5 fl oz per 100 lbs seed	For suppression of soil borne diseases apply this product as a seed treatment at the rate of 1 5 – 2 5 fl oz per 100 lbs seed

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI)** of 0 days
 Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI)** of 4 hours

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
------	----------------	--------------------	----------------------------------	--------------------------

Mint and Other Herbs/Spices Angelica Balm Basil Borage Burnet Chamomile Catnip Chervil Chive Clary Coriander Costmary Cilantro Curry Dillweed Horehound Hyssop Lavender Lemongrass Lovage Marjoram Nasturtium Parsley (dried) Peppermint Rosemary Sage Savory (summer and winter) Sweet Bay Tansy Tarragon Thyme Wintergreen Woodruff Wormwood and other herbs/spices	Downy Mildew (<i>Peronospora</i> spp)	Foliar (Ground)	1 – 4 quarts per acre	For ground applications apply this product preventatively in a minimum of 50 gallons of water per acre Repeat applications at 7 t 14 day intervals
	Powdery Mildew (<i>Erysiphe</i> spp)	Rust (<i>Puccinia menthae</i>)	Foliar (Aerial)	0.5 – 1 quart per acre

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI)** of 0 days
 Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI)** of 4 hours

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
------	----------------	--------------------	----------------------------------	--------------------------

Oil Seed Crops (not including cotton peanut or soybean) Canola Castor Flax Rapeseed Safflower Sesame Sunflower And other oilseed crops	Bacterial Pustule (<i>Xanthomonas</i> spp) Bacterial Speck (<i>Pseudomonas syringae pv glycinea</i>) Brown Spot (<i>Septoria glycines</i>) Cercospora Leaf Spot (<i>Cercospora</i> spp)	Foliar (Ground)	0.5 – 2 quarts per acre	For ground applications to optimize disease control and to maximize yields apply this product preventatively in 15-40 gallons of water per acre For improved performance apply this product in a tank mix program with another registered fungicide Consult your local Extension Specialist or Crop Consultant regarding the optimum timing of fungicide applications
	Downy Mildew (<i>Peronospora manshurica</i>) Pod and Stem Blight (<i>Diaporthe phaseolorum var sojæ</i>) (<i>Phomopsis longicola</i>) White Mold/Sclerotinia Stem Rot (<i>Sclerotinia sclerotiorum</i>)	Foliar (Aerial)	0.5 – 1 quart per acre	For aerial applications apply this product in a minimum of 3 gallons per acre For improved performance apply this product in a tank mix program with another registered fungicide Consult your local Extension Specialist or Crop Consultant regarding the optimum timing of fungicide applications
	<i>Fusarium</i> spp <i>Phytophthora</i> spp <i>Pythium</i> spp <i>Rhizoctonia</i> spp <i>Verticillium</i> spp	Seed Treatment	1.5 – 2.5 fl oz per 100 lbs seed	For suppression of soil borne diseases apply this product as a seed treatment at the rate of 1.5 – 2.5 fl oz per 100 lbs seed

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI)** of 0 days
 Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI)** of 4 hours

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
------	----------------	--------------------	----------------------------------	--------------------------

<p>Olive</p>	<p>Olive Knot (<i>Pseudomonas savastanoi</i>)</p>	<p>Foliar</p>	<p>1 – 4 quarts per acre</p>	<p>Apply this product preventatively in 50 100 gallons of water per acre</p> <p>Repeat applications at 7 14 day intervals</p> <p>Dilute applications this product can be applied by ground equipment to tree crops in dilute applications of 100 400 gallons of water Apply this product at a rate of 2 4 quarts per acre when applied alone or at 1 4 quarts per acre when tank mixed with another fungicide Avoid excessive amounts of water that result in the runoff of spray material</p>
---------------------	---	---------------	------------------------------	--

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI)** of **0 days**
 Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI)** of **4 hours**

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
<p>Peanut</p>	<p>Aspergillus Crown Rot (<i>Aspergillus niger</i>)</p> <p>Damping off (<i>Aspergillus flavus</i>) (<i>Fusarium</i> spp) (<i>Pythium</i> spp) (<i>Rhizoctonia</i> spp)</p> <p>Early Leaf Spot (<i>Cercospora arachidicola</i>)</p> <p>Late Leaf Spot (<i>Cerosporidium personatum</i>)</p> <p>Rhizoctonia Foliar Blight Peg and Root Rot (<i>Rhizoctonia solani</i>)</p> <p>White Mold (<i>Sclerotium rolfsii</i>)</p>	<p>Foliar</p>	<p>1 – 4 quarts per acre</p>	<p>For foliar applications apply this product preventatively in 20 50 gallons of water per acre</p> <p>Repeat applications at 7 14 day intervals</p> <p>Tank mix this product with another fungicide labeled for the target disease</p>

	Aspergillus Crown Rot (<i>Aspergillus niger</i>) <i>Fusarium</i> spp <i>Phytophthora</i> spp <i>Pythium</i> spp <i>Rhizoctonia</i> spp <i>Verticillium</i> spp	Soil Drench	1 – 3 quarts per 100 gallons	For soil drench applications apply this product at a concentration of 1 3 quarts per 100 gallons of water and at a sufficient rate to thoroughly soak the growing media and root zone Make an initial application of this product during or shortly after transplant to reduce transplant shock suppress soil borne diseases and improve root growth Multiple drench applications can be made on a 10 14 day interval
	White Mold (<i>Sclerotium rolfsii</i>)	In Furrow	1 – 4 quarts per acre 2 2 8 8 per 1000 ft row	For in furrow applications at planting apply this product as an in furrow spray at the rate of 1 4 quarts per acre or 2 2 8 8 fluid ounces per 1000 feet of row according to the chart in the SOIL TREATMENT USE DIRECTIONS section Apply this product in 5 15 gallons of water so as the spray is directed into the seed furrow just before the seeds are covered
		Seed Treatment	1 5 – 2 5 fl oz per 100 lbs seed	For suppression of soil borne diseases apply this product as a seed treatment at the rate of 1 5 – 2 5 fl oz per 100 lbs seed

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI) of 0 days**
Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI) of 4 hours**

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
Pome Fruits Apple Crabapple Loquat Oriental Pear Pear Quince Mayhaw and other pome fruit crops	Powdery Mildew (<i>Podosphaera leucotricha</i>) Alternaria Blotch (<i>Alternaria mali</i>) Apple Scab (<i>Venturia inaequalis</i>) Suppression only Bitter Rot (<i>Colletotrichum</i> spp)	Foliar	1 – 4 quarts per acre	For foliar applications apply this product in 50 100 gallons of water per acre Begin applications when conditions are conducive to disease development but not prior to petal fall Repeat applications on 7 10 day intervals Additional sprays beyond second cover may be needed on susceptible varieties or when environmental conditions are conducive to rapid disease development Use high label rate and shorter spray intervals when conditions are conducive to rapid disease development Fire Blight – For suppression apply 1 2

<p>Black Rot/Frogeye Leaf Spot (<i>Botryosphaeria obtusa</i>)</p> <p>Bot Rot (<i>Botryosphaeria dothidea</i>)</p> <p>Brooks Spot (<i>Mycosphaerella pomi</i>)</p> <p>Bull s Eye Rot (<i>Neofabraea spp</i>)</p> <p>Cedar Apple Rust (<i>Gymnosporangium juniperi virginianae</i>) Suppression only</p> <p>Fire Blight (<i>Erwinia amylovora</i>) Suppression only</p> <p>Flyspeck (<i>Zygophiala jamaicensis</i>)</p> <p>Scab (<i>Venturia spp</i>)</p> <p>Sooty Blotch (<i>Geastrum polystigmati</i>) (<i>Leptodontium elatus</i>) (<i>Peltaster fructicola</i>)</p> <p>White Rot (<i>Botryosphaeria dothidea</i>)</p>				<p>quarts of this product in 50 100 gallons of water per acre beginning at petal fall For maximum control use this product prior to infection events During periods of rapid development and frequent infection periods use spray intervals of 3 7 days</p> <p>Apply in sufficient water to provide full coverage For improved performance use this product in a rotational program with antibiotics registered for Fire Blight control such as but not limited to oxytetracycline or streptomycin</p> <p>Proper orchard cultural practices are essential to eliminate Fire Blight infected tissue from the orchard to assure good performance of any crop protection product Care must be taken to remove and destroy dead and diseased wood from the orchard prior to and during the growing season</p> <p>Scab – For suppression apply 1 quart of this product in 50 100 gallons of water per acre at green tip and through bloom when environmental conditions become favorable for primary Scab development and repeat on a 7 10 day interval or as needed Use this product in a tank mix or rotational program with other fungicides labeled for Scab control Following bloom this product can be applied at 2 4 quarts per acre</p> <p>Use caution when selecting spray adjuvants Select only those adjuvants which through prior experience do not affect fruit finish when combined with this product</p> <p>Dilute applications this product can be applied by ground equipment to tree crops in dilute applications of 100 400 gallons of water Apply this product at a rate of 2 4 quarts per acre when applied alone or at 1 4 quarts per acre when tank mixed with another fungicide Avoid excessive amounts of water that result in the runoff of spray material</p>
---	--	--	--	---

	<i>Phytophthora</i> spp <i>Pythium</i> spp	Plant Dip (bare root)	1 - 4 quarts per 100	For plant dip applications for improved plant growth and suppression of soil borne diseases apply this product in a 0.25 - 1% v/v suspension (1 - 4 quarts this product per 100 gallons water) as a pre plant dip immediately prior to transplanting
--	---	--------------------------	-------------------------	--

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI)** of 0 days
Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI)** of 4 hours

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
Root Tuber and Corm Crops Potato Beet Carrot Cassava Ginger Ginseng Horseradish Radish Sweet Potato Turnip and other root crops (including those for seed production)	Bacterial Leaf Blight <i>(Xanthomonas campestris)</i>	Foliar	1 - 4 quarts per acre	For foliar applications apply this product in 25 - 100 gallons of water per acre sufficient to provide thorough coverage Begin application soon after emergence or transplant and when conditions are conducive to disease development Repeat on a 7 to 10 day interval or as needed Use shorter intervals when conditions are conducive to rapid disease development For suppression of Early Blight Black Root Rot/Black Crown Rot and Late Blight begin application of this product in 25 - 100 gallons of water per acre soon after emergence when conditions are conducive to disease development Repeat on a 5 to 7 day interval or as needed For improved performance use this product in a tank mix with other registered fungicides
	Black Root Rot / Black Crown Rot <i>(Alternaria spp)</i>			
	Downy Mildew <i>(Peronospora spp)</i>			
	Early Blight <i>(Alternaria solani)</i>			
	Gray Mold <i>(Botrytis spp)</i>			
	Late Blight <i>(Phytophthora infestans)</i>			
	Powdery Mildew <i>(Erysiphe spp)</i>			
White Mold <i>(Sclerotinia sclerotiorum)</i>				

	Clubroot (<i>Plasmiodiophora brassicae</i>)	Soil Drench	1 – 3 quarts per 100 gallons	For soil drench applications apply this product at a concentration of 1 – 3 quarts per 100 gallons of water and at a sufficient rate to thoroughly soak the growing media and root zone. Make an initial application of this product during or shortly after transplant to reduce transplant shock, suppress soil borne diseases and improve root growth. Multiple drench applications can be made on a 10 – 14 day interval.
	Common Scab (<i>Streptomyces scabies</i>)			
	<i>Fusarium</i> spp			
	<i>Phytophthora</i> spp			
	<i>Pythium</i> spp			
	<i>Rhizoctonia</i> spp	In Furrow	1 – 4 quarts per acre 2.2 – 8.8 per 1000 ft row	For in furrow applications at planting apply this product as an in furrow spray at the rate of 1 – 4 quarts per acre or 2.2 – 8.8 fluid ounces per 1000 feet of row according to the chart in the SOIL TREATMENT USE DIRECTIONS section. Apply this product in 5 – 15 gallons of water so as the spray is directed into the seed furrow just before the seeds are covered.
	<i>Verticillium</i> spp			
		Seed Piece Dip	1 – 4 quarts per 100 gallons of water	For seed piece dip applications for improved plant growth and suppression of soil borne diseases apply this product in a 0.25 – 1% v/v suspension (1 – 4 quarts this product per 100 gallons water) as a pre plant dip to transplants or seed pieces immediately prior to transplanting.
		Chemigation	1 – 4 quarts per acre	For chemigation applications for improved plant growth and suppression of soil borne diseases apply this product through drip irrigation at the rate of 1 – 4 quarts per acre immediately after transplant and at 14 day intervals or begin 14 days after transplant when plant dip or soil drench applications are used.
		Seed Treatment	1.5 – 2.5 fl oz per 100 lbs seed	For suppression of soil borne diseases apply this product as a seed treatment at the rate of 1.5 – 2.5 fl oz per 100 lbs seed.

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI)** of 0 days
Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI)** of 4 hours

Crop	Target Disease	Application	Application
------	----------------	-------------	-------------

		Method	Product Use Rate per Application	Instructions
Soybean	Aerial Web Blight (<i>Rhizoctonia solani</i>) Alternaria Leafspot (<i>Alternaria</i> spp) Anthracnose (<i>Colletotrichum truncatum</i>) Asian Soybean Rust (<i>Phakopsora pachyrhizi</i>)	Foliar (Ground)	0.5 - 2 quarts per acre	For ground applications to optimize disease control and to maximize yields apply 1 - 2 quarts of this product preventatively in 15 - 40 gallons of water per acre For improved performance apply 0.5 - 2 quarts of this product in a tank mix with another registered fungicide Consult your local Extension Specialist or Crop Consultant regarding the optimum timing of fungicide applications
	Brown Spot (<i>Septoria glycines</i>) Cercospora Blight (<i>Cercospora kikuchii</i>) Frog eyed Leaf Spot (<i>Cercospora sojina</i>) Pod and Stem Blight (<i>Diaporthe</i> spp) Septoria Brown Spot (<i>Septoria glycines</i>) White Mold (<i>Sclerotinia sclerotiorum</i>)	Foliar (Aerial)	0.5 - 1 quart per acre	For aerial applications apply this product in a minimum of 3 gallons of water per acre For improved performance apply this product in a tank mix with another registered fungicide Consult your local Extension Specialist or Crop Consultant regarding the optimum timing of fungicide applications
	<i>Fusarium</i> spp <i>Phytophthora</i> spp <i>Pythium</i> spp <i>Rhizoctonia</i> spp	In Furrow	1 - 4 quarts per acre 2.2 - 8.8 per 1000 ft row	For in furrow applications at planting apply this product as an in furrow spray at the rate of 1 - 4 quarts per acre or 2.2 - 8.8 fluid ounces per 1000 feet of row according to the chart in the SOIL TREATMENT USE DIRECTIONS section Apply this product in 5 - 15 gallons of water so as the spray is directed into the seed furrow just before the seeds

				are covered
		Seed Treatment	1.5 – 2.5 fl oz per 100 lbs seed	For suppression of soil borne diseases apply this product as a seed treatment at the rate of 1.5 – 2.5 fl oz per 100 lbs seed

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI)** of **0 days**
 Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI)** of **4 hours**

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
Stone Fruits Apricot Cherry (sweet and tart) Nectarine Peach Plum Plumcot Prune and other stone fruit crops	Alternaria Spot/Fruit Rot (<i>Alternaria alternata</i>) Anthracnose (<i>Colletotrichum</i> spp) Bacterial Canker (<i>Pseudomonas</i> spp) Bacterial Spot (<i>Xanthomonas pruni</i>) Brown Rot Blossom Blight (<i>Monilinia laxa</i>) Brown Rot Fruit Rot (<i>Monilinia fruticola</i>) Cercospora Leaf Spot (<i>Cercospora</i> spp) Cherry Leaf Spot (<i>Blumeriella jaapi</i>) Gray Mold	Foliar	1 – 4 quarts per acre	For foliar applications apply this product preventatively in 50 – 100 gallons of water per acre Bacterial Blight – Apply this product in 50 – 100 gallons of water per acre postharvest before Fall rains Brown Rot Blossom Blight – Begin application of this product in 50 – 100 gallons of water per acre at early bloom and repeat through petal fall on a 7 day interval or as needed Powdery Mildew – Begin application of this product in 50 – 100 gallons of water per acre at popcorn stage and repeat on a 7 day interval or as needed For improved performance use this product in a tank mix or rotational program with other registered fungicides for powdery mildew control Scab – Begin application of this product in 50 – 100 gallons of water per acre at petal fall and repeat on a 7 to 10 day interval or as needed For improved performance tank mix this product with another fungicide labeled for Scab control For all other diseases – Begin application prior to disease development when environmental conditions and plant stage are conducive to rapid disease

	<p>(<i>Botrytis cinerea</i>)</p> <p>Powdery Mildew (<i>Podosphaera</i> spp) (<i>Sphaerotheca pannosa</i>)</p> <p>Rust (<i>Tranzschelia discolor</i>)</p> <p>Rusty Spot (<i>Podosphaera leucotricha</i>)</p> <p>Scab (<i>Cladosporium carpophilum</i>)</p> <p>Shot Hole (<i>Wilsonomyces carpophilus</i>)</p>			<p>development and repeat on a 7 - 10 day interval or as needed Use in a tank mix or rotational program when disease conditions are severe</p> <p>Dilute applications this product can be applied by ground equipment to tree crops in dilute applications of 100 - 400 gallons of water Apply this product at a rate of 2 - 4 quarts per acre when applied alone or at 1 - 4 quarts per acre when tank mixed with another fungicide Avoid excessive amounts of water that result in the runoff of spray material</p>
	<p><i>Fusarium</i> spp</p> <p><i>Phytophthora</i> spp</p> <p><i>Pythium</i> spp</p> <p><i>Rhizoctonia</i> spp</p> <p><i>Verticillium</i> spp</p>	<p>Plant Dip (bare root)</p>	<p>1 - 4 quarts per 100 gallons of water</p>	<p>For plant dip applications for improved plant growth and suppression of soil borne diseases apply this product in a 0.25 - 1% v/v suspension (1 - 4 quarts this product per 100 gallons water) as a pre plant dip immediately prior to transplanting</p>

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI)** of **0 days**
 Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI)** of **4 hours**

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
Strawberry	<p>Anthracoese (<i>Collectotrichum</i> spp) Suppression only</p> <p>Botrytis (<i>Botrytis cinerea</i>)</p> <p>Leaf Spot</p>	Foliar	1 - 3 quarts per acre	<p>For foliar applications apply this product preventatively in 50 - 100 gallons of water per acre at 7 - 14 day spray intervals or as soon as first symptoms of disease appear</p> <p>Anthracoese For suppression apply this product preventatively in 50 - 100 gallons of water per acre and repeat on a</p>

	<p><i>(Mycosphaerella fragariae)</i></p> <p>Phomopsis Leaf Blight <i>(Phomopsis obscurans)</i></p> <p>Powdery Mildew <i>(Sphaerotheca macularis)</i></p>			<p>7 to 10 day interval or as needed For best performance tank mix this product with other registered fungicides for Anthracnose control</p> <p>Dilute applications this product can be applied by ground equipment to strawberries in dilute applications of 100 to 200 gallons of water Apply this product at a rate of 2 to 3 quarts per acre when applied alone or at 1 to 3 quarts per acre when tank mixed with another fungicide Avoid excessive amounts of water that result in the runoff of spray material</p>
	<p>Black Root Rot <i>(Rhizoctonia spp)</i> <i>(Pythium spp)</i> <i>(Fusarium spp)</i> <i>(Cylindrocarpon spp)</i></p> <p>Colletotrichum Crown Rot <i>(Colletotrichum spp)</i></p> <p>Phytophthora Root Rot and Crown Rot <i>(Phytophthora spp)</i></p> <p>Verticillium Wilt <i>(Verticillium spp)</i></p> <p><i>Fusarium spp</i></p> <p><i>Pythium spp</i></p> <p><i>Phytophthora spp</i></p> <p><i>Rhizoctonia spp</i></p> <p><i>Verticillium spp</i></p>	Plant Dip	1 to 4 quarts per 100 gallons of water	For plant dip applications for improved plant growth and suppression of soil borne diseases apply this product in a 0.25 to 1% v/v suspension (1 to 4 quarts per 100 gallons water) as a pre plant dip to strawberry plants roots and crowns immediately prior to transplanting
		Soil Drench	1 to 3 quarts per 100 gallons	For soil drench applications apply this product at a concentration of 1 to 3 quarts per 100 gallons of water and at a sufficient rate to thoroughly soak the growing media and root zone Make an initial application of this product during or shortly after transplant to reduce transplant shock suppress soil borne diseases and improve root growth Multiple drench applications can be made on a 10 to 14 day interval
		Chemigation	1 to 4 quarts per acre	For chemigation applications for improved plant growth and suppression of soil borne diseases apply this product through drip irrigation at the rate of 1 to 4 quarts per acre immediately after transplant and at 14 day intervals or begin 14 days after transplant when plant dip or soil drench applications are used

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI)** of 0 days
Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI)** of 4 hours

Crop	Target Disease	Application	Application
------	----------------	-------------	-------------

		Method	Product Use Rate per Application	Instructions
Sugar Beets (includes crop for seed production)	Powdery Mildew (<i>Erysiphe betae</i>) (<i>Erysiphe polygoni</i>) Leaf Spot (<i>Cercospora beticola</i>) Ramularia (<i>Ramularia</i> spp) Rust (<i>Uromyces betae</i>)	Foliar	1 – 2 quarts per acre	To optimize disease control and to maximize yields apply this product preventatively in 15 – 40 gallons of water per acre by ground or aerial application For improved performance apply this product in a tank mix program with another registered fungicide Consult your local Extension Specialist or Crop Consultant regarding the optimum timing of fungicide applications
	<i>Fusarium</i> spp <i>Phytophthora</i> spp <i>Pythium</i> spp <i>Rhizoctonia</i> spp <i>Verticillium</i> spp	Seed Treatment	1.5 – 2.5 fl oz per 100 lbs seed	For suppression of soil borne diseases apply this product as a seed treatment at the rate of 1.5 – 2.5 fl oz per 100 lbs seed

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI)** of 0 days
 Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI)** of 4 hours

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
Sugarcane	Brown Rust (<i>Puccinia melanocephala</i>) Orange Rust (<i>Puccinia kuehni</i>)	Foliar (Ground)	1 – 2 quarts per acre	For ground applications to optimize disease control and to maximize yields apply this product preventatively in 15 – 40 gallons of water per acre by ground application Consult your local Extension Specialist or Crop Consultant regarding the optimum timing of fungicide applications For improved performance apply this

				product in a tank mix program with another registered fungicide
		Foliar (Aerial)	0.5 - 1 quart per acre	For aerial applications apply this product in a minimum of 3 gallons of water per acre Consult your local Extension Specialist or Crop Consultant regarding the optimum timing of fungicide applications For improved performance apply this product in a tank mix program with another registered fungicide
	<i>Fusarium</i> spp <i>Phytophthora</i> spp <i>Pythium</i> spp <i>Rhizoctonia</i> spp	In Furrow	1 - 4 quarts per acre 2.2 - 8.8 per 1000 ft row	For in furrow applications at planting apply this product as an in furrow spray at the rate of 1 - 4 quarts per acre or 2.2 - 8.8 fluid ounces per 1000 feet of row according to the chart in the SOIL TREATMENT USE DIRECTIONS section Apply this product in 5 - 15 gallons of water so as the spray is directed into the seed furrow just before the seeds are covered
		Seed Treatment	1.5 - 2.5 fl oz per 100 lbs seed	For suppression of soil borne diseases apply this product as a seed treatment at the rate of 1.5 - 2.5 fl oz per 100 lbs seed
<p>Regalia® Bioprotectant Concentrate has a pre harvest interval (PHI) of 0 days Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours</p>				

Crop	Target Disease	Application Method	Product Use	Application Instructions
------	----------------	--------------------	-------------	--------------------------

			Rate per Application	
Tobacco	Blue Mold (<i>Peronospora tabacina</i>)	Foliar	1 – 4 quarts per acre	For foliar applications apply this product at a rate of 2 – 4 quarts per acre when applied alone or at 1 – 4 quarts per acre when tank mixed with another fungicide preventatively in a minimum of 50 gallons of water per acre Avoid excessive amounts of water that result in spray material dripping from the foliage If necessary repeat applications at a 7 day interval
	<i>Fusarium</i> spp <i>Phytophthora</i> spp <i>Pythium</i> spp <i>Rhizoctonia</i> spp <i>Verticillium</i> spp	Plant Dip	1 – 4 quarts per 100 gallons	For plant dip applications for improved plant growth and suppression of soil borne diseases apply this product in a 0.25 – 1% v/v suspension (1 – 4 quarts per 100 gallons water) as a pre plant dip to tobacco roots and plants immediately prior to transplanting

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI)** of 0 days
Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI)** of 4 hours

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
Tree Nut Crops Walnut (Black and English) Almond Beech nut Brazil nut Butternut Cashew Chestnut Chinquapin Filbert Hickory nut Macadamia nut Pecan Pistachio and other tree	Walnut Blight (<i>Xanthomonas campestris</i>) Alternaria Late Blight Alternaria Leaf Spot (<i>Alternaria</i> spp) Anthracnose (<i>Collectotrichum</i> spp) (<i>Gnomonia leptostyla</i>) Bacterial Canker	Foliar (Ground)	1 – 4 quarts per acre	For ground applications apply this product in 50 – 100 gallons of water per acre this product can be tank mixed at the lower rate with another registered fungicide under heavy disease pressure Walnut Blight For preventative control apply this product in 50 – 100 gallons of water per acre Repeat applications at 7 to 10 day intervals Under conditions of heavy disease pressure tank mix this product with a copper based fungicide Dilute applications this product can be applied by ground equipment to tree

nut crops	(<i>Erwinia nigrifluens</i>) (<i>Pseudomonas syringae</i>) Botryosphaeria Blight (<i>Botryosphaeria dothidea</i>)			crops in dilute applications of 100 - 400 gallons of water. Apply this product at a rate of 2 - 4 quarts per acre when applied alone or at 1 - 4 quarts per acre when tank mixed with another fungicide. Avoid excessive amounts of water that result in the runoff of spray material.
	Brown Rot (<i>Monilinia</i> spp) Eastern Filbert Blight (<i>Anisogramma anomala</i>) Green Fruit Rot (<i>Botrytis cinerea</i>) Leaf Rust (<i>Tranzschelia discolor</i>) Scab (<i>Cladosporium carpophilum</i>) (<i>Sphaceloma perseae</i>) Shot Hole (<i>Wilsonomyces carpophilus</i>)	Foliar (Aerial)	0.5 - 1 quart per acre	For aerial applications apply this product in a minimum of 10 gallons per acre.
	<i>Fusarium</i> spp <i>Phytophthora</i> spp <i>Pythium</i> spp <i>Rhizoctonia</i> spp <i>Verticillium</i> spp	Plant Dip (bare root)	1 - 4 quarts per 100 gallons of water	For plant dip applications for improved plant growth and suppression of soil borne diseases apply this product in a 0.25 - 1% v/v suspension (1 - 4 quarts this product per 100 gallons water) as a pre plant dip immediately prior to transplanting.

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI)** of 0 days
Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI)** of 4 hours

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
------	----------------	--------------------	----------------------------------	--------------------------

Tropical Fruits Avocado Banana Kiwi Mango Papaya Plantain Pineapple Pomegranate and other tropical fruit crops	Anthracnose (<i>Colletotrichum gloeosporioides</i>)	Foliar (Ground)	1 – 4 quarts per acre	For ground applications apply this product preventatively in 50 100 gallons of water per acre Repeat applications at 7 t 14 day intervals Dilute applications this product can be applied by ground equipment to tree crops in dilute applications of 100 400 gallons of water Apply this product at a rate of 2 4 quarts per acre when applied alone or at 1 4 quarts per acre when tank mixed with another fungicide Avoid excessive amounts of water that result in the runoff of spray material Sigatoka Initiate applications when leaves first appear and repeat on a 7 10 day schedule Apply in sufficient water by ground or air to obtain thorough coverage of foliage For improved disease control this product may be tank mixed with oil or other fungicides registered for Sigatoka control at label rates
	Bacterial Blight (<i>Pseudomonas syringae</i>) Bacterial Canker (<i>Xanthomonas campestris</i>) Botrytis Fruit Rot (<i>Botrytis cinerea</i>) Scab (<i>Elsinoe mangiferae</i>) Sigatoka (<i>Mycosphaerella fijiensis</i>)			Foliar (Aerial)
	<i>Fusarium</i> spp <i>Phytophthora</i> spp <i>Pythium</i> spp <i>Rhizoctonia</i> spp <i>Verticillium</i> spp	Plant Dip	1 4 quarts per 100 gallons of water	For plant dip applications for improved plant growth and suppression of soil borne diseases apply this product in a 0.25 – 1% v/v suspension (1 4 quarts per 100 gallons water) as a pre plant dip immediately prior to transplanting

Regalia® Bioprotectant Concentrate has a **pre harvest interval (PHI) of 0 days**
 Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI) of 4 hours**

INTEGRATED PEST MANAGEMENT (IPM)

Many conventional fungicides have been tested in an IPM regime with Regalia® Bioprotectant Concentrate with very satisfactory results. One of the major objectives of IPM has been to reduce the probability of disease resistance development to a particular active ingredient.

The alternate use of (1-2 sprays) followed by a conventional registered fungicide (1-2 sprays) has been successfully used in many crops. In addition, the use of tank mixes with a conventional fungicide has also been successful.

Follow label instructions of the particular registered product. Do not exceed amounts or treatment intervals on the label.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place. Avoid freezing.

Pesticide Disposal: To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

Container Handling (under 5 gallons): Non-refillable container. Do not reuse or refill this container. Triple-rinse container (or equivalent) promptly after emptying. Triple-rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill or by incineration. Do not burn unless allowed by state and local ordinances.

Container Handling (over 5 gallons): Non-refillable container. Do not reuse or refill this container. Triple-rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill or by incineration. Do not burn unless allowed by state and local ordinances.

Marrone Bio Innovations is a member of the Ag Container Recycling Council.
Visit <http://www.acrecycle.org/contact.html> for information on how to arrange pick up of this empty pesticide container.

WARRANTY

To the extent consistent with applicable law, the seller makes no warranty, expressed or implied, of merchantability, fitness, or otherwise concerning use of this product. The user assumes all risks of use, storage, or handling that are not in strict accordance with the accompanying directions.

Label date

Made in the U.S.A.

US Patents No. 4,863,734 and No. 5,989,429

Regalia® is a trademark of Marrone Bio Innovations, Inc.

Marrone Bio Innovations name and logo are registered trademarks of Marrone Bio Innovations, Inc.

© Marrone Bio Innovations, Inc.

2121 Second St., Suite B 107, Davis, CA 95618

47/65

1 877 664 4476
www.marronebio.com
info@marronebio.com

Sublabel B Turf & Professional Landscape Use

Regalia® Bioprotectant Concentrate

A plant extract to boost the plants' defense mechanisms to protect against certain fungal and bacterial diseases and to improve plant health

Active ingredient	Extract of <i>Reynoutria sachalinensis</i>	5 %
Other ingredients		95 %
Total		100 %

EPA Reg No 84059 3

EPA Est No 085970 FL 001
 EPA Est No 047857 CA 001
 EPA Est No 61842 CA 001

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID	
IF SWALLOWED	Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth to mouth if possible. Call a poison control center or doctor for further treatment advice.
IF IN EYES	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or if going for treatment.	



LOT # _____

Net Contents 1 pint 1 quart 1 gallon 2.5 gallon 5 gallon 55 gallon drum
 Marrone Bio Innovations Inc 2121 Second St Suite B 107 Davis CA 95618

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION Causes moderate eye irritation Avoid contact with eyes or clothing Wear goggles or safety glasses Wash thoroughly with soap and water after handling and before eating drinking chewing gum using tobacco or using the toilet

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear

Long sleeved shirt and long pants

Shoes plus socks

Waterproof gloves

Protective eyewear

Follow manufacturer s instructions for cleaning and maintaining PPE If no such instructions for washables use detergent and hot water Keep and wash PPE separately from other laundry

ENVIRONMENTAL HAZARDS

For terrestrial uses Do not apply directly to water to areas where surface water is present or to intertidal areas below the mean high water mark Do not contaminate water when disposing of equipment washwater or rinsate

(Use the following additional statement for containers of 5 gallons or more Do not discharge effluent containing this product into lakes streams ponds estuaries oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority For guidance contact your State Water Board or Regional Office of the EPA)

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling Do not apply this product in a way that will contact workers or other persons either directly or through drift Only protected handlers may be in the area during application For any requirements specific to your State or Tribe consult the State or Tribal agency responsible for pesticide regulation

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170 This standard contains requirements for the protection of agricultural workers on farms forests nurseries and greenhouses and handlers of agricultural pesticides It contains requirements for training decontamination notification and emergency assistance It also contains specific instructions and exemptions pertaining to the statements on this label about personal protective equipment (PPE) and the restricted entry interval (REI) The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants soil or water is

Coveralls

Waterproof gloves

Shoes plus socks

Protective eyewear

NON AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are **not** within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170) The WPS applies when this product is used to produce agricultural plants on farms forests nurseries or greenhouses

Keep unprotected persons out of treated areas until sprays have dried

GENERAL INFORMATION

Regalia® Bioprotectant Concentrate is an extract from the plant *Reynoutria* spp for use on ornamental plants and turf Regalia® Bioprotectant Concentrate applied to actively growing plants (see DIRECTIONS FOR USE) will improve plant health and will help make the treated portions resistant to certain plant diseases Use Regalia® Bioprotectant Concentrate as a preventative rather than a curative application Apply prior to disease infestation to protect the growing leaf tissue See specific information for diseases controlled and use rates on ornamental plants and turf

Regalia® Bioprotectant Concentrate can be used as a seed treatment plant dip soil drench in furrow spray or applied through drip irrigation to control or suppress certain soil borne diseases and to promote healthy root growth See specific information for diseases controlled and use rates on treating seeds with Regalia® Bioprotectant Concentrate

MODE OF ACTION

The extract obtained from *Reynoutria* spp plant material contains active chemical compounds The extract when applied to the host plant increases the plant s defense system due to a five fold increase in phenolics and antioxidants and strengthens cell walls This induced resistance against important diseases is not systemic but provides some translaminar protection Repeat foliar applications at 7 to 14 day intervals to maintain induction and to protect new plant growth The resistance induction takes place within one to two days

Use Regalia® Bioprotectant Concentrate therefore as a preventative treatment

MIXING AND APPLICATION INSTRUCTIONS

- SHAKE WELL PRIOR TO USE -

Regalia® Bioprotectant Concentrate is a micro emulsion concentrate consisting of certain ingredients extracted from *Reynoutria* spp Use 50 mesh nozzle screens or larger

See **AERIAL APPLICATION** section for aerial application use directions

See **CHEMIGATION** section for chemigation use directions

See **PRE PLANT DIP** section for pre plant dip use directions

See **SEED TREATMENT** section for seed treatment use directions

See **SOIL TREATMENT** section for soil application use directions

Use higher water volumes with larger sized crops and extensive foliage to secure thorough coverage

Regalia® Bioprotectant Concentrate alone Add ½ of the required amount of water to the mix tank With the agitator running add the Regalia® Bioprotectant Concentrate to the mix tank Continue agitation while adding the remainder of the water Begin application of the solution after the Regalia® Bioprotectant Concentrate has completely dispersed into the mix water Maintain agitation until all the mixture has been applied

Regalia® Bioprotectant Concentrate + tank mixtures Add ½ ¼ of the required amount of water to the mix tank Start the agitation before adding any tank mix partners In general tank mix partners should be added in this order wettable powders dry flowable formulations liquid flowable formulations and emulsifiable formulations such as Regalia® Bioprotectant Concentrate Always allow each tank mix partner to become completely dispersed before adding the next component Maintain continuous agitation until all components have been dispersed and throughout the application process After all components are completely dispersed add the remainder of the water Regalia® Bioprotectant Concentrate cannot be mixed with another product with a prohibition against mixing Use of the tank mix must be in accordance with the more restrictive label limitations and precautions **Do not pre mix Regalia® Bioprotectant Concentrate with any other tank mix component prior to adding to the spray tank**

Compatibility Do not combine Regalia® Bioprotectant Concentrate in the spray tank with pesticides, adjuvants, or fertilizers if there has been no previous experience or use of the combination to show it is physically compatible, effective, and non-injurious under your use conditions.

Regalia® Bioprotectant Concentrate is compatible with many commonly used pesticides, fertilizers, adjuvants, and surfactants, but has not been evaluated with all potential combinations. To ensure compatibility of the tank mix combinations, evaluate prior to use as follows: Using a suitable container, add the proportional amounts of product to water. Add wettable powders first, then water dispersible granules, then liquid flowables, and lastly emulsifiable concentrates. Mix thoroughly and let stand for at least five minutes. If the combination stays mixed or can be remixed, it is physically compatible. Test the mix on a small portion of the crop to be treated to ensure that a phytotoxic response will not occur as a result of the application.

AERIAL APPLICATION INSTRUCTIONS

Apply Regalia® Bioprotectant Concentrate by aerial application to the Edible Crops listed at the rate of 0.5 – 1 quart per acre in a minimum of 5 gallons of water per acre unless specified differently in the APPLICATION RATES section. Increasing the amount of water applied per acre may improve product performance. Follow all instructions to reduce aerial drift.

AERIAL DRIFT REDUCTION ADVISORY INFORMATION

GENERAL Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. Where states have more stringent regulations, they should be observed. Note: This section is advisory in nature and does not supersede the mandatory label requirements.

INFORMATION ON DROPLET SIZE The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply droplets large enough to provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

CONTROLLING DROPLET SIZE Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets. Pressure – Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When high flow rates are needed, use higher flow rate nozzles instead of increasing pressure. Number of Nozzles – Use the minimum number of nozzles that provide uniform coverage. Nozzle Orientation – Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential. Nozzle Type – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

BOOM WIDTH For aerial applications, the boom width must not exceed 75% of the wingspan or 90% of the rotary blade. Use upwind swath displacement and apply only when wind speed is 3-10 mph as measured by an anemometer. Use medium or coarser spray according to ASAE 572 definition for standard nozzles or VMD for spinning atomizer nozzles. If application includes a no-spray zone, do not release spray at a height greater than 10 feet above the ground or crop canopy.

APPLICATION HEIGHT Do not make application at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind

SWATH ADJUSTMENT When applications are made with a crosswind the swath will be displaced downward Therefore on the up and downwind edges of the field the applicator must compensate for this displacement by adjusting the path of the aircraft upwind Swath adjustment distance should increase with increasing drift potential (higher wind smaller drops etc)

WIND Drift potential is lowest between wind speeds of 2 10 mph However many factors including droplet size and equipment type determine drift potential at any given speed Application should be avoided below 2 mph due to variable wind direction and high inversion potential NOTE Local terrain can influence wind patterns Every applicator should be familiar with local wind patterns and how they affect spray drift

TEMPERATURE AND HUMIDITY When making applications in low relative humidity set up equipment to produce larger droplets to compensate for evaporation Droplet evaporation is most severe when conditions are both hot and dry

TEMPERATURE INVERSIONS Do not apply during a temperature inversion because drift potential is high Temperature inversions restrict vertical air mixing which causes small suspended droplets to remain in a concentrated cloud This cloud can move in unpredictable directions due to the light variable winds common during inversions Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind They begin to form as the sun sets and often continue into the morning Their presence can be indicated by ground fog however if fog is not present inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion while smoke that moves upward and rapidly dissipates indicates good vertical air mixing

SENSITIVE AREAS The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e g residential areas bodies of water known habitat for threatened or endangered species non target crops) is minimal (e g when wind is blowing away from the sensitive areas) Do not allow spray to drift from the application site and contact people structures people occupy at any time and the associated property parks and recreation areas non target crops aquatic and wetland areas woodlands pastures rangelands or animals

CHEMIGATION USE DIRECTIONS

Apply Regalia® Bioprotectant Concentrate at 1 4 quarts per acre according to the instructions below unless specified differently in the APPLICATION RATES section

CHEMIGATION

General Requirements

- 1) Apply this product only through a drip system or sprinkler including center pivot lateral move end tow side (wheel) roll traveler big gun solid set hand move flood (basin) furrow border or drip (trickle) irrigation systems Do not apply this product through any other type of irrigation system
- 2) Crop injury lack of effectiveness or illegal pesticide residues in the crop can result from non uniform distribution of treated water
- 3) If you have questions about calibration you should contact State Extension Service specialists equipment manufacturers or other experts
- 4) Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place

- 5) A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise

Specific Requirements for Chemigation Systems Connected to Public Water Systems

- 1) Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year
- 2) Chemigation systems connected to public water systems must contain a functional reduced pressure zone backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe
- 3) The pesticide injection pipeline must contain a functional automatic quick closing check valve to prevent the flow of fluid back toward the injection pump
- 4) The pesticide injection pipeline must contain a functional normally closed solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down
- 5) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or in cases where there is no water pump when the water pressure decreases to the point where pesticide distribution is adversely affected
- 6) Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock
- 7) Do not apply when wind speed favors drift beyond the area intended for treatment

Specific Requirements for Sprinkler Chemigation

- 1) The system must contain a functional check valve vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow
- 2) The pesticide injection pipeline must contain a functional automatic quick closing check valve to prevent the flow of fluid back toward the injection pump
- 3) The pesticide injection pipeline must also contain a functional normally closed solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down
- 4) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops
- 5) The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected
- 6) Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock
- 7) Do not apply when wind speed favors drift beyond the area intended for treatment

Specific Requirements for Flood (Basin) Furrow and Border Chemigation

- 1) Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as a drop structure or weir box to decrease potential for water source contamination from backflow if water flow stops
- 2) The systems utilizing a pressurized water and pesticide injection system must meet the following requirements
 - a) The system must contain a functional check valve vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow
 - b) The pesticide injection pipeline must contain a functional automatic quick closing check valve to prevent the flow of fluid back toward the injection pump

- c The pesticide injection pipeline must also contain a functional normally closed solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down
- d The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops
- e The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected
- f Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock

Specific Requirements for Drip (Trickle) Chemigation

- 1) The system must contain a functional check valve vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow
- 2) The pesticide injection pipeline must contain a functional automatic quick closing check valve to prevent the flow of fluid back toward the injection pump
- 3) The pesticide injection pipeline must also contain a functional normally closed solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down
- 4) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops
- 5) The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected
- 6) Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock

Application Instructions for All Types of Chemigation

- 1) Remove scale pesticide residues and other foreign matter from the chemical supply tank and entire injector system Flush with clean water Failure to provide a clean tank void of scale or residues may cause product to lose effectiveness or strength
- 2) Determine the treatment rates as indicated in the directions for use and make proper dilutions Product can be applied continuously or at any time during the water application
- 3) Prepare a solution in the chemical tank by filling the tank with the required water and then adding product as required

PRE PLANT DIP USE DIRECTIONS

Regalia® Bioprotectant Concentrate can be applied as a pre plant dip for improved plant health and suppression of certain soil borne diseases Apply Regalia® Bioprotectant Concentrate in 1 - 4 quarts product per 100 gallons of water as a pre plant dip immediately prior to transplanting unless specified differently in the APPLICATION RATES section

SEED TREATMENT USE DIRECTIONS

Regalia® Bioprotectant Concentrate can be applied as a seed dressing for suppression of soil borne diseases to improve early season root growth Regalia® Bioprotectant Concentrate may be applied as a water based slurry with other registered seed treatment insecticides and fungicides through standard slurry or mist type commercial seed treatment equipment

Mixing instructions Prepare no more mixture than is required for the immediate operation Agitate the solution continuously during mixing and application Mechanical mixing is recommended for proper mixing of Regalia® Bioprotectant Concentrate mixtures

Regalia® Bioprotectant Concentrate alone Add ½ of the required amount of water to the mix tank With the agitator running add the Regalia® Bioprotectant Concentrate to the mix tank Continue agitation while adding the remainder of the water Begin application of the solution after the Regalia® Bioprotectant Concentrate has completely dispersed into the mix water Maintain agitation until all the mixture has been applied

Regalia® Bioprotectant Concentrate+ tank mixtures Add ½ of the required amount of water to the mix tank Start the agitation before adding any tank mix partners In general tank mix partners should be added in this order wettable powders dry flowable formulations liquid flowable formulations and emulsifiable formulations such as Regalia® Bioprotectant Concentrate Always allow each tank mix partner to become completely dispersed before adding the next component Maintain continuous agitation until all components have been dispersed and throughout the application process

Note When using Regalia® Bioprotectant Concentrate in tank mixtures all products in water soluble packaging should be added to the tank before any other tank mix partner including Regalia® Bioprotectant Concentrate Allow the water soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank mix partner to the tank

If using Regalia® Bioprotectant Concentrate in a tank mixture with other seed treatment products observe all directions for use crops/sites use rates dilution ratios precautions and limitations which appear on the tank mix partner label No label dosage may be exceeded and the most restrictive label precautions and limitations must be followed This product should not be mixed with any product which prohibits such mixing

Note Federal law requires that bags containing treated seeds shall be labeled with the following information This seed has been treated with Regalia® Bioprotectant Concentrate fungicide Do not use for food feed or oil purposes Store away from feed and foodstuffs **Treated seed bagged for later use must contain an EPA approved dye or colorant that imparts an unnatural color to the seed**

SOIL TREATMENT USE DIRECTIONS

Regalia® Bioprotectant Concentrate can be applied by soil drench in furrow spray or soil injection to improve plant health and to protect against certain soil borne diseases

In general Regalia® Bioprotectant Concentrate can be applied by the following methods unless specified differently in the APPLICATION RATES section

Soil Drench Applications

Apply Regalia® Bioprotectant Concentrate at a concentration of 1 3 quarts per 100 gallons of water and at a sufficient rate to thoroughly soak the growing media and root zone Make an initial application of Regalia® Bioprotectant Concentrate during or shortly after transplant to reduce transplant shock suppress soil borne diseases and improve root growth Multiple drench applications can be made on a 10 14 day interval

Shanked In and Injected Applications

Regalia® Bioprotectant Concentrate can be shanked in or injected into the soil alone or with most types of liquid nutrients

In Furrow Applications

At planting apply Regalia® Bioprotectant Concentrate as an in furrow spray at the rate of 1 4 quarts per acre or 2 2 8 8 fluid ounces per 1000 feet of row according to the chart below Apply Regalia® Bioprotectant Concentrate in 5 to 15 gallons of water so as the spray is directed into the seed furrow just before the seeds are covered

Rate	In Furrow Application Rates Product per Acre (fl oz)					
	30 Rows	32 Rows	34 Rows	36 Rows	38 Rows	40 Rows
2 2 fl oz per 1000 ft row	38 3	36 0	33 8	32 0	30 3	28 7

8.8 fl oz per 1000 ft row	153.2	144.0	135.2	128.0	121.2	114.8
---------------------------------	-------	-------	-------	-------	-------	-------

30 = 17 424 row ft /acre 32 = 16 315 row ft /acre 34 = 15 374 row ft /acre 36 = 14 520 row ft /acre 38 = 13 754 row ft /acre 40 = 13 068 row ft /acre

APPLICATION RATES

Regalia® Bioprotectant Concentrate used as specified will improve plant health and induce the defense system of the treated plants listed below towards the diseases specified below

The general recommended use rate for Regalia® Bioprotectant Concentrate applied alone or as an alternate spray is 2.4 quarts per 100 gallons of water (0.5:1.0% v/v dilution of Regalia® Bioprotectant Concentrate concentrate) applied at 50-100 gallons of water per acre. When tank mixed with another fungicide the use rate for Regalia is 1.4 quarts in 100 gallons of water applied at 50-100 gallons of water per acre. Use higher water volumes with larger sized crops and extensive foliage in order to secure thorough coverage. See specific application recommendations for additional details.

For greenhouse application the recommended use rate for Regalia® Bioprotectant Concentrate is 2.4 quarts in 100 gallons of water (0.5:1.0% v/v dilution of Regalia® Bioprotectant Concentrate concentrate) sprayed until just before point of runoff. When tank mixed with another fungicide the use rate for Regalia® Bioprotectant Concentrate is 1.4 quarts in 100 gallons of water. Repeat at 7 to 14 day intervals as needed. See specific application recommendations for additional details.

Crop	Target Disease	Application Method	Product Use Rate per Application	Application Instructions
Ornamentals Herbaceous Ornamentals Flowering Plants Foliage Plants Woody Ornamentals Broadleaves Shrubs and Trees Conifers Shrubs and Trees	Anthracnose (<i>Colletotrichum</i> spp.) Bacteria (<i>Erwinia</i> spp.) (<i>Pseudomonas</i> spp.) (<i>Xanthomonas</i> spp.) Black Spot of Rose (<i>Diplocarpon</i> <i>rosae</i>) Blossom Blight (<i>Monilinia</i> spp.) Downy Mildew (<i>Peronospora</i> spp.) (<i>Plasmopara</i> <i>viburni</i>) Gray Mold (<i>Botrytis cinerea</i>)	Foliar	1 – 4 quarts per acre	For foliar applications mix this product concentrate with water at a concentration of 2.4 quarts per 100 gallons of water when used alone or 1.4 quarts per 100 gallons of water when tank mixed with another fungicide. Begin applications preventatively (before disease symptoms become visible) at the 4 to 6 leaf stage and treat at 7 to 14 day intervals as needed prior to sale or harvest. Spray until just before point of runoff. This product may be used to control certain diseases of container bench flat plug bed or field grown ornamentals in greenhouses shade houses outdoor nurseries retail nurseries and other landscape areas .

<p>Leaf Spot (<i>Alternaria</i> spp) (<i>Cercospora</i> spp) (<i>Entomosporium</i> spp) (<i>Myrothecium</i> spp) (<i>Septoria</i> spp)</p> <p>Powdery Mildew (<i>Erysiphe</i> spp) (<i>Oidium</i> spp) (<i>Podosphaera</i> spp) (<i>Sphaerotheca</i> spp)</p> <p>Rust (<i>Puccinia</i> spp)</p> <p>Scab (<i>Venturia</i> spp)</p>			
<p><i>Fusarium</i> spp <i>Phytophthora</i> spp <i>Pythium</i> spp <i>Rhizoctonia</i> spp <i>Verticillium</i> spp</p>	<p>Soil Drench</p>	<p>1 –3 quarts per 100 gallons</p>	<p>For soil drench applications apply this product at a concentration of 1 – 3 quarts per 100 gallons of water and at a sufficient rate to thoroughly soak the growing media and root zone. Make an initial application of this product during or shortly after transplant to reduce transplant shock, suppress soil borne diseases and improve root growth. Multiple drench applications can be made on a 10-14 day interval.</p>
	<p>Plant Dip</p>	<p>1 – 4 quarts per 100 gallons</p>	<p>For plant dip applications for improved plant growth and suppression of soil borne diseases apply this product in a 0.25 – 1% v/v suspension (1 – 4 quarts per 100 gallons water) as a pre plant dip immediately prior to transplanting.</p>

		Chemigation	1 – 4 quarts per acre	For chemigation applications for improved plant growth apply this product through drip irrigation immediately after transplant and at 14 day intervals or begin 14 days after transplant when soil drench applications are used
--	--	-------------	-----------------------	---

Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI)** of **4 hours**

The following plant species have been treated with Regalia® Bioprotectant Concentrate to prevent disease

Plants investigated

Annual and Perennial Flowering Plants

Begonias Freesias Geraniums Gerbera Impatiens *Lamium* Lisianthus Petunias Poinsettias Roses Salvias Snapdragons Zinnias

Trees and Shrubs

Azalea Boxwood Crape Myrtle Dogwood Indian Hawthorne Jumbo Azalea Lilac Loropetalum Japanese Maple Japanese Privet *Photinia* Rhododendron *Rosaceae* Soft Touch Holly Spirea *Viburnum*

Tropical Foliage

Aglaonema Dieffenbachia Dracaena English Ivy *Hibiscus* Leatherleaf Fern *Spathiphyllum*

Since it is not possible to test all ornamental species or varieties grown in the greenhouse test Regalia® Bioprotectant Concentrate on a few plants prior to large scale usage

Crop	Target Disease	Application Method	Product Use Rate per Application (per 1 000 sq ft)	Product Use Rate per Application (per Acre)	Application Instructions
Turfgrass	Anthracnose (<i>Colletotrichum graminicola</i>)	Foliar	1 – 3 fl oz per 1000 sq ft in a minimum of 1.5 gallons of water	3 – 8 pints per acre in a minimum of 50 gallons of water	This product aids in control of turf diseases and improves turf quality For improved performance under moderate to severe disease pressure reduce spray intervals or use this product in a tank mix or rotational
Bluegrass	Bentgrass/Bermudagrass				
Bentgrass	Dead Spot (<i>Ophiophaerella agrostis</i>)				
Bermudagrass					
Dichondra					
Fescue					
Orchardgrass					
<i>Poa annua</i>	Bermudagrass Decline (<i>Gaeumannomyces graminis</i> var <i>graminis</i>)				
Ryegrass					
St Augustine					
Zoysia					

<p>mixtures and other grasses</p>	<p>Brown patch (<i>Rhizoctonia solani</i>)</p>				<p>program with other registered fungicides</p>
<p>Ornamental Grasses</p>	<p>Copper Spot (<i>Gloeocercospora sorghi</i>)</p>				<p>Begin applications preventatively (before disease symptoms become visible) and treat at 7 14 day intervals as needed Spray water volumes must be of at least 1 5 gallons of water per 1000 sq ft Under moderate to high disease pressure tank mix with other registered fungicides</p>
	<p>Dichondra Rust (<i>Puccinia dichondrae</i>)</p>				
	<p>Dollar Spot (<i>Lanzia spp</i>) (<i>Moellerodiscus spp</i> formerly <i>Sclerotinia homeocarpa</i>)</p>				
	<p>Fusarium Patch (<i>Fusarium nivale</i>)</p>				
	<p>Gray Leaf Spot (<i>Pyricularia grisea</i>)</p>				
	<p>Melting Out Leaf Spot (<i>Bipolaris spp</i>) (<i>Drechslera spp</i>)</p>				
	<p>Necrotic Ring Spot (<i>Leptosphaeria korrae</i>)</p>				
	<p>Pink Patch (<i>Limonomyces roseipellis</i>)</p>				
	<p>Powdery Mildew (<i>Erysiphe graminis</i>)</p>				
	<p>Pythium Blight Pythium Root Rot (<i>Pythium aphanidermatum</i>) (<i>Pythium spp</i>)</p>				
	<p>Red Thread (<i>Laetisaria fuciformis</i>)</p>				
	<p>Rust (<i>Puccinia spp</i>)</p>				
	<p>Rhizoctonia Large Patch (<i>Rhizoctonia solani</i>)</p>				
	<p>Snowmold Gray (<i>Typhula spp</i>)</p>				
	<p>Snowmold Pink (<i>Microdochium nivale</i>)</p>				

	<p>Southern Blight (<i>Sclerotium rolfsii</i>)</p> <p>Spring Dead Spot (<i>Leptosphaeria korrae</i>) (<i>Leptosphaeria narmari</i>) (<i>Ophiosphaerella herpotricha</i>) (<i>Gaeumannomyces graminis</i>)</p> <p>Stripe Smut (<i>Ustilago striiformis</i>) (<i>Urocystis agropyri</i>)</p> <p>Summer Bentgrass Decline</p> <p>Summer Patch Poa Patch (<i>Magnaporthe poae</i>)</p> <p>Take All Patch (<i>Gaeumannomyces graminis</i>)</p> <p>Yellow Patch (<i>Rhizoctonia cerealis</i>)</p> <p>Yellow Tuft/Downy Mildew (<i>Sclerophthora macrospora</i>)</p> <p>Zoysia Patch (<i>Rhizoctonia solani</i>)</p>				
	<p><i>Fusarium</i> spp</p> <p><i>Phytophthora</i> spp</p> <p><i>Pythium</i> spp</p> <p><i>Rhizoctonia</i> spp</p> <p><i>Verticillium</i> spp</p>	<p>Seed Treatment</p>	<p>1.5 – 2.5 fl oz per 100 lbs seed</p>		<p>For suppression of soil borne diseases apply this product as a seed treatment at the rate of 1.5 – 2.5 fl oz per 100 lbs seed</p>

11/65

Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI) of 4 hours**

STORAGE AND DISPOSAL

Do not contaminate water food or feed by storage or disposal

Pesticide Storage Store in a cool dry place Avoid freezing

Pesticide Disposal To avoid wastes use all material in this container by application according to label directions If wastes cannot be avoided offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry)

Container Handling Non refillable container Do not reuse or refill this container Triple rinse container (or equivalent) promptly after emptying Triple rinse as follows Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip Fill the container ¼ full with water and recap Shake for 10 seconds Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal Drain for 10 seconds after the flow begins to drip Repeat this procedure two more times Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration Do not burn unless allowed by state and local ordinances

Container Handling (over 5 gallons) Non refillable container Do not reuse or refill this container Triple rinse as follows Empty the remaining contents into application equipment or a mix tank Fill the container ¼ full with water Replace and tighten closures Tip container on its side and roll it back and forth ensuring at least one complete revolution for 30 seconds Stand the container on its end and tip it back and forth several times Turn the container over onto its other end and tip it back and forth several times Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal Repeat this procedure two more times Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration Do not burn unless allowed by state and local ordinances

Marrone Bio Innovations is a member of the Ag Container Recycling Council
Visit [http //www acrecycle org/contact html](http://www.acrecycle.org/contact.html) for information on how to arrange pick up of this empty pesticide container

WARRANTY

To the extent consistent with applicable law the seller makes no warranty expressed or implied of merchantability fitness or otherwise concerning use of this product The user assumes all risks of use storage or handling that are not in strict accordance with the accompanying directions

Label date

Made in the U S A

US Patents No 4 863 734 and No 5 989 429

Regalia® is a trademark of Marrone Bio Innovations Inc
Marrone Bio Innovations name and logo are registered trademarks of Marrone Bio Innovations Inc

© **Marrone Bio Innovations, Inc**
2121 Second St Suite B 107 Davis CA 95618
1 877 664 4476
www.marronebio.com
info@marronebio.com

62/65

Sublabel C Home & Garden Use

Regalia® Bioprotectant Concentrate

A plant extract to boost the plants defense mechanisms to protect against certain fungal and bacterial diseases and to improve plant health

Active ingredient Extract of *Reynoutria sachalinensis* 5 %
Other ingredients 95 %
Total 100 %

EPA Reg No 84059 3

EPA Est No 085970 FL 001
EPA Est No 047857 CA 001
EPA Est No 61842 CA 001

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID	
IF SWALLOWED	Call poison control center or doctor immediately for treatment advice Have person sip a glass of water if able to swallow Do not induce vomiting unless told to do so by the poison control center or doctor Do not give anything by mouth to an unconscious person
IF ON SKIN OR CLOTHING	Take off contaminated clothing Rinse skin immediately with plenty of water for 15 20 minutes Call a poison control center or doctor for treatment advice
IF INHALED	Move person to fresh air If person is not breathing call 911 or an ambulance then give artificial respiration preferably by mouth to mouth if possible Call a poison control center or doctor for further treatment advice
IF IN EYES	Hold eye open and rinse slowly and gently with water for 15 20 minutes Remove contact lenses if present after the first 5 minutes then continue rinsing eye Call a poison control center or doctor for treatment advice
Have the product container or label with you when calling a poison control center or doctor or if going for treatment	



CAN BE USE IN ORGANIC GARDENING



LOT # _____

Net Contents 1 pint 1 quart 1 gallon 2 5 gallon
Marrone Bio Innovations, Inc 2121 Second St Suite B 107 Davis CA 95618



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal law to apply this product in a manner inconsistent with its labeling.

HOME AND GARDEN USE DIRECTIONS

Regalia® Bioprotectant Concentrate is a broad spectrum fungicide used for the control or suppression of a broad range of foliar diseases. Regalia® Bioprotectant Concentrate may be used on vegetable crops, roses, fruits, nuts, flowers, foliage, houseplants, trees and shrubs located in residential landscapes. Regalia® Bioprotectant Concentrate can be applied up to and on the day of harvest on all fruits and vegetables.

WHEN TO USE

For best results, apply Regalia® Bioprotectant Concentrate prior to disease development or at the first sign of diseases and continue applying on a 7 day schedule or as needed.

BEFORE YOU USE

- Read and follow these directions when using.
- Do not allow spray to drift from application site.
- Use only with pressurized hand held sprayers or spray trigger bottles.
- Do not allow spray mixture to stand overnight or for prolonged periods.

Regalia® Bioprotectant Concentrate can be applied in commonly used pressurized hand held sprayers, spray trigger bottles and hose end sprayers.

HOW TO USE FOR HAND HELD SPRAYERS AND SPRAY TRIGGER BOTTLES

- Shake well before use.
- Fill sprayer or bottle with appropriate amount of water and concentrate.
- Mix the spray solution thoroughly.
- Keep the spray solution agitated during application.

HOW TO USE FOR HOSE END SPRAYERS

- Shake well before use.
- Follow hose end sprayer instructions to determine how to fill, set dial, clean and disconnect from hose.
- Set dial on sprayer to deliver rate as directed below.

HOW MUCH TO USE FOR ALL APPLICATIONS

1 fluid ounce (2 TBSP) of Regalia® Bioprotectant Concentrate per gallon of water. Spray plants to complete wetness, covering both top and bottom leaf surfaces to ensure complete coverage.

Some pesticides can cause phytotoxic effects ranging from slight burning or browning of leaves to distorted leaves, fruit, flowers or stems. Damage symptoms may vary with the type of plant that has been treated. It is impossible to test all plant species for phytotoxicity. To assure that the plants to be treated are not sensitive to the treatment, apply

a small amount of the product to a few leaves or the above ground portion of the plant and check back in 2 4 days for signs of phytotoxicity Use product according to label directions

DISEASES CONTROLLED OR SUPPRESSED ON VEGETABLES FRUITS NUTS ORNAMENTAL PLANTS TREES SHRUBS FLOWERS FOLIAGE AND TROPICAL PLANTS

- Anthracoise (*Colletotrichum* spp)
- Bacteria (*Erwinia* spp *Pseudomonas* spp *Xanthomonas* spp)
- Bacterial Leaf Blight (*Xanthomonas campestris*)
- Bacterial Speck (*Pseudomonas syringae* pv Tomato)
- Bacterial Spot (*Xanthomonas* spp)
- Bean Rust (*Uromyces appendiculatus*)
- Black Mold (*Alternaria alternata*)
- Black Rot/Black Crown Rot (*Alternaria* spp)
- Black Spot of Rose (*Diplocarpon rosea*)
- Botrytis (*Botrytis* spp)
- Botrytis Leaf Blight (*Botrytis squamosa*)
- Botrytis Neck Rot (*Botrytis* spp)
- Downy Mildew (*Bremia lactucae* *Peronospora* spp and *Plasmopara viticola*) suppression
- Early Blight (*Alternaria solani*)
- Fire Blight (*Erwinia amylovora*) – suppression
- Gray Mold (*Botrytis cinerea*)
- Greasy Spot (*Mycosphaerella citri*)
- Late Blight (*Phytophthora infestans*) – suppression
- Leaf Spots (*Alternaria* spp *Cercospora* spp *Septoria* spp)
- Onion Downy Mildew (*Peronospora destructor*)
- Onion Purple Blotch (*Alternaria porri*)
- Pin Rot (*Alternaria/Xanthomonas* complex)
- Powdery Mildew (*Uncinula necator* *Erysiphe* spp *Sphaerotheca* spp *Oidiopsis taurica* *Leveillula taurica* *Podosphaera leucotricha* *Oidium* spp *Podosphaera* spp)
- Rust (*Puccinia* spp)
- Scab (*Venturia* spp) suppression
- Sclerotinia Head and Leaf Drop (*Sclerotinia* spp) – suppression
- Sour Rot (*Alternaria tenuis* *Aspergillus* spp *Botrytis cinerea* *Cladosporium herbarum* *Penicillium* spp *Rhizopus arrhizus*)
- Target Spot (*Corynespora cassicola*)
- Walnut Blight (*Xanthomonas campestris*)

STORAGE AND DISPOSAL

Do not contaminate water food or feed by storage or disposal

Pesticide Storage Store in a cool dry place Avoid freezing

Pesticide Disposal and Container Handling If empty Nonrefillable container Do not reuse or refill this container Place in trash and offer for recycling if available

If partially filled Call your local solid waste agency or (800) 858 7378 (National Pesticide Information Center) for disposal instructions Never place unused product down any indoor or outdoor drain

WARRANTY

To the extent consistent with applicable law the seller makes no warranty expressed or implied of merchantability fitness or otherwise concerning use of this product The user assumes all risks of use storage or handling that are not in strict accordance with the accompanying directions

Label date

Made in the U S A

US Patents No 4 863 734 and No 5 989 429

65/65

Regalia® is a trademark of Marrone Bio Innovations Inc
Marrone Bio Innovations name and logo are registered trademarks of Marrone Bio Innovations Inc

© **Marrone Bio Innovations Inc**
2121 Second St Suite B 107 Davis CA 95618
1 877 664 4476
www.marronebio.com
info@marronebio.com