



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

February 26, 2025

Maggie Rodriguez
Director of Global Regulatory and Government Affairs
ProFarm Group, Inc.
1540 Drew Ave.
Davis CA 95618

Subject: Non-PRIA (Pesticide Registration Improvement Act) Labeling Amendment – Addition of an OMRI Claim, Amended Application Directions and Marketing Claims
Product Name: Regalia Bioprotectant Concentrate
EPA Registration Number: 84059-3
EPA Receipt Date: 7/13/2023
Action Case Number: 00491253

Dear Ms. Rodriguez:

The amended labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable.

This approval does not affect any terms or conditions that were previously imposed on this registration. You continue to be subject to existing terms or conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release this product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR § 152.3.


Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the U.S. Environmental Protection Agency (EPA). If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains statements or claims substantially differing from statements or claims made in connection with obtaining

a FIFRA section 3 registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Chris Pfeifer of my team by phone at (703) 244-7991 or via email at pfeifer.chris@epa.gov.

Sincerely,

A handwritten signature in blue ink, appearing to read "James Parker", is written over a horizontal line.

James Parker, Team Leader
Biochemical Pesticides Branch
Biopesticides and Pollution
Prevention Division (7511M)
Office of Pesticide Programs

Enclosure

MASTER LABEL

REGALIA[®] Bioprotectant Concentrate

Alternate Brand Names: REGALIA[®] Biofungicide Concentrate, REGALIA[®] Biofungicide, REGALIA[®] PTO Biofungicide, REGALIA[®] Rx Biofungicide, REGALIA[®] Rx, REGALIA[®] Garden, REGALIA[®] Garden Biofungicide, REGALIA[®] Home & Garden, REGALIA[®] Home & Garden Biofungicide, Regalia[®] CG, Regalia[®] CG Biofungicide, Regalia ST, Regalia Rx Plant Health, REGALIA[®] Rx Bioprotectant Concentrate; REGALIA[®] Bioprotectant Concentrate, REGALIA[®]

Sublabel A: Specialty Crops; Row Crops; Greenhouse; Turf (All combinations of terrestrial, greenhouse, indoor, and forest; food, feed, and nonfood)

Sublabel B: Professional Landscape Use (Terrestrial nonfood)

Sublabel C: Home & Garden Use (Residential Indoor and Residential Outdoor)

EPA Registration No. 84059-3

[Editorial guide/note to reviewer:

- *Optional/Alternate text appears within parentheses, editorial text appears within brackets and is not intended for final printed label.*
- *Where "Biopesticide" appears in this master label, one or more of Biofungicide, Biological Fungicide, Bioprotectant, and/or Fungicide may be used on the Final printed label.*
- *Final product label product use rate/interval may change but will still be within stated Master Label range.*
- *Alternate units (e.g. fl. oz per gallon, quarts per 100 gallons, Imperial to metric, etc) may be given on Final product label that do not affect the dilution ratio or the minimum or maximum use dilutions stated on the Master Label.]*

Sublabel A: Specialty Crops; Row Crops; Greenhouse; Turf (All combinations of terrestrial, greenhouse, indoor, and forest; food, feed, and nonfood)

REGALIA® Bioprotectant Concentrate

A plant extract to boost the plants' defense mechanisms to protect against certain fungal and bacterial diseases, and to improve plant health.

Active ingredient: Extract of *Reynoutria sachalinensis* 5 %
Other ingredients: 95 %
Total 100 %

EPA Reg. No. 84059-3

EPA Est. No. xxxxx-xx-xxx

GROUP P5 FUNGICIDE

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID	
IF SWALLOWED:	Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED:	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
IF IN EYES:	Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or if going for treatment. You may also contact 1-800-222-1222 for emergency medical treatment information.	

 CAN BE USED IN ORGANIC PRODUCTION)



(Batch)(LOT) No: xxxx (Printed on container)

Net Contents: xxxx

REGALIA® Bioprotectant Concentrate; EPA Reg. No. 84059-3

MASTER LABEL – MBI-LBL-4 r23 v4 January 28, 2025

Page 2 of 66

(Manufactured [for] [by]:) Pro Farm Group, Inc. 1530 Drew Ave., Davis, CA 95618 (1-877-664-4476;
info@profarmgroup.com)
(Name and logo of Pro Farm Group are registered trademarks of Pro Farm Group, Inc.)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear goggles or safety glasses. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Waterproof gloves
- Protective eyewear

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS

For terrestrial uses: do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exemptions pertaining to the statements on this label about personal protective equipment (PPE) and the restricted-entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. The REI does not apply when this product is used for seed treatment at planting or in hopper box treatments.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

For commercial seed treatment uses: Keep unprotected persons out of treatment area until seeds have dried or been packaged.

For all other non-WPS uses: Keep unprotected persons out of treated areas until sprays have dried.

GENERAL INFORMATION

REGALIA® Bioprotectant Concentrate is an extract from the plant *Reynoutria sachalinensis* for use on ornamental plants, turf, row, greenhouse, and edible and field crops. REGALIA® Bioprotectant Concentrate applied to the above ground portions of actively growing plants (see DIRECTIONS FOR USE) will help protect treated portions from certain plant diseases and will improve plant health. Plant health benefits often result in greater yields at harvest, especially when crops are stressed by pathogens or environmental conditions. Use REGALIA® Bioprotectant Concentrate as a preventative rather than a curative application. Apply prior to disease infestation to protect the growing buds, leaf tissue, flowers and above ground fruit and vegetables. See specific information below for diseases controlled and use rates on ornamental plants, turf, row, greenhouse, field and edible crops.

REGALIA® Bioprotectant Concentrate can be used in multiple application methods to the soil or plant roots to control or suppress certain soil-borne or foliar diseases and to promote healthy plant growth. See below for specific information on diseases controlled and use rates.

MIXING AND APPLICATION INSTRUCTIONS

– SHAKE WELL PRIOR TO USE –

Mixing instructions: Prepare no more mixture than is required for the immediate operation. Agitate the solution continuously during mixing and application. Mechanical mixing is recommended for proper mixing of REGALIA® Bioprotectant Concentrate mixtures.

REGALIA® Bioprotectant Concentrate alone: Add ½ of the required amount of water to the mix tank. With the agitator running, add the REGALIA® Bioprotectant Concentrate to the mix tank. Continue agitation while adding the remainder of the water. Begin application of the solution after the REGALIA® Bioprotectant Concentrate has completely dispersed into the mix water. Maintain agitation until all the mixture has been applied.

REGALIA® Bioprotectant Concentrate + tank mixtures: Add ½–¾ of the required amount of water to the mix tank. Start the agitation before adding any tank mix ingredients. In general, tank mix ingredients should be added in this order: wettable powders, dry flowable formulations, liquid flowable formulations, and emulsifiable formulations such as REGALIA® Bioprotectant Concentrate. Always allow each tank mix ingredient to become completely dispersed before adding the next component. Maintain continuous agitation until all components have been dispersed and throughout the application process. After all components are completely dispersed add the remainder of the water. REGALIA® Bioprotectant Concentrate cannot be mixed with another product with a prohibition against mixing. Use of the tank mix must be in accordance with the most restrictive label limitations and precautions. **Do not pre-mix REGALIA® Bioprotectant Concentrate with any other tank mix component prior to adding to the spray tank.**

Note: When using REGALIA® Bioprotectant Concentrate in tank-mixtures, all products in water soluble packaging should be added to the tank before any other tank-mix ingredient, including REGALIA® Bioprotectant Concentrate. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank-mix ingredient to the tank.

Compatibility: Do not combine REGALIA® Bioprotectant Concentrate in the spray tank with pesticides, adjuvants, or fertilizers if there has been no previous experience or use of the combination to show it is physically compatible, effective, and non-injurious under your use conditions. REGALIA® Bioprotectant Concentrate is compatible with many commonly used pesticides, fertilizers, adjuvants, and surfactants, but has not been evaluated with all potential combinations. To ensure compatibility of the tank mix combinations, evaluate prior to use as follows: Using a suitable container, add the proportional amounts of product to water. Add wettable powders first, then water dispersible granules, then liquid flowables, and lastly, emulsifiable concentrates. Mix thoroughly and let stand for at least five minutes. If the combination stays mixed or can be remixed, it is physically compatible. Test the mix on a small portion of the crop to be treated to ensure that a phytotoxic response will not occur as a result of the application.

Application Instructions: REGALIA® Bioprotectant Concentrate is a micro-emulsion concentrate consisting of certain ingredients extracted from *Reynoutria sachalinensis*. Use 50-mesh nozzle screens or larger. Use higher water volumes with larger sized crops and extensive foliage to obtain thorough coverage. Do not use carrier volumes and/or adjuvants that create spray runoff or drip-accumulation at the leaf margins or at the base of fruit or on the harvested commodity.

See FOLIAR AERIAL and FOLIAR GROUND APPLICATION section for use directions.

See BACKPACK/HANDHELD SPRAYER section for use directions.

See CHEMIGATION section for use directions.

See PRE-PLANT DIP section for use directions.

See SEED TREATMENT section for use directions.

See SOIL TREATMENT section for use directions.

FOLIAR AERIAL APPLICATION INSTRUCTIONS

0.5–1 quart per acre for FOLIAR (AERIAL) applications

- (For aerial applications, apply this product in a minimum of (2 to 10) gallons of water per acre.)
- (Apply this product preventatively or when the first disease symptoms are visible and reapply every 7–14 days)
- (For improved performance, use this product in a tank mix or rotational program with other registered fungicides.) (Tank mix this product with other registered fungicides for improved disease control under (heavy) (high disease) pressure.)
- (Repeat applications at (5-14) (7-10) (7–14) day intervals) (depending upon crop growth and disease pressure)
- (Under moderate to heavy disease pressure, tank mix this product with another fungicide.)
- (Consult your local Extension Specialist or Crop Consultant regarding the optimum timing of fungicide applications)
- (For grasses including forage, fodder, hay, and cereals, and for non-grass animal feed:) (It is important to apply this product at the flag leaf stage to maximize yield.)
- (Increasing the amount of water applied per acre will improve product performance.)
- (Follow all instructions to reduce aerial drift.)

AERIAL DRIFT REDUCTION ADVISORY INFORMATION

GENERAL: Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. Where states have more stringent regulations, they should be observed. Note: This section is advisory in nature and does not supersede the mandatory label requirements.

INFORMATION ON DROPLET SIZE: The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply droplets large enough to provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

CONTROLLING DROPLET SIZE: Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets. Pressure – Do not exceed the nozzle manufacturer’s recommended pressures. For many nozzle types, lower pressure produces larger droplets. When high flow rates are needed, use higher flow rate nozzles instead of increasing pressure. Number of Nozzles – Use the minimum number of nozzles that provide uniform coverage. Nozzle Orientation – Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential. Nozzle Type – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

BOOM WIDTH: For aerial applications, the boom width must not exceed 75% of the wingspan or 90% of the rotary blade. Use upwind swath displacement and apply only when wind speed is 3–10 mph as measured by an anemometer. Use medium or coarser spray according to ASAE 572 definition for standard nozzles or VMD for spinning atomizer nozzles. If application includes a no-spray zone, do not release spray at a height greater than 10 feet above the ground or crop canopy.

APPLICATION HEIGHT: Do not make application at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

SWATH ADJUSTMENT: When applications are made with a crosswind, the swath will be displaced downward. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.).

WIND: Drift potential is lowest between wind speeds of 2–10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY: When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

TEMPERATURE INVERSIONS: Do not apply during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small, suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SENSITIVE AREAS: The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas). Do not allow spray to drift from the application site and contact people, structures people occupy at any time and the associated property, parks and recreation areas, non-target crops, aquatic and wetland areas, woodlands, pastures, rangelands, or animals.

FOLIAR (GROUND) (EQUIPMENT) APPLICATION USE DIRECTIONS

REGALIA® Bioprotectant Concentrate can be applied in most commonly-used ground application equipment, such as tractor-mounted boom, airblast, high clearance, hose-end, backpack, and other pressurized sprayers; or hand-held sprayers; foggers or mist blowers. Thorough coverage is necessary to provide good disease control.

0.5–4 quarts per acre (0.5 – 6 tbsp. per 1000 sq. ft.) for FOLIAR (GROUND) (EQUIPMENT) applications

- (For foliar applications)(to optimize disease control and to maximize yields,) (apply ([specify volume])(of) (this product) (preventatively) in (a minimum of) 15–100 gallons of water per acre) (0.5 – 2.5 gallons of water per 1000 sq. ft.) (by (ground) (equipment)) (prior to disease development using) (sufficient volume for thorough coverage.) (or when the first symptoms of disease are visible) (at 7 day spray intervals or) (as soon as the first symptoms of disease appear) (or when environmental conditions are conducive to rapid disease development.) (Begin application soon after emergence or transplant, and when conditions are conducive to disease development.) (when applied alone, or at 1–4 quarts per acre (1.5 – 6 tbsp. per 1000 sq. ft.) when tank mixed with another fungicide preventatively in a minimum of 25 gallons of water per acre (0.5 gallons per 1000 square feet))
- (Additional sprays beyond second cover may be needed on susceptible varieties, or when environmental conditions are conducive to rapid disease development. Use high label rate and shorter spray intervals when conditions are conducive to rapid disease development.) (Use shorter intervals when conditions are conducive to rapid disease development.)
- (Avoid excessive amounts of water that result in the runoff of spray material)
- (Continue sprays at 7-day intervals or as needed) (For foliar applications, mix this product concentrate with water at a concentration of 2–4 quarts per 100 gallons of water (1.5 – 2.5 tbsp. per gallon of water) when used alone or 1–4 quarts per 100 gallons of water (1 – 2.5 tbsp. per gallon of water) when tank mixed with another fungicide) (If necessary, repeat applications at a 7-day interval.)
- (Begin applications preventatively (before disease symptoms become visible) at the 4 to 6-leaf stage and treat at 7–10 day intervals as needed prior to sale or harvest. Spray until just before point of runoff.)
- (Increase water volume as plant size increases.)
- (Do not exceed 1.0% v/v of the applied solution.)
- (Apply this product preventatively or when the first disease symptoms are visible) (and reapply every 7–14 days.)
- (Repeat applications at (5-14) (7–10) (7-14) day intervals) (depending upon crop growth and disease pressure)
- (Under moderate to heavy disease pressure, tank-mix this product with another fungicide.)(When the plants are) (under high disease pressure, tank-mix this product with another fungicide for more effective control.)(Tank mix this product with other registered fungicides for improved disease control under heavy pressure.)(For improved performance,(apply 0.5–2 quarts per acre (0.5 – 3 tbsp. per 1000 sq. ft.) of) (use) this product in a tank mix (or rotational program) with other registered fungicides)(When tank mixed with other fungicides, use 1–2 quarts of REGALIA® Bioprotectant Concentrate per acre (1.5 – 3 tbsp. per 1000 sq. ft.)).(This product can be tank mixed at the lower rate with another registered fungicide under heavy disease pressure)
- (For trees and vines:)(Dilute applications: this product can be applied by ground equipment to (vine and)tree crops in dilute applications of 100–400 gallons of water per acre (2.5 – 9 gallons of water per 1000 square feet). Apply this product at a rate of 2–4 quarts per acre (3 – 6 tbsp. per 1000 sq. ft.) when applied alone, or at 1–4 quarts per acre (1.5 – 6 tbsp. per 1000 sq. ft.) when tank mixed with another fungicide. Avoid excessive amounts of water that result in the runoff of spray material)
- (Consult your local Extension Specialist or Crop Consultant regarding the optimum timing of fungicide applications.)
- (For grasses including cereals, forage, fodder, hay, and for non-grass animal feed:) (It is important to apply this product at the flag leaf stage to maximize yield.)
- (Apply product at 1-4 quarts per acre (1.5 – 6 tbsp. per 1000 sq. ft.) when applied in a tank mix, or at 2-4 quarts per acre (3 – 6 tbsp. per 1000 sq. ft.) when applied alone.)
- (For concentrated ground applications, apply this product at 0.5–1.5 quarts per acre (0.5 – 2.5 tbsp. per 1000 sq. ft.) in a minimum of 10 gallons of water per acre (1 quart of water per 1000 sq. ft.)).
- (For plant health applications, apply (10-14 days) (two to three weeks) prior to cutting) (Foliar applications of Regalia to improve plant health can be made during the period from layby to contact sprays)
- (This product may be used to control certain diseases of container, bench, flat, plug, bed, or field-grown ornamentals in **greenhouses, shade-houses, outdoor nurseries, retail nurseries, and other landscape areas.**)

BACKPACK/HAND-HELD SPRAYER USE DIRECTIONS

The use rate for REGALIA® Bioprotectant Concentrate when applied alone or as an alternate spray in a backpack or hand-held sprayer is 1.2 – 2.5 tablespoons (Tbsp.) (0.6 – 1.5 fluid ounces [fl. oz.]) per gallon of water (0.5 – 1.0% v/v dilution of REGALIA® Bioprotectant Concentrate) applied at 1 – 2.5 gallons per 1000 square feet (sq. ft.) (25 – 100 gallons of water per acre). When tank mixed with another fungicide, the use rate for REGALIA® Bioprotectant Concentrate in a backpack or hand-held sprayer is 0.6 – 2.5 tablespoons (0.3 – 1.3 fluid ounces) per gallon of water applied at 1– 2.5 gallons per 1000 square feet (25 –100 gallons of water per acre). Use higher water volumes with larger sized crops and extensive foliage in order to secure thorough coverage. Do not use carrier volumes and/or adjuvants that create spray runoff or drip-accumulation at the base of fruit or on the harvested commodity. See specific application recommendations pertaining to each crop for additional details.

Use rate for REGALIA® Bioprotectant Concentrate when applied in small volumes

	Tablespoons REGALIA® Bioprotectant Concentrate per Gallon of Water	Fluid Ounces REGALIA® Bioprotectant Concentrate per Gallon of Water	v/v Dilution of REGALIA® Bioprotectant Concentrate	Applied (diluted) Gallons per 1,000 sq. ft.
Applied alone or as an alternate spray	1.2 – 2.5	0.6 – 1.3	0.5 – 1.0% v/v	1 – 2.5
Tank mixed with another fungicide	0.6 – 2.5	0.3 – 1.3	0.25 – 1.0% v/v	1 – 2.5

CHEMIGATION USE DIRECTIONS

(For (control) (suppression) of listed foliar diseases and to improve plant health, apply this product through overhead sprinkler, center pivot sprinkler systems, lateral move, end tow, side (wheel) roll, traveler, solid set, or hand move irrigation systems Do not use reclaimed water for application of this product.)

(For (control) (suppression) of listed soil-borne diseases and to improve plant health, apply this product only through flood (basin), furrow, border or drip (trickle) irrigation systems. Do not use reclaimed water for application of this product.)

0.5–4 quarts per acre (0.7 – 6 tbsp. per 1000 sq. ft.) for CHEMIGATION applications

- For chemigation applications (for improved plant growth) (and suppression of soil-borne diseases), apply this product through (drip) irrigation immediately after transplant and at 14-day intervals or begin 14 days after transplant when soil drench applications are used.

Spray preparation

First prepare a suspension of REGALIA® Bioprotectant Concentrate in a mix tank. Fill tank ½ to ¾ the desired amount of water. Start mechanical or hydraulic agitation. Add the required amount of REGALIA® Bioprotectant Concentrate, and then the remaining volume of water. Then set the irrigation system to deliver a minimum of 0.1 to 0.3 inch of water per acre. Start irrigation system and uniformly inject the suspension of REGALIA® Bioprotectant Concentrate into the irrigation water line so as to deliver the desired rate per acre. Inject the suspension of REGALIA® Bioprotectant Concentrate with a positive displacement pump into the main line after the filter, and ahead of a right angle turn to insure adequate mixing. Any questions on calibration should be directed to your State Extension Service Specialists, to equipment manufacturers or other experts.

Do not combine REGALIA® Bioprotectant Concentrate with pesticides, surfactants or fertilizers for application through chemigation equipment unless prior experience has shown the combination physically compatible, effective and non-injurious under conditions of use. REGALIA® Bioprotectant Concentrate has not been fully evaluated for compatibility with all adjuvants or surfactants. Conduct a spray compatibility test if a mixture with adjuvants or surfactants is planned.

CHEMIGATION

General Requirements –

- 1) Apply this product only through a drip or trickle system or center pivot sprinkler system, lateral move, end tow, side (wheel) roll, traveler, solid set, hand move, flood (basin), furrow, border or drip (trickle) irrigation systems.
- 2) Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- 3) If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.
- 4) Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- 5) A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Application Instructions for All Types of Chemigation –

- 1) Remove scale, pesticide residues, and other foreign matter from the chemical supply tank and entire injector system. Flush with clean water. Failure to provide a clean tank, void of scale or residues may cause product to lose effectiveness or strength.
- 2) Determine the treatment rates as indicated in the directions for use and make proper dilutions. Product can be applied continuously or at any time during the water application.
- 3) Prepare a solution in the chemical tank by filling the tank with the required water and then adding product as required
- 4) The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- 5) The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- 6) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- 7) Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Specific Requirements for Chemigation Systems Connected to Public Water Systems –

- 1) Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- 2) Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- 3) Do not apply when wind speed favors drift beyond the area intended for treatment.

Specific Requirements for Sprinkler Chemigation –

- 1) The system must contain a functional check valve, vacuum relief valve and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- 2) The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- 3) Do not apply when wind speed favors drift beyond the area intended for treatment.

Specific Requirements for Flood (Basin), Furrow and Border Chemigation –

- 1) Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as a drop structure or weir box to decrease potential for water source contamination from backflow if water flow stops.
- 2) The systems utilizing a pressurized water and pesticide injection system must meet the following requirements:
 - a. The system must contain a functional check valve, vacuum relief valve and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
 - b. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Specific Requirements for Drip (Trickle) Chemigation –

- 1) The system must contain a functional check valve, vacuum relief valve and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- 2) The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

PRE-PLANT DIP USE DIRECTIONS

Apply REGALIA® Bioprotectant Concentrate as a pre-plant dip for improved plant health and suppression of certain soil-borne diseases.

0.5–4 quarts per 100 gallons (1 tsp. – 2.5 tbsp. per gallon) of water for PLANT DIP (bare root) applications

- For plant dip applications for improved plant growth and suppression of soil-borne diseases, apply this product in a 0.25–0.5 % v/v suspension (1–4 quarts of this product per 100 gallons (or 1 – 2.5 tbsp. per gallon) of water as a pre-plant dip immediately prior to transplanting. Do not allow roots or plugs to be submerged in excess of 15 minutes prior to removal from the REGALIA® Bioprotectant Concentrate treatment.

SEED TREATMENT USE DIRECTIONS

REGALIA® Bioprotectant Concentrate can be applied as a seed (or seed piece) dressing for suppression of soil-borne diseases to improve early-season root growth. REGALIA® Bioprotectant Concentrate may be applied with other registered seed treatment insecticides and fungicides through standard slurry- or mist-type commercial seed treatment equipment. (REGALIA® Bioprotectant Concentrate can also be used in on-farm hopper-box or planter-box treatments.)

If using REGALIA® Bioprotectant Concentrate in a tank mixture with other seed treatment products, observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank-mix ingredient label. Do not exceed label rates and the most restrictive label precautions and limitations must be followed. Do not mix this product with any product which prohibits such mixing.

0.5–15 fluid ounces per 100 lbs. seed for SEED TREATMENT applications

- For suppression of soil-borne diseases, apply this product as a seed treatment at the rate of 0.5–15 fl. oz. per 100 lbs. seed.

(Do not apply this product through any type of irrigation system.)

Note: Federal law requires that bags containing treated seeds shall be labeled with the following information:

"This seed has been treated with Regalia® Bioprotectant Concentrate fungicide. Do not use for food, feed, or oil purposes. Store away from feed and foodstuffs."

Treated seed bagged for later use must contain an EPA-approved dye or colorant that imparts an unnatural color to the seed.

SOIL TREATMENT USE DIRECTIONS

REGALIA® Bioprotectant Concentrate can be applied by pre-plant incorporated soil application, soil drench, in-furrow spray or through drip (trickle) irrigation to improve plant health and to protect against certain soil-borne diseases.

In general, REGALIA® Bioprotectant Concentrate can be applied by the following methods, unless specified differently in the SELECTED CROPS section:

Pre-plant Incorporated Soil Applications:

- 1-4 quarts per acre in 10 – 100 gallons of water for PRE-PLANT INCORPORATED SOIL applications
- For pre-plant applications, apply this product with a broadcast spray and immediately incorporate into the soil profile prior to seeding or transplanting the crop to reduce transplant shock, suppress soil-borne diseases and improve root growth.

Soil Drench Applications:

0.5–3 quarts per 100 gallons (1 tsp. – 2 tbsp. per gallon) of water for SOIL DRENCH applications

- For soil drench applications, apply this product at a concentration of 1–3 quarts per 100 gallons (1 – 2 tbsp. per gallon) of water, and at a sufficient rate to thoroughly soak the growing media and root zone. Make an initial application of this product during or shortly after transplant to reduce transplant shock, suppress soil-borne diseases and improve root growth. (Multiple drench applications can be made on a 10–14 day interval.)

In-Furrow, Shanked-In and Injected Applications:

(REGALIA® Bioprotectant Concentrate can be shanked-in or injected into the soil alone, or with most types of liquid nutrients.)

0.5–4 quarts per acre or 1–10 fluid ounces per 1000 ft. row for IN-FURROW applications

- For in-furrow applications, at planting apply this product as an in-furrow spray at the rate of 0.5–4 quarts per acre or 1–10 fluid ounces per 1000 feet of row according to the chart (in the SOIL TREATMENT USE DIRECTIONS section)(below). Apply this product in 5–15 gallons of water so as the spray is directed into the seed furrow just before the [choose 1 or more] (seeds) (or seed pieces for potatoes) are covered.

Rate Per Acre	In-Furrow Application Rates Product per 1000 ft. row (fl. Oz.)														
	12" Rows	14" Rows	16" Rows	18" Rows	20" Rows	22" Rows	24" Rows	26" Rows	28" Rows	30" Rows	32" Rows	34" Rows	36" Rows	38" Rows	40" Rows
1 quart	0.73	0.86	0.98	1.1	1.2	1.3	1.5	1.6	1.7	1.8	2.0	2.1	2.2	2.3	2.4
2 quart	1.5	1.7	2.0	2.2	2.4	2.7	2.9	3.2	3.4	3.7	3.9	4.2	4.4	4.7	4.9
3 quart	2.2	2.6	2.9	3.3	3.7	4.0	4.4	4.8	5.1	5.1	5.9	6.2	6.6	7.0	7.3
4 quart	2.9	3.4	3.9	4.4	4.9	5.4	5.9	6.4	6.9	7.3	7.8	8.3	8.8	9.3	9.8

12" = 43,560 row ft./acre, 14" = 37,337 row ft./acre, 16" = 32,670 row ft./acre, 18" = 29,040 row ft./acre, 20" = 26,136 row ft./acre, 22" = 23,760 row ft./acre, 24" = 21,780 row ft./acre, 26" = 20,105 row ft./acre, 28" = 18,669 row ft./acre, 30" = 17,424 row ft./acre, 32" = 16,315 row ft./acre, 34" = 15,374 row ft./acre, 36" = 14,520 row ft./acre, 38" = 13,754 row ft./acre, 40" = 13,068 row ft./acre.

FOR USE ON THE FOLLOWING CROPS FOR CONTROL OR SUPPRESSION OF (THE FOLLOWING) (DISEASES) (PATHOGENS)(PESTS)

The use rate for REGALIA® Bioprotectant Concentrate when applied alone or as an alternate spray is 2–4 quarts per 100 gallons (1.5 – 2.5 tbsp. per gallon) of water (0.5–1.0% v/v dilution of REGALIA® Bioprotectant Concentrate) applied at 15–100 gallons of water per acre (0.5 – 2.5 gallons per 1000 square feet). When tank mixed with another fungicide, the use rate for REGALIA® Bioprotectant Concentrate is 1–4 quarts in 100 gallons (1 – 2.5 tbsp. per gallon) of water applied at 15–100 gallons of water per acre (0.5 – 2.5 gallons per 1000 square feet). Use higher water volumes with larger sized crops and extensive foliage in order to secure thorough coverage. Do not use carrier volumes and/or adjuvants that create spray runoff or drip-accumulation at the leaf margins or at the base of fruit or

on the harvested commodity. See specific application recommendations pertaining to each crop for additional details.

GREENHOUSE USE – This product can be used to control certain diseases of container, bench, flat, plug, bed or field-grown ornamentals and edible crops in greenhouses, shade-houses, outdoor nurseries, retail nurseries and other landscape areas. For greenhouse application on the crops and diseases listed, the recommended use rate for REGALIA® Bioprotectant Concentrate is 2–4 quarts in 100 gallons (1.5 – 2.5 tbsp. per gallon) of water (0.5–1.0% v/v dilution of REGALIA® Bioprotectant Concentrate) sprayed until just before point of runoff. When tank mixed with another fungicide, the use rate for REGALIA® Bioprotectant Concentrate is 1–4 quarts in 100 gallons (1 – 2.5 tbsp. per gallon) of water. Repeat at 7–14-day intervals as needed. See specific application recommendations for each crop for additional details. (Keep relative humidity below 70% to avoid risk of phytotoxicity.) (Since it is not possible to test all ornamental species or varieties grown in the greenhouse, test Regalia® Bioprotectant Concentrate on a few plants prior to large-scale usage.)

- Aerial Stem Rot (*Erwinia carotovora*)
- Aerial Web Blight (*Rhizoctonia solani*)
- Alfalfa Wilt (*Xylella spp.*)
- Alternaria Blight (*Alternaria cucumerina*)
- Alternaria Blotch (*Alternaria mali*)
- Alternaria Brown Spot (*Alternaria alternata*)
- Alternaria Fruit Rot (*Alternaria spp.*)
- Alternaria Leaf Blight (*Alternaria spp.*)
- Alternaria Leaf Spot (*Alternaria spp.*)
- Alternaria Leaf Spot, Boll Rot (*Alternaria spp.*)
- Alternaria Spot/Fruit Rot (*Alternaria alternata*)
- Angular Leaf Spot (*Mycosphaerella angulata*) (*Xanthomonas fragariae*)
- Anthracnose (*Colletotrichum spp.*) (*Gnomonia leptostyla*) (*Colletotrichum gloeosporioides*) (*Colletotrichum lagenarium*) (*Colletotrichum truncatum*) (*Elsinoe ampelina*)
- Anthracnose (*Colletotrichum coccodes*) (*Colletotrichum atramentarium*) (*Colletotrichum dematium*)
- Anthracnose (*Colletotrichum spp.*) – suppression only
- Anthracnose and Black Stem Rot (*Colletotrichum trifolii*)
- Anthracnose Boll Rot (*Glomeria spp.*)
- Anthracnose Fruit Rot (*Colletotrichum acutatum*)
- Anthracnose Leaf Blight (*Colletotrichum graminicola*)
- Anthracnose of Potato (*Colletotrichum coccodes*)
- Anthracnose, Boll Rot (*Glomeria spp.*)
- Apple Scab (*Venturia inaequalis*) (Suppression only)
- Ascochyta Blight, Boll Rot (*Ascochyta spp.*)
- Asian Soybean Rust (*Phakopsora pachyrhizi*)
- Aspergillus crown rot (*Aspergillus niger*)
- *Aureobasidium zeae*
- Bacteria (*Erwinia spp.*) (*Pseudomonas spp.*) (*Xanthomonas spp.*)
- Bacterial (Leaf) Spot (*Xanthomonas pruni*)
- Bacterial Blast (*Pseudomonas syringae*)
- Bacterial blight (*Pseudomonas cannabina*)
- Bacterial Blight (*Pseudomonas syringae*) (*Pseudomonas viridiflava*) (*Xanthomonas campestris* pv. *pruni*) (*Xanthomonas campestris*)
- Bacterial Blight (*Xanthomonas axonopodis*) (*Xanthomonas citri*)
- Bacterial Blight and Streak (*Xanthomonas spp.*)
- Bacterial Blight/Rot (*Xanthomonas spp.*)
- Bacterial Canker (*Erwinia nigrifluens*) (*Pseudomonas syringae*) (*Pseudomonas*

- spp.*) (*Xanthomonas campestris*)
(*Xanthomonas spp.*)
- Bacterial Leaf Blight (*Xanthomonas campestris*)
 - Bacterial Leaf Spot (*Pseudomonas spp.*)
 - Bacterial leaf streak (*Xanthomonas campestris* pv. *Holcicola*)
 - Bacterial leaf stripe (*Pseudomonas spp.*)
 - Bacterial Pustule (*Xanthomonas spp.*)
 - Bacterial rots (*Pantoea spp.*)
 - Bacterial Speck (*Pseudomonas syringae* pv. *glycinea*) (*Pseudomonas syringae*)
 - Bacterial Spot (*Xanthomonas pruni*) (*Xanthomonas spp.*) (*Xanthomonas cucurbitae*)
 - Bacterial Wilt (*Clavibacter michiganense*)
 - Barley yellow dwarf virus
 - Bentgrass/Bermudagrass Dead Spot (*Ophiosphaerella agrostis*)
 - Bermudagrass Decline (*Gaeumannomyces graminis* var. *graminis*)
 - Bitter Rot (*Colletotrichum spp.*)
 - Black dot disease (*Epicoccum nigrum*) (*Epicoccum purpurascens*)
 - Black mildew (*Schiffnerula cannabis*)
 - Black Mold (*Alternaria alternata*)
 - Black Root (*Thielaviopsis basicola*)
 - Black Root Rot / Black Crown Rot (*Alternaria spp.*)
 - Black Rot (*Guignardia bidwellii*)
 - Black Rot/Frogeye Leaf Spot (*Botryosphaeria obtusa*)
 - Black Scurf (*Rhizoctonia solani*)
 - Black shank (*Phytophthora nicotianae*)
 - Black Spot (*Guignardia citricarpa*), (*Phyllosticta citricarpa*)
 - Black Spot of Rose (*Diplocarpon rosae*)
 - Blossom Blight (*Monilinia spp.*)
 - Blue Mold (*Peronospora tabacina*)
 - Boll Rot (*Alternaria spp.*) (*Ascochyta spp.*) (*Fusarium spp.*) (*Phoma spp.*)
 - Bot Rot (*Botryosphaeria dothidea*)
 - Botryosphaeria Blight (*Botryosphaeria dothidea*)
 - Botrytis (*Botrytis cinerea*)
 - Botrytis Blight (*Botrytis cinerea*)
 - Botrytis Bud Rot (*Botrytis cinerea*)
 - Botrytis Bunch Rot (*Botrytis cinerea*)
 - Botrytis Fruit Rot (*Botrytis cinerea*)
 - Botrytis Leaf Blight (*Botrytis squamosa*)
 - Botrytis Neck Rot (*Botrytis spp.*)
 - Brooks Spot (*Mycosphaerella pomii*)
 - Brown / Hull Rot (*Monilinia spp.*)
 - Brown blight (*Alternaria alternata*) (*Alternaria tenuis*)
 - Brown leaf spot and stem canker (*Ascochyta spp.*) (*Ascochyta prasadii*) (*Phoma spp.*) (*Didymella spp.*) (*Phoma exigua*) (*Phoma glomerata*) (*Phoma herbarum*)
 - Brown patch (*Rhizoctonia solani*)
 - Brown Rot (*Monilinia spp.*)
 - Brown Rot Blossom Blight (*Monilinia laxa*)
 - Brown Rot Fruit Rot (*Monilinia fruticola*)
 - Brown Rot, Leaf Spots & Smuts (*Ceratobasidium spp.*) (*Cercospora spp.*) (*Cochliobolus spp.*) (*Drechslera spp.*)
 - Brown Rust (*Puccinia melanocephala*)
 - Brown Spot (*Alternaria spp.*) (*Septoria glycines*)
 - Brown Stripe/Gray Streak (*Cercosporidium graminis*)
 - Bull's Eye Rot (*Neofabraea spp.*)
 - Cedar-Apple Rust (*Gymnosporangium juniperi-virginianae*) – suppression only
 - Cercospora Blight (*Cercospora asparagi*) (*Cercospora kikuchii*)
 - Cercospora Blight and Leaf Spot (*Cercospora spp.*)
 - Cercospora Leaf Spot (*Cercospora citrulina*) (*Cercospora spp.*) (*Cercospora beticola*)
 - Charcoal rot (*Macrophomina phaseolina*)
 - Cherry Leaf Spot (*Blumeriella jaapii*)
 - *Cladosporium spp.*

- Cladosporium stem canker (*Cladosporium cladosporioides*)(*Cladosporium herbarum*)(*Mycosphaerella tassiana*)
- Clubroot (*Plasmodiophora brassicae*)
- Colletotrichum Crown Rot (*Colletotrichum* spp.) (*Colletotrichum graminicola*)
- Common Scab (*Streptomyces scabies*)
Suppression only
- Copper Spot (*Gloeocercospora sorghi*)
- Corn grey leaf spot (*Cercospora zeae-maydis*) (*Cercospora zeina*)
- Cotton root rot (*Phymatotrichopsis omnivora*)(*Phymatotrichum omnivorum*)
- Cranberry cotton ball (*Monilinia oxycocci*)
- Cranberry Early Rot (*Phyllosticta vacciniae*)
- Crown and Foot Rots (*Pseudocercospora herpotrichoides*, *Rhizoctonia*)
- Crown gall (*Agrobacterium tumefaciens*)
- Curvularia leaf spot (*Curvularia cymbopogonis*) (*Curvularia lunata*)(*Cochliobolus lunatus*)
- Cylindrosporium blight (*Cylindrosporium* spp.) (*Cylindrosorium cannabinum*)
- Damping off (*Aspergillus flavus*) (*Botrytis cinerea*) (*Botryotinia fuckeliana*) (*Fusarium* spp.) (*Fusarium oxysporum*) (*Fusarium solani*) (*Nectria haematococca*)(*Macrophomina phaseolina*) (*Pellicularia filamentosa*) (*Phytophthora* sp.) (*Pythium* spp.) (*Pythium aphanidermatum*)(*Pythium debaryanum*) (*Pythium ultimum*) (*Rhizoctonia* spp.) (*Rhizoctonia solani*) (*Thanatephorus cucumeris*)
- Dichondra Rust (*Puccinia dichondrae*)
- Diplodia Boll Rot (*Diplodia* spp.)
- Diplodia Ear Rot (*Stenocarpella maydis*)
- Diseases from pruning wounds including *Eutypa* (*Eutypa lata*), *Botryosphaeria rhodia*, *Phaeoacremonium aleophilum* and *P. chlamydospora*
- Dollar Spot (*Lanzia* spp.) (*Moellerodiscus* spp. formerly *Sclerotinia homeocarpa*)
- Downy Mildew (*Bremia lactucae*), (*Peronospora* spp.) (*Peronospora destructor*) (*Peronospora farinosa*) (*Peronospora mansherica*) (*Peronospora parasitica*) (*Peronospora trifoliorum*) (*Plasmopara viturni*) (*Plasmopara viticola*) (*Pseudoperonospora cubensis*) (*Pseudoperonospora humuli*)
- Downy mildew (*Pseudoperonospora cannabina*)(*Pseudoperonospora humuli*)
- Early Blight (*Alternaria solani*)
- Early Blight of celery (*Cercospora apii*)
- Early Leaf Spot (*Cercospora arachidicola*)
- Early Rot in Cranberry (*Phyllosticta vaccinii*)
- Eastern Filbert Blight (*Anisogramma anomala*)
- Eutypa (*Eutypa lata*)
- Eye Spot (*Aureobasidium zeae*)
- Fire Blight (*Erwinia amylovora*)
- Flyspeck (*Zygophiala jamaicensis*)
- Foliar Blight
- Frog-eyed Leaf Spot (*Cercospora sojae*)
- Fruit Finish
- Fungal Disease Complexes (*Bipolaris* spp., *Monographella* spp., *Phaeosphaeria* spp.)
- Fusarium foot rot and root rot (*Fusarium solani*)
- Fusarium Head Blight (*Fusarium graminearum*)
- *Fusarium oxysporum*
- Fusarium Patch (*Fusarium nivale*)
- *Fusarium proliferatum*
- Fusarium root and stem rot

- *Fusarium solani*
- *Fusarium spp.*
- Fusarium stem canker (*Fusarium sulphureum*) (*Gibberella cyanogena*) (*Gibberella saubinetii*)
- Fusarium wilt (*Fusarium oxysporum f.sp. cannabis*) (*Fusarium oxysporum f.sp. vasinfectum*) (*Fusarium oxysporum f. sp. lactucae*)
- *Fusarium wilt (Fusarium oxysporum)*
- *Glomerella tucumanensis*, also known as *Colletotrichum falcatum* (Suppression Only)
- Gray leaf spot (*Cercospora sorghi*) (*Pyricularia grisea*) (*Cercospora zeae-maydis*)
- Gray Mold (*Botrytis cinerea*) (*Botrytis spp.*)
- Greasy Spot (*Mycosphaerella citri*)
- Green Fruit Rot (*Botrytis cinerea*)
- Gummy Stem Blight (*Didymella bryoniae*)
- Hard Lock
- Hard Lock, Boll Rot (*Fusarium spp.*)
- Hemp canker (*Sclerotinia sclerotiorum*)
- Hemp Leaf Spot (*Bipolaris sp.*)
- Hull Rot (*Rhizopus stolonifer* and *Monilinia spp.*)
- Late Blight (*Phytophthora infestans*) (*Septoria apiicola*)
- Late Leaf Spot (*Cercosporidium personatum*)
- Late Rot in Cranberry
- Leaf Blight (*Pseudocercospora vitis*) (*Septosphaeria turcica*)
- Leaf Rust (*Pucciniastrum vaccinii*) (*Tranzschelia discolor*)
- Leaf Spot (*Alternaria spp.*) (*Cercospora spp.*) (*Cercospora beticola*) (*Corynespora cassicola*) (*Entomosporium spp.*) (*Mycosphaerella fragariae*) (*Myrothecium spp.*) (*Septoria spp.*)
- Leaf Spots (*Drechslera*, *Cochliobolus*, *Cercospora*)
- Leafspots and Blotches (*Pseudopeziza medicaginus*, *Stemphyllium spp.*, *Cercospora spp.*, *Stagonospora spp.*)
- Leptosphaeria blight (*Leptosphaeria cannabina*) (*Leptosphaeria woroninii*) (*Leptosphaeria acuta*)
- Melanose (*Diaporthe citri*)
- Melting Out Leaf Spot (*Bipolaris spp.*), (*Drechslera spp.*)
- Miscanthus blight
- Miscanthus streak virus
- Mosaic viruses
- Mummy Berry (*Monilinia vaccinii-corymbosi*),
- Necrotic Ring Spot (*Leptosphaeria korrae*)
- Northern Leaf Blight (*Exserohilum turcicum*)
- Northern Leaf Spot (*Cochliobolus carbonum*)
- Olive Knot (*Pseudomonas savastanoi*)
- Olive leaf spot (*Cercospora cannabis*) (*Pseudocercospora cannabina*)
- Onion Downy Mildew (*Peronospora destructor*)
- Onion Purple Blotch (*Alternaria porri*)
- Ophiobolus stem canker (*Ophiobolus cannabinus*) (*Ophiobolus anguillides*)
- Orange Rust (*Puccinia kuehnii*)
- Panicum mosaic virus
- Peg
- *Pencillium*
- Phoma Blight
- Phoma Blight, Boll Rot (*Phoma spp.*)
- Phoma stem canker (*Phoma herbarum*) (*Phoma exigua*)
- Phomopsis
- Phomopsis Fruit Rot (*Phomopsis viticola*)
- Phomopsis Leaf Blight (*Phomopsis obscurans*)
- Phomopsis Leaf Spot, Twig Blight, and Fruit Rot (*Phomopsis spp.*)
- Phomopsis stem canker (*Phomopsis cannabina*) (*Phomopsis*)

- achilleae*)(*Diaporthe arctii* var. *achilleae*)
- Phymatotrichum root rot / Phymatotrichopsis root rot (*Phymatotrichopsis omnivora*)(*Phymatotrichum omnivorum*)
- Phytophthora Blight (*Phytophthora capsici*)
- Phytophthora Root Rot and Crown Rot (*Phytophthora* spp.)
- Phytophthora (*Phytophthora* spp.)
- Pin Rot Complex (*Alternaria/Xanthomonas*)
- Pink Patch (*Limonomyces roseipellis*)
- Pink Rot (*Phytophthora erythroseptica*)(*Sclerotinia sclerotiorum*)
- Pink rot (*Trichothecium roseum*)(*Cephalothecium roseum*)
- Pithomyces blight
- Pod and Stem Blight (*Diaporthe phaseolorum* var. *sojae*) (*Phomopsis longicola*) (*Diaporthe* spp.)
- Pod and Stem Blight (*Diaporthe* spp.)
- Postbloom Fruit Drop (*Colletotrichum acutatum*)
- Powdery Mildew (*Erysiphe* spp.) (*Erysiphe betae*), (*Erysiphe cichoracearum*) (*Erysiphe cruciferarum*) (*Erysiphe graminis*) (*Erysiphe polygoni*) (*Leveillula taurica*) (*Microsphaera alni*) (*Oidium* spp.), (*Oidopsis taurica*) (*Podosphaera* spp.) (*Podosphaera leucotricha*) (*Sphaerotheca* spp.) (*Sphaerotheca fuliginea*) (*Sphaerotheca macularis*) (*Sphaerotheca pannosa*) (*Uncinula necator*)
- Powdery Mildew (*Golovinomyces*) (*Erysiphe cichoracearum*)(*Leveillula taurica*)(*Oidiopsis taurica*)(*Sphaerotheca macularis*)(*Sphaerotheca humuli*)(*Oidium* spp.)
- Powdery Mildew / Rusty Spot (*Podosphaera* spp.), (*Sphaerotheca pannosa*)
- *Puccinia* spp.
- Purple spot (*Stemphylium vesicarium*)
- Pythium (aerial blight phase) (*Pythium* spp.)
- Pythium (*Pythium* spp.)
- *Pythium acanthicum*
- *Pythium aphanidermatum*
- Pythium Blight, Pythium Root Rot (*Pythium aphanidermatum*), (*Pythium* spp.)
- *Pythium dissoticum*
- *Pythium myriotylum*
- Pythium root and damping off
- Ramularia (*Ramularia* spp.)
- Ramularia Leaf Spot (*Ramularia cynarae*)
- Red boot (*Melanospora cannabis*)
- Red Rot (*Glomerella tucumanensis*, also known as *Colletotrichum falcatum*)
- Red Thread (*Laetisaria fuciformis*)
- Rhizoctonia Foliar Blight, Peg, and Root Rot (*Rhizoctonia solani*)
- Rhizoctonia Large Patch (*Rhizoctonia solani*)
- Rhizoctonia soreshin and root rot (*Rhizoctonia solani*)
- *Rhizoctonia* spp.
- Rice Blast (*Pyricularia grisea*)
- Ripe Rot (*Colletotrichum gloeosporioides*)
- Root and collar rots (*Phytophthora*, *Pythium*, *Fusarium*, *Rhizoctonia*)
- Rot (*Rhizoctonia* spp.), (*Pythium* spp.), (*Fusarium* spp.), (*Cylindrocarpon* spp.)
- Rust (*Aecidium cannabis*)(*Uredo kriegiana*)(*Uromyces inconspicuus*)
- Rust (*Phykopsora* spp.) (*Puccinia* spp.) (*Puccinia asparagi*) (*Puccinia menthae*) (*Puccinia porri*) (*Tranzschelia discolor*) (*Uromyces appendiculatus*) (*Uromyces betae*)
- Rusty Spot (*Podosphaera leucotricha*)
- Scab (*Cladosporium carpophilum*) (*Sphaceloma perseae*) (*Elsinoe australis*)

- (*Elsinoe fawcetti*) (*Elsinoe mangiferae*)
(*Venturia spp.*)
- Sclerotinia (*Sclerotinia sclerotiorum*)
 - Sclerotinia Head and Leaf Drop (*Sclerotinia minor*) (*Sclerotinia sclerotiorum*)
 - Sclerotinia stem and crown rot (*Sclerotinia sclerotiorum*)
 - Sclerotium root and stem rot (*Sclerotium rolfsii*) (*Athella rolfsii*)
 - Seedling and Damping Off Disease Complex, including Root and Crown Rots (*Pythium*, *Phytophthora*, *Rhizoctonia*, and *Stagonospora spp.*)
 - Septoria Brown Spot (*Septoria glycines*)
 - Septoria Leaf/Speckled Leaf Spot/Blotch (*Septoria spp.*)
 - Sheath Spot and Blight (*Rhizoctonia oryzae*), (*Thanatephorus cucumeris*)
 - Shot Hole (*Wilsonomyces carpophilus*)
 - Sigatoka (*Mycosphaerella fijiensis*)
 - Smut (*Tilletia spp.*) (*Tilletia barclayana*)
 - Smuts and Bunts (*Tilletia spp.*)
 - Snowmold, Gray (*Typhula spp.*)
 - Snowmold, Pink (*Microdochium nivale*)
 - Sooty Blotch (*Geastrum polystigmati*), (*Leptodontium elatius*), (*Peltaster fructicola*)
 - Sorghum downy mildew (*Peronosclerospora sorghi*)
 - Sour Rot (*Alternaria tenuis*) (*Aspergillus spp.*) (*Botrytis cinerea*) (*Cladosporium herbarum*) (*Penicillium spp.*) (*Rhizopus arrhizus*)
 - Southern Blight (*Sclerotium rolfsii*)
 - Southern blight (*Sclerotium rolfsii*) (*Athella rolfsii*)
 - Southern leaf blight (*Bipolaris spp.*) (*Cochliobolus heterostrophus*)
 - Spring Black Stem (*Phoma medicaginis*)
 - Spring Dead Spot (*Leptosphaeria korrae*), (*Leptosphaeria narmari*), (*Ophiosphaerella herpotricha*), (*Gaeumannomyces graminis*)
 - Spur Blight (*Didymella spp.*), (*Phoma spp.*)
 - Stem Rot (*Sclerotium oryzae*)
 - Stemphylium leaf and stem spot (*Stemphylium botryosum*) (*Pleospora tarda*) (*Stemphylium cannabinum*)
 - Stemphylium Leaf Blight (*Stemphylium vesicarium*)
 - Stemphylium Leaf Spot (*Stemphylium spp.*)
 - *Striatura ulceroza* (*Pseudomonas amygdali pv. mori*)
 - Stripe Smut (*Ustilago striiformis*), (*Urocystis agropyri*)
 - Summer Bentgrass Decline
 - Summer Patch, Poa Patch (*Magnaporthe poae*)
 - Switchgrass Mosaic Virus
 - Take-All Patch (*Gaeumannomyces graminis*)
 - Tan Spot (*Pyrenophora tritici-repentis*)
 - Tar spot (*Phyllachora cannabidis*)
 - Target Spot (*Corynespora cassiicola*) (*Rhizoctonia solani*)
 - Tropical rot (*Lasiodiplodia theobromae*) (*Botryodiplodia theobromae*)
 - Twig blight (*Dendrophoma marconii*) (*Botryosphaeria marconii*)
 - Verticillium wilt (*Verticillium spp.*) (*Verticillium albo-atrum*)
 - *Verticillium spp.*
 - Verticillium wilt (*Verticillium albo-atrum*) (*Verticillium dahliae*)
 - Walnut Blight (*Xanthomonas campestris*)
 - White leaf spot (*Phomopsis ganjae*)
 - White Mold (*Sclerotinia sclerotiorum*) (*Sclerotium rolfsii*) (*Sclerotinia minor*) (*Sclerotinia trifoliorum*)
 - white mold stem rot
 - White Mold/ Sclerotinia Stem Rot (*Sclerotinia sclerotiorum*)
 - White Rot (*Botryosphaeria dothidea*)
 - White Rust (*Albugo occidentalis*)

- *Xanthomonas campestris*
- Xanthomonas leaf spot (*Xanthomonas campestris* pv. *cannabis*)
- Xanthomonas Leaf Spot (*Xanthomonas campestris*)
- *Xanthomonas* spp.
- Yellow leaf spot (*Septoria cannabis*)(*Septoria cannabina*)
- Yellow Patch (*Rhizoctonia cerealis*)
- Yellow Tuft/Downy Mildew (*Sclerophthora macrospora*)
- Zoysia Patch (*Rhizoctonia solani*)

Pre-harvest Interval (PHI) = 0 days

ROOT, TUBER AND CORM CROPS

Potato, Beet, Carrot, Cassava, Ginger, Ginseng, Horseradish, Radish, Sugar beet, Sweet Potato, Turnip (including those for seed production) and other root and tuber crops

For suppression of Early Blight, Black Root Rot/Black Crown Rot, and Late Blight, begin application of this product in 25–100 gallons of water per acre (0.5 – 2.5 gallons of water per 1000 square feet) soon after emergence when conditions are conducive to disease development. Repeat on a 5–7-day interval or as needed. For improved performance, use this product in a tank mix with other registered fungicides.

(Apply at 1 pint per acre for Plant Health Applications.)

1-16 fl oz. per 100 pounds of seed pieces for SEED PIECE DIP applications

- For seed piece dip applications for improved plant growth and suppression of soil-borne diseases, apply this product to seed pieces prior to planting.

LEAVES OF ROOT AND TUBER VEGETABLES

Beet, Chervil, and other leaves of roots and tubers

BULB VEGETABLES

Onion (Bulb and Green), Garlic, Leek, Shallot and other bulb vegetables

LEAFY VEGETABLE CROPS (except *Brassica* vegetables)

Arugula, Celery, Cilantro, Corn Salad, Cress, Dandelion, Dock, Edible Chrysanthemum, Endive, Head Lettuce, Leaf Lettuce, Parsley, Purslane, Radicchio, Rhubarb, Spinach, Swiss Chard, Turnip, Watercress and other leafy vegetable crops

- West of the Rocky Mountains – For aerial applications, apply this product at 0.5–1.5 quarts per acre in a minimum of 10 gallons of water per acre
- East of the Rocky Mountains – For aerial applications, apply this product at 0.5–1 quarts per acre in a minimum of 5 gallons of water per acre.
- For California – For aerial application apply REGALIA® Bioprotectant Concentrate at 1–3 pints per acre in 10–25 gallons of water per acre.

Restrictions:

Do not apply REGALIA® Bioprotectant Concentrate when extended/unseasonably cold or cold and cloudy conditions are expected. Regalia should **not** be applied when night-time temperatures will fall below 45°F and relative humidity is predicted to be above 80%. Applications during daylight hours are preferred over night-time applications.

(For concentrated applications, apply this product at 0.5 to 1.5 quarts per acre in a minimum of 10 gallons of water per acre.)

BRASSICAS (COLE) LEAFY VEGETABLES

Broccoli, Broccoli Rabe, Brussels Sprouts, Cabbage, Chinese Broccoli, Chinese Cabbage (Bok Choy), Chinese Cabbage (Napa), Chinese Mustard Cabbage (Gai Choy), Cauliflower, Cavalo, Collards, Kale, Kohlrabi, Mizuna, Mustard Greens, Mustard Spinach, Rape Greens, and other cole crops

LEGUME VEGETABLES, succulent or dried (not including soybeans and peanuts)(not including peanuts)

Chick Peas, Dry Beans, Green Beans, Lentils, Lima Beans, Peas, Shell Beans, Snap Beans, Split Peas (including those grown for seed or oil production), Soybeans (and other legume vegetables)

To maximize activity against *Sclerotinia* white mold in soybeans, the best timing would be V3 to R3 and should be tank mixed with a fungicide partner also labeled for white mold activity.

Repeat applications on 7–10 day intervals depending upon crop growth and disease pressure.

FOLIAGE OF LEGUME VEGETABLES

Garden peas and other foliage of legume vegetables

FRUITING VEGETABLES

Tomato, Pepper, Eggplant, Ground Cherry, Okra, Tomatillo and other fruiting vegetables

- Phytophthora Blight – Apply this product in combination with labeled rates of a copper fungicide (or with another fungicide labeled for Phytophthora Blight control).

CUCURBITS

Includes all types and hybrids of: Acorn squash, Balsam apple, Balsam pear, Bitter melon, Butternut squash, Calabaza, Cantaloupe, Casaba, Chayote, Chinese cucumber, Chinese okra, Chinese waxgourd, Citron melon, Crenshaw melon, Crookneck squash, Cucumber, Cucuzza, Gherkin, Golden pershaw melon, Honeydew melon, Honey balls, Hubbard squash, Hyotan, Mango melon, Persian melon, Pineapple melon, Pumpkin, Santa Claus melon, Scallop squash, Snake melon, Spaghetti squash, Straightneck squash, Vegetable marrow, Watermelon, Zucchini and other cucurbits

- When greenhouse cucurbits are under high disease conditions, use the shorter spray interval.
- Downy Mildew – Tank mix this product with another fungicide labeled for Downy Mildew control and re-apply at a 7-day interval or according to the label directions of the tank mix (ingredient) (partner)
- Phytophthora Blight – Apply this product in combination with labeled rates of a copper fungicide or with another fungicide labeled for Phytophthora Blight control.

CITRUS CROPS

Orange, Grapefruit, Kumquat, Lemon, Tangelo, Tangerine, Pummelo and other citrus crops

POME FRUITS

Apple, Crabapple, Loquat, Oriental Pear, Pear, Quince, Mayhaw and other pome fruits

- Use caution when selecting spray adjuvants. Select only those adjuvants which through prior experience do not affect fruit finish when combined with this product.
- Fire Blight – For suppression, apply 1-2 quarts of this product in 50–100 gallons of water per acre (1 – 2.5 gallons of water per 1000 square feet) at green tip through bloom. Do not exceed 0.5% v/v. Following bloom, this product can be applied at 2–4 quarts per acre (3 – 6 tbsp. per 1000 sq. ft.). For maximum control, use this product prior to infection events. During periods of rapid development and frequent infection periods, use spray intervals of 3–7 days.

- Apply in sufficient water to provide full coverage. For improved performance, use this product in a rotational program with copper or antibiotics registered for Fire Blight control such as but not limited to oxytetracycline or streptomycin.
- Proper orchard cultural practices are essential to eliminate Fire Blight-infected tissue from the orchard to assure good performance of any crop protection product. Remove and destroy dead and diseased wood from the orchard prior to and during the growing season.
- Scab – For suppression, apply 1-2 quarts of this product in 50–100 gallons of water per acre (1 – 2.5 gallons of water per 1000 square feet) at green tip and through bloom when environmental conditions become favorable for primary Scab development and repeat on a 7–10 day interval or as needed. Do not exceed 0.5% v/v. Use this product in a tank mix or rotational program with other fungicides labeled for Scab control. Following bloom, this product can be applied at 2–4 quarts per acre (3 – 6 tbsp. per 1000 sq. ft.).

Some sensitive tree fruit varieties have exhibited petal staining and/or necrosis after application of higher use rates. To minimize petal staining and/or necrosis:

- Use adjuvants that improve coverage, not penetration; follow the manufacturer's mixing instructions.
- Use adjuvants that through prior experience do not affect petal integrity when combined with this product.
- Apply 1-2 quarts of this product in 50–100 gallons (1 – 2.5 tbsp. per gallon) of water per acre (1.5 tbsp. in 1 gallon to 2.5 gallons of water per 1000 sq. ft.) in Pome Fruit, from 10% bloom to full bloom. Do not exceed 0.5% v/v.

STONE FRUITS

Apricot[†], Cherry (sweet and tart), Nectarine, Peach, Plum, Plumcot, Prune and other stone fruits

- Bacterial Blight – Apply this product in 50–100 gallons of water per acre (1 – 2.5 gallons of water per 1000 square feet) postharvest before fall rains.
- Brown Rot Blossom Blight – Begin application of this product in 50–100 gallons of water per acre (1 – 2.5 gallons of water per 1000 square feet) at early bloom, and repeat through petal fall on a 7-day interval or as needed.
- Powdery Mildew – Begin application of this product in 50–100 gallons of water per acre (1 – 2.5 gallons of water per 1000 square feet) at popcorn stage, and repeat on a 7-day interval or as needed. For improved performance, use this product in a tank mix or rotational program with other registered fungicides for powdery mildew control.
- Scab – Begin application of this product in 50–100 gallons of water per acre (1 – 2.5 gallons of water per 1000 square feet) at petal fall, and repeat on a 7–10-day interval or as needed. For improved performance, tank mix this product with another fungicide labeled for Scab control.
- For all other diseases – Begin application prior to disease development when environmental conditions and plant stage are conducive to rapid disease development, and repeat on a 7–10 day interval or as needed. Use in a tank mix or rotational program when disease conditions are severe.

[†]Some sensitive apricot varieties have exhibited fruit spotting as a result of application. Spray a test strip to confirm your variety is not susceptible to spotting before spraying.

Some sensitive tree fruit varieties have exhibited petal staining and/or necrosis after application of higher use rates. To minimize petal staining and/or necrosis:

- Use adjuvants that improve coverage, not penetration; follow the manufacturer's mixing instructions.
- Use adjuvants that through prior experience do not affect petal integrity when combined with this product.
- Apply 1 quart of this product in 50–100 gallons of water per acre (1.5 tbsp. in 1 gallon to 2.5 gallons of water per 1000 sq. ft.) in:
 - Cherries, from white bud (first white, popcorn) to full bloom,
 - Stone fruit, from 10% bloom to full bloom.

BERRIES (AND SMALL FRUIT)

Blueberry[†], Blackberry (all varieties), Cranberry, Currant, Elderberry, Gooseberry, Huckleberry, Juneberry, Ligonberry, Loganberry, Raspberry (red and black), Salal and cultivars and/or hybrids of these and other berries

Grape (wine, table, raisin and juice varieties), Strawberry and other smallfruit

- Mummy Berry – Initiate application at bud break stage of development. Apply this product preventatively and repeat on a 7–10 day interval or as needed. For best performance, tank mix this product with other registered fungicides for Mummy Berry control.
- Botrytis Blight – Apply this product preventatively or when the first disease symptoms are visible and reapply every 7–14 days.
- Bacterial Canker – Apply this product prior to Fall rains and repeat applications during dormancy before spring growth. This product can be tank mixed with another registered fungicide for improved control of bacterial canker. REGALIA® Bioprotectant Concentrate is compatible with fungicide/bactericides containing the active ingredients peroxyacetic acid and hydrogen peroxide such as Jet-Ag® 5.
- Anthracnose Fruit Rot and Alternaria Fruit Rot on blueberries – Initiate application at green tip and continue applications on a 7–10 day interval.
- Anthracnose on strawberry – For suppression, apply this product preventatively in 50–100 gallons of water per acre (1 – 2.5 gallons of water per 1000 square feet) and repeat on a 7-day interval or as needed. For best performance, tank mix this product with other registered fungicides for Anthracnose control.

[†] Some sensitive blueberry varieties have exhibited fruit spotting as a result of application. Spray a test strip to confirm your variety is not susceptible to spotting before spraying. Do not exceed 1% v/v.

(Grape - Do not use carrier volumes and/or adjuvants that create spray runoff or drip-accumulation at the leaf margins or at the base of fruit or on the harvested commodity. Complex tank-mixes that include REGALIA® Bioprotectant Concentrate may increase the risk of visible spray residues on grape clusters. Test the mix on a small portion of the crop to be treated to ensure that a phytotoxic response will not occur as a result of the application. Do not apply REGALIA® Bioprotectant Concentrate within 14 days of applying a sulfur fungicide during high temperatures.)

TREE NUT CROPS

Almond, Beech nut, Brazil nut, Bitternut, Cashew, Chestnut, Chinquapin, Filbert, Hickory nut, Macadamia nut, Pecan, Pistachio, Walnut (Black and English)

Some tree nut varieties have exhibited petal staining and/or necrosis after application of higher use rates.

To minimize petal staining and/or necrosis:

- Use adjuvants that improve coverage, not penetration; follow the manufacturer's mixing instructions.
- Use adjuvants that through prior experience do not affect petal integrity when combined with this product.
- Apply 1 quart in 50 gallons or 2 quarts in 100 gallons of water per acre (1.5 tbs. in 1 gallon or 3 tbs. in 2.5 gallons of water per 1000 sq. ft.) from pink bud through bloom.

(For Pecans: For Phymatotrichopsis root rot (*Phymatotrichopsis omnivore*) apply through Chemigation by drip irrigation or micro-sprinklers at 2-4 quarts/acre. Begin applications when trees begin to leaf out in the spring, repeat applications monthly.)

CEREAL GRAINS

Barley, Buckwheat, Grain Amaranth, Milo, Oat, Millets, Rice, Rye, Sorghum, Triticale, Wheat and other cereal grains

It is important to apply this product at the Feekes 4 to Feekes 10.5 stage to maximize yield. Apply this product preventatively or when the first disease symptoms appear. Repeat applications in 7–14-day intervals depending upon crop growth and disease pressure

(For (Barley) (Buckwheat) (Rye) (Amaranth) (Oat) (Millets) (Rice) (Wheat) (Sorghum)): (Apply at 1 pint per acre for foliar (ground) (aerial) Plant Health Applications.)

Corn (Sweet Corn, Field Corn, Popcorn, Silage Corn, Seed Corn)

Apply this product preventatively starting at V1-V10 or VT or when the first disease symptoms appear. (Repeat applications on 7–10 day intervals depending upon crop growth and disease pressure.)

(Apply this product in 1 pint per acre by (foliar (ground)) (in-furrow) (for) (improved plant health potential.) (increased quality and/or yields at harvest.)

FORAGE, FODDER AND STRAW OF CEREAL GRAINS

Corn, Wheat, and any other cereal grain crop

(For (Corn) (Wheat):) Apply at 1 pint per acre for (foliar (ground) (aerial)) (in-furrow) Plant Health Applications.)

GRASS FORAGE, FODDER, AND HAY

Bermuda grass, Bluegrass, Bromegrass, Fescue, Pasture and range grasses grown for hay or silage, Sudangrass, Timothy and other grass forage, fodder, and hay

(Apply at 0.5 to 1 quart per acre for foliar (ground) (aerial) Plant Health Applications.)

NON-GRASS ANIMAL FEED

Alfalfa, Clover, Kudzu, Lespedeza, Lupin, Sainfoin, Trefoil, Vetch and other non-grass animal feed

(Apply at 1 pint per acre for foliar (ground) Plant Health Applications.)

HERBS (DRIED OR FRESH) (FIELD AND GREENHOUSE)

SPICES (FIELD AND GREENHOUSE)

OIL SEED CROPS (not including peanut or soybean)

Canola, Castor, Flax, Jojoba, Rapeseed, Safflower, Sesame, and Sunflower and other oil seeds

For white mold disease control: apply at 10% bloom with a second application 7-10 days later.

(Apply at 1 pint per acre for Plant Health Applications in Canola.)

Cotton

For ground applications for foliar and boll rot disease control, apply this product preventatively in 15–40 gallons of water per acre (0.5 – 1 gallon of water per 1000 square feet) prior to disease development using sufficient volume for thorough coverage

(Apply at 0.5 to 1 quart per acre for foliar (ground) (aerial) Plant Health Applications.)

STALK, STEM, AND LEAF PETIOLE VEGETABLES

Asparagus, Celery, Agave and other stalk, stem and leaf petiole vegetables

TROPICAL AND SUBTROPICAL FRUITS, EDIBLE

Olive and other tropical and subtropical fruits with edible peel

TROPICAL AND SUBTROPICAL FRUITS, INEDIBLE

Avocado, Banana, Kiwi, Mango, Papaya, Plantain, Pineapple, Pomegranate and other tropical and subtropical fruits with inedible peel

Sigatoka – Initiate applications when leaves first appear and repeat on a 7–10 day schedule. Apply in sufficient water by ground or air to obtain thorough coverage of foliage. For improved disease control, this product may be tank-mixed with oil or other fungicides registered for Sigatoka control at label rates

(Other crops [outside crop groups])

ARTICHOKE

HOPS

- Minimum spray volumes for hop growth stages are as follows:
 - Emergence to Training: Apply 1–2 quarts this product per acre (1.5 – 3 tbsp. per 1000 sq. ft.) using a minimum spray volume of 20 gallons per acre (0.5 gallon per 1000 sq. ft.). Coverage will vary with the size of the vines and the type of spray equipment. Apply adequate spray volume to achieve complete spray coverage.
 - Training to Wire-Touch: Apply 1–2 quarts this product per acre (1.5 – 3 tbsp. per 1000 sq. ft.) using a minimum spray volume of 50 gallons per acre (1 gallon of water per 1000 sq. ft.). Coverage will vary with the size of the vines and the type of spray equipment. Apply adequate spray volume to achieve complete spray coverage.
 - Wire-Touch through Harvest: Apply 2–4 quarts (3 – 6 tbsp.) of this product using a minimum of 100 gallons of water per acre (2.5 gallons of water per 1000 sq. ft.). Higher water volumes may be necessary to achieve thorough coverage after side arms develop. Do not apply more than 4 quarts of product per acre (6 tbsp. per 1000 sq. ft.) per application. Apply adequate spray volume to achieve complete spray coverage. Use the higher rates when moderate to high disease pressure is present or expected.
 - For control of downy mildew, tank mix this product with another fungicide labeled for Downy Mildew control and re-apply at a 7-day interval or according to the label directions of the tank mix (ingredient) (partner).
-

PEANUT

(Apply at 1 pint per acre for plant health.)

HEMP

QUINOA

SUGARCANE

(Apply at 1 pint per acre for Plant Health Applications.)

CROTALARIA, SESSBANIA, KENAF

FLOWERING PLANTS

BEDDING PLANTS

GRASS GROWN FOR SEED

0.5–2.5 fluid ounces per 100 lbs. seed for SEED TREATMENT applications

ORNAMENTALS

SWITCHGRASS, MISCANTHUS

TOBACCO

For Target Spot: Apply this product at a rate of 2 quarts per acre in a minimum of 50 gallons of water per acre. (Apply at 1 pint per acre for Plant Health Applications.)

TURFGRASS AND ORNAMENTAL GRASSES: Bluegrass, Bentgrass, Bermudagrass, Dichondra, Fescue, Orchardgrass, *Poa annua*, Ryegrass, St. Augustine, Zoysia, mixtures

1–3 fluid ounces per 1000 square feet in a minimum of 1.5 gallons of water for FOLIAR applications, or

3–8 pints per acre in a minimum of 50 gallons of water for FOLIAR applications

This product aids in control of turf diseases and improves turf quality. For improved performance under moderate to severe disease pressure, reduce spray intervals or use this product in a tank mix or rotational program with other registered fungicides.

INTEGRATED PEST MANAGEMENT (IPM)

Many conventional fungicides have been tested in an IPM regime with REGALIA® Bioprotectant Concentrate with very satisfactory results. One of the major objectives of IPM has been to reduce the probability of disease resistance development to a particular active ingredient.

The alternate use of (1–2 sprays) followed by a conventional, registered fungicide (1–2 sprays) has been successfully used in many crops. In addition, the use of tank mixes with a conventional fungicide has also been successful.

Follow label instructions of the particular registered product: Do not exceed amounts or treatment intervals on the label.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place. Avoid freezing.

Pesticide Disposal: To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

Container Handling (5 gallons or less): Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill or by incineration. Do not burn unless allowed by state and local ordinances.

Container Handling (over 5 gallons): Non-refillable container. Do not reuse or refill this container Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill or by incineration. Do not burn unless allowed by state and local ordinances.

Container Handling (refillable containers – 265 gallon tote):

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. When empty, return to point of sale or offer for recycling if available or reconditioning if appropriate.

Pro Farm Group is a member of the Ag Container Recycling Council.

Visit <http://www.acrecycle.org/contact.html> for information on how to arrange pick-up of this empty pesticide container.



WARRANTY

To the extent consistent with applicable law, the seller makes no warranty, expressed or implied, of merchantability, fitness or otherwise concerning use of this product. To the extent permitted by applicable law, the user assumes all risks of use, storage or handling that are not in strict accordance with the accompanying directions.

Label date:

Made in the U.S.A.

US Patents No. 4,863,734 and No. 5,989,429

REGALIA® is a trademark of Pro Farm Group, Inc.

Name and logo of Pro Farm Group are registered trademarks of Pro Farm Group, Inc.

© Pro Farm Group, Inc.

REGALIA® Bioprotectant Concentrate; EPA Reg. No. 84059-3

MASTER LABEL – MBI-LBL-4 r23 v4 January 28, 2025

Page 26 of 66

SUBLABEL A OPTIONAL CLAIMS

1. Boost plants' defenses [for stronger healthier plants]
2. Strengthens plants' immunity
3. Improves plant health
4. Controls/Prevents powdery mildew, leaf spot and rust
5. Controls/Prevents black spot on rose
6. Labeled Fungal and bacterial disease control
7. Use on labeled fruits, vegetables and ornamentals
8. Can be applied as a pre-plant dip [for improved plant health]
9. Can be applied as a soil drench application
10. For use on ornamental plants and edible crops/fruits/vegetables.
11. For use on vegetables, roses, fruits, berries, nuts, flowers, bedding plants, (ornamental) trees and shrubs
12. REGALIA® Bioprotectant Concentrate may be used on vegetable(s) [crops], roses, fruits, berries, nuts, flowers, foliage, (ornamental) trees and shrubs.
13. REGALIA® Bioprotectant Concentrate is a broad spectrum fungicide used for the control or suppression of a broad range of labeled foliar [fungal and bacterial] diseases.
14. (Active ingredient is) a plant extract to boost the plants' defense mechanisms to protect against labeled fungal and bacterial diseases, and to improve plant health.
15. Active ingredient (is) made from a plant extract (botanical extract)(plant-based)
16. REGALIA® Bioprotectant Concentrate is an extract from the plant *Reynoutria sachalinensis*
17. REGALIA® Bioprotectant Concentrate can be applied up to and on the day of harvest [on all fruits and vegetables].
18. Made in the U.S.A.
19. This container is made with X% recycled material
20. Guaranteed results(*)
21. Label date:
22. US Patents No. 4,863,734 and No. 5,989,429
23. REGALIA® is a registered trademark of Pro Farm Group, Inc.
24. Name and logo of Pro Farm Group are registered trademarks of Pro Farm Group, Inc.
25. © insert company copyright information
26. World rights reserved
27. Distributed by: insert company name and address
28. [For] questions/comments



- 29.
30. GENERAL INFORMATION: REGALIA® Bioprotectant Concentrate is an extract from the plant *Reynoutria sachalinensis*. REGALIA® Bioprotectant Concentrate applied to actively growing plants will improve plant health, and will help make the treated portions resistant to labeled plant diseases. (Plant health benefits often result in greater yields at harvest, especially when crops/plants are stressed by pathogens or environmental conditions).
31. MODE OF ACTION: REGALIA® Bioprotectant Concentrate contains an extract obtained from *Reynoutria sachalinensis* plant material composed of active chemical compounds. Application of the extract activates the immune response, increasing the plant's defense system through a five-fold increase in phenolics and antioxidants, and cell wall strengthening. These immune responses, sometimes referred to as induced systemic resistance or systemic acquired resistance, protects treated plants from numerous diseases.
When applied at rates and timing for disease control, the translaminar activity of REGALIA® Bioprotectant Concentrate provides induced resistance against important diseases and acts within one to two days of application.
REGALIA® Bioprotectant Concentrate should be used as a preventative treatment prior to the development of disease

symptoms. Repeat foliar applications per label instructions, following a 7–14-day interval to maintain plant immune response induction and to protect new plant growth.

(In addition to foliar applications, REGALIA® Bioprotectant Concentrate can be used in multiple application methods as a plant dip, soil drench, pre-plant incorporated soil application, in-furrow spray, or applied through drip irrigation to control or suppress certain soil-borne diseases and to promote healthy root growth and improve above ground plant growth.)

(When applied at rates and timing for plant health effects, the induced plant defense responses minimize the impacts of stress and disease, resulting in optimized yields at harvest. Applying Regalia® Bioprotectant Concentrate has been shown to increase leaf chlorophyll content and increase soluble protein content in some crops. These effects often lead to improved crop quality and/or yields.)

32. REGALIA® Bioprotectant Concentrate is (an extract from the plant *Reynoutria sachalinensis*) for use on labeled (ornamental plants), (turf), (agricultural crops), (edible crops), (non-edible crops), (food crops), (non-food crops), (feed crops), or (non-feed crops).
33. Optional Language: (*) and (*= Not for use in California)
34. Repackaging or relabeling of this product without express written permission from Pro Farm Group is prohibited.
35. Biofungicide
36. [UPC code]
37. [RF code]
38. Bio with Bite™
39. For (labeled) disease control
40. For (labeled) fungal control
41. Can be used in sustainable production
42. For use in sustainable production
43. For maximum harvest
44. US MRL exempt
45. Minimal PPE
46. Protection from Multiple Labeled Diseases
47. Leaves no detectable residues
48. Product(s) thoroughly tested
49. Proven results, since 2007
50. Trial Tested
51. (number)+ trials
52. Read full label before use
53. Prevents (labeled disease) and (labeled fungal) build up
54. Protection from labeled bacterial and fungal disease
55. REGALIA® Bioprotectant Concentrate can be applied by (any labeled use pattern) to protect against (labeled diseases) and (labeled fungal pathogens).
56. 2-10 tablespoons per 1,000 sq. ft. [must be consistent with rates to be listed on label]
57. For turf
58. For recreational turf and landscapes [when crop is listed]
59. (Specify pest:) (Tank-mix) (or rotate) with a (fungicide) for improved control.
60. Refer to the table in the SOIL TREATMENT USE DIRECTIONS (In-Furrow Applications) section to determine the proper rate per 1000 foot of row.
61. REGALIA® Bioprotectant Concentrate should be used as part of an Integrated Pest Management System.
62. REGALIA® Bioprotectant Concentrate can be applied following a soil fumigant.
63. Use the high(er) labeled rate when high(er) (disease)(fungal pathogen) pressure is expected.
64. Optional Language: (*) and (*= Not labeled for this use in California)
65. [optional listing of all pests, alternatively list specific pests for each crop]
66. Bioprotectant Concentrate
67. Outdoor
68. See back of booklet [or other specific location] for precautionary statements and first aid
69. See attached booklet for complete directions for use and precautionary statements

70. Indoor/Outdoor
71. See attached booklet for full label
72. *rate may require more than one package
73. (see (side) (back) (inside) (other) panel for additional (first aid) (and) (precautionary) statements)
74. (Refer to inside of label booklet for additional precautionary information and Directions for Use including First Aid and Storage and Disposal.)
75. May be used in hydroponic crop production.
76. Peel here (for (additional))(more)(complete) label instructions)
77. Plant-Based Actives
78. Results Done Right
79. Starts Working Immediately
80. Proven Effective
81. Proven Efficacy
82. Highly Effective Protection
83. Field Tested
84. 20% more vs. XX lb size [XX lbs]
85. 20% more vs XX size [[XX [ounce] [lbs]]
86. Xx% more vs xx lb size
87. TWIN [PACK/PAK]
88. Value [Pack/Pak]
89. VALUE [SIZE/SIZED]
90. Great Value
91. Club [Pack/Pak/Size]
92. Combo [Pack/Pak/Size]
93. Bonus [Size/Pack/Pak]
94. A [Insert dollar amount] Value
95. Treats up to XX sq. ft. (XX size)
96. Covers XX sq. ft (all pests) (XX pound size)
97. 100% Satisfaction Guaranteed
98. Guaranteed Results [or your money back]
99. {100%} Satisfaction [Guaranteed] or your money back
100. Money Back Guarantee
101. © [copyrighted material]
102. TM [trademarked]
103. Indoor/Outdoor
104. See back of booklet for precautionary statements and first aid
105. See attached booklet for complete directions for use and precautionary statements
106. Grow a better world
107. Take root in something bigger
108. The power Science meets the force of nature
109. A new wave of sustainable plant wellness products
110. Help your plants live their best life
111. BioUnite
112. BioFocus
113. BioTrust
114. BioFusion
115. Terraconnect
116. Bottle (body) contains XX% (post-consumer) recycled plastic (excluding dye) (excluding colorant)
117. Bottle (body contains XX% (post-consumer) recycled content (excluding dye) (excluding colorant)
118. Recyclable plastic bottle (body) made of XX% (post-consumer) recycled plastic (excluding dye) (excluding colorant)
119. Recyclable plastic bottle (body) made of XX% (post-consumer) content (excluding dye) (excluding colorant)

Sublabel B: Professional Landscape Use (Terrestrial nonfood)

REGALIA® Bioprotectant Concentrate

A plant extract to boost the plants' defense mechanisms to protect against certain fungal and bacterial diseases, and to improve plant health.

Active ingredient: Extract of *Reynoutria sachalinensis* 5 %
Other ingredients: 95 %
Total 100 %

EPA Reg. No. 84059-3

EPA Est. No. xxxxx-xx-xxx

GROUP P5 FUNGICIDE

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID	
IF SWALLOWED:	Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED:	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
IF IN EYES:	Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or if going for treatment. You may also contact 1-800-222-1222 for emergency medical treatment information.	

()( CAN BE USED IN ORGANIC PRODUCTION)

) (For Organic Production)( For Organic Use - OMRI.org)()

(Batch)(LOT) No: xxxx (Printed on container)

Net Contents: xxxx

(Manufactured [for] [by]): Pro Farm Group, Inc. 1530 Drew Ave., Davis, CA 95618 (1-877-664-4476;
info@profarmgroup.com)

REGALIA® Bioprotectant Concentrate; EPA Reg. No. 84059-3

MASTER LABEL – MBI-LBL-4 r23 v4 January 28, 2025

Page 30 of 66

(Name and logo of Pro Farm Group are registered trademarks of Pro Farm Group, Inc.)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear goggles or safety glasses. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Waterproof gloves
- Protective eyewear

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS

For terrestrial uses: do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exemptions pertaining to the statements on this label about personal protective equipment (PPE) and the restricted-entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. The REI does not apply when this product is used for seed treatment at planting or in hopper box treatments.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

For commercial seed treatment uses: Keep unprotected persons out of treatment area until seeds have dried or been packaged.

For all other non-WPS uses: Keep unprotected persons out of treated areas until sprays have dried.

GENERAL INFORMATION

REGALIA® Bioprotectant Concentrate is an extract from the plant *Reynoutria sachalinensis* for use on ornamental plants, turf, and edible crops. REGALIA® Bioprotectant Concentrate applied to the above ground portions of actively growing plants (see DIRECTIONS FOR USE) will help protect treated portions from certain plant diseases and will improve plant health. Plant health benefits often result in greater yields at harvest, especially when crops are stressed by pathogens or environmental conditions. Use REGALIA® Bioprotectant Concentrate as a preventative rather than a curative application. Apply prior to disease infestation to protect the growing leaf tissue, flowers and above ground fruit and vegetables. See specific information below for diseases controlled and use rates on ornamental plants, turf, and edible crops.

REGALIA® Bioprotectant Concentrate can be used in multiple application methods to the soil or plant roots to control or suppress certain soil-borne or foliar diseases and to promote healthy plant growth. See below for specific information on diseases controlled and use rates.

MODE OF ACTION

The extract obtained from *Reynoutria sachalinensis* plant material contains bioactive compounds. The extract, when applied to the host plant, activates the plant's defense system to increase phenolics and antioxidants, and strengthen cell walls. This mode of action is classified as induced systemic resistance or systemic acquired resistance.

When applied at rates and timing for disease control, the induced resistance against important diseases provides translaminar activity, which takes place within one to two days of application. Repeat foliar applications per label instructions. Use REGALIA® Bioprotectant Concentrate, therefore, as a preventative treatment. In addition to foliar applications, REGALIA® Bioprotectant Concentrate can be used in multiple application methods as a plant dip, soil drench, pre-plant incorporated soil application, in-furrow spray, or applied through drip irrigation to control or suppress certain soil-borne diseases and to promote healthy root growth and improve above ground plant growth.

When applied at rates and timing for plant health effects, the improved plant defense responses minimize the impacts of stress and disease, resulting in optimized yields at harvest. Applying Regalia® Bioprotectant Concentrate has been shown to increase leaf chlorophyll content and increase soluble protein content in some crops. These effects often lead to improved crop quality and/or yields.

MIXING AND APPLICATION INSTRUCTIONS

– SHAKE WELL PRIOR TO USE –

Mixing instructions: Prepare no more mixture than is required for the immediate operation. Agitate the solution continuously during mixing and application. Mechanical mixing is recommended for proper mixing of REGALIA® Bioprotectant Concentrate mixtures.

REGALIA® Bioprotectant Concentrate alone: Add ½ of the required amount of water to the mix tank. With the agitator running, add the REGALIA® Bioprotectant Concentrate to the mix tank. Continue agitation while adding the remainder of the water. Begin application of the solution after the REGALIA® Bioprotectant Concentrate has completely dispersed into the mix water. Maintain agitation until all the mixture has been applied.

REGALIA® Bioprotectant Concentrate + tank mixtures: Add ½–¾ of the required amount of water to the mix tank. Start the agitation before adding any tank mix ingredients. In general, tank mix ingredients should be added in this order: wettable powders, dry flowable formulations, liquid flowable formulations, and emulsifiable formulations such as REGALIA® Bioprotectant Concentrate. Always allow each tank mix ingredient to become completely dispersed before adding the next component. Maintain continuous agitation until all components have been dispersed and throughout the application process. After all components are completely dispersed add the remainder of the water. REGALIA® Bioprotectant Concentrate cannot be mixed with another product with a prohibition against mixing. Use of the tank mix must be in accordance with the most restrictive label limitations and precautions. **Do not pre-mix REGALIA® Bioprotectant Concentrate with any other tank mix component prior to adding to the spray tank.**

Note: When using REGALIA® Bioprotectant Concentrate in tank-mixtures, all products in water soluble packaging should be added to the tank before any other tank-mix ingredient, including REGALIA® Bioprotectant Concentrate. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank-mix ingredient to the tank.

Compatibility: Do not combine REGALIA® Bioprotectant Concentrate in the spray tank with pesticides, adjuvants, or fertilizers if there has been no previous experience or use of the combination to show it is physically compatible, effective, and non-injurious under your use conditions. REGALIA® Bioprotectant Concentrate is compatible with many commonly used pesticides, fertilizers, adjuvants, and surfactants, but has not been evaluated with all potential combinations. To ensure compatibility of the tank mix combinations, evaluate prior to use as follows: Using a suitable container, add the proportional amounts of product to water. Add wettable powders first, then water dispersible granules, then liquid flowables, and lastly, emulsifiable concentrates. Mix thoroughly and let stand for at least five minutes. If the combination stays mixed or can be remixed, it is physically compatible. Test the mix on a small portion of the crop to be treated to ensure that a phytotoxic response will not occur as a result of the application.

Application Instructions: REGALIA® Bioprotectant Concentrate is a micro-emulsion concentrate consisting of certain ingredients extracted from *Reynoutria sachalinensis*. Use 50-mesh nozzle screens or larger. Use higher water volumes with larger sized crops and extensive foliage to obtain thorough coverage. Do not use carrier volumes and/or adjuvants that create spray runoff or drip-accumulation at the leaf margins or at the base of fruit or on the harvested commodity.

See FOLIAR AERIAL and FOLIAR GROUND APPLICATION section for use directions.

See BACKPACK/HANDHELD SPRAYER section for use directions.

See CHEMIGATION section for use directions.

See PRE-PLANT DIP section for use directions.

See SEED TREATMENT section for use directions.

See SOIL TREATMENT section for use directions.

FOLIAR AERIAL APPLICATION INSTRUCTIONS

0.5–1 quart per acre for FOLIAR (AERIAL) applications

- (For aerial applications, apply this product in a minimum of (2 to 10) gallons of water per acre.)
- (Apply this product preventatively or when the first disease symptoms are visible and reapply every 7–14 days)
- (For improved performance, use this product in a tank mix or rotational program with other registered fungicides.) (Tank mix this product with other registered fungicides for improved disease control under (heavy) (high disease) pressure.)
- (Repeat applications at (5-14) (7-10) (7–14) day intervals) (depending upon crop growth and disease pressure)
- (Under moderate to heavy disease pressure, tank mix this product with another fungicide.)

- (Consult your local Extension Specialist or Crop Consultant regarding the optimum timing of fungicide applications)
- (For grasses including forage, fodder, hay, and cereals, and for non-grass animal feed:) (It is important to apply this product at the flag leaf stage to maximize yield.)
- (Increasing the amount of water applied per acre will improve product performance.)
- (Follow all instructions to reduce aerial drift.)

AERIAL DRIFT REDUCTION ADVISORY INFORMATION

GENERAL: Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. Where states have more stringent regulations, they should be observed. Note: This section is advisory in nature and does not supersede the mandatory label requirements.

INFORMATION ON DROPLET SIZE: The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply droplets large enough to provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

CONTROLLING DROPLET SIZE: Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets. Pressure – Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When high flow rates are needed, use higher flow rate nozzles instead of increasing pressure. Number of Nozzles – Use the minimum number of nozzles that provide uniform coverage. Nozzle Orientation – Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential. Nozzle Type – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

BOOM WIDTH: For aerial applications, the boom width must not exceed 75% of the wingspan or 90% of the rotary blade. Use upwind swath displacement and apply only when wind speed is 3–10 mph as measured by an anemometer. Use medium or coarser spray according to ASAE 572 definition for standard nozzles or VMD for spinning atomizer nozzles. If application includes a no-spray zone, do not release spray at a height greater than 10 feet above the ground or crop canopy.

APPLICATION HEIGHT: Do not make application at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

SWATH ADJUSTMENT: When applications are made with a crosswind, the swath will be displaced downward. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.).

WIND: Drift potential is lowest between wind speeds of 2–10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY: When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

TEMPERATURE INVERSIONS: Do not apply during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small, suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SENSITIVE AREAS: The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas). Do not allow spray to drift from the application site and contact people, structures people occupy at any time and the associated property, parks and recreation areas, non-target crops, aquatic and wetland areas, woodlands, pastures, rangelands, or animals.

FOLIAR (GROUND) (EQUIPMENT) APPLICATION USE DIRECTIONS

REGALIA® Bioprotectant Concentrate can be applied in most commonly-used ground application equipment, such as tractor-mounted boom, airblast, high clearance, hose-end, backpack, and other pressurized sprayers; or hand-held sprayers; foggers or mist blowers. Thorough coverage is necessary to provide good disease control.

0.5–4 quarts per acre (0.5 – 6 tbsp. per 1000 sq. ft.) for FOLIAR (GROUND) (EQUIPMENT) applications

- (For foliar applications)(to optimize disease control and to maximize yields,)(apply ([specify volume])(of) (this product) (preventatively) in (a minimum of) 15–100 gallons of water per acre) (0.5 – 2.5 gallons of water per 1000 sq. ft.) (by (ground) (equipment)) (prior to disease development using) (sufficient volume for thorough coverage.) (or when the first symptoms of disease are visible) (at 7 day spray intervals or) (as soon as the first symptoms of disease appear) (or when environmental conditions are conducive to rapid disease development.) (Begin application soon after emergence or transplant, and when conditions are conducive to disease development.) (when applied alone, or at 1–4 quarts per acre (1.5 – 6 tbsp. per 1000 sq. ft.) when tank mixed with another fungicide preventatively in a minimum of 50 gallons of water per acre (1 gallons per 1000 square feet))
- (Additional sprays beyond second cover may be needed on susceptible varieties, or when environmental conditions are conducive to rapid disease development. Use high label rate and shorter spray intervals when conditions are conducive to rapid disease development.) (Use shorter intervals when conditions are conducive to rapid disease development.)
- (Avoid excessive amounts of water that result in the runoff of spray material)
- (Continue sprays at 7-day intervals or as needed) (For foliar applications, mix this product concentrate with water at a concentration of 2–4 quarts per 100 gallons of water (1.5 – 2.5 tbsp. per gallon of water) when used alone or 1–4 quarts per 100 gallons of water (1 – 2.5 tbsp. per gallon of water) when tank mixed with another fungicide) (If necessary, repeat applications at a 7-day interval.)
- (Begin applications preventatively (before disease symptoms become visible) at the 4 to 6-leaf stage and treat at 7–10 day intervals as needed prior to sale or harvest. Spray until just before point of runoff.)
- (Increase water volume as plant size increases.)
- (Do not exceed 1.0% v/v of the applied solution.)
- (Apply this product preventatively or when the first disease symptoms are visible) (and reapply every 7–14 days.)
- (Repeat applications at (5-14) (7–10) (7-14) day intervals) (depending upon crop growth and disease pressure)
- (Under moderate to heavy disease pressure, tank-mix this product with another fungicide.)(When the plants are) (under high disease pressure, tank-mix this product with another fungicide for more effective control.)(Tank mix this product with other registered fungicides for improved disease control under heavy pressure.)(For improved performance,(apply 0.5–2 quarts per acre (0.5 – 3 tbsp. per 1000 sq. ft.) of) (use) this product in a tank mix (or rotational program) with other registered fungicides)(When tank mixed with other fungicides, use 1–2 quarts of REGALIA® Bioprotectant Concentrate per acre (1.5 – 3 tbsp. per 1000 sq.

- ft.).) (This product can be tank mixed at the lower rate with another registered fungicide under heavy disease pressure)
- (For trees and vines:) (Dilute applications: this product can be applied by ground equipment to (vine and) tree crops in dilute applications of 100–400 gallons of water per acre (2.5 – 9 gallons of water per 1000 square feet). Apply this product at a rate of 2–4 quarts per acre (3 – 6 tbsp. per 1000 sq. ft.) when applied alone, or at 1–4 quarts per acre (1.5 – 6 tbsp. per 1000 sq. ft.) when tank mixed with another fungicide. Avoid excessive amounts of water that result in the runoff of spray material)
 - (Consult your local Extension Specialist or Crop Consultant regarding the optimum timing of fungicide applications.)
 - (For grasses including cereals, forage, fodder, hay, and for non-grass animal feed:) (It is important to apply this product at the flag leaf stage to maximize yield.)
 - (Apply product at 1-4 quarts per acre (1.5 – 6 tbsp. per 1000 sq. ft.) when applied in a tank mix, or at 2-4 quarts per acre (3 – 6 tbsp. per 1000 sq. ft.) when applied alone.)
 - (For concentrated ground applications, apply this product at 0.5–1.5 quarts per acre (0.5 – 2.5 tbsp. per 1000 sq. ft.) in a minimum of 10 gallons of water per acre (1 quart of water per 1000 sq. ft.).)
 - (For plant health applications, apply (10-14 days) (two to three weeks) prior to cutting) (Foliar applications of Regalia to improve plant health can be made during the period from layby to contact sprays)
 - (This product may be used to control certain diseases of container, bench, flat, plug, bed, or field-grown ornamentals in **shade-houses, outdoor nurseries, retail nurseries, and other landscape areas.**)

BACKPACK/HAND-HELD SPRAYER USE DIRECTIONS

The use rate for REGALIA® Bioprotectant Concentrate when applied alone or as an alternate spray in a backpack or hand-held sprayer is 1.2 – 2.5 tablespoons (Tbsp.) (0.6-1.3 fluid ounces [fl. oz.]) per gallon of water (0.5 – 1.0% v/v dilution of REGALIA® Bioprotectant Concentrate) applied at 1 – 2.5 gallons per 1000 square feet (sq. ft.) (25 – 100 gallons of water per acre). When tank mixed with another fungicide, the use rate for REGALIA® Bioprotectant Concentrate in a backpack or hand-held sprayer is 0.6 – 2.5 tablespoons (0.3 – 1.3 fluid ounces) per gallon of water applied at 1 – 2.5 gallons per 1000 square feet (50 –100 gallons of water per acre). Use higher water volumes with larger sized crops and extensive foliage in order to secure thorough coverage. Do not use carrier volumes and/or adjuvants that create spray runoff or drip-accumulation at the base of fruit or on the harvested commodity. See specific application recommendations pertaining to each crop for additional details.

Use rate for REGALIA® Bioprotectant Concentrate when applied in small volumes

	Tablespoons REGALIA® Bioprotectant Concentrate per Gallon of Water	Fluid Ounces REGALIA® Bioprotectant Concentrate per Gallon of Water	v/v Dilution of REGALIA® Bioprotectant Concentrate	Applied (diluted) Gallons per 1,000 sq. ft.
Applied alone or as an alternate spray	1.2 – 2.5	0.6 – 1.3	0.5 – 1.0% v/v	1 – 2.5
Tank mixed with another fungicide	0.6 – 2.5	0.3 – 1.3	0.25 – 1.0% v/v	1 – 2.5

CHEMIGATION USE DIRECTIONS

(For (control) (suppression) of listed foliar diseases and to improve plant health apply this product through overhead sprinkler, center pivot sprinkler systems, lateral move, end tow, side (wheel) roll, traveler, solid set, or hand move irrigation systems. Do not use reclaimed water for application of this product.)

(For (control) (suppression) of listed soil-borne diseases and to improve plant health, apply this product only through flood (basin), furrow, border or drip (trickle) irrigation systems. Do not use reclaimed water for application of this product.)

0.5–4 quarts per acre (0.7 – 6 tbsp. per 1000 sq. ft.) for CHEMIGATION applications

- For chemigation applications (for improved plant growth) (and suppression of soil-borne diseases), apply this product through (drip) irrigation immediately after transplant and at 14-day intervals or begin 14 days after transplant when soil drench applications are used.

Spray preparation

First prepare a suspension of REGALIA® Bioprotectant Concentrate in a mix tank. Fill tank ½ to ¾ the desired amount of water. Start mechanical or hydraulic agitation. Add the required amount of REGALIA® Bioprotectant Concentrate, and then the remaining volume of water. Then set the irrigation system to deliver a minimum of 0.1 to 0.3 inch of water per acre. Start irrigation system and uniformly inject the suspension of REGALIA® Bioprotectant Concentrate into the irrigation water line so as to deliver the desired rate per acre. Inject the suspension of REGALIA® Bioprotectant Concentrate with a positive displacement pump into the main line after the filter, and ahead of a right angle turn to insure adequate mixing. Any questions on calibration should be directed to your State Extension Service Specialists, to equipment manufacturers or other experts.

Do not combine REGALIA® Bioprotectant Concentrate with pesticides, surfactants or fertilizers for application through chemigation equipment unless prior experience has shown the combination physically compatible, effective and non-injurious under conditions of use. REGALIA® Bioprotectant Concentrate has not been fully evaluated for compatibility with all adjuvants or surfactants. Conduct a spray compatibility test if a mixture with adjuvants or surfactants is planned.

CHEMIGATION

General Requirements –

- 1) Apply this product only through a drip or trickle system or center pivot sprinkler system, lateral move, end tow, side (wheel) roll, traveler, solid set, hand move, flood (basin), furrow, border or drip (trickle) irrigation systems.
- 2) Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- 3) If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.
- 4) Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- 5) A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Application Instructions for All Types of Chemigation –

- 1) Remove scale, pesticide residues, and other foreign matter from the chemical supply tank and entire injector system. Flush with clean water. Failure to provide a clean tank, void of scale or residues may cause product to lose effectiveness or strength.
- 2) Determine the treatment rates as indicated in the directions for use and make proper dilutions. Product can be applied continuously or at any time during the water application.
- 3) Prepare a solution in the chemical tank by filling the tank with the required water and then adding product as required
- 4) The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- 5) The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- 6) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

- 7) Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Specific Requirements for Chemigation Systems Connected to Public Water Systems –

- 1) Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- 2) Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- 3) Do not apply when wind speed favors drift beyond the area intended for treatment.

Specific Requirements for Sprinkler Chemigation –

- 1) The system must contain a functional check valve, vacuum relief valve and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- 2) The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- 3) Do not apply when wind speed favors drift beyond the area intended for treatment.

Specific Requirements for Flood (Basin), Furrow and Border Chemigation –

- 1) Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as a drop structure or weir box to decrease potential for water source contamination from backflow if water flow stops.
- 2) The systems utilizing a pressurized water and pesticide injection system must meet the following requirements:
 - a. The system must contain a functional check valve, vacuum relief valve and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
 - b. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Specific Requirements for Drip (Trickle) Chemigation –

- 1) The system must contain a functional check valve, vacuum relief valve and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- 2) The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

PRE-PLANT DIP USE DIRECTIONS

Apply REGALIA® Bioprotectant Concentrate as a pre-plant dip for improved plant health and suppression of certain soil-borne diseases.

0.5–4 quarts per 100 gallons (1 tsp. – 2.5 tbsp. per gallon) of water for PLANT DIP (bare root) applications

- For plant dip applications for improved plant growth and suppression of soil-borne diseases, apply this product in a 0.25–0.5 % v/v suspension (1–4 quarts of this product per 100 gallons (or 1 – 2.5 tbsp. per gallon) of water as a pre-plant dip immediately prior to transplanting. Do not allow roots or plugs to be submerged in excess of 15 minutes prior to removal from the REGALIA® Bioprotectant Concentrate treatment.

SEED TREATMENT USE DIRECTIONS

REGALIA® Bioprotectant Concentrate can be applied as a seed (or seed piece) dressing for suppression of soil-borne diseases to improve early-season root growth. REGALIA® Bioprotectant Concentrate may be applied with other registered seed treatment insecticides and fungicides through standard slurry- or mist-type commercial seed treatment equipment. (REGALIA® Bioprotectant Concentrate can also be used in on-farm hopper-box or planter-box treatments.)

If using REGALIA® Bioprotectant Concentrate in a tank mixture with other seed treatment products, observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank-mix ingredient label. Do not exceed label rates and the most restrictive label precautions and limitations must be followed. Do not mix this product with any product which prohibits such mixing.

0.5–15 fluid ounces per 100 lbs. seed for SEED TREATMENT applications

- For suppression of soil-borne diseases, apply this product as a seed treatment at the rate of 0.5–15 fl. oz. per 100 lbs. seed.

(Do not apply this product through any type of irrigation system.)

Note: Federal law requires that bags containing treated seeds shall be labeled with the following information:

"This seed has been treated with Regalia® Bioprotectant Concentrate fungicide. Do not use for food, feed, or oil purposes. Store away from feed and foodstuffs."

Treated seed bagged for later use must contain an EPA-approved dye or colorant that imparts an unnatural color to the seed.

SOIL TREATMENT USE DIRECTIONS

REGALIA® Bioprotectant Concentrate can be applied by pre-plant incorporated soil application, soil drench, in-furrow spray or through drip (trickle) irrigation to improve plant health and to protect against certain soil-borne diseases.

In general, REGALIA® Bioprotectant Concentrate can be applied by the following methods, unless specified differently in the SELECTED CROPS section:

Pre-plant Incorporated Soil Applications:

- 1-4 quarts per acre in 10 – 100 gallons of water for PRE-PLANT INCORPORATED SOIL applications
- For pre-plant applications, apply this product with a broadcast spray and immediately incorporate into the soil profile prior to seeding or transplanting the crop to reduce transplant shock, suppress soil-borne diseases and improve root growth.

Soil Drench Applications:

0.5–3 quarts per 100 gallons (1 tsp. – 2 tbsp. per gallon) of water for SOIL DRENCH applications

- For soil drench applications, apply this product at a concentration of 1–3 quarts per 100 gallons (1 – 2 tbsp. per gallon) of water, and at a sufficient rate to thoroughly soak the growing media and root zone. Make an initial application of this product during or shortly after transplant to reduce transplant shock, suppress soil-borne diseases and improve root growth. (Multiple drench applications can be made on a 10–14 day interval.)

In-Furrow, Shank-In and Injected Applications:

(REGALIA® Bioprotectant Concentrate can be shanked-in or injected into the soil alone, or with most types of liquid nutrients.)

0.5–4 quarts per acre or 1–10 fluid ounces per 1000 ft. row for IN-FURROW applications

- For in-furrow applications, at planting apply this product as an in-furrow spray at the rate of 0.5–4 quarts per acre or 1–10 fluid ounces per 1000 feet of row according to the chart (in the SOIL TREATMENT USE DIRECTIONS section)(below). Apply this product in 5–15 gallons of water so as the spray is directed into the seed furrow just before the [choose 1 or more] (seeds) (or seed pieces for potatoes) are covered.

Rate	In-Furrow Application Rates Product per 1000 ft. row (fl. oz.)
------	--

Per Acre	12" Rows	14" Rows	16" Rows	18" Rows	20" Rows	22" Rows	24" Rows	26" Rows	28" Rows	30" Rows	32" Rows	34" Rows	36" Rows	38" Rows	40" Rows
1 quart	0.73	0.86	0.98	1.1	1.2	1.3	1.5	1.6	1.7	1.8	2.0	2.1	2.2	2.3	2.4
2 quart	1.5	1.7	2.0	2.2	2.4	2.7	2.9	3.2	3.4	3.7	3.9	4.2	4.4	4.7	4.9
3 quart	2.2	2.6	2.9	3.3	3.7	4.0	4.4	4.8	5.1	5.1	5.9	6.2	6.6	7.0	7.3
4 quart	2.9	3.4	3.9	4.4	4.9	5.4	5.9	6.4	6.9	7.3	7.8	8.3	8.8	9.3	9.8

12" = 43,560 row ft./acre, 14" = 37,337 row ft./acre, 16" = 32,670 row ft./acre, 18" = 29,040 row ft./acre, 20" = 26,136 row ft./acre, 22" = 23,760 row ft./acre, 24" = 21,780 row ft./acre, 26" = 20,105 row ft./acre, 28" = 18,669 row ft./acre, 30" = 17,424 row ft./acre, 32" = 16,315 row ft./acre, 34" = 15,374 row ft./acre, 36" = 14,520 row ft./acre, 38" = 13,754 row ft./acre, 40" = 13,068 row ft./acre.

**FOR USE ON THE FOLLOWING CROPS FOR CONTROL OR SUPPRESSION OF (THE FOLLOWING) (DISEASES)
(PATHOGENS)(PESTS)**

The use rate for REGALIA® Bioprotectant Concentrate when applied alone or as an alternate spray is 2–4 quarts per 100 gallons (1.5 – 2.5 tbsp. per gallon) of water (0.5–1.0% v/v dilution of REGALIA® Bioprotectant Concentrate) applied at 15–100 gallons of water per acre (0.5 - 2.5 gallons per 1000 square feet). When tank mixed with another fungicide, the use rate for REGALIA® Bioprotectant Concentrate is 1–4 quarts in 100 gallons (1 – 2.5 tbsp. per gallon) of water applied at 15–100 gallons of water per acre (0.5 – 2.5 gallons per 1000 square feet). Use higher water volumes with larger sized crops and extensive foliage in order to secure thorough coverage. Do not use carrier volumes and/or adjuvants that create spray runoff or drip-accumulation at the leaf margins or at the base of fruit or on the harvested commodity. See specific application recommendations pertaining to each crop for additional details.

(This product can be used to control certain diseases of container, bench, flat, plug, bed or field-grown ornamentals and edible crops in shade-houses, outdoor nurseries, retail nurseries and other landscape areas. For application on the crops and diseases listed, the recommended use rate for REGALIA® Bioprotectant Concentrate is 2–4 quarts in 100 gallons (1.5 – 2.5 tbsp. per gallon) of water (0.5–1.0% v/v dilution of REGALIA® Bioprotectant Concentrate) sprayed until just before point of runoff. When tank mixed with another fungicide, the use rate for REGALIA® Bioprotectant Concentrate is 1–4 quarts in 100 gallons (1 – 2.5 tbsp. per gallon) of water. Repeat at 7–14-day intervals as needed. See specific application recommendations for each crop for additional details. (Keep relative humidity below 70% to avoid risk of phytotoxicity.) (Since it is not possible to test all ornamental species or varieties, test Regalia® Bioprotectant Concentrate on a few plants prior to large-scale usage.))

- Aerial Stem Rot (*Erwinia carotovora*)
- Aerial Web Blight (*Rhizoctonia solani*)
- Alfalfa Wilt (*Xylella spp.*)
- Alternaria Blight (*Alternaria cucumerina*)
- Alternaria Blotch (*Alternaria mali*)
- Alternaria Brown Spot (*Alternaria alternata*)
- Alternaria Fruit Rot (*Alternaria spp.*)
- Alternaria Leaf Blight (*Alternaria spp.*)
- Alternaria Leaf Spot (*Alternaria spp.*)
- Alternaria Leaf Spot, Boll Rot (*Alternaria spp.*)
- Alternaria Spot/Fruit Rot (*Alternaria alternata*)
- Angular Leaf Spot (*Mycosphaerella angulata*) (*Xanthomonas fragariae*)
- Anthracnose (*Collectotrichum spp.*) (*Gnomonia leptostyla*) (*Collectotrichum gloeosporioides*) (*Collectotrichum lagenarium*) (*Collectotrichum truncatum*) (*Elsinoe ampelina*)

- Anthracnose (*Colletotrichum coccodes*)
(*Colletotrichum atramentarium*)
(*Colletotrichum dematium*)
- Anthracnose (*Colletotrichum spp.*) – suppression only
- Anthracnose and Black Stem Rot (*Colletotrichum trifolii*)
- Anthracnose Boll Rot (*Glomeria spp.*)
- Anthracnose Fruit Rot (*Colletotrichum acutatum*)
- Anthracnose Leaf Blight (*Colletotrichum graminicola*)
- Anthracnose of Potato (*Colletotrichum coccodes*)
- Anthracnose, Boll Rot (*Glomeria spp.*)
- Apple Scab (*Venturia inaequalis*) (Suppression only)
- Ascochyta Blight, Boll Rot (*Ascochyta spp.*)
- Asian Soybean Rust (*Phakopsora pachyrhizi*)
- Aspergillus crown rot (*Aspergillus niger*)
- *Aureobasidium zeae*
- Bacteria (*Erwinia spp.*) (*Pseudomonas spp.*) (*Xanthomonas spp.*)
- Bacterial (Leaf) Spot (*Xanthomonas pruni*)
- Bacterial Blast (*Pseudomonas syringae*)
- Bacterial blight (*Pseudomonas cannabina*)
- Bacterial Blight (*Pseudomonas syringae*) (*Pseudomonas viridiflava*) (*Xanthomonas campestris* pv. *pruni*) (*Xanthomonas campestris*)
- Bacterial Blight (*Xanthomonas axonopodis*) (*Xanthomonas citri*)
- Bacterial Blight and Streak (*Xanthomonas spp.*)
- Bacterial Blight/Rot (*Xanthomonas spp.*)
- Bacterial Canker (*Erwinia nigrifluens*) (*Pseudomonas syringae*) (*Pseudomonas spp.*) (*Xanthomonas campestris*) (*Xanthomonas spp.*)
- Bacterial Leaf Blight (*Xanthomonas campestris*)
- Bacterial Leaf Spot (*Pseudomonas spp.*)
- Bacterial leaf streak (*Xanthomonas campestris* pv. *Holcicola*)
- Bacterial leaf stripe (*Pseudomonas spp.*)
- Bacterial Pustule (*Xanthomonas spp.*)
- Bacterial rots (*Pantoea spp.*)
- Bacterial Speck (*Pseudomonas syringae* pv. *glycinea*) (*Pseudomonas syringae*)
- Bacterial Spot (*Xanthomonas pruni*) (*Xanthomonas spp.*) (*Xanthomonas cucurbitae*)
- Bacterial Wilt (*Clavibacter michiganense*)
- Barley yellow dwarf virus
- Bentgrass/Bermudagrass Dead Spot (*Ophiosphaerella agrostis*)
- Bermudagrass Decline (*Gaeumannomyces graminis* var. *graminis*)
- Bitter Rot (*Colletotrichum spp.*)
- Black dot disease (*Epicoccum nigrum*) (*Epicoccum purpurascens*)
- Black mildew (*Schiffnerula cannabis*)
- Black Mold (*Alternaria alternata*)
- Black Root (*Thielaviopsis basicola*)
- Black Root Rot / Black Crown Rot (*Alternaria spp.*)
- Black Rot (*Guignardia bidwellii*)
- Black Rot/Frogeye Leaf Spot (*Botryosphaeria obtusa*)
- Black Scurf (*Rhizoctonia solani*)
- Black shank (*Phytophthora nicotianae*)
- Black Spot (*Guignardia citricarpa*), (*Phyllosticta citricarpa*)
- Black Spot of Rose (*Diplocarpon rosae*)
- Blossom Blight (*Monilinia spp.*)
- Blue Mold (*Peronospora tabacina*)
- Boll Rot (*Alternaria spp.*) (*Ascochyta spp.*) (*Fusarium spp.*) (*Phoma spp.*)
- Bot Rot (*Botryosphaeria dothidea*)
- Botryosphaeria Blight (*Botryosphaeria dothidea*)
- Botrytis (*Botrytis cinerea*)
- Botrytis Blight (*Botrytis cinerea*)
- Botrytis Bud Rot (*Botrytis cinerea*)

- Botrytis Bunch Rot (*Botrytis cinerea*)
- Botrytis Fruit Rot (*Botrytis cinerea*)
- Botrytis Leaf Blight (*Botrytis squamosa*)
- Botrytis Neck Rot (*Botrytis* spp.)
- Brooks Spot (*Mycosphaerella pomii*)
- Brown / Hull Rot (*Monilinia* spp.)
- Brown blight (*Alternaria alternata*) (*Alternaria tenuis*)
- Brown leaf spot and stem canker (*Ascochyta* spp.) (*Ascochyta prasadii*) (*Phoma* spp.) (*Didymella* spp.) (*Phoma exigua*) (*Phoma glomerata*) (*Phoma herbarum*)
- Brown patch (*Rhizoctonia solani*)
- Brown Rot (*Monilinia* spp.)
- Brown Rot Blossom Blight (*Monilinia laxa*)
- Brown Rot Fruit Rot (*Monilinia fruticola*)
- Brown Rot, Leaf Spots & Smuts (*Ceratobasidium* spp.) (*Cercospora* spp.) (*Cochliobolus* spp.) (*Drechslera* spp.)
- Brown Rust (*Puccinia melanocephala*)
- Brown Spot (*Alternaria* spp.) (*Septoria glycines*)
- Brown Stripe/Gray Streak (*Cercosporidium graminis*)
- Bull's Eye Rot (*Neofabraea* spp.)
- Cedar-Apple Rust (*Gymnosporangium juniperi-virginianae*) – suppression only
- Cercospora Blight (*Cercospora asparagi*) (*Cercospora kikuchii*)
- Cercospora Blight and Leaf Spot (*Cercospora* spp.)
- Cercospora Leaf Spot (*Cercospora citrulina*) (*Cercospora* spp.) (*Cercospora beticola*)
- Charcoal rot (*Macrophomina phaseolina*)
- Cherry Leaf Spot (*Blumeriella jaapii*)
- *Cladosporium* spp
- Cladosporium stem canker (*Cladosporium cladosporioides*) (*Cladosporium herbarum*) (*Mycosphaerella tassiana*)
- Clubroot (*Plasmodiophora brassicae*)
- Colletotrichum Crown Rot (*Colletotrichum* spp.) (*Colletotrichum graminicola*)
- Common Scab (*Streptomyces scabies*)
Suppression only
- Copper Spot (*Gloeocercospora sorghi*)
- Corn grey leaf spot (*Cercospora zeae-maydis*) (*Cercospora zeina*)
- Cotton root rot (*Phymatotrichopsis omnivora*) (*Phymatotrichum omnivorum*)
- Cranberry cotton ball (*Monilinia oxycocci*)
- Cranberry Early Rot (*Phyllosticta vaccinia*)
- Crown and Foot Rots (*Pseudocercospora herpotrichoides*, *Rhizoctonia*)
- Crown gall (*Agrobacterium tumefaciens*)
- Curvularia leaf spot (*Curvularia cymbopogonis*) (*Curvularia lunata*) (*Cochliobolus lunatus*)
- Cyindrosporium blight (*Cylindrosporium* spp.) (*Cylindrosorium cannabinum*)
- Damping off (*Aspergillus flavus*) (*Botrytis cinerea*) (*Botryotinia fuckeliana*) (*Fusarium* spp.) (*Fusarium oxysporum*) (*Fusarium solani*) (*Nectria haematococca*) (*Macrophomina phaseolina*) (*Pellicularia filamentosa*) (*Phytophthora* sp.) (*Pythium* spp.) (*Pythium aphanidermatum*) (*Pythium debaryanum*) (*Pythium ultimum*) (*Rhizoctonia* spp.) (*Rhizoctonia solani*) (*Thanatephorus cucumeris*)
- Dichondra Rust (*Puccinia dichondrae*)
- Diplodia Boll Rot (*Diplodia* spp.)
- Diplodia Ear Rot (*Stenocarpella maydis*)
- Diseases from pruning wounds including *Eutypa* (*Eutypa lata*), *Botryosphaeria rhodia*, *Phaeoacremonium aleophilum* and *P. chlamydospora*

- Dollar Spot (*Lanzia spp.*)
(*Moellerodiscus spp.* formerly
Sclerotinia homeocarpa)
- Downy Mildew (*Bremia lactucae*),
(*Peronospora spp.*) (*Peronospora
farinosa*) (*Peronospora destructor*)
(*Peronospora mansherica*)
(*Peronospora parasitica*) (*Peronospora
trifoliorum*) (*Plasmopara viburni*)
(*Plasmopara viticola*)
(*Pseudoperonospora cubensis*)
(*Pseudoperonospora humuli*)
- Downy mildew (*Pseudoperonospora
cannabina*)(*Pseudoperonospora humuli*)
- Early Blight (*Alternaria solani*)
- Early Blight of celery (*Cercospora apii*)
- Early Leaf Spot (*Cercospora
arachidicola*)
- Early Rot in Cranberry (*Phyllosticta
vaccinii*)
- Eastern Filbert Blight (*Anisogramma
anomala*)
- Eutypa (*Eutypa lata*)
- Eye Spot (*Aureobasidium zeae*)
- Fire Blight (*Erwinia amylovora*)
- Flyspeck (*Zygophiala jamaicensis*)
- Foliar Blight
- Frog-eyed Leaf Spot (*Cercospora soja*)
- Fruit Finish
- Fungal Disease Complexes (*Bipolaris
spp.*, *Monographella spp.*,
Phaeosphaeria spp.)
- Fusarium foot rot and root rot
(*Fusarium solani*)
- Fusarium Head Blight (*Fusarium
graminearum*)
- *Fusarium oxysporum*
- Fusarium Patch (*Fusarium nivale*)
- *Fusarium proliferatum*
- Fusarium root and stem rot
- *Fusarium solani*
- *Fusarium spp.*
- Fusarium stem canker (*Fusarium
sulphureum*) (*Gibberella
cyanogena*)(*Gibberella saubinetii*)
- Fusarium wilt (*Fusarium oxysporum
f.sp. cannabis*) (*Fusarium oxysporum
f.sp. vasinfectum*) (*Fusarium oxysporum
f. sp. lactucae*)
- *Fusarium wilt (Fusarium oxysporum)*
- *Glomerella tucumanensis*, also known
as *Colletotrichum falcatum*
(Suppression Only)
- Gray leaf spot (*Cercospora sorghi*)
(*Pyricularia grisea*) (*Cercospora zeae-
maydis*)
- Gray Mold (*Botrytis cinerea*) (*Botrytis
spp.*)
- Greasy Spot (*Mycosphaerella citri*)
- Green Fruit Rot (*Botrytis cinerea*)
- Gummy Stem Blight (*Didymella
bryoniae*)
- Hard Lock
- Hard Lock, Boll Rot (*Fusarium spp.*)
- Hemp canker (*Sclerotinia sclerotiorum*)
- Hemp Leaf Spot (*Bipolaris sp.*)
- Hull Rot (*Rhizopus stolonifer* and
Monilinia spp.)
- Late Blight (*Phytophthora infestans*)
(*Septoria apiicola*)
- Late Leaf Spot (*Cercosporidium
personatum*)
- Late Rot in Cranberry
- Leaf Blight (*Pseudocercospora vitis*)
(*Septosphaeria turcica*)
- Leaf Rust (*Pucciniastrum vaccinii*)
(*Tranzschelia discolor*)
- Leaf Spot (*Alternaria spp.*) (*Cercospora
spp.*) (*Cercospora beticola*)
(*Corynespora cassicola*)
(*Entomosporium spp.*) (*Mycosphaerella
fragariae*) (*Myrothecium spp.*) (*Septoria
spp.*)
- Leaf Spots (*Dreschlera*, *Cochliobolus*,
Cercospora)
- Leafspots and Blotches (*Pseudopeziza
medicaginis*, *Stemphyllium spp.*,
Cercospora spp., *Stagonospora spp.*)

- Leptosphaeria blight (*Leptosphaeria cannabina*)(*Leptosphaeria woroninii*)(*Leptosphaeria acuta*)
- Melanose (*Diaporthe citri*)
- Melting Out Leaf Spot (*Bipolaris* spp.), (*Drechslera* spp.)
- Miscanthus blight
- Miscanthus streak virus
- Mosaic viruses
- Mummy Berry (*Monilinia vaccinii-corymbosi*),
- Necrotic Ring Spot (*Leptosphaeria korrae*)
- Northern Leaf Blight (*Exserohilum turcicum*)
- Northern Leaf Spot (*Cochliobolus carbonum*)
- Olive Knot (*Pseudomonas savastanoi*)
- Olive leaf spot (*Cercospora cannabis*) (*Pseudocercospora cannabina*)
- Onion Downy Mildew (*Peronospora destructor*)
- Onion Purple Blotch (*Alternaria porri*)
- Ophiobolus stem canker (*Ophiobolus cannabinus*)(*Ophiobolus anguillides*)
- Orange Rust (*Puccinia kuehnii*)
- Panicum mosaic virus
- Peg
- *Pencillium*
- Phoma Blight
- Phoma Blight, Boll Rot (*Phoma* spp.)
- Phoma stem canker (*Phoma herbarum*) (*Phoma exigua*)
- Phomopsis
- Phomopsis Fruit Rot (*Phomopsis viticola*)
- Phomopsis Leaf Blight (*Phomopsis obscurans*)
- Phomopsis Leaf Spot, Twig Blight, and Fruit Rot (*Phomopsis* spp.)
- Phomopsis stem canker (*Phomopsis cannabina*)(*Phomopsis achilleae*)(*Diaporthe arctii* var. *achilleae*)
- Phymatotrichum root rot /Phymatotrichopsis root rot (*Phymatotrichopsis omnivora*)(*Phymatotrichum omnivorum*)
- Phytophthora Blight (*Phytophthora capsici*)
- Phytophthora Root Rot and Crown Rot (*Phytophthora* spp.)
- Phytophthora (*Phytophthora* spp.)
- Pin Rot Complex (*Alternaria/Xanthomonas*)
- Pink Patch (*Limonomyces roseipellis*)
- Pink Rot (*Phytophthora erythroseptica*)(*Sclerotinia sclerotiorum*)
- Pink rot (*Trichothecium roseum*)(*Cephalothecium roseum*)
- Pithomyces blight
- Pod and Stem Blight (*Diaporthe phaseolorum* var. *sojae*) (*Phomopsis longicola*) (*Diaporthe* spp.)
- Pod and Stem Blight (*Diaporthe* spp.)
- Postbloom Fruit Drop (*Colletotrichum acutatum*)
- Powdery Mildew (*Erysiphe* spp.) (*Erysiphe betae*), (*Erysiphe cichoracearum*) (*Erysiphe cruciferarum*) (*Erysiphe graminis*) (*Erysiphe polygoni*) (*Leveillula taurica*) (*Microsphaera alni*) (*Oidium* spp.), (*Oidopsis taurica*) (*Podosphaera* spp.) (*Podosphaera leucotricha*) (*Sphaerotheca* spp.) (*Sphaerotheca fuliginea*) (*Sphaerotheca macularis*) (*Sphaerotheca pannosa*) (*Uncinula necator*)
- Powdery Mildew (*Golovinomyces*) (*Erysiphe cichoracearum*)(*Leveillula taurica*)(*Oidiopsis taurica*)(*Sphaerotheca macularis*)(*Sphaerotheca humuli*)(*Oidium* spp.)
- Powdery Mildew / Rusty Spot (*Podosphaera* spp.), (*Sphaerotheca pannosa*)
- *Puccinia* spp.
- Purple spot (*Stemphylium vesicarium*)

- Pythium (aerial blight phase) (*Pythium* spp.)
- Pythium (*Pythium* spp.)
- *Pythium acanthicum*
- *Pythium aphanidermatum*
- Pythium Blight, Pythium Root Rot (*Pythium aphanidermatum*), (*Pythium* spp.)
- *Pythium dissoticum*
- *Pythium myriotylum*
- Pythium root and damping off
- Ramularia (*Ramularia* spp.)
- Ramularia Leaf Spot (*Ramularia cynarae*)
- Red boot (*Melanospora cannabis*)
- Red Rot (*Glomerella tucumanensis*, also known as *Colletotrichum falcatum*)
- Red Thread (*Laetisaria fuciformis*)
- Rhizoctonia Foliar Blight, Peg, and Root Rot (*Rhizoctonia solani*)
- Rhizoctonia Large Patch (*Rhizoctonia solani*)
- Rhizoctonia soreshin and root rot (*Rhizoctonia solani*)
- *Rhizoctonia* spp.
- Rice Blast (*Pyricularia grisea*)
- Ripe Rot (*Colletotrichum gloeosporioides*)
- Root and collar rots (*Phytophthora*, *Pythium*, *Fusarium*, *Rhizoctonia*)
- Rot (*Rhizoctonia* spp.), (*Pythium* spp.), (*Fusarium* spp.), (*Cylindrocarpon* spp.)
- Rust (*Aecidium cannabis*)(*Uredo kriegiana*)(*Uromyces inconspicuus*)
- Rust (*Phylopsora* spp.) (*Puccinia* spp.) (*Puccinia asparagi*) (*Puccinia menthae*) (*Puccinia porri*) (*Tranzschelia discolor*) (*Uromyces appendiculatus*) (*Uromyces betae*)
- Rusty Spot (*Podosphaera leucotricha*)
- Scab (*Cladosporium carpophilum*) (*Sphaceloma perseae*) (*Elsinoe australis*) (*Elsinoe fawcetti*) (*Elsinoe mangiferae*) (*Venturia* spp.)
- Sclerotinia (*Sclerotinia Sclerotiorum*)
- Sclerotinia Head and Leaf Drop (*Sclerotinia minor*) (*Sclerotinia sclerotiorum*)
- Sclerotinia stem and crown rot (*Sclerotinia sclerotiorum*)
- Sclerotium root and stem rot (*Sclerotium rolfsii*)(*Athella rolfsii*)
- Seedling and Damping Off Disease Complex, including Root and Crown Rots (*Pythium*, *Phytophthora*, *Rhizoctonia*, and *Stagonospora* spp.)
- Septoria Brown Spot (*Septoria glycines*)
- Septoria Leaf/Speckled Leaf Spot/Blotch (*Septoria* spp.)
- Sheath Spot and Blight (*Rhizoctonia oryzae*), (*Thanatephorus cucumeris*)
- Shot Hole (*Wilsonomyces carpophilus*)
- Sigatoka (*Mycosphaerella fijiensis*)
- Smut (*Tilletia* spp.) (*Tilletia barclayana*)
- Smuts and Bunts (*Tilletia* spp.)
- Snowmold, Gray (*Typhula* spp.)
- Snowmold, Pink (*Microdochium nivale*)
- Sooty Blotch (*Geastrum polystigmati*), (*Leptodontium elatius*), (*Peltaster fruticola*)
- Sorghum downy mildew (*Peronosclerospora sorghi*)
- Sour Rot (*Alternaria tenuis*) (*Aspergillus* spp.) (*Botrytis cinerea*) (*Cladosporium herbarum*) (*Penicillium* spp.) (*Rhizopus arrhizus*)
- Southern Blight (*Sclerotium rolfsii*)
- Southern blight (*Sclerotium rolfsii*) (*Athella rolfsii*)
- Southern leaf blight (*Bipolaris* spp.) (*Cochliobolus heterostrophus*)
- Spring Black Stem (*Phoma medicaginis*)
- Spring Dead Spot (*Leptosphaeria korrae*), (*Leptosphaeria narmari*), (*Ophiosphaerella herpotricha*), (*Gaeumannomyces graminis*)
- Spur Blight (*Didymella* spp.), (*Phoma* spp.)
- Stem Rot (*Sclerotium oryzae*)

- Stemphylium leaf and stem spot (*Stemphylium botryosum*) (*Pleospora tarda*) (*Stemphylium cannabinum*)
- Stemphylium Leaf Blight (*Stemphylium vesicarium*)
- Stemphylium Leaf Spot (*Stemphylium* spp.)
- *Striatura ulcerosa* (*Pseudomonas amygdali* pv. *mori*)
- Stripe Smut (*Ustilago striiformis*), (*Urocystis agropyri*)
- Summer Bentgrass Decline
- Summer Patch, Poa Patch (*Magnaporthe poae*)
- Switchgrass Mosaic Virus
- Take-All Patch (*Gaeumannomyces graminis*)
- Tan Spot (*Pyrenophora tritici-repentis*)
- Tar spot (*Phyllachora cannabidis*)
- Target Spot (*Corynespora cassiicola*) (*Rhizoctonia solani*)
- Tropical rot (*Lasiodiplodia theobromae*)(*Botryodiplodia theobromae*)
- Twig blight (*Dendrophoma marconii*)(*Botryosphaeria marconii*)
- Verticillium wilt (*Verticillium* spp.) (*Verticillium albo-atrum*)
- *Verticillium* spp.
- Verticillium wilt (*Verticillium albo-atrum*)(*Verticillium dahliae*)
- Walnut Blight (*Xanthomonas campestris*)
- White leaf spot (*Phomopsis ganjae*)
- White Mold (*Sclerotinia sclerotiorum*) (*Sclerotium rolfsii*) (*Sclerotinia minor*) (*Sclerotinia trifoliorum*)
- white mold stem rot
- White Mold/ Sclerotinia Stem Rot (*Sclerotinia sclerotiorum*)
- White Rot (*Botryosphaeria dothidea*)
- White Rust (*Albugo occidentalis*)
- *Xanthomonas campestris*
- Xanthomonas leaf spot (*Xanthomonas campestris* pv. *cannabidis*)
- Xanthomonas Leaf Spot (*Xanthomonas campestris*)
- *Xanthomonas* spp.
- Yellow leaf spot (*Septoria cannabidis*)(*Septoria cannabina*)
- Yellow Patch (*Rhizoctonia cerealis*)
- Yellow Tuft/Downy Mildew (*Sclerophthora macrospora*)
- Zoysia Patch (*Rhizoctonia solani*)

Pre-harvest Interval (PHI) = 0 days

ROOT, TUBER AND CORM CROPS

Potato, Beet, Carrot, Cassava, Ginger, Ginseng, Horseradish, Radish, Sugar beet, Sweet Potato, Turnip (including those for seed production) and other root and tuber crops

For suppression of Early Blight, Black Root Rot/Black Crown Rot, and Late Blight, begin application of this product in 25–100 gallons of water per acre (0.5 – 2.5 gallons of water per 1000 square feet) soon after emergence when conditions are conducive to disease development. Repeat on a 5–7-day interval or as needed. For improved performance, use this product in a tank mix with other registered fungicides.

(Apply at 1 pint per acre for Plant Health Applications.)

1-16 fl oz. per 100 pounds of seed pieces for SEED PIECE DIP applications

- For seed piece dip applications for improved plant growth and suppression of soil-borne diseases, apply this product to seed pieces prior to planting.

LEAVES OF ROOT AND TUBER VEGETABLES

Beet, Chervil, and other leaves of roots and tubers

BULB VEGETABLES

Onion (Bulb and Green), Garlic, Leek, Shallot and other bulb vegetables

LEAFY VEGETABLE CROPS (except *Brassica* vegetables)

Arugula, Celery, Cilantro, Corn Salad, Cress, Dandelion, Dock, Edible Chrysanthemum, Endive, Head Lettuce, Leaf Lettuce, Parsley, Purslane, Radicchio, Rhubarb, Spinach, Swiss Chard, Turnip, Watercress and other leafy vegetable crops

- West of the Rocky Mountains – For aerial applications, apply this product at 0.5–1.5 quarts per acre in a minimum of 10 gallons of water per acre
- East of the Rocky Mountains – For aerial applications, apply this product at 0.5–1 quarts per acre in a minimum of 5 gallons of water per acre.
- For California – For aerial application apply REGALIA® Bioprotectant Concentrate at 1–3 pints per acre in 10–20 gallons of water per acre.

Restrictions:

Do not apply REGALIA® Bioprotectant Concentrate when extended/unseasonably cold or cold and cloudy conditions are expected. Regalia should **not** be applied when night-time temperatures will fall below 45°F and relative humidity is predicted to be above 80%. Applications during daylight hours are preferred over night-time applications.

(For concentrated applications, apply this product at 0.5 to 1.5 quarts per acre in a minimum of 10 gallons of water per acre.)

BRASSICAS (COLE) LEAFY VEGETABLES

Broccoli, Broccoli Rabe, Brussels Sprouts, Cabbage, Chinese Broccoli, Chinese Cabbage (Bok Choy), Chinese Cabbage (Napa), Chinese Mustard Cabbage (Gai Choy), Cauliflower, Cavalo, Collards, Kale, Kohlrabi, Mizuna, Mustard Greens, Mustard Spinach, Rape Greens, and other cole crops

LEGUME VEGETABLES, succulent or dried (not including soybeans and peanuts)(not including peanuts)

Chick Peas, Dry Beans, Green Beans, Lentils, Lima Beans, Peas, Shell Beans, Snap Beans, Split Peas (including those grown for seed or oil production), Soybeans (and other legume vegetables)

To maximize activity against *Sclerotinia* white mold in soybeans, the best timing would be V3 to R3 and should be tank mixed with a fungicide partner also labeled for white mold activity.

Repeat applications on 7–10 day intervals depending upon crop growth and disease pressure.

FOLIAGE OF LEGUME VEGETABLES

Garden peas and other foliage of legume vegetables

FRUITING VEGETABLES

Tomato, Pepper, Eggplant, Ground Cherry, Okra, Tomatillo and other fruiting vegetables

- Phytophthora Blight – Apply this product in combination with labeled rates of a copper fungicide (or with another fungicide labeled for Phytophthora Blight control).

CUCURBITS

Includes all types and hybrids of: Acorn squash, Balsam apple, Balsam pear, Bitter melon, Butternut squash, Calabaza, Cantaloupe, Casaba, Chayote, Chinese cucumber, Chinese okra, Chinese waxgourd, Citron melon, Crenshaw melon, Crookneck squash, Cucumber, Cucuzza, Gherkin, Golden pershaw melon, Honeydew melon, Honey balls, Hubbard squash, Hyotan, Mango melon, Persian melon, Pineapple melon, Pumpkin, Santa Claus melon, Scallop squash, Snake melon, Spaghetti squash, Straightneck squash, Vegetable marrow, Watermelon, Zucchini and other cucurbits

- When cucurbits are under high disease conditions, use the shorter spray interval.
- Downy Mildew – Tank mix this product with another fungicide labeled for Downy Mildew control and re-apply at a 7-day interval or according to the label directions of the tank mix (ingredient) (partner)
- Phytophthora Blight – Apply this product in combination with labeled rates of a copper fungicide or with another fungicide labeled for Phytophthora Blight control.

CITRUS CROPS

Orange, Grapefruit, Kumquat, Lemon, Tangelo, Tangerine, Pummelo and other citrus crops

POME FRUITS

Apple, Crabapple, Loquat, Oriental Pear, Pear, Quince, Mayhaw and other pome fruits

- Use caution when selecting spray adjuvants. Select only those adjuvants which through prior experience do not affect fruit finish when combined with this product.
- Fire Blight – For suppression, apply 1-2 quarts of this product in 50–100 gallons of water per acre (1 – 2.5 gallons of water per 1000 square feet) at green tip through bloom. Do not exceed 0.5% v/v. Following bloom, this product can be applied at 2–4 quarts per acre (3 – 6 tbsp. per 1000 sq. ft.). For maximum control, use this product prior to infection events. During periods of rapid development and frequent infection periods, use spray intervals of 3–7 days.
- Apply in sufficient water to provide full coverage. For improved performance, use this product in a rotational program with copper or antibiotics registered for Fire Blight control such as but not limited to oxytetracycline or streptomycin.
- Proper orchard cultural practices are essential to eliminate Fire Blight-infected tissue from the orchard to assure good performance of any crop protection product. Remove and destroy dead and diseased wood from the orchard prior to and during the growing season.
- Scab – For suppression, apply 1-2 quarts of this product in 50–100 gallons of water per acre (1 – 2.5 gallons of water per 1000 square feet) at green tip and through bloom when environmental conditions become favorable for primary Scab development and repeat on a 7–10 day interval or as needed. Do not exceed 0.5% v/v. Use this product in a tank mix or rotational program with other fungicides labeled for Scab control. Following bloom, this product can be applied at 2–4 quarts per acre (3 – 6 tbsp. per 1000 sq. ft.).

Some sensitive tree fruit varieties have exhibited petal staining and/or necrosis after application of higher use rates. To minimize petal staining and/or necrosis:

- Use adjuvants that improve coverage, not penetration; follow the manufacturer's mixing instructions.
- Use adjuvants that through prior experience do not affect petal integrity when combined with this product.
- Apply 1-2 quarts of this product in 50–100 gallons (1 – 2.5 tbsp. per gallon) of water per acre (1.5 tbsp. in 1 gallon to 2.5 gallons of water per 1000 sq. ft.) in Pome Fruit, from 10% bloom to full bloom. Do not exceed 0.5% v/v.

STONE FRUITS

Apricot[†], Cherry (sweet and tart), Nectarine, Peach, Plum, Plumcot, Prune and other stone fruits

- Bacterial Blight – Apply this product in 50–100 gallons of water per acre (1 – 2.5 gallons of water per 1000 square feet) postharvest before fall rains.
- Brown Rot Blossom Blight – Begin application of this product in 50–100 gallons of water per acre (1 – 2.5 gallons of water per 1000 square feet) at early bloom, and repeat through petal fall on a 7-day interval or as needed.
- Powdery Mildew – Begin application of this product in 50–100 gallons of water per acre (1 – 2.5 gallons of water per 1000 square feet) at popcorn stage, and repeat on a 7-day interval or as needed. For improved

performance, use this product in a tank mix or rotational program with other registered fungicides for powdery mildew control.

- **Scab** – Begin application of this product in 50–100 gallons of water per acre (1 – 2.5 gallons of water per 1000 square feet) at petal fall, and repeat on a 7–10-day interval or as needed. For improved performance, tank mix this product with another fungicide labeled for Scab control.
- **For all other diseases** – Begin application prior to disease development when environmental conditions and plant stage are conducive to rapid disease development, and repeat on a 7–10 day interval or as needed. Use in a tank mix or rotational program when disease conditions are severe.

†Some sensitive apricot varieties have exhibited fruit spotting as a result of application. Spray a test strip to confirm your variety is not susceptible to spotting before spraying.

Some sensitive tree fruit varieties have exhibited petal staining and/or necrosis after application of higher use rates. To minimize petal staining and/or necrosis:

- Use adjuvants that improve coverage, not penetration; follow the manufacturer's mixing instructions.
- Use adjuvants that through prior experience do not affect petal integrity when combined with this product.
- Apply 1 quart of this product in 50–100 gallons of water per acre (1.5 tbs. in 1 gallon to 2.5 gallons of water per 1000 sq. ft.) in:
 - Cherries, from white bud (first white, popcorn) to full bloom,
 - Stone fruit, from 10% bloom to full bloom.

BERRIES (AND SMALL FRUIT)

Blueberry[†], Blackberry (all varieties), Cranberry, Currant, Elderberry, Gooseberry, Huckleberry, Juneberry, Ligonberry, Loganberry, Raspberry (red and black), Salal and cultivars and/or hybrids of these and other berries

Cranberry

Grape (wine, table, raisin, and juice varieties), Strawberry and other smallfruit

- **Mummy Berry** – Initiate application at bud break stage of development. Apply this product preventatively and repeat on a 7–10 day interval or as needed. For best performance, tank mix this product with other registered fungicides for Mummy Berry control.
- **Botrytis Blight** – Apply this product preventatively or when the first disease symptoms are visible and reapply every 7–14 days.
- **Bacterial Canker** – Apply this product prior to Fall rains and repeat applications during dormancy before spring growth. This product can be tank mixed with another registered fungicide for improved control of bacterial canker. REGALIA® Bioprotectant Concentrate is compatible with fungicide/bactericides containing the active ingredients peroxyacetic acid and hydrogen peroxide such as Jet-Ag® 5.
- **Anthracoze Fruit Rot and Alternaria Fruit Rot on blueberries** – Initiate application at green tip and continue applications on a 7–10 day interval.
- **Anthracoze on strawberry** – For suppression, apply this product preventatively in 50–100 gallons of water per acre (1 – 2.5 gallons of water per 1000 square feet) and repeat on a 7-day interval or as needed. For best performance, tank mix this product with other registered fungicides for Anthracnose control.

† Some sensitive blueberry varieties have exhibited fruit spotting as a result of application. Spray a test strip to confirm your variety is not susceptible to spotting before spraying. Do not exceed 1% v/v.

(Grape - Do not use carrier volumes and/or adjuvants that create spray runoff or drip-accumulation at the leaf margins or at the base of fruit or on the harvested commodity. Complex tank-mixes that include REGALIA® Bioprotectant Concentrate may increase the risk of visible spray residues on grape clusters. Test the mix on a small portion of the crop to be treated to ensure that a phytotoxic response will not occur as a result of the application. Do not apply REGALIA® Bioprotectant Concentrate within 14 days of applying a sulfur fungicide during high temperatures.)

TREE NUT CROPS

Almond, Beech nut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert, Hickory nut, Macadamia nut, Pecan, Pistachio, Walnut (Black and English)

Some tree nut varieties have exhibited petal staining and/or necrosis after application of higher use rates.

To minimize petal staining and/or necrosis:

- Use adjuvants that improve coverage, not penetration; follow the manufacturer's mixing instructions.
- Use adjuvants that through prior experience do not affect petal integrity when combined with this product.
- Apply 1 quart in 50 gallons or 2 quarts in 100 gallons of water per acre (1.5 tbsp. in 1 gallon or 3 tbsp. in 2.5 gallons of water per 1000 sq. ft.) from pink bud through bloom.

(For Pecans: For *Phymatotrichopsis* root rot (*Phymatotrichopsis omnivore*) apply through Chemigation by drip irrigation or micro-sprinklers at 2-4 quarts/acre. Begin applications when trees begin to leaf out in the spring, repeat applications monthly.)

CEREAL GRAINS

Barley, Buckwheat, Grain Amaranth, Milo, Oat, Millets, Rice, Rye, Sorghum, Triticale, Wheat and other cereal grains

It is important to apply this product at the Feekes 4 to Feekes 10.5 stage to maximize yield. Apply this product preventatively or when the first disease symptoms appear. Repeat applications in 7–14-day intervals depending upon crop growth and disease pressure

(For (Barley) (Buckwheat) (Rye) (Amaranth) (Oat) (Millets) (Rice) (Wheat) (Sorghum)): (Apply at 1 pint per acre for foliar (ground) (aerial) Plant Health Applications.)

Corn (Sweet Corn, Field Corn, Popcorn, Silage Corn, Seed Corn)

Apply this product preventatively starting at V1-V10 or VT or when the first disease symptoms appear. (Repeat applications on 7–10 day intervals depending upon crop growth and disease pressure.)

(Apply this product in 1 pint per acre by (foliar (ground)) (in-furrow) (for) (improved plant health potential.) (increased quality and/or yields at harvest.)

FORAGE, FODDER AND STRAW OF CEREAL GRAINS

Corn, Wheat, and any other cereal grain crop

(For (Corn) (Wheat):) Apply at 1 pint per acre for (foliar (ground) (aerial)) (in-furrow) Plant Health Applications.)

GRASS FORAGE, FODDER, AND HAY

Bermuda grass, Bluegrass, Bromegrass, Fescue, Pasture and range grasses grown for hay or silage, Sudangrass, Timothy and other grass forage, fodder, and hay

(Apply at 0.5 to 1 quart per acre for foliar (ground) (aerial) Plant Health Applications.)

NON-GRASS ANIMAL FEED

Alfalfa, Clover, Kudzu, Lespedeza, Lupin, Sainfoin, Trefoil, Vetch and other non-grass animal feed

(Apply at 1 pint per acre for foliar (ground) Plant Health Applications.)

HERBS (DRIED OR FRESH) (FIELD AND GREENHOUSE)

SPICES (FIELD AND GREENHOUSE)

OIL SEED CROPS (not including peanut or soybean)

Canola, Castor, Flax, Jojoba, Rapeseed, Safflower, Sesame, and Sunflower and other oil seeds

For white mold disease control: apply at 10% bloom with a second application 7-10 days later.

(Apply at 1 pint per acre for Plant Health Applications in Canola.)

Cotton

For ground applications for foliar and boll rot disease control, apply this product preventatively in 15–40 gallons of water per acre (0.5 – 1 gallon of water per 1000 square feet) prior to disease development using sufficient volume for thorough coverage

(Apply at 0.5 to 1 quart per acre for foliar (ground) (aerial) Plant Health Applications.)

STALK, STEM, AND LEAF PETIOLE VEGETABLES

Asparagus, Celery, Agave and other stalk, stem and leaf petiole vegetables

TROPICAL AND SUBTROPICAL FRUITS, EDIBLE

Olive and other tropical and subtropical fruits with edible peel

TROPICAL AND SUBTROPICAL FRUITS, INEDIBLE

Avocado, Banana, Kiwi, Mango, Papaya, Plantain, Pineapple, Pomegranate and other tropical and subtropical fruits with inedible peel

Sigatoka – Initiate applications when leaves first appear and repeat on a 7–10 day schedule. Apply in sufficient water by ground or air to obtain thorough coverage of foliage. For improved disease control, this product may be tank-mixed with oil or other fungicides registered for Sigatoka control at label rates

(Other crops [outside crop groups])

ARTICHOKE

HOPS

- Minimum spray volumes for hop growth stages are as follows:
- Emergence to Training: Apply 1–2 quarts this product per acre (1.5 – 3 tbsp. per 1000 sq. ft.) using a minimum spray volume of 20 gallons per acre (0.5 gallon per 1000 sq. ft.). Coverage will vary with the size of the vines and the type of spray equipment. Apply adequate spray volume to achieve complete spray coverage.

- Training to Wire-Touch: Apply 1–2 quarts this product per acre (1.5 – 3 tbsp. per 1000 sq. ft.) using a minimum spray volume of 50 gallons per acre (1 gallon of water per 1000 sq. ft.). Coverage will vary with the size of the vines and the type of spray equipment. Apply adequate spray volume to achieve complete spray coverage.
- Wire-Touch through Harvest: Apply 2–4 quarts (3 – 6 tbsp.) of this product using a minimum of 100 gallons of water per acre (2.5 gallons of water per 1000 sq. ft.). Higher water volumes may be necessary to achieve thorough coverage after side arms develop. Do not apply more than 4 quarts of product per acre (6 tbsp. per 1000 sq. ft.) per application. Apply adequate spray volume to achieve complete spray coverage. Use the higher rates when moderate to high disease pressure is present or expected.
- For control of downy mildew, tank mix this product with another fungicide labeled for Downy Mildew control and re-apply at a 7-day interval or according to the label directions of the tank mix (ingredient) (partner).

PEANUT

(Apply at 1 pint per acre for plant health.)

HEMP

SUGARCANE

(Apply at 1 pint per acre for Plant Health Applications.)

CROTALARIA, SESSBANIA, KENAF

FLOWERING PLANTS

BEDDING PLANTS

GRASS GROWN FOR SEED

0.5–2.5 fluid ounces per 100 lbs. seed for SEED TREATMENT applications

ORNAMENTALS

SWITCHGRASS, MISCANTHUS

TOBACCO

For Target Spot: Apply this product at a rate of 2 quarts per acre in a minimum of 50 gallons of water per acre.

(Apply at 1 pint per acre for Plant Health Applications.)

INTEGRATED PEST MANAGEMENT (IPM)

Many conventional fungicides have been tested in an IPM regime with REGALIA® Bioprotectant Concentrate with very satisfactory results. One of the major objectives of IPM has been to reduce the probability of disease resistance development to a particular active ingredient.

The alternate use of (1–2 sprays) followed by a conventional, registered fungicide (1–2 sprays) has been successfully used in many crops. In addition, the use of tank mixes with a conventional fungicide has also been successful.

Follow label instructions of the particular registered product: Do not exceed amounts or treatment intervals on the label.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place. Avoid freezing.

Pesticide Disposal: To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

Container Handling (5 gallons or less): Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill or by incineration. Do not burn unless allowed by state and local ordinances.

Container Handling (over 5 gallons): Non-refillable container. Do not reuse or refill this container Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill or by incineration. Do not burn unless allowed by state and local ordinances.

Container Handling (refillable containers – 265 gallon tote):

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. When empty, return to point of sale or offer for recycling if available or reconditioning if appropriate.



Pro Farm Group is a member of the Ag Container Recycling Council.

Visit <http://www.acrecycle.org/contact.html> for information on how to arrange pick-up of this empty pesticide container.

WARRANTY

To the extent consistent with applicable law, the seller makes no warranty, expressed or implied, of merchantability, fitness or otherwise concerning use of this product. To the extent permitted by applicable law, the user assumes all risks of use, storage or handling that are not in strict accordance with the accompanying directions.

Label date:

Made in the U.S.A.

US Patents No. 4,863,734 and No. 5,989,429

REGALIA® is a trademark of Pro Farm Group, Inc.

Name and logo of Pro Farm Group are registered trademarks of Pro Farm Group, Inc.

© Pro Farm Group, Inc.

REGALIA® Bioprotectant Concentrate; EPA Reg. No. 84059-3

MASTER LABEL – MBI-LBL-4 r23 v4 January 28, 2025

Page 54 of 66

SUBLABEL B OPTIONAL CLAIMS

1. Boost plants' defenses [for stronger healthier plants]
2. Strengthens plants' immunity
3. Improves plant health
4. Controls/Prevents powdery mildew, leaf spot and rust
5. Controls/Prevents black spot on rose
6. Labeled Fungal and bacterial disease control
7. Use on labeled fruits, vegetables and ornamentals
8. Can be applied as a pre-plant dip [for improved plant health]
9. Can be applied as a soil drench application
10. For use on ornamental plants and edible crops/fruits/vegetables.
11. For use on vegetables, roses, fruits, berries, nuts, flowers, bedding plants, (ornamental) trees and shrubs
12. REGALIA® Bioprotectant Concentrate may be used on vegetable(s) [crops], roses, fruits, berries, nuts, flowers, foliage, (ornamental) trees and shrubs.
13. REGALIA® Bioprotectant Concentrate is a broad spectrum fungicide used for the control or suppression of a broad range of labeled foliar [fungal and bacterial] diseases.
14. (Active ingredient is) a plant extract to boost the plants' defense mechanisms to protect against labeled fungal and bacterial diseases, and to improve plant health.
15. Active ingredient (is) made from a plant extract (botanical extract)(plant-based)
16. REGALIA® Bioprotectant Concentrate is an extract from the plant *Reynoutria sachalinensis*
17. REGALIA® Bioprotectant Concentrate can be applied up to and on the day of harvest [on all fruits and vegetables].
18. Made in the U.S.A.
19. This container is made with X% recycled material
20. Guaranteed results(*)
21. Label date:
22. US Patents No. 4,863,734 and No. 5,989,429
23. REGALIA® is a registered trademark of Pro Farm Group, Inc.
24. Name and logo of Pro Farm Group are registered trademarks of Pro Farm Group, Inc.
25. © insert company copyright information
26. World rights reserved
27. Distributed by: insert company name and address
28. [For] questions/comments



- 29.
30. GENERAL INFORMATION: REGALIA® Bioprotectant Concentrate is an extract from the plant *Reynoutria sachalinensis*. REGALIA® Bioprotectant Concentrate applied to actively growing plants will improve plant health, and will help make the treated portions resistant to labeled plant diseases. (Plant health benefits often result in greater yields at harvest, especially when crops/plants are stressed by pathogens or environmental conditions).
31. MODE OF ACTION: REGALIA® Bioprotectant Concentrate contains an extract obtained from *Reynoutria sachalinensis* plant material composed of active chemical compounds. Application of the extract activates the immune response, increasing the plant's defense system through a five-fold increase in phenolics and antioxidants, and cell wall strengthening. These immune responses, sometimes referred to as induced systemic resistance or systemic acquired resistance, protects treated plants from numerous diseases.
When applied at rates and timing for disease control, the translaminar activity of REGALIA® Bioprotectant Concentrate provides induced resistance against important diseases and acts within one to two days of application.
REGALIA® Bioprotectant Concentrate should be used as a preventative treatment prior to the development of disease

symptoms. Repeat foliar applications per label instructions, following a 7–14-day interval to maintain plant immune response induction and to protect new plant growth.

(In addition to foliar applications, REGALIA® Bioprotectant Concentrate can be used in multiple application methods as a plant dip, soil drench, pre-plant incorporated soil application, in-furrow spray, or applied through drip irrigation to control or suppress certain soil-borne diseases and to promote healthy root growth and improve above ground plant growth.)

(When applied at rates and timing for plant health effects, the induced plant defense responses minimize the impacts of stress and disease, resulting in optimized yields at harvest. Applying Regalia® Bioprotectant Concentrate has been shown to increase leaf chlorophyll content and increase soluble protein content in some crops. These effects often lead to improved crop quality and/or yields.)

32. REGALIA® Bioprotectant Concentrate is (an extract from the plant *Reynoutria sachalinensis*) for use on labeled (ornamental plants), (turf), (agricultural crops), (edible crops), (non-edible crops), (food crops), (non-food crops), (feed crops), or (non-feed crops).
33. Optional Language: (*) and (*= Not for use in California)
34. Repackaging or relabeling of this product without express written permission from Pro Farm Group is prohibited.
35. Biofungicide
36. [UPC code]
37. [RF code]
38. Bio with Bite™
39. For (labeled) disease control
40. For (labeled) fungal control
41. Can be used in sustainable production
42. For use in sustainable production
43. For maximum harvest
44. US MRL exempt
45. Minimal PPE
46. Protection from Multiple Labeled Diseases
47. Leaves no detectable residues
48. Product(s) thoroughly tested
49. Proven results, since 2007
50. Trial Tested
51. (number)+ trials
52. Read full label before use
53. Prevents (labeled disease) and (labeled fungal) build up
54. Protection from labeled bacterial and fungal disease
55. REGALIA® Bioprotectant Concentrate can be applied by (any labeled use pattern) to protect against (labeled diseases) and (labeled fungal pathogens).
56. 2-10 tablespoons per 1,000 sq. ft. [must be consistent with rates to be listed on label]
57. For turf
58. For recreational turf and landscapes [when crop is listed]
59. (Specify pest:) (Tank-mix) (or rotate) with a (fungicide) for improved control.
60. Refer to the table in the SOIL TREATMENT USE DIRECTIONS (In-Furrow Applications) section to determine the proper rate per 1000 foot of row.
61. REGALIA® Bioprotectant Concentrate should be used as part of an Integrated Pest Management System.
62. REGALIA® Bioprotectant Concentrate can be applied following a soil fumigant.
63. Use the high(er) labeled rate when high(er) (disease)(fungal pathogen) pressure is expected.
64. Optional Language: (*) and (*= Not labeled for this use in California)
65. [optional listing of all pests, alternatively list specific pests for each crop]
66. Bioprotectant Concentrate
67. Outdoor
68. See back of booklet [or other specific location] for precautionary statements and first aid
69. See attached booklet for complete directions for use and precautionary statements

70. Indoor/Outdoor
71. See attached booklet for full label
72. *rate may require more than one package
73. (see (side) (back) (inside) (other) panel for additional (first aid) (and) (precautionary) statements)
74. (Refer to inside of label booklet for additional precautionary information and Directions for Use including First Aid and Storage and Disposal.)
75. May be used in hydroponic crop production.
76. Peel here (for (additional))(more)(complete) label instructions)
77. Plant-Based Actives
78. Results Done Right
79. Starts Working Immediately
80. Proven Effective
81. Proven Efficacy
82. Highly Effective Protection
83. Field Tested
84. 20% more vs. XX lb size [XX lbs]
85. 20% more vs XX size [[XX [ounce] [lbs]]
86. Xx% more vs xx lb size
87. TWIN [PACK/PAK]
88. Value [Pack/Pak]
89. VALUE [SIZE/SIZED]
90. Great Value
91. Club [Pack/Pak/Size]
92. Combo [Pack/Pak/Size]
93. Bonus [Size/Pack/Pak]
94. A [Insert dollar amount] Value
95. Treats up to XX sq. ft. (XX size)
96. Covers XX sq. ft (all pests) (XX pound size)
97. 100% Satisfaction Guaranteed
98. Guaranteed Results [or your money back]
99. {100%} Satisfaction [Guaranteed] or your money back
100. Money Back Guarantee
101. © [copyrighted material]
102. TM [trademarked]
103. Indoor/Outdoor
104. See back of booklet for precautionary statements and first aid
105. See attached booklet for complete directions for use and precautionary statements
106. Grow a better world
107. Take root in something bigger
108. The power Science meets the force of nature
109. A new wave of sustainable plant wellness products
110. Help your plants live their best life
111. BioUnite
112. BioFocus
113. BioTrust
114. BioFusion
115. Terraconnect
116. Bottle (body) contains XX% (post-consumer) recycled plastic (excluding dye) (excluding colorant)
117. Bottle (body contains XX% (post-consumer) recycled content (excluding dye) (excluding colorant)
118. Recyclable plastic bottle (body) made of XX% (post-consumer) recycled plastic (excluding dye) (excluding colorant)
119. Recyclable plastic bottle (body) made of XX% (post-consumer) content (excluding dye) (excluding colorant)

Sublabel C: Home & Garden Use (Residential Indoor and Residential Outdoor)

REGALIA® Bioprotectant Concentrate

A plant extract to boost the plants' defense mechanisms to protect against certain fungal and bacterial diseases, and to improve plant health.

Active ingredient: Extract of *Reynoutria sachalinensis*..... 5 %
Other ingredients: 95 %
Total 100 %

EPA Reg. No. 84059-3

EPA Est. No. 085970-FL-001

EPA Est. No. 084059-MI-001

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back/side/top/bottom [panel/label] for additional precautionary statements.

FIRST AID	
IF SWALLOWED:	Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED:	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
IF IN EYES:	Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER Have the product container or label with you when calling a poison control center or doctor, or if going for treatment. You may also contact 1-800-222-1222 for emergency medical treatment information.	



(CAN BE USED IN ORGANIC GARDENING)



) (For Organic Gardening)(

LOT #: Printed on container

Net Contents: xxxx

Pro Farm Group, Inc. 1530 Drew Ave., Davis, CA 95618

Made in the USA

REGALIA® Bioprotectant Concentrate; EPA Reg. No. 84059-3

MASTER LABEL – MBI-LBL-4 r23 v4 January 28, 2025

Page 58 of 66

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS – CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS: To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal law to apply this product in a manner inconsistent with its labeling.

OPTION 1

HOME AND GARDEN USE DIRECTIONS

REGALIA® Bioprotectant Concentrate is a broad spectrum (biological/bio) fungicide used for the control or suppression of a broad range of foliar diseases. REGALIA® Bioprotectant Concentrate may be used on vegetable crops, roses, fruits, nuts, flowers, bedding plants, foliage, houseplants, trees and shrubs located in residential landscapes. REGALIA® Bioprotectant Concentrate can be applied up to and on the day of harvest on all fruits and vegetables.

WHEN TO USE

For best results, apply REGALIA® Bioprotectant Concentrate prior to disease development or at the first sign of diseases and continue applying on a 7-day schedule or as needed.

BEFORE YOU USE

Read and follow these directions when using:

Do not allow spray to drift from application site.

Use only with pressurized hand-held sprayers, spray trigger bottles, or hose-end sprayers.

Do not allow spray mixture to stand overnight or for prolonged periods.

REGALIA® Bioprotectant Concentrate can be applied in commonly used pressurized hand-held sprayers, spray trigger bottles and hose-end sprayers.

HOW TO USE FOR HAND-HELD SPRAYERS AND SPRAY TRIGGER BOTTLES

Shake well before use.

Fill sprayer or bottle with appropriate amount of water and concentrate.

Mix the spray solution thoroughly.

Keep the spray solution agitated during application.

HOW TO USE FOR HOSE-END SPRAYERS

Shake well before use.

Follow hose-end sprayer instructions to determine how to fill, set dial, clean and disconnect from hose.

Set dial on sprayer to deliver rate as directed below.

HOW MUCH TO USE FOR ALL APPLICATIONS:

1 fluid ounce (2 TBSP) of REGALIA® Bioprotectant Concentrate per gallon of water.

Spray plants to complete wetness, covering both top and bottom leaf surfaces to ensure complete coverage.

Some pesticides (garden care products)(plant protection products) can cause phytotoxic effects ranging from slight burning or browning of leaves to distorted leaves, fruit, flowers or stems. Damage symptoms may vary with the type of plant that has been treated. It is impossible to test all plant species for phytotoxicity. To assure that the plants to

REGALIA® Bioprotectant Concentrate; EPA Reg. No. 84059-3

MASTER LABEL – MBI-LBL-4 r23 v4 January 28, 2025

Page 59 of 66

be treated are not sensitive to the treatment, apply a small amount of the product to a few leaves or the above ground portion of the plant and check back in 2–4 days for signs of phytotoxicity. Use product according to label directions.

OPTION 2

WHEN TO USE

For best results, apply REGALIA® Bioprotectant Concentrate prior to disease development or at the first sign of diseases and continue applying on a 7-day schedule or as needed.

[As a preventative, apply every 7 to 14 days until the potential for disease has passed. To control disease that is already present, apply on a 7-day schedule until disease symptoms are gone. Then continue spraying every 14 days to prevent disease recurrence.]

BEFORE YOU USE

Read label [before use]

Do not allow spray to drift from application site.

Use only with pressurized hand-held sprayers/[hand-held] trigger-spray bottles/hose-end sprayers.

Do not allow spray mixture to stand overnight or for prolonged periods.

HOW TO USE [for pressurized hand-held sprayers/[hand-held] trigger-sprayers]

Shake [concentrate] before use

Fill sprayer with 1fl.oz. (2 TBSP) of REGALIA® Bioprotectant Concentrate per gallon of water [(or 1 1/2 tsp. per 32 fl. oz. of water)]

Mix thoroughly

Spray both tops and bottoms of leaves/leaf surfaces [thoroughly] until dripping wet.

Shake sprayer occasionally during application [to keep product well mixed]

HOW TO USE [for hose-end sprayers]

Shake [concentrate] before use

Follow hose-end sprayer instructions to determine how to fill, set dial, clean and disconnect from hose.

Set dial on sprayer to deliver 1fl.oz. (2 TBSP) of REGALIA® Bioprotectant Concentrate per gallon of water

Spray both tops and bottoms of leaves/leaf surfaces [thoroughly] until dripping wet.

HOW TO USE [for Pre-Plant Dip Applications]

REGALIA® Bioprotectant Concentrate can be applied as a pre-plant dip for improved health and suppression of certain soil-borne diseases when transplanting. Shake [concentrate] before use. Mix 1oz. (2TBSP) REGALIA® Bioprotectant Concentrate per gallon of water, briefly submerge roots or plugs in mixture (ensuring full coverage, then remove) before transplanting.

HOW TO USE [for Soil Drench Applications]

Shake [concentrate] before use. Mix REGALIA® Bioprotectant Concentrate at a concentration of 1oz (2TBSP) per gallon of water and apply at a sufficient rate to thoroughly soak the soil and root zone. Make an initial application of REGALIA® Bioprotectant Concentrate during or shortly after transplant to reduce transplant shock, suppress soil-borne diseases and improve root growth. Multiple drench applications can be made on a 10-14-day interval.

[HOW IT WORKS/MODE OF ACTION]

REGALIA® Bioprotectant Concentrate contains an extract from the plant *Reynoutria sachalinensis*, that when applied according to label directions triggers an immune response that protects treated plants from numerous diseases. The induced resistance provided by REGALIA® is not systemic, therefore thorough coverage of above-ground plant parts is important to achieving optimal control. Repeat foliar applications at 7–14 day intervals to maintain induction and to protect new plant growth. Use REGALIA® Bioprotectant Concentrate as a preventative treatment prior to the development of disease symptoms.

[Phytotoxicity Note:]

or

(Some garden care products can cause adverse effects such as (yellowing) (browning) (slight burning of leaves.) (Some sensitive plant varieties may be slightly burned by garden care products.) (In general, plant damage can be avoided by following labeled rates. Plants to be treated can be checked for sensitivity to REGALIA® Bioprotectant Concentrate by applying the product to a few leaves and checking back in 2–4 days for signs of damage. Complex tank-mixes that include REGALIA® Bioprotectant Concentrate may increase the risk of visible spray residues on grape clusters. Test the mix on a small portion of the plants to be treated to ensure that a phytotoxic response will not occur as a result of the application. Do not apply REGALIA® Bioprotectant Concentrate within 14 days of applying a sulfur fungicide during periods of high temperatures. Do not use water volumes and/or adjuvants that create spray runoff or drip-accumulation at the leaf margins or at the base of fruit or on the harvested commodity.)

DISEASES CONTROLLED OR SUPPRESSED[†] [ON VEGETABLES, FRUITS, TREE NUTS, ORNAMENTAL PLANTS, TREES, SHRUBS, FLOWERS, BEDDING PLANTS, HEMP, FOLIAGE AND TROPICAL PLANTS]

Anthracoise (*Colletotrichum* spp.)

Bacteria (*Erwinia* spp., *Pseudomonas* spp., *Xanthomonas* spp.)

Bacterial Leaf Blight (*Xanthomonas campestris*)

Bacterial Speck (*Pseudomonas syringae* pv. Tomato)

Bacterial Spot (*Xanthomonas* spp.)

Bean Rust (*Uromyces appendiculatus*)

Black Mold (*Alternaria alternata*)

Black Rot/Black Crown Rot (*Alternaria* spp.)

Black Spot of Rose (*Diplocarpon rosea*)

Botrytis (*Botrytis* spp.)

Botrytis Leaf Blight (*Botrytis squamosa*)

Botrytis Neck Rot (*Botrytis* spp.)

Brown Rot (*Monilinia* spp.)

Downy Mildew[†] (*Bremia lactucae*, *Peronospora* spp., and *Plasmopara viticola*)

Early Blight (*Alternaria solani*)

Fire Blight (*Erwinia amylovora*)

Gummy Stem Blight (*Didymella bryoniae*)

Gray Mold (*Botrytis cinerea*)

Greasy Spot (*Mycosphaerella citri*)

Late Blight[†] (*Phytophthora infestans*)

Leaf Spots (*Alternaria* spp., *Bipolaris* spp., *Cercospora* spp. *Septoria* spp.)

Onion Downy Mildew (*Peronospora destructor*)

Onion Purple Blotch (*Alternaria porri*)

Pin Rot (*Alternaria/Xanthomonas* complex)

Phytophthora spp.

Powdery Mildew (*Uncinula necator*, *Erysiphe* spp., *Sphaerotheca* spp., *Oidiopsis taurica*, *Leveillula taurica*,

Podosphaera leucotricha, *Oidium* spp., *Podosphaera* spp.)

Rust (*Puccinia* spp.)

Scab[†] (*Venturia* spp.)

Sclerotinia Head and Leaf Drop[†] (*Sclerotinia* spp.)

Sour Rot (*Alternaria tenuis*, *Aspergillus* spp., *Botrytis cinerea*, *Cladosporium herbarum*, *Penicillium* spp., *Rhizopus arrhizus*)

Target Spot (*Corynespora cassiicola*)

Walnut Blight (*Xanthomonas campestris*)

White mold (*Sclerotinium rolfsii*, *Sclerotinium sclerotium*, *Sclerotinium* spp. *Sclerotinium trifolorium*)

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

REGALIA® Bioprotectant Concentrate; EPA Reg. No. 84059-3

MASTER LABEL – MBI-LBL-4 r23 v4 January 28, 2025

Page 61 of 66

Pesticide Storage: Store in a cool, dry place. Avoid freezing.

Pesticide Disposal and Container Handling: If empty: Nonrefillable container. Do not reuse or refill this container. Place in trash and offer for recycling if available.

If partially filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

WARRANTY

To the extent consistent with applicable law, the seller makes no warranty, expressed or implied, of merchantability, fitness or otherwise concerning use of this product. To the extent permitted by applicable law, the user assumes all risks of use, storage or handling that are not in strict accordance with the accompanying directions.



-or-

(*)The (insert company name) Guarantee – If for any reason you are not satisfied with this product, send proof of purchase to the address shown and we will gladly refund your purchase price.

SUBLABEL C OPTIONAL CLAIMS

1. Boost plants' defenses [for stronger healthier plants]
2. Strengthens plants' immunity
3. Improves plant health
4. Controls/Prevents common garden diseases [and improves plant health]
5. Controls/Prevents powdery mildew, leaf spot and rust
6. Controls/Prevents black spot on rose
7. Labeled Fungal and bacterial disease control
8. Defends gardens by boosting plant defenses
9. Defending gardens against diseases...one plant at a time!
10. Use on labeled fruits, vegetables and ornamentals
11. Can be applied as a pre-plant dip [for improved plant health]
12. Can be applied as a soil drench application
13. [Can be] use(d) as a preventative to protect labeled [growing] plants from common garden diseases
14. For use on labeled ornamental plants and edible crops/fruits/vegetables.
15. For use on labeled vegetables, roses, fruits, berries, nuts, flowers, bedding plants, houseplants, (ornamental) trees and shrubs [located in residential landscapes].
16. REGALIA® Bioprotectant Concentrate may be used on labeled vegetable(s) [crops], roses, fruits, berries, nuts, flowers, foliage, houseplants, (ornamental) trees and shrubs [located in residential landscapes].
17. REGALIA® Bioprotectant Concentrate is a broad spectrum fungicide used for the control or suppression of a broad range of labeled foliar [fungal and bacterial] diseases.
18. (Active ingredient is) a plant extract to boost the plants' defense mechanisms to protect against certain labeled fungal and bacterial diseases, and to improve plant health.
19. Active ingredient (is) made from a plant extract (botanical extract)(plant-based)
20. REGALIA® Bioprotectant Concentrate is an extract from the plant *Reynoutria sachalinensis*
21. REGALIA® Bioprotectant Concentrate can be applied up to and on the day of harvest [on all labeled fruits and vegetables].
22. Made in the U.S.A.
23. This container is made with X% recycled material
24. Guaranteed results(*)
25. Label date:
26. US Patents No. 4,863,734 and No. 5,989,429
27. REGALIA® is a trademark of Pro Farm Group, Inc.
28. Name and logo of Pro Farm Group are registered trademarks of Pro Farm Group, Inc.
29. © insert company copyright information
30. World rights reserved
31. Distributed by: insert company name and address
32. [For] questions/comments



- 33.
34.  Can be used for organic gardening
35.  For (use in) organic gardening
36. GENERAL INFORMATION: REGALIA® Bioprotectant Concentrate is an extract from the plant *Reynoutria sachalinensis*. REGALIA® Bioprotectant Concentrate applied to actively growing plants will improve plant health, and will help make the treated portions resistant to certain plant diseases. [Plant health benefits often result in greater yields at harvest, especially when crops/plants are stressed by pathogens or environmental conditions].
37. MODE OF ACTION: REGALIA® Bioprotectant Concentrate contains an extract obtained from *Reynoutria sachalinensis* plant material composed of active chemical compounds. Application of the extract activates the immune response,

increasing the plant's defense system through a five-fold increase in phenolics and antioxidants, and cell wall strengthening. These immune responses, sometimes referred to as induced systemic resistance or systemic acquired resistance, protects treated plants from numerous diseases.

When applied at rates and timing for disease control, the translaminar activity of REGALIA® Bioprotectant Concentrate provides induced resistance against important diseases and acts within one to two days of application. REGALIA® Bioprotectant Concentrate should be used as a preventative treatment prior to the development of disease symptoms. Repeat foliar applications per label instructions, following a 7–14-day interval to maintain plant immune response induction and to protect new plant growth.

(In addition to foliar applications, REGALIA® Bioprotectant Concentrate can be used in multiple application methods as a plant dip, soil drench, pre-plant incorporated soil application, in-furrow spray, or applied through drip irrigation to control or suppress certain soil-borne diseases and to promote healthy root growth and improve above ground plant growth.)

(When applied at rates and timing for plant health effects, the induced plant defense responses minimize the impacts of stress and disease, resulting in optimized yields at harvest. Applying Regalia® Bioprotectant Concentrate has been shown to increase leaf chlorophyll content and increase soluble protein content in some crops. These effects often lead to improved crop quality and/or yields.)

38. REGALIA® Bioprotectant Concentrate is (an extract from the plant *Reynoutria sachalinensis*) for use on labeled (ornamental plants), (turf), (agricultural crops), (edible crops), (non-edible crops), (food crops), (non-food crops), (feed crops), or (non-feed crops).
39. Optional Language: (*) and (*= Not for use in California)
40. Repackaging or relabeling of this product without express written permission from Pro Farm Group is prohibited.
41. Biofungicide
42. [UPC code]
43. [RF code]
44. Bio with Bite™
45. For labeled disease control
46. For labeled fungal control
47. For maximum harvest
48. US MRL exempt
49. Minimal PPE
50. Protection from Multiple Labeled Diseases
51. Leaves no detectable residues
52. Product(s) thoroughly tested
53. Proven results, since 2007
54. Trial Tested
55. (number)+ trials
56. Read full label before use
57. Prevents labeled (disease) and (fungal) build up
58. Protection from labeled bacterial and fungal disease
59. REGALIA® Bioprotectant Concentrate can be applied by (any labeled use pattern) to protect against (labeled) (diseases) and (fungal pathogens).
60. 2-10 tablespoons per 1,000 sq. ft. [must be consistent with rates to be listed on label]
61. For (turf) (grass)
62. For recreational turf (grass) and landscapes [when crop is listed]
63. (Specify pest:) (Tank-mix) (or rotate) with a (fungicide) for improved control.
64. Refer to the table in the SOIL TREATMENT USE DIRECTIONS (In-Furrow Applications) section to determine the proper rate per 1000 foot of row.
65. REGALIA® Bioprotectant Concentrate should be used as part of an Integrated Pest Management System.
66. REGALIA® Bioprotectant Concentrate can be applied following a soil fumigant.
67. Use the high(er) labeled rate when high(er) (disease)(fungal pathogen) pressure is expected.
68. Optional Language: (*) and (*= Not labeled for this use in California)
69. [optional listing of all pests, alternatively list specific pests for each crop]

70. Bioprotectant Concentrate
71. Outdoor
72. See back of booklet [or other specific location] for precautionary statements and first aid
73. See attached booklet for complete directions for use and precautionary statements
74. Indoor commercial/Outdoor
75. See attached booklet for full label
76. (see (side) (back) (inside) (other) panel for additional (first aid) (and) (precautionary) statements)
77. (Refer to inside of label booklet for additional precautionary information and Directions for Use including First Aid and Storage and Disposal.)
78. May be used in hydroponic crop production.
79. Peel here (for (additional)(more)(complete) label instructions)
80. READY TO USE
81. Combo Product
82. Effective Home Perimeter Treatment
83. Plant-Based Actives
84. Results Done Right
85. Protects your (grass) (lawn) [from top to bottom]
86. Starts Working Immediately
87. Proven Effective
88. Proven Efficacy
89. Highly Effective Protection
90. Field Tested
91. 20% more vs. XX lb size [XX lbs]
92. 20% more vs XX size [[XX [ounce] [lbs]]
93. Xx% more vs xx lb size
94. TWIN [PACK/PAK]
95. Value [Pack/Pak]
96. VALUE [SIZE/SIZED]
97. Great Value
98. Club [Pack/Pak/Size]
99. Combo [Pack/Pak/Size]
100. Bonus [Size/Pack/Pak]
101. A [Insert dollar amount] Value
102. Treats up to XX sq. ft. (XX size)
103. Covers XX sq. ft (all pests) (XX pound size)
104. 100% Satisfaction Guaranteed
105. Guaranteed Results [or your money back]
106. {100%} Satisfaction [Guaranteed] or your money back
107. Money Back Guarantee
108. © [copyrighted material]
109. TM [trademarked]
110. Indoor/Outdoor
111. See back of booklet for precautionary statements and first aid
112. See attached booklet for complete directions for use and precautionary statements
113. Grow a better world
114. Take root in something bigger
115. The power Science meets the force of nature
116. A new wave of sustainable plant wellness products
117. Help your plants live their best life
118. For residential use only.
119. BioUnite
120. BioFocus

- 121. BioTrust
- 122. BioFusion
- 123. Terraconnect
- 124. Bottle (body) contains XX% (post-consumer) recycled plastic (excluding dye) (excluding colorant)
- 125. Bottle (body contains XX% (post-consumer) recycled content (excluding dye) (excluding colorant)
- 126. Recyclable plastic bottle (body) made of XX% (post-consumer) recycled plastic (excluding dye) (excluding colorant)
- 127. Recyclable plastic bottle (body) made of XX% (post-consumer) content (excluding dye) (excluding colorant)
- 128. "3 in 1" optional claims (disease control, immunity, plant health)
- 129. Covers up to x sq ft
- 130. Hose end sprayer directions
- 131. Add capful equivalent conversion
- 132. For Listed Houseplants, Fruits, Vegetables and Flowers
- 133. Controls [insert disease],[insert disease],[insert disease], and other labeled diseases
- 134. Concentrate makes up to X gallons
- 135. Increased harvest and yield
- 136. Doubles your harvest
- 137. Bigger blooms
- 138. Proven results
- 139. Highly effective when used as directed
- 140. Visible results
- 141. Fast acting
- 142. Long lasting results
- 143. Won't harm plants, blooms or lawns
- 144. Year round
- 145. Stops and cures labeled diseases on contact
- 146. For Roses
- 147. For Flowers
- 148. For Lawns
- 149. Rainproof in hours
- 150. Doesn't create resistance
- 151. effective
- 152. Effective long lasting results
- 153. Super concentrate(d)
- 154. 5x the value
- 155. New technology
- 156. Patented ingredients
- 157. Patented technology