

84229-2

02/28/2008

1/25



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg.

Number:

84229-2

Date of Issuance:

Feb 28, 2008

NOTICE OF PESTICIDE:

☒ Registration☐ Reregistration

(under FIFRA, as amended)

Term of Issuance: conditional

Name of Pesticide Product:

Tide Aba 0.15EC Ag
Insecticide/Miticide

Name and Address of Registrant (include ZIP Code):

Tide International, USA, Inc.
c/o Pyxis Regulatory Consulting, Inc.
4116 136th St., NW
Gig Harbor, WA 98332

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) and (B) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Per the indicated EPA Guidelines, submit the following within 18 months measured from the effective date of this registration.
 - a. one year storage stability (830.6317)
 - b. corrosion characteristics (830.6320)
3. Make the following label changes before you release the product for shipment:
 - a. Revise the EPA Registration Number to read, "EPA Reg. No. 84229-2"
 - b. Celeriac, restrictions - change "do not apply more than 20 fl. oz." to "... 16 fl. oz."
 - c. Citrus, restrictions - add "Do not allow livestock to graze in treated orchards."
 - d. Cotton, restrictions - add "Do not apply through any type of irrigation system."
 - e. Mint, rate - change "8-16 fl. oz./A" to 8-12 fl. oz./A."
 - f. Potatoes, restrictions - change similar text to "Do not apply more than 32 fl. oz./A Aba 0.15 EC to control spider mites or Colorado potato beetle. Do not apply more than 48 fl. oz./A Aba 0.15 EC to control leafminers."

(continued next page)

Signature of Approving Official:

151

Thomas C. Harris
Biologist
Insecticide Rodenticide Branch
Registration Division (7505P)

Date:

FEB 28 2008

2/25

4. Submit two copies of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label stamped "accepted with comments" is enclosed for your records.

3/25

RESTRICTED USE PESTICIDE

Toxic to Fish, Mammals and Aquatic Organisms

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicators certification.

Tide Aba 0.15 EC Ag

Miticide/Insecticide

ACTIVE INGREDIENT:

Abamectin: 1.9 %

OTHER INGREDIENTS: 98.1 %

TOTAL: 100.0 %

Tide Aba 0.15 EC contains 0.15 lbs active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN

Warning Aviso

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none">• Move person to fresh air• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact Chemtrec at 1-800-424-9300 for emergency medical treatment information.	

ACCEPTED
with COMMENTS
in EPA Letter Dated:
2/28/2008
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
As amended, for the pesticide
Registered under EPA Reg. No.
84229-2

4/25

NOTE TO PHYSICIAN

Early signs of intoxication include dilation of pupils, muscular incoordination, and muscular tremors. Toxicity following accidental ingestion of this product can be minimized by early administration of chemical adsorbents (e.g., activated charcoal).

If toxicity from exposure has progressed to cause severe vomiting, the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parental fluid replacement therapy should be given, along with other required supportive measures (such as maintenance of blood pressure levels and proper respiratory functionality) as indicated by clinical signs, symptoms, and measurements.

In severe cases, observations should continue for at least several days until clinical condition is stable and normal. Since abamectin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic abamectin exposure.

**EPA Reg. No. 84229-
EPA Est. No.**

Manufactured for:
Tide International USA Inc.
21 Hubble
Irvine, CA 92618

Net Contents:

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING

Causes substantial but temporary eye injury. Do not get in eyes. Harmful if absorbed through skin or if swallowed. Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category E on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Socks and shoes.
- Chemical-resistant gloves (such as barrier laminate, nitrile rubber, neoprene rubber, viton).
- Protective eyewear.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from target areas. Do not contaminate water when disposing of equipment wash water or rinsate.

5/25

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Use of this product may pose a risk to threatened and endangered species of fish, amphibians, crustaceans (including fresh water shrimp), and insects. All use of this product in the state of California should comply with the recommendations of the California Endangered Species Project. Before using this product in California, consult with your country agriculture commissioner to determine use limitations that apply in your area.

PHYSICAL OR CHEMICAL HAZARDS

Combustible: Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

IMPORTANT – READ BEFORE USE

Read the entire Directions for Use, Limited Warranty and Disclaimer before using this product.

If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following warranty disclaimer, inherent risks of use and limitation of remedies.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls over short-sleeved shirt and short pants.
- Chemical-resistant gloves – EPA chemical-resistance category E (e.g., barrier laminate, nitrile rubber, neoprene rubber, viton).
- Chemical-resistant shoes and socks.
- Protective eyewear.

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR INSECT CONTROL, AND/OR ILLEGAL RESIDUES.

6/25

GENERAL INFORMATION

Aba 0.15 EC is an insecticide that will control certain insect pests. The product must be mixed with water and applied to labeled crops according to the specific crop use directions for effective control of these insects. For good mite and insect control, Aba 0.15 EC needs to be applied so that the foliage is thoroughly covered by the spray solution.

Phytotoxicity: Aba 0.15 EC has been tested for phytotoxicity and has been proven to be safe on a variety of crops. Aba 0.15 EC is also compatible with many commonly used pesticides, crop oils, and nutritional sprays. However, it is impossible to test all crop varieties and combination of spray mixtures under all possible growing conditions. Prior to mixing, users should test the compatibility of Aba 0.15 EC with any combination product using physical compatibility jar tests. Before applying Aba 0.15 EC and combination products to large areas of the crop, test small areas for phytotoxicity.

New York State Restriction: Do not apply Aba 0.15 EC by aerial application in New York State.

Chemigation: Do not apply this product through any type of irrigation system.

Spray Drift Precautions for Aba 0.15 EC Use on Celeriac, Cotton, Cucurbits, Fruiting Vegetables, Grapes, Hops, Leafy Vegetables, Potatoes, and Strawberries:

- Do not apply with ground application equipment within 25 ft. or with aircraft within 150 ft. of lakes, reservoirs, rivers, permanent streams, marshes, potholes, natural ponds, and estuaries, of commercial fish farm ponds.
- All aerial and ground application equipment must be properly calibrated, using appropriate carriers, and maintained. For aerial applications, mount the spray boom on the aircraft in a manner that minimizes drift caused by wing tip vortices. Use the minimum practical boom length but do not exceed 75% of wingspan or rotor diameter.
- Use the largest droplet size consistent with pest control. Minimize the formation of very small droplets by selecting appropriate nozzles, orienting nozzles away from the air stream as much as possible, and avoiding excessive spray boom pressure. Release the spray at the lowest height consistent with pest control and flight safety. Avoid application more than 10 ft. above the crop canopy.
- Make aerial or ground applications when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoiding applications when wind direction is toward the aquatic area can reduce risk of exposure to aquatic areas. Do not cultivate within 25 ft. (10 ft. for cotton) of the aquatic area to allow growth of a vegetative filter strip.
- Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore increase the likelihood of spray drifting to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.
- Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and temperatures that increase with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Resistance Management: Aba 0.15 EC contains the active ingredient abamectin. There are inherent risks of pests developing resistance to any product, therefore, it is strongly advised that Aba 0.15 EC be used in a sound resistance management program (including practices such as non-chemical alternatives (beneficial arthropods), rotation of Aba 0.15 EC with insecticides that have different modes of action, rotation of susceptible and non-susceptible plants, etc.). Once insect or mite-tolerant strains develop, Aba 0.15 EC will not be effective against labeled pests. In crops that have pests with multiple generations per crop per year, resistance management practices help reduce the risk of resistance development to Aba 0.15 EC. Additional information is available on implementation of these or other resistance management practices from your local agricultural advisor or company representative.

CROP USE DIRECTIONS

ALMONDS AND WALNUTS

Aba 0.15 EC will control the following pests in almonds and walnuts:

European red mite
Pacific Spider mite
Strawberry spider mite
Twospotted spider mite

Rate of Aba 0.15 EC	Directions for Application	Restrictions
<p>For Dilute Sprays^a: 2.5 – 5.0 fl. oz./100 gals</p> <p>For Concentrate Sprays^b: 10 – 20 fl. oz./A</p> <p>PLUS</p> <p>A Horticultural Spray Oil that is not a dormant oil</p> <p>For Dilute Sprays: 0.25% or 1 gal./A</p> <p>For Concentrate Sprays: Minimum of 1 gal./A</p>	<p>Apply Aba 0.15 EC using conventional dilute or concentrate sprays using ground application equipment. Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage. The size and number of trees per acre as well as foliage density will determine the gallons of spray required. Apply only in 40 gals. or more of water/A. Control of these mites/insects requires a thorough coverage of the spray solution on the tree foliage, and residual spider mite control increases if sprays are made to younger leaves.</p> <p>Time Aba 0.15 EC applications to occur when spider mites first appear. Repeat applications following the directions in the Restrictions column of this table.</p>	<p>Do not apply by air.</p> <p>Do not apply Aba 0.15 EC without a horticultural spray oil that is not a dormant oil (but one that is approved for use on almonds and walnuts).</p> <p>Do not apply more than 20 fl. oz Aba 0.15 EC per application, and no more than 40 fl. oz. per growing season.</p> <p>Make only 2 applications per growing season 21 days or more apart.</p> <p>Do not allow livestock to graze in treated orchards.</p> <p>Preharvest Interval: 21 days</p>
<p>^a Dilute Spray Rate is based on a volume of 400 gals./A dilute spray.</p> <p>^b Concentrate Spray Rate: determine the amount of Aba 0.15 EC required in a full cover dilute spray. Then use the same amount of product per acre in concentrate sprays that would be required for the dilute sprays to the same orchard/grove. Note that for small trees, the amount may be less than 10 fl. oz./A.</p>		
<p>Buffer Zone: To reduce the risk to aquatic organisms, maintain a 25-yard buffer zone around all water bodies containing aquatic life (streams, ponds, lakes, rivers, springs, irrigation canals containing water year-round, swamps, bogs and marshes).</p>		

8/25

APPLES

Aba 0.15 EC will control the following pests in almonds and walnuts:

European red mite
McDaniel spider mite
Tentiform leafminer
Twospotted spider mite
White apple leafhopper

Rate of Aba 0.15 EC	Directions for Application	Restrictions
<p>For Dilute Sprays^a: 2.5 – 5.0 fl. oz./100 gals</p> <p>For Concentrate Sprays^b: 10 – 20 fl. oz./A</p> <p>And add (for best results)</p> <p>A Horticultural Spray Oil that is not a dormant oil</p> <p>For Dilute Sprays: 0.25% or 1 gal./A</p> <p>For Concentrate Sprays: Minimum of 1 gal./A</p>	<p>Apply Aba 0.15 EC using conventional dilute or concentrate sprays using ground application equipment. Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage. The size and number of trees per acre as well as foliage density will determine the gallons of spray required. Apply only in 40 gals. or more of water/A. Control of these mites/insects requires a thorough coverage of the spray solution on the tree foliage.</p> <p>Time Aba 0.15 EC applications to occur when spider mites first appear or when insect thresholds are reached. Repeat applications following the directions in the Restrictions column of this table. Spider Mites: Apply before a threshold of 5 spider mites per leaf is reached. For residual spider mite control with Aba 0.15 EC, apply in combination with horticultural spray oil (not a dormant oil) to newer leaves of trees. Apply Aba 0.15 EC for spider mite control from petal fall through 6 weeks following petal fall.</p> <p>Tentiform leafminer: Apply onto eggs for control of new hatches and to early sap-feeder stages of first and second-generation tentiform leafminers when locally established thresholds have been reached. Do not apply during bloom.</p> <p>White Apple Leafhopper (Not for use west of the Rocky Mountains): Apply Aba 0.15 EC only to the first generation of white apple leafhoppers, just after petal fall and in combination with horticultural spray oil (not a dormant oil).</p>	<p>Do not apply by air.</p> <p>Reduced efficacy and/or residual control may occur if Aba 0.15 EC is not applied in combination with a horticultural spray oil (not a dormant oil). Follow all directions on the spray oil label or in spray guides. Use caution when applying Aba 0.15 EC with a horticultural spray oil or injury may occur with certain apple varieties (such as russetting on light-skinned Golden Delicious apples). In some cases, phytotoxicity and crop loss has been observed when Aba 0.15 EC plus a horticultural spray oil was applied less than 14 days before or after a captan application.</p> <p>Do not apply more than 20 fl. oz. Aba 0.15 EC per application, and no more than 40 fl. oz. per growing season.</p> <p>Make only 2 applications per growing season 21 days or more apart.</p> <p>Do not allow livestock to graze in treated orchards.</p> <p>Preharvest Interval: 28 days</p>

^a Dilute Spray Rate is based on a volume of 400 gals./A dilute spray.

^b Concentrate Spray Rate: determine the amount of Aba 0.15 EC required in a full cover dilute spray. Then use the same amount of product per acre in concentrate sprays that would be required for the dilute sprays to the same orchard/grove. Note that for small trees, the amount may be less than 10 fl. oz./A.

Spray Drift Precautions: Observe the following precautions when spraying in the vicinity of aquatic areas such as lakes, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farms:

- Spray last 3 rows upwind of aquatic areas using nozzles on one side only, with spray directed away from aquatic areas
- Avoid spray going over tops of trees by adjusting or turning off top nozzles. Shut off nozzles on the side away from the grove/orchard when spraying the outside row. Shut off nozzles when turning at ends of row and passing tree gaps in rows.
- Do not apply Aba 0.15 EC when weather conditions favor drift to aquatic areas.
- Do not apply within 110 feet upwind of aquatic areas or when wind speed is above 8 miles per hour.
- Do not apply during a temperature inversion.

AVOCADOS

Aba 0.15 EC will control the following pests in avocados:

Avocado thrips (Scirtothrips perseae)

Persea Mite (Oligonychus perseae)

Rate of Aba 0.15 EC	Directions for Application	Restrictions
<p>10 – 20 fl. oz./A (Do NOT use less than 10 fl. oz. of Aba 0.15 EC/A)</p> <p>And add (for best results): a Horticultural Spray Oil that is not a dormant oil</p> <p>Minimum of 1 to 4% v/v (Follow all directions on the spray oil label. Test small areas with oil concentrations greater than 2.0% before making applications to large areas.)</p>	<p>Apply Aba 0.15 EC using conventional dilute or concentrate sprays using ground or air application equipment. Although aerial application is permitted, best control of thrips is achieved with ground applications which provide better spray coverage. The user accepts all liability for thrip level and duration of control if Aba 0.15 EC is applied by air.</p> <p>Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage. The size and number of trees per acre as well as foliage density will determine the gallons of spray required. Apply only in 100 gals. or more of water/A by ground and 50 gals. of water by air application. [For ground applications where the spray volume is greater than 400 gals./A, use 2.5 fl. oz. Aba 0.15 EC per 100 gals. of water in combination with 1.0-4.0% of an NR (narrow range) 415 oil.] Control of these pests requires a thorough coverage of the spray solution on the tree foliage.</p> <p>Time Aba 0.15 EC applications to occur when immature thrips first appear but before 5 or more immature thrips per leaf/fruit are detected. Repeat applications following the directions in the Restrictions column of this table.</p> <p>Apply Aba 0.15 EC at 10-15 fl. oz/A for low (1-2 thrips per leaf/fruit) to moderate (3-4 thrips per leaf/fruit). Use the higher rate of 15-20 fl. oz./A for severe (more than 5 thrips per leaf/fruit) outbreaks. Apply Aba 0.15 EC in combination with a 1.0-4.0% NR 415 oil in combination with a Horticultural Spray Oil (not a dormant oil) which is approved for use on avocados.</p>	<p>Do not apply more than 20 fl. oz Aba 0.15 EC per application, and no more than 40 fl. oz. per growing season.</p> <p>Make only 2 applications per growing season 30 days or more apart.</p> <p>Preharvest Interval: 14 days</p>

CELERIAC (*Apium graveollens*)

Aba 0.15 EC will control the following pests in celeriac:

Twospotted spider mite

Rate of Aba 0.15 EC	Directions for Application	Restrictions
16.0 fl. oz./A	<p>Apply Aba 0.15 EC using conventional dilute or concentrate sprays using ground application equipment.</p> <p>Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage of the crop foliage. Apply only in 20 gals. or more of water/A by ground application. Control of this mite requires a thorough coverage of the spray solution on the crop foliage.</p> <p>Time Aba 0.15 EC applications to occur when mites first appear. Repeat applications following the directions in the Restrictions column of this table.</p> <p>Optimum results are obtained if Aba 0.15 EC is used in combination with a non-ionic surfactant.</p>	<p>Refer to the section on Spray Drift Precautions in the General Information section of this label.</p> <p>Do not apply by air.</p> <p>Do not apply more than 20 fl. oz Aba 0.15 EC per application, and no more than 48 fl. oz. after transplanting.</p> <p>To prevent resistance, make only 2 sequential applications per growing season 7 days or more apart.</p> <p>Preharvest Interval: 7 days</p>

11/25

CITRUS FRUIT CROP GROUP

Aba 0.15 EC will control the following pests in citrus (which includes but is not limited to citrus citron, citrus hybrids, citrus species (includes chironja, tangelo, tangor), grapefruit, kumquat, lemon, lime, mandarin (tangerine), sour orange, and sweet orange):

Asian citrus psylla
Broad mite
Citrus bud mite
Citrus leafminer
Citrus rust mite
Citrus thrips
Twospotted spider mite

Rate of Aba 0.15 EC	Directions for Application	Restrictions
<p>For Dilute Sprays^a: 1.0 – 2.0 fl. oz./100 gals. for all insects (except Asian citrus psylla, citrus leafminers and citrus thrips—use the concentrate spray rate)</p> <p>For Concentrate Sprays^{b, c}: 10 – 20 fl. oz./A for all listed insects/mites</p> <p>And add (for best results) a Horticultural Spray Oil that is not a dormant oil - 0.20% or 1 gal./A (see footnotes for additional rates)^c</p>	<p>Apply Aba 0.15 EC using conventional dilute or concentrate sprays using ground application equipment. Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage (outside coverage for Asian citrus psylla, citrus leafminer or citrus thrips). The size and number of trees per acre as well as foliage density will determine the gallons of spray required.</p> <p>Asian citrus psylla: Applications made in the spring, summer or fall should be directed at newly expanding foliage and will only control adults and nymphs present at the time of application.</p> <p>Mites: Applications made in the spring, summer or fall should be made when mites first appear.</p> <p>Citrus bud mite: Time application to occur at "bud swell".</p> <p>Citrus leafminer: Applications in the spring, summer or fall should be directed at newly expanding foliage.</p> <p>Citrus thrips: Correct timing is required. Apply after egg hatch has begun (typically, early to mid-hatch is best) but only the current generation will be controlled.</p>	<p>Do not apply by air.</p> <p>Do not apply more than 40 fl. oz. Aba 0.15 EC per growing season.</p> <p>Do not apply more than 3 applications per growing season 30 days or more apart.</p> <p>To prevent pest resistance to Aba 0.15 EC, do not apply in citrus nurseries.</p> <p>Preharvest Interval: 7 days</p>

^a Dilute Spray Rate is based on a volume of 1,000 gals./A dilute spray.

^b Concentrate Spray Rate: determine the amount of Aba 0.15 EC that would be required in a full cover dilute spray.

^c For optimum control of **Asian citrus psylla**, **broad mites**, **citrus rust mites** and **citrus leafminers**, apply 150-300 gals spray mix per acre plus a minimum of 3 gal./A of a horticultural oil (not a dormant oil) and apply using ground equipment at 1 to 1.5 miles per hour. For optimum control of **citrus bud mite**, apply Aba 0.15 EC in a minimum of 500 gals. per acre plus a minimum of 2 qts. (0.5%) per acre of a horticultural oil (not a dormant oil). For optimum control of **citrus thrips**, use 100-250 gals. per acre spray mix for best coverage of outside foliage.

12/25

Spray Drift Precautions: Observe the following precautions when spraying in the vicinity of aquatic areas such as lakes, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farms:

- Spray last 3 rows upwind of aquatic areas using nozzles on one side only, with spray directed away from aquatic areas
- Avoid spray going over tops of trees by adjusting or turning off top nozzles. Shut off nozzles on the side away from the grove/orchard when spraying the outside row. Shut off nozzles when turning at ends of row and passing tree gaps in rows.
- Do not apply Aba 0.15 EC when weather conditions favor drift to aquatic areas.
- Do not apply within 110 feet upwind of aquatic areas or when wind speed is above 8 miles per hour.
- Do not apply during a temperature inversion.

COTTON – WEST OF THE MISSISSIPPI ONLY

Aba 0.15 EC will control the following pests in cotton (**West of the Mississippi Only**):

Carmines spider mites
Pacific spider mites
Strawberry spider mites
Twospotted spider mites

Rate of Aba 0.15 EC	Directions for Application	Restrictions
<p>Early season: 4 – 6 fl. oz./A</p> <p>Mid-season/Lay-by: 8 – 16 fl. oz./A</p>	<p>Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage. Control of mites requires a thorough coverage of the spray solution on the crop foliage.</p> <p>Early season: Apply Aba 0.15 EC when the cotton is no taller than 10 inches. Apply only using ground application equipment. Do not apply less than 4 fl. oz. Aba 0.15 EC per acre.</p> <p>Mid-season and lay-by: Use Aba 0.15 EC at the specified rates 8 - 16 fl. oz./A) depending on the size of the plants and foliage density. Apply using ground application or by air in fixed-wing aircraft or by helicopter. Apply by air in a minimum of 5 gals. of water /A.</p> <p>Time Aba 0.15 EC applications to occur when mites first appear. Repeat applications following the directions in the Restrictions column of this table.</p>	<p>Refer to the section on Spray Drift Precautions in the General Information section of this label.</p> <p>Do not apply Aba 0.15 EC east of the Mississippi River.</p> <p>Do not apply more than 32 fl. oz. Aba 0.15 EC per growing season.</p> <p>Make only 2 applications per growing season 21 days or more apart.</p> <p>Do not feed foliage to livestock; do not allow livestock to graze treated cotton.</p> <p>Preharvest Interval: 20 days</p>

13/25

CUCURBITS CROP GROUP

Aba 0.15 EC will control the following pests in the Cucurbit Crop Group (which includes but is not limited to citron melon, cucumbers, gherkin, edible gourds (includes hyotan, cucuzza, hechima, Chinese okra); *Momordica* spp. (includes balsam apple, balsam pear, bittermelon, Chinese cucumber); muskmelon, cantaloupe, pumpkin, summer squash, winter squash (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); and watermelon):

Spider mites

Leafminers

Rate of Aba 0.15 EC	Directions for Application	Restrictions
8 – 16 fl. oz./A	<p>Apply Aba 0.15 EC using ground or air application equipment. Although aerial application is permitted, best control of insect pests is achieved with ground applications which proved better spray coverage. The user accepts all liability for mite level and duration of control if Aba 0.15 EC is applied by air.</p> <p>Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage. Control of these mites/insects requires a thorough coverage of the spray solution on the crop foliage.</p> <p>Do not apply in less than 20 gals. of water/A by ground and in less than 5 gals. of water by air application. To ensure thorough spray coverage, use a greater spray volume for high pest pressure, dense crop canopies, and adverse application conditions (high temperatures).</p> <p>Time Aba 0.15 EC applications to occur when insects first appear. Use 8 – 12 fl. oz./A for low to moderate insect pressure use 16 fl. oz./A for heavy insect infestations. Repeat applications following the directions in the Restrictions column of this table.</p>	<p>Refer to the section on Spray Drift Precautions in the General Information section of this label.</p> <p>Do not apply Aba 0.15 EC by air in the State of New York.</p> <p>Do not apply more than 48 fl. oz Aba 0.15 EC per growing season.</p> <p>Make only 2 sequential applications 7 days or more apart.</p> <p>Preharvest Interval: 7 days</p>

14/25

FRUITING VEGETABLE CROP GROUP

Aba 0.15 EC will control the following pests in the Fruiting Vegetable Crop Group (which includes but is not limited to tomato; eggplant; peppers (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper); and tomatillo):

Broad mite
Colorado potato beetle
Liriomyza leafminers
Spider mites
Thrips palmi
Tomato pinworm
Tomato russet mite
Potato psyllid (peppers only; provides suppression only)

Rate of Aba 0.15 EC	Directions for Application	Restrictions
<p>8 – 16 fl. oz./A (use the 16 fl. oz./A rate for tomato pinworm)</p> <p>Note: Although Aba 0.15 EC can be used without a wetting agent, improved spreading and penetration for better insect control is achieved using a combination of Aba 0.15 EC and a surfactant. Nonionic type surfactants are recommended (not binder or sticker-type surfactants).</p>	<p>Apply Aba 0.15 EC using ground or air application equipment. Although aerial application is permitted, best control of mites is achieved with ground applications which proved better spray coverage. The user accepts all liability for mite level and duration of control if Aba 0.15 EC is applied by air.</p> <p>Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage. Control of these mites/insects requires a thorough coverage of the spray solution on the crop foliage.</p> <p>Do not apply in less than 20 gals. of water/A by ground and in less than 5 gals. of water by air application. To ensure thorough spray coverage, use a greater spray volume for high pest pressure, dense crop canopies, and adverse application conditions (high temperatures).</p> <p>Repeat applications following the directions in the Restrictions column of this table.</p> <p>Mites and thrips : Apply when insects first appear using 8 fl. oz. for low to moderate insect populations and 16 fl. oz./A for heavy infestations.</p> <p>Tomato pinworm: Make application when moth activity is first observed and not later than when larvae emerge. Apply using the 16 fl. oz./A rate.</p> <p>Colorado potato beetle: Apply 8 fl. oz. for low to moderate insect populations and 16 fl. oz./A for heavy infestations.</p> <p>Potato psyllid (Peppers only): Aba 0.15 EC provides suppression only. Apply when insects first appear. Use 8 – 12 fl. oz. for low to moderate insect populations and 16 fl. oz./A for heavy infestations. Make sequential applications at 10-14 day intervals when heavy egg and nymph populations are found. Use of a spreading or penetration adjuvant or horticultural oil is recommended.</p>	<p>Refer to the section on Spray Drift Precautions in the General Information section of this label.</p> <p>Do not apply Aba 0.15 EC by air in the State of New York.</p> <p>Do not apply more than 48 fl. oz Aba 0.15 EC per growing season.</p> <p>Make only 2 sequential applications 7 days or more apart.</p> <p>To prevent insect resistance to Aba 0.15 EC, do not apply to fruiting vegetables grown for transplants.</p> <p>Preharvest Interval: 7 days</p>

15/25

GRAPES

Aba 0.15 EC will control the following pests in grapes:

Pacific spider mite
Twospotted spider mite
Western grapeleaf skeletonizer
Willamette spider mite.
*Western grape leafhopper**
*Varigated leafhopper**

*Aba 0.15 EC provides knock-down control only of these pests.

Rate of Aba 0.15 EC	Directions for Application	Restrictions
8 – 16 fl. oz./A PLUS a nonionic type surfactant is recommended (not binder or sticker-type surfactants).	<p>Apply Aba 0.15 EC using ground application equipment. Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage. Control of these mites/insects requires a thorough coverage of the spray solution on the crop foliage. Do not spray alternate rows: apply Aba 0.15 EC on both sides of the row to ensure maximum coverage of the crop.</p> <p>Do not apply in less than 50 gals. of water/A by ground. If electro-static sprayers are used, the volume may be less than 50 gals./A but do not use less than 5 gals. of water/A.</p> <p>Always use Aba 0.15 EC with a nonionic surfactant. This combination improves the wetting of leaves and provides an even coverage of spray. Not all combinations of nonionic surfactants and Aba 0.15 EC have been tested on every grapes variety or all types of growing conditions. Before using a nonionic surfactant, carefully read and follow the surfactant product's label.</p> <p>Apply when insects first appear using 8 - 12 fl. oz. for low to moderate insect populations and 16 fl. oz./A for heavy infestations. Repeat applications following the directions in the Restrictions column of this table.</p> <p>Mites : time application to occur when mites are first detected but before motiles exceed 5 per leaf. Western grapeleaf skeletonizer: Make application when larvae appear. Best control is achieved if application is made just after egg hatch. Western grape leafhopper, Varigated leafhopper: For knock-down control only, apply when pests first appear.</p>	<p>Refer to the section on Spray Drift Precautions in the General Information section of this label.</p> <p>Do not apply by air.</p> <p>Do not apply more than 16 fl. oz. per application or more than 32 fl. oz Aba 0.15 EC per growing season.</p> <p>Make only 2 applications per growing season. One repeat application may be applied 21 day or more after the first application if needed to maintain control.</p> <p>Do not allow livestock to graze in treated vineyards.</p> <p>Preharvest Interval: 28 days</p>

16/25

HERB CROP SUBGROUP (EXCEPT CHIVES)

Aba 0.15 EC will control the following pests in the Herb Crop Subgroup (except chives) (which includes but is not limited to basil; chamomile; catnip; coriander (leaf); dill weed; lavender; marigold; marjoram (*Oiganum* species); nasturtium; parsley (dried); rosemary; sage; summer and winter savory; tarragon; thyme; and wintergreen):

Liriomyza leafminers

Rate of Aba 0.15 EC	Directions for Application	Restrictions
<p>8 – 16 fl. oz./A</p> <p>Note: To improve the wetting of leaves and provide an even coverage of spray of the foliage, use Aba 0.15 EC inc combination with a wetting agent, such as a nonionic surfactant.</p>	<p>Apply Aba 0.15 EC using conventional dilute or concentrate ground application equipment. Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage. Control of leafminers requires a thorough coverage of the spray solution on the upper and lower leaves.</p> <p>Do not apply in less than 30 gals. of water/A by ground.</p> <p>Use the low rate for low to moderate insect pressure and the higher rate for heavy insect pressure. Time application to occur when adult flies are first detected. Repeat applications following the directions in the Restrictions column of this table.</p> <p>Not all combinations of nonionic surfactants and Aba 0.15 EC have been tested on every herb types and variety or under all types of growing conditions. Before using a nonionic surfactant, carefully read and follow the surfactant product's label.</p>	<p>Do not apply by air.</p> <p>Make only 2 applications per single cutting.</p> <p>Do not apply more than 6 applications per cropping season at the 8 fl. oz./A rate or more than 3 applications at the 16 fl. oz./A rate.</p> <p>Repeat applications may be made 7 days or more after the last application.</p> <p>Preharvest Interval: 14 days</p>

17/25

HOPS (EXCEPT IN CALIFORNIA)

Aba 0.15 EC will control the following pests in hops (not approved for use in California):

Twospotted spider mite

Rate of Aba 0.15 EC	Directions for Application	Restrictions
<p>8 – 16 fl. oz./A</p> <p>Note: Although Aba 0.15 EC can be used without a wetting agent, improved spreading and penetration for better insect control is achieved using a combination of Aba 0.15 EC and a surfactant. Nonionic type surfactants are recommended (not binder or sticker-type surfactants).</p>	<p>Apply Aba 0.15 EC using ground application equipment. Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage. Control of mites requires a thorough coverage of the spray solution on the upper and lower leaves of the crop.</p> <p>The size and foliage density of plants will determine the gallons of spray required for thorough coverage. Use the same amount of Aba 0.15 EC for concentrate or dilute sprays. For plants 6 – 8 feet in height (1/2 trellis growth), apply 8 – 16 fl. oz./A of Aba 0.15 EC in a minimum of 40 gals. of water per acre. For plants taller than 6-8 feet, do not use less than 16 fl. oz./A in a minimum of 100 gals. of water/A.</p> <p>Time application to occur when mites are first detected. One repeat application may be made following the directions in the Restrictions column of this table.</p>	<p>Refer to the section on Spray Drift Precautions in the General Information section of this label.</p> <p>Do not apply by air.</p> <p>Make only 2 applications per season.</p> <p>Do not apply more than 32 fl. oz./A per growing season.</p> <p>Repeat applications may be made 21 days or more after the last application. To prevent insect resistance; use other miticides with different modes of action before making a repeat application of Aba 0.15 EC.</p> <p>Do not allow livestock to graze in treated hop yards.</p> <p>Preharvest Interval: 28 days</p>

18/25

LEAFY VEGETABLE CROP GROUP

Aba 0.15 EC will control the following pests in the Leafy Vegetable Crop Group (which includes but is not limited to celery, lettuce (head and leaf), amaranth (Chinese spinach), garden cress, upland cress, dandelion, endive, fennel, parsley, radicchio, rhubarb, spinach, and Swiss chard):

Carmine spider mite
Liriomyza leafminers
Twospotted spider mite

Rate of Aba 0.15 EC	Directions for Application	Restrictions
<p>8 – 16 fl. oz./A</p> <p>Note: Although Aba 0.15 EC can be used without a wetting agent, improved spreading and penetration for better insect control is achieved using a combination of Aba 0.15 EC and a surfactant. Nonionic type surfactants are recommended (not binder or sticker-type surfactants).</p>	<p>Apply Aba 0.15 EC using ground or air application equipment. Although aerial application is permitted, best control of mites is achieved with ground applications which proved better spray coverage. The user accepts all liability for mite level and duration of control if Aba 0.15 EC is applied by air.</p> <p>Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage. Control of these mites/insects requires a thorough coverage of the spray solution on the crop foliage.</p> <p>Do not apply in less than 20 gals. of water/A by ground and in less than 5 gals. of water by air application. To ensure thorough spray coverage, use a greater spray volume for high pest pressure, dense crop canopies, and adverse application conditions (high temperatures).</p> <p>Apply Aba 0.15 EC at 8 – 12 fl. oz./A for low to moderate insect populations and at 16 fl. oz./A for heavy infestations. Make repeat applications if needed to maintain control according to the Restrictions column of this table.</p> <p>Mites : Apply when insects first appear. Leafminers: Time application to occur when adult flies are first detected.</p>	<p>Refer to the section on Spray Drift Precautions in the General Information section of this label.</p> <p>Do not apply Aba 0.15 EC by air in the State of New York.</p> <p>Do not apply more than 48 fl. oz. Aba 0.15 EC per growing season.</p> <p>Make only 2 sequential applications 7 days or more apart.</p> <p>To prevent insect resistance to Aba 0.15 EC, do not apply to leafy vegetables grown for transplants.</p> <p>Preharvest Interval: 7 days</p>

19/25

MINT (PEPPERMINT AND SPEARMINT)

Aba 0.15 EC will control the following pests in peppermint and spearmint:

Twospotted spider mite

Rate of Aba 0.15 EC	Directions for Application	Restrictions
<p>8 – 16 fl. oz./A</p> <p>Note: Although Aba 0.15 EC can be used without a wetting agent, improved spreading and penetration for better insect control is achieved using a combination of Aba 0.15 EC and either a nonionic surfactant or organosilicone-based surfactant at the manufacturer's recommended use rates.</p>	<p>Apply Aba 0.15 EC using ground or air application equipment. Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage. Control of these mites/insects requires a thorough coverage of the spray solution on the crop foliage.</p> <p>Do not apply in less than 20 gals. of water/A by ground and in less than 5 gals. of water by air application. To ensure thorough spray coverage, use a greater spray volume for high pest pressure, dense crop canopies, and adverse application conditions (high temperatures).</p> <p>Time applications to occur when insects first appear. Make repeat applications if needed to maintain control according to the Restrictions column of this table.</p>	<p>Do not apply Aba 0.15 EC by air in the State of New York.</p> <p>Do not apply more than 12 fl. oz. Aba 0.15 EC per application or more than 36 fl. oz. per growing season.</p> <p>Make only 2 sequential applications 7 days or more apart. Do not apply more than 3 applications per season.</p> <p>Do not allow livestock to graze or feed treated foliage to livestock.</p> <p>Preharvest Interval: 28 days</p>

20/25

PEARS (INCLUDING ORIENTAL PEARS)

Aba 0.15 EC will control the following pests in pears and Oriental pears:

European red mite
McDaniel spider mite
Pear psylla
Pear rust mite
Twospotted spider mite
Yellow mite

Rate of Aba 0.15 EC	Directions for Application	Restrictions
<p>For Dilute Sprays^a: 2.5 – 5.0 fl. oz./100 gals</p> <p>For Concentrate Sprays^b: 10 – 20 fl. oz./A</p> <p>And add (for best results)</p> <p>A Horticultural Spray Oil that is not a dormant oil</p> <p>For Dilute Sprays: 0.25% or 1 gal./A</p> <p>For Concentrate Sprays: Minimum of 1 gal./A</p>	<p>Apply Aba 0.15 EC using conventional dilute or concentrate sprays using ground application equipment. Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage. The size and number of trees per acre as well as foliage density will determine the gallons of spray required. Apply only in 40 gals. or more of water/A. Control of these mites/insects requires a thorough coverage of the spray solution on the tree foliage.</p> <p>Time Aba 0.15 EC applications to occur when spider mites first appear or when insect thresholds are reached. Repeat applications following the directions in the Restrictions column of this table.</p>	<p>Do not apply by air.</p> <p>Reduced efficacy and/or residual control may occur if Aba 0.15 EC is not applied with a horticultural spray oil (not a dormant oil). Follow all directions on the spray oil label or spray guides. In some cases, phytotoxicity and crop loss has been observed when Aba 0.15 EC plus a horticultural spray oil was applied in less than 14 days before or after a captan application.</p> <p>Do not apply more than 20 fl. oz. Aba 0.15 EC per application, and no more than 40 fl. oz. per growing season.</p> <p>Make only 2 applications per growing season 21 days or more apart.</p> <p>Do not allow livestock to graze in treated orchards.</p> <p>Preharvest Interval: 28 days</p>
<p>^a Dilute Spray Rate is based on a volume of 400 gals./A dilute spray.</p> <p>^b Concentrate Spray Rate: determine the amount of Aba 0.15 EC required in a full cover dilute spray. Then use the same amount of product per acre in concentrate sprays that would be required for the dilute sprays to the same orchard/grove. Note that for small trees, the amount may be less than 10 fl. oz./A.</p>		
<p>Spray Drift Precautions: Observe the following precautions when spraying in the vicinity of aquatic areas such as lakes, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farms:</p> <ul style="list-style-type: none"> --Spray last 3 rows upwind of aquatic areas using nozzles on one side only, with spray directed away from aquatic areas --Avoid spray going over tops of trees by adjusting or turning off top nozzles. Shut off nozzles on the side away from the grove/orchard when spraying the outside row. Shut off nozzles when turning at ends of row and passing tree gaps in rows. --Do not apply Aba 0.15 EC when weather conditions favor drift to aquatic areas. --Do not apply within 110 feet upwind of aquatic areas or when wind speed is above 8 miles per hour. --Do not apply during a temperature inversion. 		

21/25

PLUMS AND PRUNES

Aba 0.15 EC will control the following pests in plums and prunes:

European red mite

Pacific spider mite

Twospotted spider mite

Rate of Aba 0.15 EC	Directions for Application	Restrictions
<p>For Dilute Sprays^a: 2.5 – 5.0 fl. oz./100 gals</p> <p>For Concentrate Sprays^b: 10 – 20 fl. oz./A</p> <p>PLUS</p> <p>A Horticultural Spray Oil that is not a dormant oil</p> <p>For Dilute Sprays: 0.25% or 1 gal./A</p> <p>For Concentrate Sprays: Minimum of 1 gal./A</p> <p>OR</p> <p>Nonionic surfactant (at the manufacturer's recommended label rate).</p>	<p>Apply Aba 0.15 EC using conventional dilute or concentrate sprays using ground application equipment. Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage. The size and number of trees per acre as well as foliage density will determine the gallons of spray required. Apply only in 40 gals. or more of water/A. Control of these mites/insects requires a thorough coverage of the spray solution on the tree foliage.</p> <p>Always use Aba 0.15 EC with either a nonionic surfactant or a horticultural spray oil (not a dormant oil). The combination of Aba 0.15 EC and the nonionic surfactant helps to provide an even coverage of spray on the foliage and to improve the wetting and penetration of leaves for better insect control. Not all combinations of nonionic surfactants and Aba 0.15 EC have been tested on every plum/prune variety or under all types of growing conditions. Before using a nonionic surfactant, carefully read and follow the surfactant product's label.</p> <p>Efficacy may be Improved with a combination of Aba 0.15 EC and a horticultural spray oil (not a dormant oil). However, the chances of phytotoxicity to fruit and foliage increase with this combination. Apply at the recommended rates in this table.</p> <p>Time Aba 0.15 EC applications to occur when mites first appear. Repeat applications following the directions in the Restrictions column of this table.</p>	<p>Do not apply by air.</p> <p>Do not apply more than 20 fl. oz. Aba 0.15 EC per application, and no more than 40 fl. oz. per growing season.</p> <p>Make only 2 applications per growing season 21 days or more apart.</p> <p>Do not allow livestock to graze in treated orchards.</p> <p>Preharvest Interval: 21 days</p>
<p>^a Dilute Spray Rate is based on a volume of 400 gals./A dilute spray.</p> <p>^b Concentrate Spray Rate: determine the amount of Aba 0.15 EC required in a full cover dilute spray. Then use the same amount of product per acre in concentrate sprays that would be required for the dilute sprays to the same orchard/grove. Note that for small trees, the amount may be less than 10 fl. oz./A.</p>		

Spray Drift Precautions: Observe the following precautions when spraying in the vicinity of aquatic areas such as lakes, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farms:

- Spray last 3 rows upwind of aquatic areas using nozzles on one side only, with spray directed away from aquatic areas
- Avoid spray going over tops of trees by adjusting or turning off top nozzles. Shut off nozzles on the side away from the grove/orchard when spraying the outside row. Shut off nozzles when turning at ends of row and passing tree gaps in rows.
- Do not apply Aba 0.15 EC when weather conditions favor drift to aquatic areas.
- Do not apply within 100 feet upwind of aquatic areas or when wind speed is above 8 miles per hour.
- Do not apply during a temperature inversion.

23/25

POTATOES

Aba 0.15 EC will control the following pests in potatoes:

Colorado potato beetle
Liriomyza leafminers
Spider mites

Rate of Aba 0.15 EC	Directions for Application	Restrictions
<p>8 – 16 fl. oz./A</p> <p>Note: Although Aba 0.15 EC can be used without a wetting agent, improved spreading and penetration for better insect control is achieved using a combination of Aba 0.15 EC and either a nonionic surfactant or organosilicone-based surfactant at the manufacturer's recommended use rates.</p>	<p>Apply Aba 0.15 EC using ground or air application equipment. Although aerial application is permitted, best control of mites is achieved with ground applications which proved better spray coverage. The user accepts all liability for mite level and duration of control if Aba 0.15 EC is applied by air.</p> <p>Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage. Control of these mites/insects requires a thorough coverage of the spray solution on the crop foliage.</p> <p>Do not apply in less than 20 gals. of water/A by ground and in less than 5 gals. of water by air application. To ensure thorough spray coverage, use a greater spray volume for high pest pressure, dense crop canopies, and adverse application conditions (high temperatures).</p> <p>Use of Aba 0.15 EC in combination with a sticker or binder type product such as Bravo® Weather Stik® may reduce control of insects and spider mites.</p> <p>Apply Aba 0.15 EC at 8 – 12 fl. oz./A for low to moderate insect populations and at 16 fl. oz./A for heavy infestations. Make repeat applications if needed to maintain control according to the Restrictions column of this table.</p> <p>Spider mites: Time applications to occur when insects are first detected.</p> <p>Colorado potato beetle: Apply Aba 0.15 EC when about 50% of the egg masses have hatched and the larvae are present. When 2 applications are required, do not make the second application to the same Colorado potato beetle generation.</p> <p>Liriomyza leafminers: Time application to occur when adult flies are first detected.</p>	<p>Refer to the section on Spray Drift Precautions in the General Information section of this label.</p> <p>Do not apply Aba 0.15 EC by air in the State of New York.</p> <p>Do not apply more than 32 fl. oz./A Aba 0.15 EC to control spider mites or more than 48 fl. oz./A to control Colorado potato beetle.</p> <p>Make only 2 sequential applications 7 days or more apart.</p> <p>Do not allow livestock to graze or feed treated foliage to livestock.</p> <p>Preharvest Interval: 14 days</p>

24/25

STRAWBERRIES

Aba 0.15 EC will control the following pests in strawberries:

Strawberry spider mite
Twospotted spider mite.

Rate of Aba 0.15 EC	Directions for Application	Restrictions
16 fl. oz./A Note: Although Aba 0.15 EC can be used without a wetting agent, improved spreading and penetration for better insect control is achieved using a combination of Aba 0.15 EC and a nonionic surfactant at the manufacturer's recommended use rate.	<p>Apply Aba 0.15 EC using ground application equipment. Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage. Control of these mites requires a thorough coverage of the spray solution on the crop foliage. Apply Aba 0.15 EC to the tops and undersides of the leaves.</p> <p>Do not apply in less than 50 gals. of water/A by ground. If electro-static sprayers are used, the volume may be less than 50 gals./A but do not use less than 10 gals. of water/A.</p> <p>Apply Aba 0.15 EC when mites are first detected. A second repeat application may be made 7 – 10 days later. This sequence of two applications may be repeated if needed but follow the directions in the Restrictions column of this table.</p>	<p>Refer to the section on Spray Drift Precautions in the General Information section of this label.</p> <p>Do not apply by air.</p> <p>Do not apply more than 16 fl. oz. per application or more than 64 fl. oz Aba 0.15 EC per growing season.</p> <p>After the first 2 applications, wait 21 day or more before making additional applications if needed to maintain control.</p> <p>To prevent insect resistance to Aba 0.15 EC, do not apply in strawberry nurseries.</p> <p>Preharvest Interval: 3 days</p>

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

Pesticide Storage

Store in a tightly closed container in a cool, dry place.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal

Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures allowed by State or local authorities.

Bravo and Weather Stik are registered trademarks of a Syngenta Group Company

LIMITED WARRANTY AND DISCLAIMER

Tide International USA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below.

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such use or the results to be obtained if not used in accordance with printed directions and established safe practice. To the extent consistent with applicable law, buyer's exclusive remedy and manufacturer's or seller's exclusive liability for any and all claims, losses, damages or injuries resulting from the use or handling of this product, whether or not based in contract, negligence, strict liability in tort or otherwise shall be limited, at the manufacturer's option to replacement of, or the repayment of the purchase price for, the quantity of product with respect to which damages are claimed.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Tide International USA or the seller. All such risks shall be assumed by buyer.

Terms and Conditions of Use

If terms of the Limited Warranty & Disclaimer and Inherent Risks of Use are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer and Inherent Risks of Use.