



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

September 24, 2015

Nicole Cochran
Agent for Tide International USA, Inc.
c/o Pyxis Regulatory Consulting, Inc.
4110 136th St. NW
Gig Harbor, WA 98332

Subject: Label Amendment – Adding chives, bulb onions, dry beans, and tuberous and
corn vegetables to the label as well as condensing other sections to fit current crop groupings
Product Name: Timectin 0.15EC Ag Insecticide/Miticide
EPA Registration Number: 84229-2
Application Date: June 16, 2015
Decision Number: 506209

Dear Nicole Cochran:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2
EPA Reg. No. 84229-2
Decision No. 506209

with FIFRA section 6. If you have any questions, please contact Jessica Rogala by phone at 703-347-0263, or via email at rogala.jessica@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to be 'MS' with a long horizontal stroke extending to the right.

Mark Suarez
Product Manager 07
Invertebrate & Vertebrate Branch 3
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

RESTRICTED USE PESTICIDE

Toxic to Fish, Mammals and Aquatic Organisms

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicators certification.

Timectin™ 0.15 EC Ag

Insecticide/Miticide

[Alternate Brand Names: AgMectin 0.15EC Insecticide/Miticide, Tide Timectin™ 0.15 EC Ag Insecticide/Miticide]

ACTIVE INGREDIENT:

Abamectin: 1.9 %

OTHER INGREDIENTS: 98.1 %

TOTAL: 100.0 %


Timectin 0.15 EC Ag Insecticide/Miticide contains 0.15 lb. active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If in eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	• Move person to fresh air • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact Chemtrec at 1-800-424-9300 for emergency medical treatment information.	



NOTE TO PHYSICIAN

Early signs of intoxication include dilation of pupils, muscular incoordination, and muscular tremors. Toxicity following accidental ingestion of this product can be minimized by early administration of chemical adsorbents (e.g., activated charcoal).

If toxicity from exposure has progressed to cause severe vomiting, the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parental fluid replacement therapy should be given, along with other required supportive measures (such as maintenance of blood pressure levels and proper respiratory functionality) as indicated by clinical signs, symptoms, and measurements.

In severe cases, observations should continue for at least several days until clinical condition is stable and normal. Since abamectin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic abamectin exposure.

EPA Reg. No. 84229-2

EPA Est. No.

Manufactured for:

Tide International USA, Inc.

21 Hubble

Irvine, CA 92618

Net Contents:

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Causes substantial but temporary eye injury. Do not get in eyes. Harmful if absorbed through skin or if swallowed. Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category E on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Socks and shoes
- Chemical-resistant gloves (such as barrier laminate, nitrile rubber, neoprene rubber, viton)
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from target areas. Do not contaminate water when disposing of equipment wash water or rinsate.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Use of this product may pose a risk to threatened and endangered species of fish, amphibians, crustaceans (including fresh water shrimp), and insects. All use of this product in the state of California should comply with the recommendations of the California Endangered Species Project. Before using this product in California, consult with your county agriculture commissioner to determine use limitations that apply in your area.

PHYSICAL OR CHEMICAL HAZARDS

Combustible: Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

IMPORTANT – READ BEFORE USE

Read the entire Directions for Use, Limited Warranty and Disclaimer before using this product.

If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following warranty disclaimer, inherent risks of use and limitation of remedies.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves – EPA chemical-resistance category E (e.g., barrier laminate, nitrile rubber, neoprene rubber, viton)
- Chemical-resistant shoes and socks
- Protective eyewear

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR INSECT CONTROL, AND/OR ILLEGAL RESIDUES.

Timectin 0.15 EC Ag Insecticide/Miticide is an insecticide that will control certain insect pests. The product must be mixed with water and applied to labeled crops according to the specific crop use directions for effective control of these insects. For good mite and insect control, Timectin 0.15 EC Ag Insecticide/Miticide needs to be applied so that the foliage is thoroughly covered by the spray solution.

Phytotoxicity: Timectin 0.15 EC Ag Insecticide/Miticide has been tested for phytotoxicity and has been proven to be safe on a variety of crops. Timectin 0.15 EC Ag Insecticide/Miticide is also compatible with many commonly used pesticides, crop oils, and nutritional sprays. However, it is impossible to test all crop varieties and combination of spray mixtures under all possible growing conditions. Prior to mixing, users should test the compatibility of Timectin 0.15 EC Ag Insecticide/Miticide with any combination product using physical compatibility jar tests. Before applying Timectin 0.15 EC Ag Insecticide/Miticide and combination products to large areas of the crop, test small areas for phytotoxicity.

New York State Restriction: Do not apply Timectin 0.15 EC Ag Insecticide/Miticide by aerial application in New York State.

Chemigation: Do not apply this product through any type of irrigation system.

SPRAY DRIFT

Responsibility

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator is responsible for considering all of these factors when making application decisions.

NOTE: When states have more stringent regulations, they must be observed.

Spray Drift Precautions for Application with Aircraft or Ground Application Equipment

- Apply Timectin 0.15 EC Ag Insecticide/Miticide only when wind velocity favors on target product deposition (approximately 3 to 10 mph).
- Do not apply with ground application equipment within 25 ft. of or with aircraft within 150 ft. of lakes, reservoirs, rivers, permanent streams, marshes, pot holes, natural ponds, estuaries, or commercial fish farm ponds.
- Do not cultivate within 25 ft. of the aquatic area to allow growth of a vegetative filter strip.
- Do not allow this product to drift onto non-target areas. Drift may result in illegal residues or injury to nontarget species. Risk of exposure to sensitive areas can be reduced by applying this product when the wind direction is away from the sensitive area.
- Do not apply when the weather conditions may cause drift.
- Avoid application when the temperature is high and/or the humidity is low. These conditions increase the evaporation of spray droplets and the likelihood of drift to aquatic areas.
- Do not apply when wind speed or wind gusts are greater than 15 mph.
- Do not apply when wind speed is below 2 mph because wind direction will vary and there is a high potential for inversion.
- Observe the following precautions when using ground application to spray tree crops or hops in the vicinity of aquatic areas such as lakes, reservoirs, permanent streams, marshes, potholes, natural ponds, estuaries, or commercial fish ponds:
 - Do not apply Timectin 0.15 EC Ag Insecticide/Miticide when weather conditions favor drift to aquatic areas.
 - Do not apply within 110 ft. upwind of aquatic areas or when wind speed is above 8 mph.
 - Spray last 3 rows upwind of aquatic areas using nozzles on one side only, with spray directed away from the aquatic areas.
 - Avoid spray going over tops of trees by adjusting or turning off top nozzles. Shut off nozzles on the side away from the grove/orchard when spraying the outside row. Shut off nozzles when turning at ends of row and passing tree gaps in rows.

Spray Drift Precautions for Aerial Application

Drift Management Requirements

The following drift management requirements must be followed to avoid off-target movement from aerial applications to agricultural field crops.

Outermost Nozzle Distance

The distance of the outermost nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.

Nozzle Directions

Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Maximum Wind Speed

Do not apply when wind speed is greater than 15 mph.

Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see **Wind, Temperature and Humidity, and Temperature Inversions**).

Controlling Droplet Size

Volume – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure – Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of Nozzles – Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation – Orienting nozzles so that the spray is released parallel to the air stream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.

Nozzle Type – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

Boom Length – For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

Application Height – Applications should not be made at a height greater than 10 ft. above the top of the target plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment – When applications are made with a cross wind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind.

Wind – Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application must be avoided below 2 mph due to variable wind direction and high inversion potential.

NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity – To compensate for evaporation when applying Timectin 0.15 EC Ag Insecticide/Miticide in low relative humidity, set up equipment to produce larger droplets. Evaporation of droplets is most severe when conditions are both hot and dry.

Temperature Inversions - Timectin 0.15 EC Ag Insecticide/Miticide must not be applied during a temperature inversion because the potential for drift is high. Temperature inversions restrict vertical air mixing, and this causes small, suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds that are common during inversions. Temperature inversions are characterized by temperatures that increase with altitude and are common on nights with limited cloud cover and light to no wind. Inversions begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, the movement of smoke from a ground source or an aircraft smoke generator can also identify inversions. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates, indicates good vertical air mixing.

Resistance Management: Timectin 0.15 EC Ag Insecticide/Miticide contains the active ingredient abamectin. Abamectin is a Group 6 insecticide. There are inherent risks of pests developing resistance to any product, therefore, it is strongly advised that Timectin 0.15 EC Ag Insecticide/Miticide be used in a sound resistance management program (including practices such as non-chemical alternatives (beneficial arthropods), rotation of Timectin 0.15 EC Ag Insecticide/Miticide with insecticides that have different modes of action, rotation of susceptible and non-susceptible plants, etc.). Once insect or mite-tolerant strains develop, Timectin 0.15 EC Ag Insecticide/Miticide will not be effective against labeled pests. In crops that have pests with multiple generations per crop per year, resistance management practices help reduce the risk of resistance development to Timectin 0.15 EC Ag Insecticide/Miticide. Additional information is available on implementation of these or other resistance management practices from your local agricultural advisor or company representative.

CROP USE DIRECTIONS

APPLES

Timectin 0.15 EC Ag Insecticide/Miticide will control the following pests in apples:

European red mite
McDaniel spider mite
Tentiform leafminer
Twospotted spider mite
White apple leafhopper

Rate of Timectin 0.15 EC Ag Insecticide/Miticide	Directions for Application	Restrictions
For Dilute Sprays^a: 2.5 – 5.0 fl. oz./100 gals For Concentrate Sprays^b: 10 – 20 fl. oz./A And add (for best results) A Horticultural Spray Oil that is not a dormant oil such as VINTRE. Horticultural Spray Oil: For Dilute Sprays: 0.25% or 1 gal./A For Concentrate Sprays: Minimum of 1 gal./A VINTRE (only during non-fruit bearing stages) Use 16 - 64 fl. oz. per acre at rates no greater than 0.3% v/v.	Apply Timectin 0.15 EC Ag Insecticide/Miticide using conventional dilute or concentrate sprays using ground application equipment. Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage. The size and number of trees per acre as well as foliage density will determine the gallons of spray required. Apply only in 40 gals. or more of water/A. Control of these mites/insects requires a thorough coverage of the spray solution on the tree foliage. Time Timectin 0.15 EC Ag Insecticide/Miticide applications to occur when spider mites first appear or when insect thresholds are reached. Repeat applications following the directions in the Restrictions column of this table. Spider Mites: Apply before a threshold of 5 spider mites per leaf is reached. For residual spider mite control with Timectin 0.15 EC Ag Insecticide/Miticide, apply in combination with horticultural spray oil (not a dormant oil) to newer leaves of trees. Apply Timectin 0.15 EC Ag Insecticide/Miticide for spider mite control from petal fall through 6 weeks following petal fall. Tentiform leafminer: Apply onto eggs for control of new hatches and to early sap-feeder stages of first and second-generation tentiform leafminers when locally established thresholds have been reached. Do not apply during bloom. White Apple Leafhopper (Not for use west of the Rocky Mountains): Apply Timectin 0.15 EC Ag Insecticide/Miticide only to the first generation of white apple leafhoppers, just after petal fall and in combination with horticultural spray oil (not a dormant oil).	Do not apply by air. Reduced efficacy and/or residual control may occur if Timectin 0.15 EC Ag Insecticide/Miticide is not applied in combination with VINTRE or a horticultural spray oil (not a dormant oil). Follow all directions on the VINTRE or spray oil label or in spray guides. Use caution when applying Timectin 0.15 EC Ag Insecticide/Miticide with VINTRE or a horticultural spray oil or injury may occur with certain apple varieties (such as russetting on light-skinned Golden Delicious apples). In some cases, phytotoxicity and crop loss has been observed when Timectin 0.15 EC Ag Insecticide/Miticide plus VINTRE or a horticultural spray oil was applied less than 14 days before or after a captan application. Do not apply more than 20 fl. oz. Timectin 0.15 EC Ag Insecticide/Miticide per application, and no more than 40 fl. oz. per growing season. Make only 2 applications per growing season 21 days or more apart. Do not allow livestock to graze in treated orchards. Preharvest Interval: 28 days

^a Dilute Spray Rate is based on a volume of 400 gals./A dilute spray.

^b Concentrate Spray Rate: determine the amount of Timectin 0.15 EC Ag Insecticide/Miticide required in a full cover dilute spray. Then use the same amount of product per acre in concentrate sprays that would be required for the dilute sprays to the same orchard/grove. Note that for small trees, the amount may be less than 10 fl. oz./A.

AVOCADOS

Timectin 0.15 EC Ag Insecticide/Miticide will control the following pests in avocados:

Avocado thrips (Scirtothrips perseae)

Persea Mite (Oligonychus perseae)

Rate of Timectin 0.15 EC Ag Insecticide/Miticide	Directions for Application	Restrictions
10 – 20 fl. oz./A (Do NOT use less than 10 fl. oz. of Timectin 0.15 EC Ag Insecticide/Miticide/A) And add (for best results): a Horticultural Spray Oil that is not a dormant oil such as VINTRE. Horticultural Spray Oil: Minimum of 1 to 4% v/v (Follow all directions on the spray oil label. Test small areas with oil concentrations greater than 2.0% before making applications to large areas.) VINTRE Use 16-32 fl. oz. per acre at rates no greater than 0.3% v/v.	Apply Timectin 0.15 EC Ag Insecticide/Miticide using conventional dilute or concentrate sprays using ground or air application equipment. Although aerial application is permitted, best control of thrips is achieved with ground applications which provide better spray coverage. The user accepts all liability for thrip level and duration of control if Timectin 0.15 EC Ag Insecticide/Miticide is applied by air. Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage. The size and number of trees per acre as well as foliage density will determine the gallons of spray required. Apply only in 100 gals. or more of water/A by ground and 50 gals. of water by air application. [For ground applications where the spray volume is greater than 400 gals./A, use 2.5 fl. oz. Timectin 0.15 EC Ag Insecticide/Miticide per 100 gals. of water in combination with VINTRE or 1.0-4.0% of an NR (narrow range) 415 oil.] Control of these pests requires a thorough coverage of the spray solution on the tree foliage. Time Timectin 0.15 EC Ag Insecticide/Miticide applications to occur when immature thrips first appear but before 5 or more immature thrips per leaf/fruit are detected. Repeat applications following the directions in the Restrictions column of this table. Apply Timectin 0.15 EC Ag Insecticide/Miticide at 10-15 fl. oz/A for low (1-2 thrips per leaf/fruit) to moderate (3-4 thrips per leaf/fruit). Use the higher rate of 15-20 fl. oz./A for severe (more than 5 thrips per leaf/fruit) outbreaks. Apply Timectin 0.15 EC Ag Insecticide/Miticide in combination with VINTRE or a 1.0-4.0% NR 415 oil in combination with a Horticultural Spray Oil (not a dormant oil) which is approved for use on avocados.	Do not apply more than 20 fl. oz. Timectin 0.15 EC Ag Insecticide/Miticide per application, and no more than 40 fl. oz. per growing season. Make only 2 applications per growing season 30 days or more apart. Preharvest Interval: 14 days

CELERIAC (*Apium graveollens*)

Timectin 0.15 EC Ag Insecticide/Miticide will control the following pests in celeriac:

Twospotted spider mite

Rate of Timectin 0.15 EC Ag Insecticide/Miticide	Directions for Application	Restrictions
16.0 fl. oz./A	Apply Timectin 0.15 EC Ag Insecticide/Miticide using conventional dilute or concentrate sprays using ground application equipment. Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage of the crop foliage. Apply only in 20 gals. or more of water/A by ground application. Control of this mite requires a thorough coverage of the spray solution on the crop foliage. Time Timectin 0.15 EC Ag Insecticide/Miticide applications to occur when mites first appear. Repeat applications following the directions in the Restrictions column of this table. Optimum results are obtained if Timectin 0.15 EC Ag Insecticide/Miticide is used in combination with a non-ionic surfactant such as WETCIT.	Refer to the section on Spray Drift Precautions of this label. Do not apply by air. Do not apply more than 16 fl. oz Timectin 0.15 EC Ag Insecticide/Miticide per application, and no more than 48 fl. oz. after transplanting. To prevent resistance, make only 2 sequential applications per growing season 7 days or more apart. Preharvest Interval: 7 days

CITRUS FRUITS (CROP GROUP 10)

Timectin 0.15 EC Ag Insecticide/Miticide will control the following pests in citrus: calamondin, citrus citron, citrus hybrids (including chironja, tangelo, tangor), grapefruit, kumquat, lemon, lime, mandarin (tangerine), sour orange, sweet orange, pummelo, Satsuma mandarin:

Asian citrus psylla
Broad mite
Citrus bud mite
Citrus leafminer
Citrus rust mite
Citrus thrips
Twospotted spider mite

Rate of Timectin 0.15 EC Ag Insecticide/Miticide	Directions for Application	Restrictions
For Dilute Sprays^a: 1.0 – 2.0 fl. oz./100 gals. for all insects (except Asian citrus psylla, citrus leafminers and citrus thrips—use the concentrate spray rate) For Concentrate Sprays^{b, c}: 10 – 20 fl. oz./A for all listed insects/mites And add (for best results) a Horticultural Spray Oil that is not a dormant oil such as OROCIT, VINTRE or CITRI-KING. Horticultural Spray Oil 0.20% or 1 gal./A OROCIT or VINTRE or CITRI-KING Use 16-64 fl. oz. per acre in sufficient water at rates no greater than 0.3% v/v. (see footnotes for additional rates)^c	Apply Timectin 0.15 EC Ag Insecticide/Miticide using conventional dilute or concentrate sprays using ground application equipment. Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage (outside coverage for Asian citrus psylla, citrus leafminer or citrus thrips). The size and number of trees per acre as well as foliage density will determine the gallons of spray required. Asian citrus psylla: Applications made in the spring, summer or fall should be directed at newly expanding foliage and will only control adults and nymphs present at the time of application. Mites: Applications made in the spring, summer or fall should be made when mites first appear. Citrus bud mite: Time application to occur at “bud swell”. Citrus leafminer^d: Applications in the spring, summer or fall should be directed at newly expanding foliage. Citrus thrips: Correct timing is required. Apply after egg hatch has begun (typically, early to mid-hatch is best) but only the current generation will be controlled.	Aerial application is permitted only for control of citrus leafminer. Aerial application to citrus is not approved in California. Do not apply more than 40 fl. oz. Timectin 0.15 EC Ag Insecticide/Miticide per growing season. Do not apply more than 3 applications per growing season 30 days or more apart. To prevent pest resistance to Timectin 0.15 EC Ag Insecticide/Miticide, do not apply in citrus nurseries. Do not allow livestock to graze in treated orchards. Preharvest Interval: 7 days

^a Dilute Spray Rate is based on a volume of 1,000 gals./A dilute spray.

^b Concentrate Spray Rate: determine the amount of Timectin 0.15 EC Ag Insecticide/Miticide that would be required in a full cover dilute spray.

^c For optimum control of **Asian citrus psylla, broad mites, citrus rust mites and citrus leafminers**, apply 150-300 gals spray mix per acre plus OROCIT, VINTRE or CITRI-KING or a minimum of 3 gal./A of a horticultural oil (not a dormant oil) and apply using ground equipment at 1 to 1.5 miles per hour. For optimum control of **citrus bud mite**, apply Timectin 0.15 EC Ag Insecticide/Miticide in a minimum of 500 gals. per acre plus OROCIT, VINTRE or CITRI-KING or a minimum of 2 qts. (0.5%) per acre of a horticultural oil (not a dormant oil). For optimum control of **citrus thrips**, use 100-250 gals. per acre spray mix for best coverage of outside foliage.

^d **Citrus leafminer** – Ground or aerial application is permitted; however, do not apply with aircraft to citrus in California. For aerial application use a minimum of 10 gallons of finished spray volume per acre. Under conditions such as high pest population, dense foliage, or adverse application conditions (such as high temperatures) use a greater volume of water to ensure adequate coverage.

COTTON

Timectin 0.15 EC Ag Insecticide/Miticide will control the following pests in cotton:

Carmine spider mites

Pacific spider mites

Strawberry spider mites

Twospotted spider mites

Rate of Timectin 0.15 EC Ag Insecticide/Miticide	Directions for Application	Restrictions
Early season: 4 – 6 fl. oz./A Mid-season/Lay-by: 8 – 16 fl. oz./A	Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage. Control of mites requires a thorough coverage of the spray solution on the crop foliage. Early season: Apply Timectin 0.15 EC Ag Insecticide/Miticide when the cotton is no taller than 10 inches. Apply only using ground application equipment. Do not apply less than 4 fl. oz. Timectin 0.15 EC Ag Insecticide/Miticide per acre. Mid-season and lay-by: Use Timectin 0.15 EC Ag Insecticide/Miticide at the specified rates 8 - 16 fl. oz./A) depending on the size of the plants and foliage density. Apply using ground application or by air in fixed-wing aircraft or by helicopter. Apply by air in a minimum of 5 gals. of water /A. Time Timectin 0.15 EC Ag Insecticide/Miticide applications to occur when mites first appear. Repeat applications following the directions in the Restrictions column of this table.	Refer to the section on Spray Drift Precautions of this label. Do not apply through any type of irrigation system. Do not apply more than 32 fl. oz. Timectin 0.15 EC Ag Insecticide/Miticide per growing season. Make only 2 applications per growing season 21 days or more apart. Do not feed foliage to livestock; do not allow livestock to graze treated cotton. Preharvest Interval: 20 days

CUCURBIT VEGETABLES (CROP GROUP 9)

Timectin 0.15 EC Ag Insecticide/Miticide will control the following pests in the Cucurbit Crop Group chayote (fruit), Chinese waxgourd (Chinese preserving melon), citron melon, cucumber, gherkin, edible gourds (includes hyotan, cucuzza, hechima, Chinese okra), *Momordica* spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber), muskmelon (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon and snake melon), pumpkin, summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini), winter squash (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash), watermelon:

Spider mites

Leafminers

Rate of Timectin 0.15 EC Ag Insecticide/Miticide	Directions for Application	Restrictions
8 – 16 fl. oz./A	Apply Timectin 0.15 EC Ag Insecticide/Miticide using ground or air application equipment. Although aerial application is permitted, best control of insect pests is achieved with ground applications which provide better spray coverage. The user accepts all liability for mite level and duration of control if Timectin 0.15 EC Ag Insecticide/Miticide is applied by air. Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage. Control of these mites/insects requires a thorough coverage of the spray solution on the crop foliage. Do not apply in less than 20 gals. of water/A by ground and in less than 5 gals. of water by air application. To ensure thorough spray coverage, use a greater spray volume for high pest pressure, dense crop canopies, and adverse application conditions (high temperatures). Time Timectin 0.15 EC Ag Insecticide/Miticide applications to occur when insects first appear. Use 8 – 12 fl. oz./A for low to moderate insect pressure use 16 fl. oz./A for heavy insect infestations. Repeat applications following the directions in the Restrictions column of this table.	Refer to the section on Spray Drift Precautions of this label. Do not apply Timectin 0.15 EC Ag Insecticide/Miticide by air in the State of New York. Do not apply more than 48 fl. oz Timectin 0.15 EC Ag Insecticide/Miticide per growing season. Make only 2 sequential applications 7 days or more apart. Preharvest Interval: 7 days

DRY BEANS

Timectin 0.15 EC Ag Insecticide/Miticide will control the following pests in: *Cicer arietinum* (chickpea, garbanzo bean); *Lupinus* spp. (including sweet lupine, white sweet lupine, white lupine, and grain lupine); *Phaseolus* spp. (including kidney bean, lima bean, mung bean, navy bean, pinto bean, snap bean, and waxbean); *Vicia faba* (broad bean, fava bean); *Vigna* spp. (including adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea*, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, and yardlong bean):

Liriomyza leafminers

Spider mites

*For use on cowpeas that are grown only for dry seed. Do not allow livestock to graze cowpea forage and do not harvest cowpea forage or hay for use as livestock feed.

Rate of Timectin 0.15 EC Ag Insecticide/Miticide	Directions for Application	Restrictions
Low to moderate infestations: 8.0 – 12.0 fl. oz./A High infestations: 13 – 16 fl. oz./A	Apply with ground application equipment or aircraft. Thorough coverage of the crop canopy is essential for optimum results. Inadequate coverage can result in reduced control. For best control, apply with ground application equipment. With aerial application, the resulting control of leafminers and spider mites could be less than with ground application. Apply when adult leafminer flies or spider mites are first observed and repeat application, as needed, to maintain control with constraints of a sound resistance management program. (see Restrictions)	Do not apply Timectin 0.15 EC Ag Insecticide/Miticide by air in the State of New York. Do not apply more than 16 fl. oz. (or 0.019 lb. ai./A) of Timectin 0.15 EC Ag Insecticide/Miticide, or any other foliarly applied abamectin containing product, per application. Do not apply more than 48 fl. oz. (or 0.056 lb. ai./A) of Timectin 0.15 EC Ag Insecticide/Miticide, or any other foliarly applied abamectin containing product, per growing season. Make a maximum of 2 applications of Timectin 0.15 EC Ag Insecticide/Miticide, or any other foliarly applied abamectin containing product per growing season. Make repeat applications at least 6 days apart. Preharvest Interval: 7 days Do not apply in less than 10 gals. of water/A with ground application equipment. Do not apply in less than 5 gal. of water/A with aircraft. Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to insure adequate coverage.

FRUITING VEGETABLES (Except Cucurbits) (CROP GROUP 8)

Timectin 0.15 EC Ag Insecticide/Miticide will control the following pests in the Fruiting Vegetable Crop Group: eggplant, groundcherry, pepino, pepper (includes bell pepper, chili pepper, cooking pepper, pimiento, sweet pepper), tomatillo, tomato:

Broad mite
Colorado potato beetle
Liriomyza leafminers
Spider mites
Thrips palmi
Tomato pinworm
Tomato russet mite
Potato psyllid (peppers only; provides suppression only)

Rate of Timectin 0.15 EC Ag Insecticide/Miticide	Directions for Application	Restrictions
8 – 16 fl. oz./A (use the 16 fl. oz./A rate for tomato pinworm) Note: Although Timectin 0.15 EC Ag Insecticide/Miticide can be used without a wetting agent, improved spreading and penetration for better insect control is achieved using a combination of Timectin 0.15 EC Ag Insecticide/Miticide and a surfactant such as WETCIT. Nonionic type surfactants are recommended (not binder or sticker–type surfactants).	Apply Timectin 0.15 EC Ag Insecticide/Miticide using ground or air application equipment. Although aerial application is permitted, best control of mites is achieved with ground applications which provide better spray coverage. The user accepts all liability for mite level and duration of control if Timectin 0.15 EC Ag Insecticide/Miticide is applied by air. Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage. Control of these mites/insects requires a thorough coverage of the spray solution on the crop foliage. Do not apply in less than 20 gals. of water/A by ground and in less than 5 gals. of water by air application. To ensure thorough spray coverage, use a greater spray volume for high pest pressure, dense crop canopies, and adverse application conditions (high temperatures). Repeat applications following the directions in the Restrictions column of this table. Mites and thrips : Apply when insects first appear using 8 fl. oz. for low to moderate insect populations and 16 fl. oz./A for heavy infestations. Tomato pinworm: Make application when moth activity is first observed and not later than when larvae emerge. Apply using the 16 fl. oz./A rate. Colorado potato beetle: Apply 8 fl. oz. for low to moderate insect populations and 16 fl. oz./A for heavy infestations. Potato psyllid (Peppers only): Timectin 0.15 EC Ag Insecticide/Miticide provides suppression only. Apply when insects first appear. Use 8 – 12 fl. oz. for low to moderate insect populations and 16 fl. oz./A for heavy infestations. Make sequential applications at 10-14 day intervals when heavy egg and nymph populations are found. Use of a spreading or penetration adjuvant such as WETCIT or horticultural oil is recommended.	Refer to the section on Spray Drift Precautions of this label. Do not apply Timectin 0.15 EC Ag Insecticide/Miticide by air in the State of New York. Do not apply more than 48 fl. oz Timectin 0.15 EC Ag Insecticide/Miticide per growing season. Make only 2 sequential applications 7 days or more apart. To prevent insect resistance to Timectin 0.15 EC Ag Insecticide/Miticide, do not apply to fruiting vegetables grown for transplants. Preharvest Interval: 7 days

GRAPES

Timectin 0.15 EC Ag Insecticide/Miticide will control the following pests in grapes:

Pacific spider mite
Twospotted spider mite
Western grapeleaf skeletonizer
Willamette spider mite.
*Western grape leafhopper**
*Varigated leafhopper**

* Timectin 0.15 EC Ag Insecticide/Miticide provides knock-down control only of these pests.

Rate of Timectin 0.15 EC Ag Insecticide/Miticide	Directions for Application	Restrictions
8 – 16 fl. oz./A PLUS VINTRE (Use 16-32 fl. oz. per acre at rates no greater than 0.3% v/v) OR a nonionic type surfactant (not binder or sticker–type surfactants).	Apply Timectin 0.15 EC Ag Insecticide/Miticide using ground application equipment. Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage. Control of these mites/insects requires a thorough coverage of the spray solution on the crop foliage. Do not spray alternate rows: apply Timectin 0.15 EC Ag Insecticide/Miticide on both sides of the row to ensure maximum coverage of the crop. Do not apply in less than 50 gals. of water/A by ground. If electro-static sprayers are used, the volume may be less than 50 gals./A but do not use less than 5 gals. of water/A. Always use Timectin 0.15 EC Ag Insecticide/Miticide with VINTRE or a nonionic surfactant. This combination improves the wetting of leaves and provides an even coverage of spray. Not all combinations of VINTRE or nonionic surfactants and Timectin 0.15 EC Ag Insecticide/Miticide have been tested on every grape variety or all types of growing conditions. Before using VINTRE or a nonionic surfactant, carefully read and follow the surfactant product's label. Apply when insects first appear using 8 - 12 fl. oz. for low to moderate insect populations and 16 fl. oz./A for heavy infestations. Repeat applications following the directions in the Restrictions column of this table. Mites : time application to occur when mites are first detected but before motiles exceed 5 per leaf. Western grapeleaf skeletonizer: Make application when larvae appear. Best control is achieved if application is made just after egg hatch. Western grape leafhopper, Varigated leafhopper: For knock-down control only, apply when pests first appear.	Refer to the section on Spray Drift Precautions of this label. Do not apply by air. Do not apply more than 16 fl. oz. per application or more than 32 fl. oz Timectin 0.15 EC Ag Insecticide/Miticide per growing season. Make only 2 applications per growing season. One repeat application may be applied 21 days or more after the first application if needed to maintain control. Do not allow livestock to graze in treated vineyards. Preharvest Interval: 28 days

HERB CROP SUBGROUP (EXCEPT CHIVES) (CROP SUBGROUP 19A)

Timectin 0.15 EC Ag Insecticide/Miticide will control the following pests in the Herb Crop Subgroup (except chives): angelica, balm, basil, borage, burnet, chamomile, catnip, chervil (dried), chives, chives (Chinese), clary, coriander (leaf), costmary, cilantro (leaf), curry (leaf), dillweed, horehound, hyssop, lavender, lemongrass, lovage (leaf), marigold, marjoram (*Origanum* spp.), nasturtium, parsley (dried), pennyroyal, rosemary, rue, sage, savory (summer and winter), sweet bay, tansy, tarragon, thyme, wintergreen, woodruff, wormwood:

Liriomyza leafminers

Rate of Timectin 0.15 EC Ag Insecticide/Miticide	Directions for Application	Restrictions
8 – 16 fl. oz./A Note: To improve the wetting of leaves and provide an even coverage of spray of the foliage, use Timectin 0.15 EC Ag Insecticide/Miticide in combination with a wetting agent, such as WETCIT or a nonionic surfactant.	Apply Timectin 0.15 EC Ag Insecticide/Miticide using conventional dilute or concentrate ground application equipment. Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage. Control of leafminers requires a thorough coverage of the spray solution on the upper and lower leaves. Do not apply in less than 30 gals. of water/A by ground. Use the low rate for low to moderate insect pressure and the higher rate for heavy insect pressure. Time application to occur when adult flies are first detected. Repeat applications following the directions in the Restrictions column of this table. Not all combinations of WETCIT or nonionic surfactants and Timectin 0.15 EC Ag Insecticide/Miticide have been tested on every herb type and variety or under all types of growing conditions. Before using WETCIT or a nonionic surfactant, carefully read and follow the surfactant product's label.	Do not apply by air. Make only 2 applications per single cutting. Do not apply more than 6 applications per cropping season at the 8 fl. oz./A rate or more than 3 applications at the 16 fl. oz./A rate. Repeat applications may be made 7 days or more after the last application. Preharvest Interval: 14 days

HOPS (EXCEPT IN CALIFORNIA)

Timectin 0.15 EC Ag Insecticide/Miticide will control the following pests in hops (not approved for use in California):

Twospotted spider mite

Rate of Timectin 0.15 EC Ag Insecticide/Miticide	Directions for Application	Restrictions
8 – 16 fl. oz./A Note: Although Timectin 0.15 EC Ag Insecticide/Miticide can be used without a wetting agent, improved spreading and penetration for better insect control is achieved using a combination of Timectin 0.15 EC Ag Insecticide/Miticide and a surfactant such as WETCIT. Nonionic type surfactants are recommended (not binder or sticker–type surfactants).	Apply Timectin 0.15 EC Ag Insecticide/Miticide using ground application equipment. Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage. Control of mites requires a thorough coverage of the spray solution on the upper and lower leaves of the crop. The size and foliage density of plants will determine the gallons of spray required for thorough coverage. Use the same amount of Timectin 0.15 EC Ag Insecticide/Miticide for concentrate or dilute sprays. For plants 6 – 8 feet in height (1/2 trellis growth), apply 8 – 16 fl. oz./A of Timectin 0.15 EC Ag Insecticide/Miticide in a minimum of 40 gals. of water per acre. For plants taller than 6-8 feet, do not use less than 16 fl. oz./A in a minimum of 100 gals. of water/A. Time application to occur when mites are first detected. One repeat application may be made following the directions in the Restrictions column of this table.	Refer to the section on Spray Drift Precautions of this label. Do not apply by air. Make only 2 applications per season. Do not apply more than 32 fl. oz./A per growing season. Repeat applications may be made 21 days or more after the last application. To prevent insect resistance, use other miticides with different modes of action before making a repeat application of Timectin 0.15 EC Ag Insecticide/Miticide. Do not allow livestock to graze in treated hop yards. Preharvest Interval: 28 days

LEAFY VEGETABLES (Except BRASSICA Vegetables) (CROP GROUP 4)

Timectin 0.15 EC Ag Insecticide/Miticide will control the following pests in the Leafy Vegetables (except brassica vegetables) Crop Group: amaranth (leafy amaranth, Chinese spinach, tampala), arugula (Roquette), cardoon, celery, Chinese celery, celtuce, chervil, Chrysanthemum (edible-leaved), Chrysanthemum (garland), corn salad, garden cress, upland cress (yellow rocket, winter cress), dandelion, dock (sorrel), endive (escarole), Florence fennel (finocchio), head and leaf lettuce, orach, parsley, garden purslane, winter purslane, radicchio (red chicory), rhubarb, spinach, New Zealand spinach, vine spinach (Malabar spinach, Indian spinach), Swiss chard:

Carmine spider mite
Liriomyza leafminers
Twospotted spider mite

Rate of Timectin 0.15 EC Ag Insecticide/Miticide	Directions for Application	Restrictions
8 – 16 fl. oz./A Note: Although Timectin 0.15 EC Ag Insecticide/Miticide can be used without a wetting agent, improved spreading and penetration for better insect control is achieved using a combination of Timectin 0.15 EC Ag Insecticide/Miticide and a surfactant such as WETCIT. Nonionic type surfactants are recommended (not binder or sticker–type surfactants).	Apply Timectin 0.15 EC Ag Insecticide/Miticide using ground or air application equipment. Although aerial application is permitted, best control of mites is achieved with ground applications which provide better spray coverage. The user accepts all liability for mite level and duration of control if Timectin 0.15 EC Ag Insecticide/Miticide is applied by air. Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage. Control of these mites/insects requires a thorough coverage of the spray solution on the crop foliage. Do not apply in less than 20 gals. of water/A by ground and in less than 5 gals. of water by air application. To ensure thorough spray coverage, use a greater spray volume for high pest pressure, dense crop canopies, and adverse application conditions (high temperatures). Apply Timectin 0.15 EC Ag Insecticide/Miticide at 8 – 12 fl. oz./A for low to moderate insect populations and at 16 fl. oz./A for heavy infestations. Make repeat applications if needed to maintain control according to the Restrictions column of this table. Mites: Apply when insects first appear. Leafminers: Time application to occur when adult flies are first detected.	Refer to the section on Spray Drift Precautions of this label. Do not apply Timectin 0.15 EC Ag Insecticide/Miticide by air in the State of New York. Do not apply more than 48 fl. oz. Timectin 0.15 EC Ag Insecticide/Miticide per growing season. Make only 2 sequential applications 7 days or more apart. To prevent insect resistance to Timectin 0.15 EC Ag Insecticide/Miticide, do not apply to leafy vegetables grown for transplants. Preharvest Interval: 7 days

MINT (PEPPERMINT AND SPEARMINT)

Timectin 0.15 EC Ag Insecticide/Miticide will control the following pests in peppermint and spearmint:

Twospotted spider mite

Rate of Timectin 0.15 EC Ag Insecticide/Miticide	Directions for Application	Restrictions
8 – 12 fl. oz./A Note: Although Timectin 0.15 EC Ag Insecticide/Miticide can be used without a wetting agent, improved spreading and penetration for better insect control is achieved using a combination of Timectin 0.15 EC Ag Insecticide/Miticide and either a nonionic surfactant such as WETCIT or organosilicone-based surfactant at the manufacturer's recommended use rates.	Apply Timectin 0.15 EC Ag Insecticide/Miticide using ground or air application equipment. Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage. Control of these mites/insects requires a thorough coverage of the spray solution on the crop foliage. Do not apply in less than 20 gals. of water/A by ground and in less than 5 gals. of water by air application. To ensure thorough spray coverage, use a greater spray volume for high pest pressure, dense crop canopies, and adverse application conditions (high temperatures). Time applications to occur when insects first appear. Make repeat applications if needed to maintain control according to the Restrictions column of this table.	Do not apply Timectin 0.15 EC Ag Insecticide/Miticide by air in the State of New York. Do not apply more than 12 fl. oz. Timectin 0.15 EC Ag Insecticide/Miticide per application or more than 36 fl. oz. per growing season. Make only 2 sequential applications 7 days or more apart. Do not apply more than 3 applications per season. Do not allow livestock to graze or feed treated foliage to livestock. Preharvest Interval: 28 days

ONION, BULB (CROP SUBGROUP 3-07A)

Timectin 0.15 EC Ag Insecticide/Miticide will control the following pests in the Onion, Bulb Subgroup: onion, bulb including daylily, bulb; fritillaria, bulb; garlic, bulb; garlic, great-headed, bulb; garlic, serpent, bulb; lily, bulb; onion, Chinene, bulb; onion, peal; onion, potato, bulb; shallot, bulb; cultivars, varieties, and/or hybrids of these:

Liriomyza leafminers

Thrips

Rate of Timectin 0.15 EC Ag Insecticide/Miticide	Directions for Application	Restrictions
Low to moderate infestations: 8.0 – 12.0 fl. oz./A High infestations: 13 – 16 fl. oz./A	Apply with ground application equipment or aircraft. Thorough coverage of the crop canopy is essential for optimum results. Inadequate coverage can result in reduced control. For best control, apply with ground application equipment. With aerial application, the resulting control of leafminers and thrips could be less than with ground application. Leafminer: Apply when adult leafminer flies are first observed and repeat application, as needed, to maintain control within constraints of a sound resistance management program. (see Restrictions) Thrips: Apply in a thrips management program and when thrips are at economic threshold. Repeat application, as needed, to maintain control within constraints of a sound resistance management program (see Restrictions). Do not use Timectin 0.15 EC Ag Insecticide/Miticide as a rescue treatment for thrips control. Note: Add a non-ionic activator type wetting, spreading and/or penetrating adjuvant approved for use on bulb onion. Do not use binder sticker type adjuvants. Tank mixing with products that have binder sticker type formulations can reduce Timectin 0.15 EC Ag Insecticide/Miticide performance. (see Restrictions)	Do not apply Timectin 0.15 EC Ag Insecticide/Miticide by air in the State of New York. Do not apply more than 16 fl. oz. (or 0.019 lb. ai./A) of Timectin 0.15 EC Ag Insecticide/Miticide, or any other foliarly applied abamectin containing product, per application. Do not apply more than 48 fl. oz. (or 0.056 lb. ai./A) of Timectin 0.15 EC Ag Insecticide/Miticide, or any other foliarly applied abamectin containing product, per growing season. Wait at least 7 days before repeating application. Make 2 consecutive applications of Timectin 0.15 EC Ag Insecticide/Miticide then rotate to a chemistry with a different mode of action. Make at least 2 applications of a chemistry with a different mode of action before making additional Timectin 0.15 EC Ag Insecticide/Miticide applications. Do not make more than 2 sequential applications of Timectin 0.15 EC Ag Insecticide/Miticide or any other foliarly applied abamectin containing product. Preharvest Interval: 30 days Do not apply in less than 20 gals. of water/A with ground application equipment. Do not apply in less than 5 gal. of water/A with aircraft. Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to insure adequate coverage.

PEARS (INCLUDING ORIENTAL PEARS)

Timectin 0.15 EC Ag Insecticide/Miticide will control the following pests in pears and Oriental pears:

European red mite
McDaniel spider mite
Pear psylla
Pear rust mite
Twospotted spider mite
Yellow mite

Rate of Timectin 0.15 EC Ag Insecticide/Miticide	Directions for Application	Restrictions
For Dilute Sprays^a: 2.5 – 5.0 fl. oz./100 gals For Concentrate Sprays^b: 10 – 20 fl. oz./A And add (for best results) A Horticultural Spray Oil that is not a dormant oil For Dilute Sprays: 0.25% or 1 gal./A For Concentrate Sprays: Minimum of 1 gal./A	Apply Timectin 0.15 EC Ag Insecticide/Miticide using conventional dilute or concentrate sprays using ground application equipment. Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage. The size and number of trees per acre as well as foliage density will determine the gallons of spray required. Apply only in 40 gals. or more of water/A. Control of these mites/insects requires a thorough coverage of the spray solution on the tree foliage. Time Timectin 0.15 EC Ag Insecticide/Miticide applications to occur when spider mites first appear or when insect thresholds are reached. Repeat applications following the directions in the Restrictions column of this table.	Do not apply by air. Reduced efficacy and/or residual control may occur if Timectin 0.15 EC Ag Insecticide/Miticide is not applied with a horticultural spray oil (not a dormant oil). Follow all directions on the spray oil label or spray guides. In some cases, phytotoxicity and crop loss has been observed when Timectin 0.15 EC Ag Insecticide/Miticide plus a horticultural spray oil was applied in less than 14 days before or after a captan application. Do not apply more than 20 fl. oz. Timectin 0.15 EC Ag Insecticide/Miticide per application, and no more than 40 fl. oz. per growing season. Make only 2 applications per growing season 21 days or more apart. Do not allow livestock to graze in treated orchards. Preharvest Interval: 28 days
^a Dilute Spray Rate is based on a volume of 400 gals./A dilute spray. ^b Concentrate Spray Rate: determine the amount of Timectin 0.15 EC Ag Insecticide/Miticide required in a full cover dilute spray. Then use the same amount of product per acre in concentrate sprays that would be required for the dilute sprays to the same orchard/grove. Note that for small trees, the amount may be less than 10 fl. oz./A.		

STONE FRUIT CROP GROUP (CROP GROUP 12)

Timectin 0.15 EC Ag Insecticide/Miticide will control the following pests in the Stone Fruit Crop Group: apricot; cherry (sweet and tart), nectarine, peach, plum, plum (chickasaw, damson, and Japanese), plumcot, and prune (fresh):

European spider mite

Pacific spider mite

Twospotted spider mite

Rate of Timectin 0.15 EC Ag Insecticide/Miticide	Directions for Application	Restrictions
For Dilute Sprays^a: 2.5 – 5.0 fl. oz./100 gals For Concentrate Sprays^b: 10 – 20 fl. oz./A Horticultural Spray Oil: For Dilute Sprays: 0.25% or 1 gal./A For Concentrate Sprays: Minimum of 1 gal./A Nonionic Surfactant Labeled rate	Apply when mites first appear. A second application may be made, if needed to maintain control. (see Restrictions) Always apply Timectin 0.15 EC Ag Insecticide/Miticide in combination with a nonionic surfactant (at the manufacturer's specified label rate) that spreads and/or penetrates the leaf cuticle, or apply with horticultural Spray oil (not a dormant oil). The combination of Timectin 0.15 EC Ag Insecticide/Miticide and the nonionic surfactant helps to provide an even coverage of spray on the foliage and to improve the wetting and penetration of leaves for better insect control. A nonionic surfactant that spreads on and or/penetrates the leaf cuticle can improve insect control. Although Timectin 0.15 EC Ag Insecticide/Miticide has been tested in combination with nonionic surfactants for safety to Stone Fruit, it is impossible to test on all Stone Fruit varieties under the variety of conditions that may cause crop injury. Therefore, when using Timectin 0.15 EC Ag Insecticide/Miticide in combination with a nonionic surfactant, carefully follow the Directions for Use and Precautions on the surfactant label and in official spray guides. A horticultural oil (not a dormant oil) may improve efficacy and can be used but it may increase the possibility of phytotoxicity to foliage and fruit. Use nonionic surfactants at the manufacturers recommended rate. Use a minimum of 0.25% horticultural spray oil (not a dormant oil) in the dilute spray mixture and not less than 1.0 gal. of horticultural spray oil per acre in the final finished spray. Apply Timectin 0.15 EC Ag Insecticide/Miticide using conventional dilute or concentrate sprays using ground application equipment. Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage. The size and number of trees per acre as well as foliage density will determine the gallons of spray required. Apply only in 40 gals. or more of water/A. Control of these mites/insects requires a thorough coverage of the spray solution on the tree foliage.	Do not apply by air. Do not apply more than 20 fl. oz. (or 0.023 lb. ai./A) of Timectin 0.15 EC Ag Insecticide/Miticide, or any other foliarly applied abamectin containing product, per application. Do not apply more than 40 fl. oz. (or 0.047 lb. ai./A) of Timectin 0.15 EC Ag Insecticide/Miticide, or any other foliarly applied abamectin containing product, per growing season. Make a maximum of 2 applications of Timectin 0.15 EC Ag Insecticide/Miticide, or any other foliarly applied abamectin containing product per growing season 21 days or more apart. Do not allow livestock to graze in treated orchards. Preharvest Interval: 21 days Do not apply in less than 40 gals. of water/A.

^a Dilute Spray Rate is based on a volume of 400 gals./A dilute spray.

^b Concentrate Spray Rate: determine the amount of Timectin 0.15 EC Ag Insecticide/Miticide required in a full cover dilute spray. Then use the same amount of product per acre in concentrate sprays that would be required for the dilute sprays to the same orchard/grove. Note that for small trees, the amount may be less than 10 fl. oz./A.

STRAWBERRIES

Timectin 0.15 EC Ag Insecticide/Miticide will control the following pests in strawberries:

Carmine mites
Cyclamen mites
Aphids
Thrips
Whiteflies
Strawberry spider mite
Twospotted spider mite

Rate of Timectin 0.15 EC Ag Insecticide/Miticide	Directions for Application	Restrictions
16 fl. oz./A Note: Although Timectin 0.15 EC Ag Insecticide/Miticide can be used without a wetting agent, improved spreading and penetration for better insect control is achieved using a combination of Timectin 0.15 EC Ag Insecticide/Miticide and a nonionic surfactant such as WETCIT at the manufacturer's recommended use rate.	Apply Timectin 0.15 EC Ag Insecticide/Miticide using ground application equipment. Calibrate equipment prior to application to ensure sufficient water is used for thorough coverage. Control of these mites requires a thorough coverage of the spray solution on the crop foliage. Apply Timectin 0.15 EC Ag Insecticide/Miticide to the tops and undersides of the leaves. Do not apply in less than 50 gals. of water/A by ground. If electro-static sprayers are used, the volume may be less than 50 gals./A but do not use less than 10 gals. of water/A. Apply Timectin 0.15 EC Ag Insecticide/Miticide when mites are first detected. A second repeat application may be made 7 – 10 days later. This sequence of two applications may be repeated if needed but follow the directions in the Restrictions column of this table.	Refer to the section on Spray Drift Precautions of this label. Do not apply by air. Do not apply more than 16 fl. oz. per application or more than 64 fl. oz Timectin 0.15 EC Ag Insecticide/Miticide per growing season. After the first 2 applications, wait 21 days or more before making additional applications if needed to maintain control. To prevent insect resistance to Timectin 0.15 EC Ag Insecticide/Miticide, do not apply in strawberry nurseries. Preharvest Interval: 3 days

TREE NUTS CROP GROUP (CROP GROUP 14) and PISTACHIO

Timectin 0.15 EC Ag Insecticide/Miticide will control the following pests in the Tree Nut Crop Group: almond, beech nut, Brazil nut, butternut, cashew, chestnut, chinquapin, filbert (hazelnut), hickory nut, macadamia nut, pecan, and walnuts (black and English):

European red mite

Pacific spider mite

Strawberry spider mite

Twospotted spider mite

Rate of Timectin 0.15 EC Ag Insecticide/Miticide	Directions for Application	Restrictions
For Dilute Sprays^a: 2.5 – 5.0 fl. oz./100 gals For Concentrate Sprays^b: 10 – 20 fl. oz./A Horticultural Spray Oil that is not a dormant oil: For Dilute Sprays: 0.25% or 1 gal./A For Concentrate Sprays: Minimum of 1 gal./A	Apply when spider mites first appear. Residual spider mite control is greater from spray deposits on newer leaves compared to older leavers. Always apply Timectin 0.15 EC Ag Insecticide/Miticide in combination with a horticultural spray oil (not a dormant oil) approved for use on tree nuts. A second application may be made, if needed, to maintain control. (see Restrictions)	Do not apply by air. Do not apply more than 20 fl. oz. (or 0.023 lb. ai./A) of Timectin 0.15 EC Ag Insecticide/Miticide, or any other foliarly applied abamectin containing product, per application. Do not apply more than 40 fl. oz. (or 0.047 lb. ai./A) of Timectin 0.15 EC Ag Insecticide/Miticide, or any other foliarly applied abamectin containing product, per growing season. Make a maximum of 2 applications of Timectin 0.15 EC Ag Insecticide/Miticide, or any other foliarly applied abamectin containing product per growing season. If a second application is necessary, wait at least 21 days before repeating application. Do not allow livestock to graze in treated groves/orchards. Preharvest Interval: 21 days Do not apply in less than 40 gals. of water/A.
^a Dilute Spray Rate is based on a volume of 400 gals./A dilute spray. ^b Concentrate Spray Rate: First determine the amount of Timectin 0.15 EC Ag Insecticide/Miticide required in a full cover dilute spray. Then use the same amount of product per acre in concentrate sprays that would be required for the dilute sprays to the same orchard/grove. This may result in use of less than 10 fl. oz./A on small trees.		

TUBEROUS AND CORM VEGETABLES CROP SUBGROUP (CROP SUBGROUP 1C)

Timectin 0.15 EC Ag Insecticide/Miticide will control the following pests in the Tuberous and Corm Vegetables Crop Subgroup: arracha; arrowroot; artichoke, Chinese; artichoke, Jerusalem; canna, edible; cassava (bitter and sweet); chayote (root); chufa; dasheen; ginger; leren; potato; sweet potato; taniar; turmeric; yam bean; and yam, true:

Colorado potato beetle

Liriomyza leafminers

Potato psyllid

Spider mites

Rate of Timectin 0.15 EC Ag Insecticide/Miticide	Directions for Application	Restrictions
Low to moderate infestations: 8.0 – 12.0 fl. oz./A Severe infestations: 16 fl. oz./A	Thorough coverage of the crop canopy is essential for optimal results. Inadequate coverage may result in reduced control. Apply Timectin 0.15 EC Ag Insecticide/Miticide using ground or air application equipment. Although aerial application is permitted, best control of mites is achieved with ground applications which provide better spray coverage. The user accepts all liability for mite level and duration of control if Timectin 0.15 EC Ag Insecticide/Miticide is applied by air. The addition of a nonionic surfactant or organosilicone-based surfactant is suggested for optimum spider mite and insect control. Use nonionic surfactant or organosilicone-based surfactant at the manufacturers recommended rate. Use of Timectin 0.15 EC Ag Insecticide/Miticide in combination with a sticker or binder type product such as Bravo® Weather Stik® may reduce control of insects and spider mites. Spider mites: Time applications to occur when mites first appear. Repeat application as needed to maintain control. (see Restrictions) Colorado potato beetle: Apply Timectin 0.15 EC Ag Insecticide/Miticide when about 50% of the egg masses have hatched and the larvae are present. If 2 applications are needed, limit them to a single Colorado potato beetle generation per crop. Do not make more than 2 applications per crop. (see Restrictions) Liriomyza leafminers: Time application to occur when adult flies are first observed. Repeat applications as needed to maintain control. (see Restrictions)	Do not apply Timectin 0.15 EC Ag Insecticide/Miticide by air in the State of New York. Do not apply more than 16 fl. oz./A (or 0.019 lb. a.i./A) per application. Do not apply in less than 20 gal. of water/A with ground equipment. Do not apply in less than 5 gal. of water/A with aircraft. Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to insure adequate coverage. Do not apply more than 32 fl. oz. (or 0.038 lb. a.i./A) of Timectin 0.15 EC Ag Insecticide/Miticide, or any other foliarly applied abamectin containing product, per crop for control of spider mite or Colorado potato beetle. Do not apply more than 48 fl. oz. (or 0.056 lb. a.i./A) of Timectin 0.15 EC Ag Insecticide/Miticide, or any other foliarly applied abamectin containing product, per crop for control of leafminer. Make only 2 sequential applications. Applications of Timectin 0.15 EC Ag Insecticide/Miticide (or other abamectin containing products) must be 7 days or more apart. Do not allow livestock to graze or feed treated foliage to livestock. Preharvest Interval: 14 days
^a Dilute Spray Rate is based on a volume of 400 gals./A dilute spray. ^b Concentrate Spray Rate: First determine the amount of Timectin 0.15 EC Ag Insecticide/Miticide required in a full cover dilute spray. Then use the same amount of product per acre in concentrate sprays that would be required for the dilute sprays to the same orchard/grove. This may result in use of less than 10 fl. oz./A on small trees.		

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

Pesticide Storage

Store in a tightly closed container in a cool, dry place.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal

Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

LIMITED WARRANTY AND DISCLAIMER

Tide International USA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below.

Our directions for use of this product are based upon tests believed to be reliable. The use of this product being beyond control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such use or the results to be obtained if not used in accordance with printed directions and established safe practice. To the extent consistent with applicable law, buyer's exclusive remedy and manufacturer's or seller's exclusive liability for any and all claims, losses, damages or injuries resulting from the use or handling of this product, whether or not based in contract, negligence, strict liability in tort or otherwise shall be limited, at the manufacturer's option to replacement of, or the repayment of the purchase price for, the quantity of product with respect to which damages are claimed.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Tide International USA or the seller. All such risks shall be assumed by buyer.

Terms and Conditions of Use

If terms of the Limited Warranty & Disclaimer and Inherent Risks of Use are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer and Inherent Risks of Use.

Bravo® and Weather Stik® are registered trademarks of a Syngenta Group Company.

Timectin™ is a trademark of Tide International USA, Inc.

OROCIT®, VINTRE® and CITRI-KING® are registered trademarks of ORO AGRI Inc.

[EPA approval date]