



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

August 6, 2025

Maryanne Kellogg
Regulatory Consultant
Tide International, USA, Inc.
c/o Pyxis Regulatory Consulting Inc.
4110 136th St. Ct. NW
Gig Harbor, WA 98332

Subject: Label Amendment - Registration Review Mitigation for Hexazinone
Product Name: Tide Hexazinone 75 WDG
EPA Registration Number: 84229-32
Application Date: April 29, 2019
Case Number: NA

Dear Maryanne Kellogg:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Hexazinone Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Caleb Carr by phone at 202-566-0636, or via email at carr.caleb@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Julie R. Javier". The signature is fluid and cursive, with the first name "Julie" being the most prominent.

Julie Javier, Team Leader
Risk Mitigation and Implementation Branch 4
Pesticide Re-Evaluation Division
Office of Pesticide Programs

ENCLOSURE: Stamped label

[Note to reviewer: [Text] in brackets indicates optional text].

HEXAZINONE	GROUP	5	HERBICIDE
------------	-------	---	-----------

Tide Hexazinone 75 WDG

[Alternate Brand Name: Tide Hexar™ 75 WDG]

Dispersible Granules

ACTIVE INGREDIENT:

Hexazinone [3-cyclohexyl-6-(dimethylamino)-1-methyl-1,3,5-triazine-2,4(1H,3H)-dione].....

By Weight

75.0%

OTHER INGREDIENTS:

25.0%

TOTAL:

100.0%

KEEP OUT OF REACH OF CHILDREN

DANGER/PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID	
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.	
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. Call CHEMTREC at 1-800-424-9300 for emergency medical treatment information.	

[See] [inside] [label][booklet] [for] [additional] [Precautionary Statements] [,] [and] [Directions for Use] [including] [Storage and Disposal] [instructions] [.]


EPA Reg. No. 84229-32

EPA Est. No.

Manufactured for:

Tide International, USA, Inc.
21 Hubble
Irvine, CA 92618

Net Weight: lbs.
Batch/lot number:



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER. Corrosive. Causes irreversible eye damage. May be fatal if swallowed. Harmful if absorbed through skin. Do not get in eyes, on skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Protective eyewear (goggles, face shield, or safety glasses)
- Long-sleeved shirt and long pants
- Shoes plus socks
- Waterproof gloves

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining personal PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Mixers and loaders supporting chemigation, groundboom, or aerial application to alfalfa, sugarcane, pineapple, or low- or high-bush blueberries, and mixers and loaders supporting application to all non-crop sites, (forestry, Christmas trees, pasture/rangeland, bermudagrass/bahiagrass, and uncultivated areas including rights of way) must wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters.

In addition, when applying to uncultivated areas (including rights of way) using aerial application equipment, mixers and loaders must use closed mixing and loading systems that meet the requirements listed in the WPS for agricultural pesticides [40 CFR 170,607(d)(2)(i) & (ii)] for inhalation protection.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product and as soon as possible, wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

The active ingredient hexazinone, in this product is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated area. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Tide Hexazinone 75 WDG must be used only in accordance with instructions on this label, or in supplemental Tide International, USA, Inc. labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

The correct use rates by crop and geographical area, specified on the label, and proper mixing/loading site considerations and application procedures must be followed to minimize potential for hexazinone movement into ground water. Users are encouraged to consult with their state Department of Agriculture, Extension Service, or other pesticide lead agency for information regarding soil permeability, aquifer vulnerability, and best management practices for their area.

For all forestry uses application by fixed-wing aircraft is prohibited.

MANDATORY SPRAY DRIFT MANAGEMENT

Aerial Applications:

- Do not release spray at a height greater than 10 feet above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to use a medium or coarser droplet size according to the most current version of the American Society of Agricultural & Biological Engineers Standard 641 (ASABE S641).
- Applicators must use ½ swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- If the wind speed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- Do not apply during temperature inversions.

Ground Boom Applications:

- Apply with the nozzle height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy. For all other ground applications, the nozzle must be no more than 4 feet from the target vegetation.
- Applicators are required to use a medium or coarser droplet size according to the most current version of the American Society of Agricultural & Biological Engineers Standard 572 (ASAE S572).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.

BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure – Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- Adjust nozzles – Follow nozzle manufactures recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT – Aircraft

Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

PRODUCT INFORMATION

Tide Hexazinone 75 WDG is a water-dispersible granule that is mixed in water and applied as a spray for weed control in certain crops, Christmas trees, forestry site preparation and release areas, and industrial areas. Tide Hexazinone 75 WDG is also used as a basal soil treatment for brush control in reforestation areas, rangeland, pastures and noncrop areas.

Tide Hexazinone 75 WDG is an effective general herbicide providing both contact and residual control of many annual and biennial weeds and woody plants. It is also effective for control of most perennial weeds.

Tide Hexazinone 75 WDG is noncorrosive to equipment.

Care must be exercised when applying Tide Hexazinone 75 WDG near desirable trees or shrubs as they can absorb Tide Hexazinone 75 WDG through roots extending into treated areas.

Apply this product to agricultural and non-agricultural sites that contain areas of temporary surface water caused by collection of water between planting beds, in equipment ruts, or in other depressions created by management activities. DO NOT make applications to natural or man-made bodies of water such as lakes, reservoirs, ponds, streams and canals.

ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY

Tide Hexazinone 75 WDG is absorbed through the roots and foliage. Moisture is required to activate Tide Hexazinone 75 WDG in the soil. Best results are obtained when the soil is moist at the time of application and ¼ - ½ inches of rainfall occurs within 2 weeks after application.

For best results, apply Tide Hexazinone 75 WDG preemergence or postemergence when weeds are less than 2 inches in height or diameter. Herbicidal activity is most effective under conditions of high temperature (above 80°F), high humidity, and good soil moisture. Reduced herbicidal activity can occur when vegetation is dormant, semi-dormant, or under stress (e.g., temperature or moisture).

Herbicidal activity will usually appear within 2 weeks after application to susceptible plants under warm, humid conditions. When weather is cool or dry, or when susceptible plants are under stress, herbicidal activity will occur 4-6 weeks after application. If rainfall after application is inadequate to activate Tide Hexazinone 75 WDG in the soil, plants may recover from contact effects and continue to grow.

On woody plants, symptoms usually appear within 3-6 weeks after sufficient rainfall has carried the herbicide into the root zone during periods of active growth. Defoliation and subsequent refoliation may occur, but susceptible plants are killed.

The degree and duration of control will depend on the following:

- Use rate
- Weed spectrum and size at time of application
- Environmental conditions at and following treatment

Where a rate range is shown, use the higher levels of the dosage range on hard-to-control species, fine-textured soils, or soils containing greater than 5% organic matter or carbon. Use the lower levels of the dosage range on coarse-textured soils and/or on soils low in organic matter. Refer to specific uses for rate ranges.

APPLICATION INFORMATION

Apply Tide Hexazinone 75 WDG by ground equipment and where permitted, with aerial equipment. Use rates, minimum spray gallonage, and other application information are described for various uses.

Dispose of the equipment washwater by applying it to a use-site listed on this label or in accordance with directions given in the STORAGE AND DISPOSAL section of this label.

Before spraying, calibrate equipment to determine the quantity of water necessary to uniformly and thoroughly cover the vegetation and soil in a measured area to be treated. Make sure the volume of water is sufficient to completely suspend the Tide Hexazinone 75 WDG.

TANK MIXTURES

Tank mix Tide Hexazinone 75 WDG with other herbicides and/or adjuvants registered for the uses (crops) specified on this label.

Refer to the label of the tank mix partner(s) for any additional use instructions or restrictions. The most restrictive label provisions apply. If other label instructions conflict with this label, do not tank mix the herbicide and/or adjuvant with Tide Hexazinone 75 WDG.

INVASIVE SPECIES MANAGEMENT

Use this product for use on public, private, and tribal lands to treat certain weed species infestations that have been determined to be invasive, consistent with the Federal Interagency Committee for the Management of Noxious and Exotic Weeds (FICMNEW) National Early Detection and Rapid Response (EDRR) System for invasive plants. Effective EDRR systems address invasions by eradicating the invader where possible, and controlling them when the invasive species is too established to be feasibly eradicated. Once an EDRR assessment has been completed and action is advised, a Rapid Response needs to be taken to quickly contain, deny reproduction, and if possible eliminate the invader. Consult your appropriate state extension service, forest service, or regional multidisciplinary invasive species management coordination team to determine the appropriate Rapid Response provisions and allowed treatments in your area.

WEED RESISTANCE MANAGEMENT

For resistance management, Tide Hexazinone 75 WDG is a Group 5 herbicide. Any weed population may contain or develop plants naturally resistant to Tide Hexazinone 75 WDG and other Group 5 herbicides. Weed species with acquired resistance to Group 5 herbicides may eventually dominate the weed population if Group 5 herbicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by Tide Hexazinone 75 WDG or other Group 5 herbicides. Users should scout before and after application.

Suspected herbicide-resistant weeds may be identified by these indicators:

- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
- A spreading patch of non-controlled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.

To delay herbicide resistance:

- Avoid the consecutive use of Tide Hexazinone 75 WDG or other target site of action Group 5 herbicides that might have a similar target site of action, on the same weed species.
- Use tank mixtures or premixes with herbicides from different target site of action Groups as long as the involved products are all registered for the same use, have different sites of action and are both effective at the tank mix or prepack rate on the weed(s) of concern (an herbicide mode of action classification by itself may not adequately address specific weeds that are resistant to specific herbicides)
- Base herbicide use on a comprehensive Integrated Pest Management (IPM) program.
- Scout fields prior to application to identify the weed species present and their growth state to determine if the intended application will be effective.
- Scout fields after application to verify that the treatment was effective.
- Contact your local extension specialist, certified crop advisors and/or manufacturer for herbicide resistance management and/or integrated weed management recommendations for specific crops and resistant weed biotypes.

Report any incidence of non-performance of this product against a particular weed species to your Tide International, USA, Inc. representative. If resistance is suspected, treat weed escapes with an herbicide having a different mechanism of action and/or use non-chemicals means to remove escapes, as practical, with the goal of preventing further seed production.

INTEGRATED PEST MANAGEMENT

This product may be used as a part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

AGRICULTURAL USES

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

ALFALFA

Tide Hexazinone 75 WDG is labeled for control of certain weeds in established alfalfa grown for hay or seed production.

- Do not apply within 30 days of harvest (cutting for hay), or feeding of forage or grazing.
- Do not exceed 1 pound Tide Hexazinone 75 WDG (0.75 lb active ingredient hexazinone) per acre per application.
- Do not exceed 2 pounds (1.5 pounds active ingredient hexazinone) per acre per year.

APPLICATION INFORMATION

NON-DORMANT AND SEMI-DORMANT VARIETIES

In the following states, make a single application of Tide Hexazinone 75 WDG during the winter months when alfalfa plants are in the least active stage of growth.

Arizona	Montana	Oklahoma	Washington
California	Nebraska	Oregon	Wyoming
Colorado	Nevada	South Dakota	
Idaho	New Mexico	Texas	
Kansas	North Dakota	Utah	

In the following states, make a single application of Tide Hexazinone 75 WDG either in the spring before new growth exceeds 2 inches in height or to alfalfa stubble after cutting, following hay removal and before regrowth exceeds 2 inches in height.

Arkansas	Maine	New Jersey	Vermont
Connecticut	Maryland	New York	Virginia
Delaware	Massachusetts	North Carolina	West Virginia
Illinois	Michigan	Ohio	Wisconsin
Indiana	Minnesota	Pennsylvania	
Iowa	Missouri	Rhode Island	
Kentucky	New Hampshire	Tennessee	

NOTE: Severe alfalfa injury may result following application, if after cutting the regrowth is more than 2 inches high, or there is significant stubble left after cutting or grazing, or the air temperature is above 90°F.

DORMANT VARIETIES

Make a single application of Tide Hexazinone 75 WDG after alfalfa becomes dormant and before new growth exceeds 2 inches in height in the spring. Where weeds have emerged, use a surfactant.

USE RATES

Use higher rates on hard-to-control species, (see **Weeds Controlled** section below) fine textured soils, soils containing greater than 5% organic matter, or under adverse environmental conditions such as temperature extremes or when weeds are stressed due to low rainfall.

For dormant alfalfa, use a surfactant approved for crops at the rate of 0.25% v/v (1 quart per 100 gallons of spray solution).

Select the appropriate rate for soil texture and organic matter content as follows:

Soils	Tide Hexazinone 75 WDG (lb/acre) Percent Organic Matter in Soil		
	<1%	1-5%	>5%
Coarse Texture Loamy sand, sandy loam	2/3 – 1 (0.5 lb – 0.75 lb ai)	2/3 – 1 (0.5 lb – 0.75 lb ai)	1 (0.75 lb ai)
Medium Texture Loam, silt loam, silt, clay loam, sandy clay loam	2/3 – 1 (0.5 lb – 0.75 lb ai)	1 (0.75 lb ai)	1 (0.75 lb ai)
Fine Texture Silty clay loam, sandy clay, silty clay, clay	1 (0.75 lb ai)	1 (0.75 lb ai)	1 (0.75 lb ai)
NOTE: <ul style="list-style-type: none">In the state of Montana (MT), do not apply to soils with less than 1.5% organic matter.In the state of Wyoming (WY):<ul style="list-style-type: none">-Do not apply to soils with less than 0.5% organic matter.-Apply to irrigated alfalfa only.			

WEEDS CONTROLLED

Tide Hexazinone 75 WDG, when applied preemergence or early postemergence at the following rates, is labeled for the control or suppression of the following species in alfalfa:

1/3 – 2/3 Lb/Acre (0.25 – 0.5 lb ai/acre)	
Tansymustard	<i>Descurainia pinnata</i>

2/3 – 1 Lb/Acre (0.5 – 0.75 lb ai/acre)	
Bluegrass, annual	<i>Poa annua</i>
Brome, downy (cheatgrass)	<i>Bromus tectorum</i>
Buckwheat, wild	<i>Polygonum convolvulus</i>
Catchfly, English	<i>Silene gallica</i>
Chamomile, mayweed (dogfennel)	<i>Anthemis cotula</i>
Chickweed, common	<i>Stellaria media</i>
Fiddleneck, tarweed	<i>Amsinckia lycopsoides</i>
Filaree	<i>Erodium sp.</i>
Flixweed	<i>Descurainia Sophia</i>
Groundsel, common	<i>Senecio vulgaris</i>
Henbit*	<i>Lamium amplexicaule</i>
Lettuce, Miner's	<i>Montia perfoliata</i>
Mustard, blue	<i>Chorispora tenella</i>
Mustard, Jim Hill (tumble)	<i>Sisymbrium altissimum</i>
Mustard, wild	<i>Brassica kaber</i>

2/3 – 1 Lb/Acre (0.5 – 0.75 lb ai/acre)	
Orchardgrass (seedling)	<i>Dactylis glomerata</i>
Pennycress, field	<i>Thlaspi arvense</i>
Pigweed, redroot	<i>Amaranthus retroflexus</i>
Radish, wild	<i>Raphanus raphanistrum</i>
Rocket, London	<i>Sisymbrium irio</i>
Rocket, common yellow	<i>Barbarea vulgaris</i>
Salsify	<i>Tragopogon spp.</i>
Shepherdspurse	<i>Capsella bursa-pastoris</i>
Speedwell, purslane	<i>Veronica peregrina</i>
Spurry, corn	<i>Spergula arvensis</i>

1 Lb/Acre (0.75 lb ai/acre)	
Alfalfa* (seedling)	<i>Medicago sativa</i>
Barley, foxtail (seedling)	<i>Hordeum jubatum</i>
Bluegrass, perennial* (spring only)	<i>Poa spp</i>
Cockle, white*	<i>Melandrium album</i>
Dandelion, common*	<i>Taraxacum officinale</i>
Dandelion, false* (spotted catsear)	<i>Hypochaeris radicata</i>
Foxtail*	<i>Setaria spp.</i>
Kochia	<i>Kochia scoparia</i>
Lambsquarters, common	<i>Chenopodium album</i>
Lettuce, prickly*	<i>Lactuca serriola</i>
Mallow, common	<i>Malva neglecta</i>
Ryegrass, Italian (annual)	<i>Lolium multiflorum</i>
Quackgrass*	<i>Elytrigia repens</i>
Speedwell, ivyleaf	<i>Veronica hederifolia</i>
Tea, Mexican*	<i>Chenopodium ambrosioides</i>
Thistle, Canada (seedling)	<i>Cirsium arvense</i>
Thistle, Russian	<i>Salsola iberica</i>

*Suppression – a visible reduction in plant population and/or plant vigor as compared to an untreated area and generally not accepted as control.

Tide Hexazinone 75 WDG, when applied late in spring or after cutting at the following rates, will control these species listed below:

2/3 – 1 Lb/Acre (0.5 – 0.75 lb ai/acre)	
Crabgrass	<i>Digitaria spp.</i>
Fleabane	<i>Conyza spp.</i>
Foxtail	<i>Setaria spp.</i>
Jimsonweed	<i>Datura stramonium</i>
Lambsquarters, common	<i>Chenopodium album</i>
Pigweed, redroot	<i>Amaranthus retroflexus</i>

SEED ALFALFA (CA, ID, MT, NV, OR, UT, WA)

Use Tide Hexazinone 75 WDG for general broadleaf weed and grass control in established alfalfa grown for seed.

DORMANT VARIETIES

Make a single application of Tide Hexazinone 75 WDG after alfalfa becomes dormant and before new growth exceeds 2 inches in height in the spring. Where weeds have emerged, use a surfactant.

NON-DORMANT AND SEMI-DORMANT VARIETIES

In the following states, make a single application of Tide Hexazinone 75 WDG during the winter months when alfalfa plants are in the least active stage of growth.

WEEDS CONTROLLED

Refer to the Alfalfa – Weeds Controlled section for specific use rates and weeds controlled.

USE PRECAUTIONS AND RESTRICTIONS

SEED ALFALFA

- Do not apply within 30 days of harvest (cutting for hay), or feeding of forage or grazing.
- Do not use Tide Hexazinone 75 WDG on fields with sandy loam or loamy sand soils having less than 1% organic matter.
- Do not exceed 2/3 pound Tide Hexazinone 75 WDG (0.5 lb ai) per acre on fields with sandy loam or loamy sand soils having 1 – 2% organic matter.
- Do not exceed 2/3 pound Tide Hexazinone 75 WDG (0.5 lb ai) per acre on seed alfalfa that has been established for only one growing season.

SEED ALFALFA

WALLA WALLA COUNTY, WASHINGTON

Use Tide Hexazinone 75 WDG for the suppression of prickly lettuce and quackgrass and control of Canada thistle (seedling), kochia, and certain other weeds in established alfalfa grown for seed.

Use Rates 1 pound (0.75 lb ai) per acre	
Kochia	<i>Kochia scoparia</i>
Lettuce, prickly*	<i>Lactuca serriola</i>
Quackgrass*	<i>Elytrigia repens</i>
Thistle, Canada (seedling)	<i>Cirsium arvense</i>

*Suppression

USE PRECAUTIONS AND RESTRICTIONS SEED ALFALFA

WALLA WALLA COUNTY WASHINGTON

- Do not apply within 30 days of harvest (cutting for hay), or feeding of forage or grazing.
- Do not exceed 1 pound (0.75 lb ai) Tide Hexazinone 75 WDG herbicide per acre per application.
- Do not exceed 2 pounds (1.5 pounds active ingredient hexazinone) per acre per year.

SPRAY EQUIPMENT

Apply Tide Hexazinone 75 WDG using a fixed boom power sprayer or aerial equipment.

For ground applications apply in a minimum of 20 gallons of spray solution per acre and by air in a minimum of 5 gallons.

CHEMIGATION ALFALFA

Apply this product only through center pivot or linear-move sprinkler irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

Severe alfalfa injury may result following application after cutting if either the regrowth is more than 2" high or significant stubble is left after alfalfa cutting.

If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments when needed.

DORMANT APPLICATIONS

Select the appropriate rate (see **Use Rates** section) for soil texture and organic matter content using 0.25" to 0.75" of sprinkler irrigation as a continuous injection during the application. Best results are obtained when soil is moist at time of application, and when weeds have not germinated or are less than 2" tall or across.

APPLICATION AFTER CUTTING

Apply Tide Hexazinone 75 WDG at 5.3 ounces (0.25 lb ai) per acre to stubble after cutting, following hay removal, and before regrowth exceeds 2" in height. Apply Tide Hexazinone 75 WDG using 0.25" to 0.75" of sprinkler irrigation as a continuous injection during the application. Best results are obtained when soil is moist at time of application and when weeds have not germinated or are less than 2" tall or across.

NOTE: Increased potential for crop injury is possible when making an application when daily temperatures are forecast to be in the mid-to-high 90 degree range within 3 to 5 days after treatment.

SPRINKLER CHEMIGATION

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

MIXING INSTRUCTIONS

1. Fill the supply tank 1/4 to 1/3 full of water
2. While agitating, add the required amount of Tide Hexazinone 75 WDG and continue agitation until the product is fully dispersed, at least 5 minutes.
3. Once the Tide Hexazinone 75 WDG is fully dispersed, maintain agitation and continue filling tank with water. Tide Hexazinone 75 WDG must be thoroughly mixed with water before adding any other material.
4. As the tank is filling, add tank mix partners (if desired). Follow use precautions and directions on the tank mix partner label.
5. After thorough mixing, the agitation system can be stopped to prevent excessive foaming in the tank. Once thoroughly mixed the solution in the supply tank does not require additional agitation unless specified on the companion products label. If foaming occurs in the injection supply tank, add a defoaming agent (defoamer) if desired.
6. Apply Tide Hexazinone 75 WDG spray mixture within 48 hours of mixing to avoid product degradation.

USE PRECAUTIONS AND RESTRICTIONS – CHEMIGATION

- Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place.
- Distributing treated water in an uneven manner can result in crop injury, lack of effectiveness, or over-tolerance pesticide residues in the crop. Therefore, to ensure that the mixture is applied evenly at the labeled rate, use sufficient water, apply the mixture for the proper length of time and ensure sprinkler produces a uniform water pattern.
- Do not permit run-off during chemigation.

POSTING OF AREAS TO BE TREATED

Posting of areas to be chemigated is required when 1) any part of a treated area is within 300 feet of sensitive areas such as residential areas, labor camps, businesses, daycare centers, hospitals, in-patient clinics, nursing homes, or any public areas such as schools, parks, playgrounds, or other public facilities

not including public roads, or 2) when the chemigated area is open to the public such as golf courses or retail greenhouses.

Posting must conform to all the following requirements:

- Treated areas shall be posted with signs at all usual points of entry and along likely routes of approach from the listed sensitive areas. When there are no usual points of entry, signs must be posted in the corners of the treated areas and in any other location affording maximum visibility to sensitive areas.
- The printed side of the sign must face away from the treated area towards the sensitive area. The signs shall be printed in English. Signs must be posted prior to application and must remain posted until foliage has dried and soil surface water has disappeared. Signs must remain in place indefinitely as long as they are composed of materials to prevent deterioration and maintain legibility for the duration of the posting period.
- All words shall consist of letters at least 2 ½ inches tall, and all letters and the symbol shall be a color which sharply contrasts with their immediate background. At the top of the sign shall be the words "KEEP OUT", followed by an octagonal stop sign symbol at least 8 inches in diameter containing the word "STOP". Below the symbol shall be the words "PESTICIDE IN IRRIGATION WATER".
- Posting required for chemigation does not replace other posting and reentry requirements for farm worker safety.

REPLANTING (FOLLOWING ALFALFA)

- Do not replant treated areas to any crop except corn, root crops, or sugarcane within two years after treatment, to prevent crop injury.
- Plant corn 12 months after the last treatment in areas of moderate to high rainfall (greater than 20 inches), provided the use rate did not exceed 1 lb. Tide Hexazinone 75 WDG (0.75 lb ai) per acre.
- Plant root crops such as potatoes, sugarbeets, radish and carrots 12 months after the last treatment, provided the use rate does not exceed 2/3 lb. Tide Hexazinone 75 WDG (0.5 lb ai) per acre. Sites with use rates higher than 2/3 lb. Tide Hexazinone 75 WDG (0.5 lb ai) per acre must not be replanted to any root crop within 2 years after application of Tide Hexazinone 75 WDG, to prevent unacceptable crop injury.
In areas where irrigation is needed to produce the crop, extend the crop rotation intervals listed if the normal irrigation amount is reduced for any reason.
- Plant sugarcane any time following treatment.
- In California, do not replant seed alfalfa to any crop within two years after treatment, to prevent crop injury.

CROP ROTATION

Field Bioassay

In arid climates (10 inches of rainfall or less per year) or areas where drought conditions have prevailed for one or more years, a field bioassay must be completed prior to planting any desired crop.

Extend the rotation intervals listed above based on the results of the bioassay.

A successful bioassay means growing to maturity a test strip of the crop(s) intended for production. The test crop(s) strip must cross the entire field including knolls, low areas, and areas where any berms were located.

**ALFALFA – IMPREGNATION ON DRY BULK FERTILIZER
(EXCEPT CALIFORNIA AND ARIZONA)**

Impregnate or coat dry bulk fertilizer with Tide Hexazinone 75 WDG for application to established alfalfa. All instructions on this label must be followed along with state regulations relating to dry bulk fertilizer blending, impregnating and labeling.

If fertilizer materials are excessively dusty, use a suitable additive to reduce dust prior to impregnation, as dusty fertilizer will result in poor distribution during application. The dry fertilizer must be properly impregnated and uniformly applied to the alfalfa to avoid crop injury and/or poor weed control.

To impregnate the fertilizer, use a system consisting of a conveyor or closed drum used to blend dry bulk fertilizer. Any commonly used fertilizer can be impregnated with Tide Hexazinone 75 WDG, except potassium nitrate or sodium nitrate. Do not use Tide Hexazinone 75 WDG on limestone.

Use a minimum of 250 lb. dry bulk fertilizer per acre and up to a maximum of 450 lb. per acre. To impregnate or coat the dry bulk fertilizer with Tide Hexazinone 75 WDG, mix 2 2/3 pounds of Tide Hexazinone 75 WDG (2 lbs ai) with sufficient water to make one gallon of suspension and thoroughly agitate. Direct the nozzles to deliver a fine spray of this suspension toward the fertilizer for thorough coverage while avoiding spray contact with mixing equipment. Uniform impregnation of Tide Hexazinone 75 WDG to dry bulk fertilizer will vary, and if the absorptivity is not adequate, use an absorptive powder to produce a dry, free-flowing mixture. "Microcel E" is the absorbent powder of choice. When another herbicide is used with Tide Hexazinone 75 WDG, mix and impregnate the fertilizer immediately.

Apply impregnated fertilizer as soon as possible after impregnation for optimum performance.

Select the rate of Tide Hexazinone 75 WDG to apply per acre from the appropriate section of this label. Then refer to the rate chart below to determine the amount of Tide Hexazinone 75 WDG that is to be impregnated on a ton of dry bulk fertilizer, based on the amount of fertilizer to be distributed in one acre.

Rate Chart for Impregnating Fertilizer with Tide Hexazinone 75 WDG

Fertilizer	Tide Hexazinone 75 WDG Rate Per Acre	
	2/3 Lbs. (0.5 lb ai)	1 Lb. (0.75 lb ai)
250 lbs.	5.3 lbs./ton	8.0 lbs./ton
300 lbs.	4.4 lbs./ton	6.6 lbs./ton
350 lbs.	3.7 lbs./ton	5.7 lbs./ton
400 lbs.	3.3 lbs./ton	5.0 lbs./ton
450 lbs.	2.9 lbs./ton	4.4 lbs./ton

For rates other than those listed, use the following formula to calculate the amounts of Tide Hexazinone 75 WDG to be impregnated per ton of dry fertilizer.

$$\text{lbs. Tide Hexazinone 75 WDG/Acre} \times 1 \text{ Ton Fertilizer} = \text{lbs. Tide Hexazinone 75 WDG per Ton of Fertilizer}$$

APPLICATION

Uniform application of Tide Hexazinone 75 WDG -impregnated dry fertilizer is essential for satisfactory weed control. Accurate calibration of the application equipment is essential for uniform distribution to the surface. The customary method of application is to apply ½ the labeled rate and overlap 50%. This results in the best distribution pattern.

USE PRECAUTIONS AND RESTRICTIONS- ALFALFA

- Best results are obtained when ½ - 1 inches of rainfall or sprinkler irrigation occurs within two weeks after application, when soil is moist at time of application, and when weeds have not germinated or are less than 2 inches in height or diameter. Heavy rainfall or excessive irrigation after application may result in crop injury or poor performance of the herbicide.
- On soils high in organic matter (greater than 5%), the effectiveness of Tide Hexazinone 75 WDG can be significantly reduced and weed control may be unsatisfactory.
- Avoid overlapping of spray swaths and shut off spray booms while starting, turning, slowing or stopping or crop injury may result.
- Crop injury, including mortality, may result in fields with restricted root growth due to non-uniform soil profiles such as gravel bases and clay lenses.
- Crop injury may result if hot weather, mid-to-high 90 degree range or higher, occurs within a few days after application.

- Do not apply to snow-covered or frozen ground.
- Crop injury to alfalfa can be influenced by several factors including alfalfa variety, soil conditions, uniformity of application and environmental conditions, etc., if no prior use history for the site or variety, treat only a small area when first using Tide Hexazinone 75 WDG.
- If abnormally dry conditions exist following application, restrict the first irrigation to no more than ½ acre inch of water.
- Temporary yellowing of alfalfa may occur following Tide Hexazinone 75 WDG applications.
- Treat only stands of alfalfa established for one year or for one growing season (except in California), provided:
 - The alfalfa stand has a well-developed tap root structure that is at least 10 inches in length (0.25 inch diameter below the crown) throughout the field and the crop is healthy, vigorous, and not under stress from weather conditions, low fertility, insects or disease damage.
 - In areas with shorter growing seasons, such as, higher elevations, adequate alfalfa tap root growth may not occur and especially when alfalfa is grown together with a cover or nurse crop. If an adequate tap root is not present, delay application of Tide Hexazinone 75 WDG until the alfalfa has gone through a minimum of two growing seasons.
- In California, treat fall planted alfalfa in the following winter months with Tide Hexazinone 75 WDG at 1/3 to 2/3 pounds (0.25 – 0.5 lb ai) per acre (use higher rate for fine textured soils) provided:
 - alfalfa root growth exceeds 6 inches in length
 - vegetative top growth of alfalfa has lateral development of secondary growth
 - alfalfa is healthy and vigorous, not growing under stress from insect, disease, winter injury or other types of stress.

Injury may result to alfalfa plants that fail to meet these growth criterion listed above.

- Do not use Tide Hexazinone 75 WDG on seedling alfalfa, alfalfa-grass mixtures, or other mixed stands as injury may result to the seedling alfalfa or companion crop.
- Do not use a surfactant with Tide Hexazinone 75 WDG when treating non-dormant alfalfa.
- Do not use Tide Hexazinone 75 WDG on gravelly or rocky soils, exposed subsoils, hardpan, sand poorly drained soil, or alkali soils.

BLUEBERRY

HIGH BUSH BLUEBERRIES

Tide Hexazinone 75 WDG is labeled for control of certain herbaceous and woody weeds in established high bush blueberry fields.

APPLICATION INFORMATION

Apply Tide Hexazinone 75 WDG to high bush blueberries that have been established for 3 or more years. Apply Tide Hexazinone 75 WDG in the spring before the lower leaves of the blueberry plant have fully expanded. Avoid contact of the leaves with the spray solution.

Using calibrated ground spray equipment, make the application in sufficient water then provide thorough and uniform coverage to the treated area (usually 20 gallons per acre). Shut off spray booms when starting, turning, slowing or stopping, or injury to the crop may result.

USE PRECAUTIONS AND RESTRICTIONS

HIGH BUSH BLUEBERRIES

- Do not apply through any type of irrigation system.
- Do not apply within 90 days of harvest.
- Do not apply to flooded field with standing water.
- Application to blueberry foliage will result in crop injury.
- Since the effect of Tide Hexazinone 75 WDG on blueberries varies with soil type, plant vigor, uniformity of applications and amount of rainfall, it is suggested that growers limit their first use to small areas.

USE RATES (Lbs./Acre) HIGH BUSH BLUEBERRIES

Soil Texture	less than or equal to 3% organic matter	greater than 3% organic matter
Coarse loamy sand, sandy loam (50-85% sand)	1.3 (0.975 lb ai)	1.6 (1.2 lbs ai)
Medium loam, silt loam, silt, clay loam, sandy clay loam	--	2.6 (1.95 lbs ai)
Fine silty clay loam, clay loam, sandy clay, silty clay, clay	1.3 – 2* (0.975 – 1.5 lbs ai)	2.6 (1.95 lbs ai)

*Use the higher rate as the soil organic matter approaches 3%.

LOW BUSH BLUEBERRIES

Use Tide Hexazinone 75 WDG for the control of certain weeds in low bush blueberries.

APPLICATION INFORMATION

Apply Tide Hexazinone 75 WDG only to pruned blueberry fields in the spring before leaf emergence. Using calibrated ground spray equipment make the application in sufficient water to provide thorough and uniform coverage to the treated area (usually 20 gallons per acre). Shut off spray booms when starting, turning, slowing or stopping, or injury to the crop may result.

USE PRECAUTIONS AND RESTRICTIONS LOWBUSH BLUEBERRIES

- Do not apply through any type of irrigation system.
- Do not apply to flooded field with standing water.
- Do not apply within 450 days of harvest.
- Do not exceed 2.4 pounds Tide Hexazinone 75 WDG (1.8 lbs ai) per acre if field has been treated with hexazinone within the past 8 years.
- Application to blueberry foliage will result in crop injury.
- Since the effect of Tide Hexazinone 75 WDG on blueberries varies with soil type, plant vigor, uniformity of applications and amount of rainfall, it is suggested that growers limit their first use to small areas. If excessive leaf drop is observed after treatment, reduce rate in future applications.
- Maintain a 50-foot buffer from any well head or water reservoir.

LOW BUSH BLUEBERRIES (LBS./ACRE)

Soil Texture	less than or equal to 3% organic matter	greater than 3% organic matter
Coarse loamy sand, sandy loam (50-85% sand)	1.2 (0.9 lb ai)	1.6 (1.2 lbs ai)
Medium loam, silt loam, silt, clay loam, sandy clay loam	--	2 (1.5 lbs ai)
Fine silty clay loam, clay loam, sandy clay, silty clay, clay	1.2 – 2.4* (0.9 – 1.8 lbs ai)	2.4 – 3.6** (1.8 – 2.7 lbs ai)

*Use the higher rate as the soil organic matter approaches 3%.

**Use the higher rate for harder to control species.

IMPREGNATION ON DRY BULK FERTILIZER

Impregnate or coat dry bulk fertilizer with Tide Hexazinone 75 WDG for application to established high bush or low bush blueberries. All instructions on this label must be followed along with state regulations relating to dry bulk fertilizer blending, impregnating and labeling.

If fertilizer materials are excessively dusty, use a suitable additive to reduce dust prior to impregnation, as dusty fertilizer will result in poor distribution during application. The dry fertilizer must be properly impregnated and uniformly applied to the alfalfa to avoid crop injury and/or poor weed control.

To impregnate the fertilizer, use a system consisting of a conveyor or closed drum used to blend dry bulk fertilizer. Any commonly used fertilizer can be impregnated with Tide Hexazinone 75 WDG, except potassium nitrate or sodium nitrate. Do not use Tide Hexazinone 75 WDG on limestone.

Use a minimum of 250 lbs. dry bulk fertilizer per acre and up to a maximum of 450 lbs. per acre. To impregnate or coat the dry bulk fertilizer with Tide Hexazinone 75 WDG, mix 2 2/3 pounds of this product (2 lbs ai) with sufficient water to make one gallon of suspension and thoroughly agitate. Direct the nozzles to deliver a fine spray of this suspension toward the fertilizer for thorough coverage while avoiding spray contact with mixing equipment. Uniform impregnation of Tide Hexazinone 75 WDG to dry bulk fertilizer will vary, and if the absorptivity is not adequate, use an absorptive powder to produce a dry, free-flowing mixture. "Microcel E" is the absorbent powder of choice. When another herbicide is used with Tide Hexazinone 75 WDG, mix and impregnate the fertilizer immediately.

Apply impregnated fertilizer as soon as possible after impregnation for optimum performance.

Select the rate of Tide Hexazinone 75 WDG to apply per acre from the appropriate section of this label. Then refer to the rate chart below to determine the amount of Tide Hexazinone 75 WDG that is to be impregnated on a ton of dry bulk fertilizer, based on the amount of fertilizer to be distributed in one acre.

Rate Chart for Impregnating Fertilizer with Tide Hexazinone 75 WDG

Fertilizer Rate/Acre	Tide Hexazinone 75 WDG Rate Per Acre			
	2/3 Lb. (0.5 lb ai)	1 Lb. (0.75 lb ai)	1 1/3 Lbs. (1 lb ai)	2 Lbs. (1.5 lbs ai)
250 lbs.	5.3 lbs./ton	8.0 lbs./ton	10.6 lbs./ton	16.0 lbs./ton
300 lbs.	4.4 lbs./ton	6.6 lbs./ton	8.8 lbs./ton	13.3 lbs./ton
350 lbs.	3.7 lbs./ton	5.7 lbs./ton	7.5 lbs./ton	11.4 lbs./ton
400 lbs.	3.3 lbs./ton	5.0 lbs./ton	6.7 lbs./ton	10.0 lbs./ton
450 lbs.	2.9 lbs./ton	4.4 lbs./ton	5.9 lbs./ton	8.9 lbs./ton

For rates other than those listed, use the following formula to calculate the amounts of Tide Hexazinone 75 WDG to be impregnated per ton of dry fertilizer.

$$\text{lbs. Tide Hexazinone 75 WDG/Acre} \times 1 \text{ Ton Fertilizer} = \text{lbs. Tide Hexazinone 75 WDG per Ton of Fertilizer}$$

APPLICATION

Uniform application of Tide Hexazinone 75 WDG-impregnated dry fertilizer is essential for satisfactory weed control. Accurate calibration of the application equipment is essential for uniform distribution to the surface. The customary method of application is to apply 1/2 the labeled rate and overlap 50%. This results in the best distribution pattern.

WEEDS CONTROLLED

Tide Hexazinone 75 WDG is labeled for the control or suppression of the following species in high and low bush blueberry crops:

Aster, heath*	<i>Aster ericoides</i>
Barnyardgrass	<i>Echinochloa crus-galli</i>
Blackberry* (briar)	<i>Rubus</i> spp.
Bluegrass, Kentucky (perennial)*	<i>Poa pratensis</i>
Brome, downy (cheatgrass)	<i>Bromus tectorum</i>
Broomsedge*	<i>Andropogon virginicus</i>
Carrot, wild*	<i>Daucus carota</i>

Catchfly, English	<i>Silene gallica</i>
Chamomile, mayweed	<i>Anthemis cotula</i>
Cherry, wild	<i>Prunus serotia</i>
Chickweed, common	<i>Stellaria media</i>
Cinquefoil	<i>Potentilla</i> spp.
Cockle, white*	<i>Melandrium album</i>
Dandelion, common*	<i>Taraxacum officinale</i>
Dandelion, false* (spotted catsear)	<i>Hypochaeris radicata</i>
Daisy, oxeye	<i>Chrysanthemum leucanthemum</i>
Dock, curly*	<i>Rumex crispus</i>
Dogfennel	<i>Eupatorium capillifolium</i>
Fescue*	<i>Festuca</i> spp.
Fiddleneck, tarweed	<i>Amsinckia lycopsoides</i>
Filaree	<i>Erodium</i> spp.
Fireweed* (willowweed)	<i>Epilobium angustifolium</i>
Fleabane, flax-leaved	<i>Conyza bonariensis</i>
Flixweed	<i>Descurainia Sophia</i>
Foxtail, yellow	<i>Setaria lutescens</i>
Goldenrod	<i>Solidago</i> spp.
Groundsel, common	<i>Senecio vulgaris</i>
Hawkweed	<i>Hieracium</i> spp.
Horseweed/marestail	<i>Conyza canadensis</i>
Jimsonweed	<i>Datura stramonium</i>
Lambsquarters, common	<i>Chenopodium album</i>
Lettuce, Miner's	<i>Montia perfoliata</i>
Lettuce, prickly*	<i>Lactuca serriola</i>
Mustard, blue	<i>Chorispora tenella</i>
Mustard, Jim Hill (tumble)	<i>Sisymbrium altissimum</i>
Orchardgrass*	<i>Dactylis glomerata</i>
Orchardgrass (seedling)	<i>Dactylis glomerata</i>
Panicgrass (witchgrass)	<i>Panicum capillare</i>
Panicum, fall	<i>Panicum dichotomiflorum</i>
Pearly everlasting	<i>Anaphalis margaritacea</i>
Pennycress, field	<i>Thlaspi arvense</i>
Pigweed, redroot	<i>Amaranthus retroflexus</i>
Quackgrass	<i>Agropyron repens</i>
Radish, wild	<i>Raphanus raphanistrum</i>
Ragweed, common	<i>Ambrosia elatior</i>
Raspberry* (briar)	<i>Rubus</i> spp.
Rocket, London	<i>Sisymbrium irio</i>
Rocket, common yellow	<i>Barbarea vulgaris</i>
Ryegrass, Italian (annual)	<i>Lolium multiflorum</i>
Ryegrass, perennial*	<i>Lolium perenne</i>
Salsify	<i>Tragopogon</i> spp.
Shepherdspurse	<i>Capsella bursa-pastoris</i>
Smartweed, Pennsylvania	<i>Polygonum pensylvanicum</i>
Sorrel, red	<i>Rumex acetosella</i>
Sorrel, sheep	<i>Rumex angiocarpus</i>
Spurry, corn	<i>Spergula arvensis</i>
Strawberry, wild	<i>Fragaria virginiana</i>
Tansymustard (pinnate)	<i>Descurainia pinnata</i>
Tea, Mexican*	<i>Chenopodium ambrosioides</i>
Velvetgrass	<i>Holcus lanatus</i>
Yarrow	<i>Achillea</i> spp.

2.4 to 3.6 Lbs./acre (1.8 – 2.7 lbs ai/acre)	
Dogbane**	<i>Apocynum</i> spp.
Meadow-sweet	<i>Filipendula ulmaria</i>
Blackberry, trailing	<i>Rubus ursinus</i>
Laurel, sheep	<i>Kalmia angustifolia</i>
Rose, wild**	<i>Rosa</i> spp.

*Suppression – a visible reduction in plant population and/or plant vigor as compared to an untreated area and generally not accepted as control.

**Harder to control species.

CHRISTMAS TREES

Tide Hexazinone 75 WDG is labeled for control of certain weeds where the following species are grown:

Fir, Douglas (western US only)	<i>Pseudotsuga menziesii</i>
Fir, Fraser	<i>Abies fraseri</i>
Fir, grand	<i>Abies grandis</i>
Fir, noble	<i>Abies procera</i>
Pine, Austrian	<i>Pinus nigra</i>
Pine, loblolly	<i>Pinus taeda</i>
Pine, ponderosa	<i>Pinus ponderosa</i>
Pine, Scotch	<i>Pinus sylvestris</i>
Spruce, Sitka	<i>Picea sitchensis</i>

Unless otherwise directed in separately published Tide instructions, do not use Tide Hexazinone 75 WDG on Christmas trees in the following states:

Alabama	Louisiana	New Jersey	Texas
Arkansas	Maine	New York	Vermont
Connecticut	Maryland	North Carolina	Virginia
Delaware	Massachusetts	Pennsylvania	West Virginia
Georgia	Mississippi	Rhode Island	
Florida	New Hampshire	South Carolina	

APPLICATION INFORMATION

EASTERN US

Apply Tide Hexazinone 75 WDG as a broadcast spray in the spring prior to bud break. If application is made after bud break, use directional spray equipment to prevent contact with foliage.

WESTERN US

Areas of greater than 20 inches annual rainfall – Apply Tide Hexazinone 75 WDG as a broadcast spray in the spring prior to conifer bud break. If application is made after bud break, use directional spray equipment to prevent contact with foliage.

Areas of less than 20 inches annual rainfall – Apply Tide Hexazinone 75 WDG in the fall before the soil freezes or in the spring after snow cover melts, but before conifer bud break occurs.

USE RATES

The rates listed below are for broadcast application. For band application, use proportionately less; for example, use ½ of the broadcast rates when treating a 3-foot band where row spacing is 6 feet. Use the higher end of the rate range on the heavier soil type.

Do not use more than one application of Tide Hexazinone 75 WDG per year.

Soils	Tide Hexazinone 75 WDG (Lbs./Acre)	
	First Year Plantings	Established Trees
Coarse Texture		
Loamy sand, sandy loam (50-85% sand)	1 1/3 (1 lb ai)	1 1/3 – 1 2/3 (1 – 1.25 lbs ai)
Medium Texture		
Loam, silt loam, silt, clay loam, sandy clay loam	1 1/3 – 1 2/3 (1 – 1.25 lbs ai)	1 2/3 – 2 1/3 (1.25 – 1.75 lbs ai)
Fine Texture		
Silty clay loam, clay loam, sandy clay, silty clay, clay	1 2/3 – 2 (1.25 – 1.5 lbs ai)	2 1/3 – 2 2/3 (1.75 – 2 lbs ai)

First year plantings – Transplant stock that is 2 years old or more (1 year old for loblolly pine). Apply Tide Hexazinone 75 WDG only if rainfall has settled the soil around the base and root systems of the transplants.

Established trees – Trees that have been planted in the plantation for 1 year or more.

WEEDS CONTROLLED

Tide Hexazinone 75 WDG is labeled for the control or suppression of the following weed species in Christmas tree crops:

Aster, heath*	<i>Aster ericoides</i>
Barnyardgrass	<i>Echinochloa crus-galli</i>
Bentgrass, common	<i>Agrostis alba</i>
Bluegrass, annual	<i>Poa annua</i>
Bromegrass	<i>Bromus</i> spp.
Burnweed, American*	<i>Erechtites hieracifolius</i>
Carrot, wild	<i>Daucus carota</i>
Crabgrass*	<i>Digitaria</i> spp.
Curly dock*	<i>Rumex crispus</i>
Daisy, oxeye	<i>Chrysanthemum leucanthemum</i>
Dandelion, common*	<i>Taraxacum officinale</i>
Dandelion, false* (spotted catsear)	<i>Hypochaeris radicata</i>
Fescue*	<i>Festuca</i> spp.
Fleabane	<i>Conyza</i> spp.
Foxtail	<i>Setaria</i> spp.
Goldenrod*	<i>Solidago</i> spp.
Groundsel, common	<i>Senecio vulgaris</i>
Horseweed/marestail	<i>Conyza canadensis</i>
Orchardgrass*	<i>Dactylis glomerata</i>
Ragweed, common	<i>Ambrosia elatior</i>
Ryegrass, Italian (annual)	<i>Lolium multiflorum</i>
Ryegrass, perennial*	<i>Lolium perenne</i>
Smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>
Velvetgrass, common	<i>Holcus lanatus</i>

*Suppression – a visible reduction in plant population and/or plant vigor as compared to an untreated area and generally not accepted as control.

SPRAY EQUIPMENT

Apply Tide Hexazinone 75 WDG by ground equipment or by air.

Select a spray volume that will ensure a through and uniform application. Apply a minimum of 5 gallons per acre by air and a minimum of 10 gallons per acre by ground equipment.

USE PRECAUTIONS AND RESTRICTIONS CHRISTMAS TREES

- Do not use Tide Hexazinone 75 WDG in nurseries, seed beds, or ornamental plantings.
- Do not add a surfactant in applications over the top of conifers.
- Weed control results from spring applications depend on sufficient moisture to activate Tide Hexazinone 75 WDG.
- Livestock may be grazed immediately following a broadcast application of Tide Hexazinone 75 WDG at rates of 1.5 pounds (1.12 lbs ai) per acre or less, and treated vegetation may be cut, dried, and fed after 38 days.
- Do not cut treated vegetation for feed, or graze livestock on treated areas for 60 days following application of Tide Hexazinone 75 WDG at broadcast rates exceeding 1.5 pounds (1.12 lbs ai) per acre.
- Poor weed and brush control may result from the following:
 - Heavy duff or slash present at the time of application.
 - Use on poorly drained sites.
 - Applications made when soil is saturated with water and rain is imminent within 24 hours.
 - Applications to soils high in organic matter (greater than 5%).
- Injury may occur when Tide Hexazinone 75 WDG is used on the following:
 - Trees that show poor vigor, insect damage, disease, winter injury, or other stress conditions.
 - Any soil containing less than 1% organic matter
 - Loamy sand or sandy loam with less than 2% organic matter (except Jeffrey Pine and Ponderosa Pine).
 - Foliage after bud break.
 - Gravelly or rocky soils, exposed subsoils, clay knobs, sand, or sandy soil with 85% or more sand.

PINEAPPLE

Tide Hexazinone 75 WDG is labeled for control of certain weeds in pineapple. See Use Precautions and Restrictions section below for information on maximum chemigation application rate.

APPLICATION INFORMATION

Mix the proper amount of Tide Hexazinone 75 WDG in water. Add a surfactant at the rate of 0.25% V/V.

Use the lower rates on coarse-textured soils or in areas where rainfall exceeds 65 inches per year. Use the higher rates on fine-textured soils or in areas where rainfall is less than 65 inches per year.

Intercrop period – Apply Tide Hexazinone 75 WDG as a broadcast spray in 100-400 gallons of water per acre at the rate of 1/3 – 2 1/3 pounds (0.25 – 1.5 lbs ai) per acre. For aerial application, use at least 10 gallons water per acre.

Post mulch, preplant – Apply Tide Hexazinone 75 WDG as a broadcast spray in 100-400 gallons of water per acre at the rate of 1/3 – 2 1/3 (0.25 – 1.5 lbs ai) pounds per acre.

Post plant, before planted cuttings start active growth – Apply Tide Hexazinone 75 WDG as a broadcast spray in 100-400 gallons of water per acre at the rate of 1/3 – 2 1/3 pounds (0.25 – 1.5 lbs ai) per acre. When weed growth has escaped control by other herbicide applications, apply a post-planting application after the planted cuttings start to grow.

Prior to forcing first ratoon – Apply Tide Hexazinone 75 WDG as a broadcast spray in 100-400 gallons of water per acre at the rate of 1/3 – 2 1/3 pounds (0.25 – 1.5 lbs ai) per acre.

Directed postemergence (pineapple and weeds) interspace application – Apply Tide Hexazinone 75 WDG as a directed spray 3-10 months after planting in 50-200 gallons of water per acre (broadcast basis) at the rate of 1/3 – 2 1/3 pounds (0.25 – 1.5 lbs ai) per acre (broadcast basis) using a stroller boom or knapsack.

Directed spot treatments for perennial grasses before floral induction – Spray perennial grasses postemergence to wet (50-200 gallons per acre depending on size) with 1 1/3 – 2 1/3 pounds Tide Hexazinone 75 WDG (1.0 – 1.5 lbs ai) per 100 gallons of water as a spot treatment.

Treatments to field edges and roadsides – Apply Tide Hexazinone 75 WDG at 2 1/3 – 4 8/10 pounds (1.5 – 3.6 lbs ai) per acre in 100-400 gallons of water.

WEEDS CONTROLLED

Tide Hexazinone 75 WDG is labeled for the control or suppression of the following species in pineapple crops:

Ageratum, tropic	<i>Ageratum conyzoides</i>
Balsamapple	<i>Momordica charantia</i>
Castorbean	<i>Ricinus communis</i>
Crabgrass	<i>Digitaria</i> spp.
Crotalaria	<i>Crotalaria</i> spp.
Dallisgrass	<i>Paspalum dilatatum</i>
Guineagrass	<i>Panicum maximum</i>
Junglerice	<i>Echinochloa colonum</i>
Kao haole*	<i>Leucaena glauca</i>
Moana loa vine*	<i>Canavalia cathartica</i>
Morningglory	<i>Ipomoea</i> spp.
Oxalis	<i>Oxalis</i> spp.
Popolo	<i>Solanum sandwicense</i>
Richardsonium	<i>Richardsonia</i> spp.
Vaseygrass	<i>Paspalum urvillei</i>

*Suppression – a visible reduction in plant population and/or plant vigor as compared to an untreated area and generally not accepted as control.

USE PRECAUTIONS AND RESTRICTIONS-PINEAPPLE

- Do not exceed 4.8 lbs. Tide Hexazinone 75 WDG (3.6 lbs ai) per acre per crop.
- For Chemigation Application, do not exceed 2 lbs. Tide Hexazinone 75 WDG (1.5 lbs ai) per acre per application.
- Do not apply Tide Hexazinone 75 WDG within 181 days of harvest.

SUGARCANE

Tide Hexazinone 75 WDG is labeled for selective weed control in sugarcane except in the State of Florida.

APPLICATION INFORMATION

Apply a single treatment of Tide Hexazinone 75 WDG per year using a fixed-boom sprayer and a minimum of 25 gallons per acre unless otherwise directed.

HAWAII

Apply Tide Hexazinone 75 WDG pre- or postemergence at the following rates for the indicated soil texture:

Soils	Tide Hexazinone 75 WDG (Lb./Acre) (Plus surfactant 0.25% by volume)
Coarse Texture	
Sand, loamy sand, sandy loam	2/3 – 1 1/10 (0.5 – 0.85 lb ai)
Medium Texture	
Loam, silt loam, silty clay loam	2/3 – 1 1/10 (0.5 – 0.85 lb ai)
Fine Texture	
Clay, gray hydromorphic clay	1 1/10 (0.85 lb ai)

Use the higher levels of the labeled rate ranges on soils high in organic matter. Do not apply more than twice the highest labeled rate for the indicated soil texture per crop (18-24 months).

Add an adjuvant for all uses.

For preemergence use only, apply Tide Hexazinone 75 WDG with aerial equipment using at least 10 gallons of spray per acre.

Apply Tide Hexazinone 75 WDG as a spot spray application for emerged weeds in sugarcane. Mix 1 to 4 pounds of Tide Hexazinone 75 WDG (0.75 – 3 lbs ai) per 100 gallons of water. Apply a sufficient volume of spray solution to thoroughly wet weed foliage but do not exceed a use rate of 1.1 pounds (0.85 lbs ai) per acre. Use the lower concentrations on coarse-textured soils that are low in organic matter, and use the higher concentrations on fine-textured soils that are high in organic matter.

LOUISIANA

Apply 2/3 – 1 1/10 pounds of Tide Hexazinone 75 WDG (0.5 – 0.85 lb ai) per acre broadcast in the fall before sugarcane emerges or in the spring before active cane tillering begins. Fall treatments of 2/3 – 1 1/10 pound Tide Hexazinone 75 WDG (0.5 – 0.85 lb ai) per acre. Do not apply more than 2 pounds (1.5 lbs ai) per year. Use the lower rates on coarse textured soils and the higher rates on fine textured soils.

PUERTO RICO

For preemergence treatments, apply 1/3 – 2/3 pound of Tide Hexazinone 75 WDG (0.25 – 0.5 lb ai) per acre to weeds after they have emerged. Use the lower rates on coarse-textured soils and the higher rates on fine-textured soils (high in clay or organic matter). Apply up to 2/3 pounds of Tide Hexazinone 75 WDG (0.5 lb ai) per ratoon per acre.

For spot treatment of emerged weeds, apply Tide Hexazinone 75 WDG with a knapsack sprayer in concentrations of 1/3 – 2/3 pound (0.25 – 0.5 lb ai) per 100 gallons of water. Apply a sufficient spray volume to wet the weed foliage. Do not exceed 100 gallons of spray per treated acre. Use the lower concentration on fine-textured soils.

Note: Since it is difficult to calibrate “spot” knapsack applications, extra care must be taken not to exceed the rate equivalent of the maximum of 2/3 pound Tide Hexazinone 75 WDG (0.5 lb ai) per acre.

Do not apply more than 1 1/3 pounds of Tide Hexazinone 75 WDG (1 lb ai) per acre per crop season.

TEXAS

Apply 2/3 – 1 1/10 pounds of Tide Hexazinone 75 WDG (0.5 - 0.85 lb ai) per acre. On plant cane, apply the herbicide before the cane emerges or as a directed layby treatment. On stubble cane, apply Tide Hexazinone 75 WDG preemergence to early postemergence (up to the 3-leaf stage) or as a directed layby treatment. A pre- or early postemergence treatment may be followed by a layby treatment, provided at least 60 days have elapsed and 3 inches of rainfall or sprinkler irrigation have occurred since the first treatment.

Do not apply more than 2 1/3 pounds of Tide Hexazinone 75 WDG (1.75 lbs ai) per acre per season.

Use the following rates according to the different soil textures:

Soils	Tide Hexazinone 75 WDG (Lb./Acre)	
	Preemergence	+ Layby
Coarse Texture*		
Sandy loam	1/3 (0.25 lb ai)	1/3 (0.25 lb ai)
Medium Texture		
Loam, silt loam	9/10 (0.67 lb ai)	9/10 (0.67 lb ai)
Fine Texture		
Clay loam	1 1/10 (0.85 lb ai)	1 1/10 (0.85 lb ai)

* With at least 2% organic matter.

On dormant cane, add a surfactant to the spray mixture to increase control of emerged weeds.

WEEDS CONTROLLED

Tide Hexazinone 75 WDG is labeled for the control or suppression of the following species in sugarcane crops:

Ageratum, tropic*	<i>Ageratum conyzoides</i>
Alexandergrass	<i>Brachiaria plantaginea</i>
Balsamapple	<i>Momordica charantia</i>
Barnyardgrass	<i>Echinochloa crus-galli</i>
Bermudagrass*	<i>Cynodon dactylon</i>
Burnweed, American (fireweed)	<i>Erechtites hieracifolius</i>
Chickweed, common	<i>Stellaria media</i>
Crabgrass, large	<i>Digitaria sanguinalis</i>
Crabgrass, smooth	<i>Digitaria ischaemum</i>
Crotalaria, fuzzy	<i>Crotalaria incana</i>
Crotalaria, showy	<i>Crotalaria spectabilis</i>
Cuphea, tarweed	<i>Cuphea carthagenensis</i>
Dallisgrass	<i>Paspalum dilatatum</i>
Fingergrass, radiate	<i>Chloris radiata</i>
Fingergrass, swollen	<i>Chloris barbata</i>
Foxtail, bristly	<i>Setaria verticillata</i>
Foxtail, yellow	<i>Setaria lutescens</i>
Geranium, Carolina	<i>Geranium carolinianum</i>
Goosegrass	<i>Elusine indica</i>
Guineagrass	<i>Panicum maximum</i>
Henbit	<i>Lamium amplexicaule</i>
Itchgrass*	<i>Rottboellia cochinchinensis</i>
Job's-tears	<i>Coix lacryma</i>
Johnsongrass (seedling)	<i>Sorghum halepense</i>
Junglerice	<i>Echinochloa colonum</i>
Lambsquarters, common	<i>Chenopodium album</i>
Millet, Texas	<i>Panicum texanum</i>
Morningglory, hairy	<i>Ipomoea pentaphylla</i>
Morningglory, threelobe	<i>Ipomoea triloba</i>
Mustard, wild	<i>Sinapis arvensis</i>
Oxalis	<i>Oxalis spp</i>
Paintbrush, Flora's	<i>Emilia sonchifolia</i>
Panicum, browntop	<i>Panicum fasciculatum</i>
Paspalum, ricegrass	<i>Paspalum orbiculare</i>
Paspalum, sour	<i>Paspalum conjugatum</i>
Pigweed, redroot	<i>Amaranthus retroflexus</i>
Pigweed, slender (green)	<i>Amaranthus viridis</i>
Pigweed, smooth	<i>Amaranthus chlorostachys</i>
Popolo	<i>Solanum sandwicense</i>
Purslane, common	<i>Portulaca oleracea</i>
Sandbur	<i>Cenchrus spp</i>
Sensitive plant (hila hila)	<i>Mimosa spp</i>
Signalgrass, broadleaf	<i>Brachiaria platyphylla</i>
Sowthistle, common	<i>Sonchus oleraceus</i>
Spanishneedles	<i>Bidens bipinnata</i>
Sprangletop	<i>Leptochloa spp</i>
Spurge, prostrate	<i>Euphorbia humistrata</i>
Spurge, graceful	<i>Chamaesyce hypericifolia</i>
Sunflower	<i>Helianthus spp</i>
Vaseygrass	<i>Paspalum urvillei</i>

Waltheria (hia loa)	<i>Waltheria</i> spp.
---------------------	-----------------------

* Suppression – a visible reduction in plant population and/or plant vigor as compared to an untreated area and generally not accepted as control.

USE PRECAUTIONS AND RESTRICTIONS – SUGARCANE

- Do not exceed 1.13 pounds Tide Hexazinone 75 WDG (0.85 pound active ingredient hexazinone) per acre per application.
- Do not plant any crop other than sugarcane following an application of Tide Hexazinone 75 WDG.
- Do not feed sugarcane forage to livestock.
- Do not apply Tide Hexazinone 75 WDG:
 - Within 180 days of harvest in Hawaii.
 - Within 234 days of harvest in Louisiana.
 - Within 288 days of harvest in Puerto Rico.
 - Within 234 days of harvest in Texas.
- To avoid injury to sugarcane, observe the following precautions:
 - Do not use Tide Hexazinone 75 WDG on cane that shows poor vigor because of insect damage, disease or winter injury, or shows symptoms of other stress conditions such as drought stress.
 - Do not add a surfactant in applications unless otherwise specified.
 - Do not use Tide Hexazinone 75 WDG on gravelly or rocky soils, thinly covered subsoils, or coarse-textured soils (sands to sandy loams) with less than 1% organic matter.
 - Temporary chlorosis of the crop may result from application over emerged cane. Applications during active cane growth must be directed to cover the weeds and soil while minimizing crop contact.
 - Do not use Tide Hexazinone 75 WDG on varieties known to be susceptible to herbicides.
 - Extremely heavy rainfall after application may result in poor weed control and/or crop injury, especially if the application is made to dry soil.

FORESTRY

SITE PREPARATION

Tide Hexazinone 75 WDG is labeled for weed and brush control in areas where the following species are grown:

EASTERN US AND LAKE STATES

Fir, balsam	<i>Abies balsamea</i>
Pine, Austrian	<i>Pinus negra</i>
Pine, loblolly	<i>Pinus taeda</i>
Pine, longleaf	<i>Pinus palustris</i>
Pine, ponderosa	<i>Pinus ponderosa</i>
Pine, red	<i>Pinus resinosa</i>
Pine, Scotch	<i>Pinus sylvestris</i>
Pine, shortleaf	<i>Pinus echinata</i>
Pine, slash	<i>Pinus elliotii</i>
Pine, Virginia	<i>Pinus virginiana</i>
Spruce, black	<i>Picea mariana</i>
Spruce, red	<i>Picea rubens</i>
Spruce, white	<i>Picea glauca</i>

WESTERN US

Fir, Douglas	<i>Pseudotsuga menziesii</i>
Fir, grand	<i>Abies grandis</i>
Fir, Noble	<i>Abies procera</i>
Fir, white	<i>Abies concolor</i>

Pine, Jeffrey	<i>Pinus jeffreyi</i>
Pine, Lodgepole	<i>Pinus contorta</i>
Pine, ponderosa	<i>Pinus ponderosa</i>
Spruce, blue	<i>Picea pungens</i>
Spruce, Engleman	<i>Picea englemannii</i>
Spruce, Sitka	<i>Picea sitchensis</i>

APPLICATION INFORMATION

EASTERN US

Apply Tide Hexazinone 75 WDG from early spring to early summer after hardwoods have broken bud and before the foliage has hardened off.

Soils	Tide Hexazinone 75 WDG (Lbs./Acre) Eastern US
Coarse Texture	
Sand, loamy sand, sandy loam	2 2/3 – 4 (2 – 3 lbs ai)
Medium Texture	
Loam, silt loam, sandy clay loam	4 – 5 1/3 (3 – 4 lbs ai)
Fine Texture	
Silty clay loam, clay loam, sandy clay, silt, silty clay, clay	5 1/3 – 6 2/3 (4 – 5 lbs ai)

The rates listed are for broadcast application. Use the lower rates on coarse textured soils and soils low in organic matter. Use the higher rates on fine textured soils and soils high in organic matter. Use the higher rates where weeds identified with an * in the Weed Controlled list predominate.

WESTERN US

For **SITE PREPARATION**, apply Tide Hexazinone 75 WDG at 1.3 to 4 pounds (0.975 – 3 lbs ai) per acre. Use the lower rates on coarse textured soils and soils low in organic matter. Use the higher rates on fine textured soils and soils high in organic matter. Use the higher rates where weeds identified in this label as “suppression” predominate.

In areas where other conifer species may be mixed in with the conifers listed above, apply Tide Hexazinone 75 WDG if the user has prior experience with Tide Hexazinone 75 WDG on the other conifer species. With no prior experience, it is advised that either a small area of plantings be tested for conifer safety prior to treating larger areas, or make no application of Tide Hexazinone 75 WDG in these areas within the site preparation area. Conifer species that are sensitive to Tide Hexazinone 75 WDG (hexazinone) such as, sugar pine and western larch, require 18 months before interplanting on treated sites.

Applications made to shelter wood sites may also result in mortality to over-story conifers. Factors that influence conifer sensitivity in these sites could include application rate, conifer species, soil characteristics, uniformity of spray distribution across the treatment swath and environmental stress.

Rain Belt (areas of high spring rainfall): For best results, apply in late winter or spring when weeds and brush are actively growing.

Snow Belt (areas of low spring rainfall): For best results, apply in the fall before soil freezes, or in the spring after snow cover melts in anticipation of rainfall. Weed and brush control results from spring applications will be dependent on sufficient rainfall following application to activate Tide Hexazinone 75 WDG.

PLANTS CONTROLLED

Tide Hexazinone 75 WDG is labeled for the control or suppression of the following species in site preparations for forestry crops:

HERBACEOUS PLANTS

Asters	
Aster, heath*	<i>Aster ericoides</i>
Barnyardgrass	<i>Echinochloa crus-galli</i>
Bentgrass	<i>Agrostis</i> spp.
Bluegrass, annual	<i>Poa annua</i>
Bromegrass	<i>Bromus</i> spp.
Carrot, wild	<i>Daucus carota</i>
Crabgrass*	<i>Digitaria</i> spp.
Daisy, oxeye	<i>Chrysanthemum leucanthemum</i>
Dandelion, common*	<i>Taraxacum officinale</i>
Dandelion, false* (spotted catsear)	<i>Hypochaeris radicata</i>
Dock, curly*	<i>Rumex crispus</i>
Elksedge	<i>Carex geyeri</i>
Fescue*	<i>Festuca</i> spp.
Fireweed* (willowweed)	<i>Epilobium angustifolium</i>
Fleabane	<i>Conyza</i> spp.
Foxtail	<i>Setaria</i> spp.
Goldenrod*	<i>Solidago</i> spp.
Groundsel, common	<i>Senecio vulgaris</i>
Horseweed/marestail	<i>Conyza canadensis</i>
Mullein, common**	<i>Verbascum thapsus</i>
Orchardgrass*	<i>Dactylis glomerata</i>
Pinegrass	<i>Calamagrostis rubescens</i>
Quackgrass*	<i>Agropyron repens</i>
Ragweed, common	<i>Ambrosia elatior</i>
Ryegrass, Italian (annual)	<i>Lolium multiflorum</i>
Ryegrass, perennial*	<i>Lolium perenne</i>
Smartweed, Pennsylvania	<i>Polygonum pensylvanicum</i>
Squawcarpet	<i>Ceanothus prostrates</i>
Thistle, Canada*	<i>Cirsium arvense</i>
Velvetgrass, common	<i>Holcus lanatus</i>

**For Western US site preparation, apply at 4 pounds (3 lbs ai) per acre.

WOODY PLANTS

Ash	<i>Fraxinus</i> spp.
Aspen, big tooth	<i>Populus grandidentata</i>
Aspen, trembling	<i>Populus tremuloides</i>
Birch	<i>Betula</i> spp.
Blackgum	<i>Nyssa sylvatica</i>
Cherry, black	<i>Prunus serotina</i>
Deerbrush	<i>Ceanothus integerrimus</i>
Dogwood, flowering*	<i>Cornus florida</i>
Elm	<i>Ulmus</i> spp.
Hawthorn	<i>Crataegus</i> spp.
Hazel	<i>Corylus</i> spp.
Hickory	<i>Carya</i> spp.
Honeysuckle*	<i>Lonicera</i> spp.
Manzanita, Greenleaf	<i>Arctostaphylos patula</i>
Maple, red*	<i>Acer rubrum</i>
Oaks	<i>Quercus</i> spp.
Poplar, balsam	<i>Populus balsamifera</i>
Snowbrush (varnishleaf)	<i>Ceanothus velutinus</i>
Sourwood*	<i>Oxydendrum arboretum</i>

Sweetgum	<i>Liquidambar</i> spp.
Willows	<i>Salix</i> spp.

*Suppression is a visible reduction in plant competition (reduced population and/or vigor) as compared to an untreated area. Degree of suppression will vary with rate applied, size of plants at application and environmental conditions following treatment. Species indicated above, especially resprouts of these species, may require a follow up treatment for acceptable control. Burning, as a follow up treatment, will enhance control of resprouts.

Within several weeks after Tide Hexazinone 75 WDG activation by rainfall, affected vegetation may be burned, if desired. This burn may further enhance control of vegetation. Burn the vegetation only after any residual stand is completely defoliated, at least twice, allowing for sufficient root uptake of Tide Hexazinone 75 WDG. In the West, results may take one to two years in areas of low rainfall.

SPRAY EQUIPMENT

When applied as a liquid spray using water as the carrier, apply Tide Hexazinone 75 WDG by ground equipment or by air (helicopter only).

For ground application, use enough water for thorough coverage, usually a minimum of 25 gallons per acre. For aerial applications (helicopter only), use at least 5 gallons of water per acre.

Restriction

When using a mechanically-pressurized handgun, do not exceed 1.5 pounds Tide Hexazinone 75 WDG (1.12 lbs a.i.) per gallon.

GRID APPLICATION

Mix 2 2/3 pounds of Tide Hexazinone 75 WDG (2 lbs ai) with sufficient water to make one gallon of suspension and thoroughly agitate. To maintain the Tide Hexazinone 75 WDG in suspension, use intermittent agitation.

Apply the Tide Hexazinone 75 WDG suspension directly to the soil surface in a grid pattern using an exact delivery handgun applicator. This equipment delivers a thin stream of predetermined volume. Tide Hexazinone 75 WDG must be applied during the period from hardwood bud break to early summer.

Application rate and grid pattern will depend on soil texture and woody plant composition. Use the lower rates on coarse textured soils and when the major components of the hardwoods are susceptible species. Use the high rates on fine-textured soils and where weeds identified in this label as "partial control or suppression" predominate.

Application Patterns and Rates for Tide Hexazinone 75 WDG Suspension			
	ML/Spot	Grid (Ft)	Lbs./Acre
Coarse	0.6	3x3	2 (1.5 lbs ai)
	2.0	4x4	4 (3 lbs ai)
	3.1	4x6	4 (3 lbs ai)
Medium/Fine	1.6	3x3	5.3 (3.97 lbs ai)
	2.8	4x4	5.3 (3.97 lbs ai)
	3.5	4x4	6.6 (4.95 lbs ai)
	5.2	4x6	6.6 (4.95 lbs ai)

BASAL (SOIL) SINGLE STEM TREATMENTS

Mix 2 2/3 pounds of Tide Hexazinone 75 WDG (2 lbs ai) with sufficient water to make one gallon of suspension and thoroughly agitate. Apply the Tide Hexazinone 75 WDG suspension with an exact-delivery handgun applicator. This equipment delivers a thin stream of predetermined volume when triggered. Apply the Tide Hexazinone 75 WDG suspension at the rate of 2 to 4 ml for each inch of stem diameter at breast height. Direct the treatment to the soil within 3 feet of the root collar of woody plants to be controlled.

For multi-stemmed and low-growing brush that have stem diameters that are difficult to determine, apply the Tide Hexazinone 75 WDG suspension at the rate of 2 to 4 ml per 3 feet of canopy width. For tall, slender (columnar) brush types, apply 4 to 8 ml per 3 feet of height. Base the rate on whichever canopy dimension is greater (width or height). Apply the lower volumes for coarse textured soils or soils with low organic matter and the higher volumes for fine textured soils or soils with high organic matter.

When treating brush that requires more than a single delivery of the Tide Hexazinone 75 WDG suspension, apply subsequent deliveries equally spaced around the target plant. If treating brush on sloping sites, apply most of the suspension on the uphill side of the stem. If treating resprouts from brush disturbed by cutting or other mechanical methods, the rate of application must be proportional to the original tree size, not just the size of sprout regrowth.

USE PRECAUTIONS AND RESTRICTIONS

SITE PREPARATION

- Where burning is desired, burn the vegetation only after any residual brush has completely defoliated, at least twice, allowing for sufficient root uptake of Tide Hexazinone 75 WDG.
- Following harvest, allow sufficient time for stumps and injured trees to adequately resprout before applying Tide Hexazinone 75 WDG.

FORESTRY – RELEASE

Tide Hexazinone 75 WDG is labeled for conifer release where the following species are grown:

EASTERN US AND LAKE STATES

Fir, balsam	<i>Abies balsamea</i>
Pine, loblolly	<i>Pinus taeda</i>
Pine, longleaf	<i>Pinus palustris</i>
Pine, red	<i>Pinus resinosa</i>
Pine, shortleaf	<i>Pinus echinata</i>
Pine, slash	<i>Pinus elliotii</i>
Pine, Virginia	<i>Pinus virginiana</i>
Spruce, black	<i>Picea mariana</i>
Spruce, Norway	<i>Picea abies</i>
Spruce, red	<i>Picea rubens</i>
Spruce, white	<i>Picea glauca</i>

WESTERN US

Fir, Douglas	<i>Pseudotsuga menziesii</i>
Fir, grand	<i>Abies grandis</i>
Fir, Noble	<i>Abies procera</i>
Fir, white	<i>Abies concolor</i>
Hemlock, Western	<i>Tsuga heterophylla</i>
Pine, Jeffrey	<i>Pinus jeffreyi</i>
Pine, lodgepole	<i>Pinus contorta</i>
Pine, ponderosa	<i>Pinus ponderosa</i>
Spruce, blue	<i>Picea pungens</i>
Spruce, Engelman	<i>Picea engelmannii</i>
Spruce, Sitka	<i>Picea sitchensis</i>

APPLICATION INFORMATION

EASTERN US

Apply Tide Hexazinone 75 WDG from early spring to early summer after hardwoods have broken bud and before full leaf expansion.

Applications made over the top of pines may result in excessive pine injury under conditions of high humidity and temperature (80 degrees F).

WESTERN US

Rainbelt (areas of high spring rainfall): For best results, apply in late winter or spring when brush is actively growing, but prior to conifer budbreak. Dormant trees are less susceptible to injury. Applications where the spray comes into direct contact with conifers after dormancy break in the spring or before the final resting bud has hardened in the fall may severely injure or kill the trees.

Snowbelt (areas of low spring rainfall): For best results, apply in the fall before soil freezes and after final resting bud has hardened on the conifers. Or, make spring applications after snow cover melts in anticipation of rainfall prior to conifer bud break. Brush control results from spring treatments will be dependent on sufficient rainfall following application to activate Tide Hexazinone 75 WDG.

USE RATES

The rates listed below are for broadcast application. Do not use more than one application of Tide Hexazinone 75 WDG per year. Use the higher rate range for harder to control* (suppression) species in the **PLANTS CONTROLLED** listings of the Site Prep and Release sections.

EASTERN US

Crop Species	Soil Description	Tide Hexazinone 75 WDG (Lbs./Acre) Established Trees
Loblolly pine Longleaf pine Shortleaf pine Virginia pine Slash pine	Loamy sand, sandy loam	1 1/3 – 2 (1 – 1.5 lbs ai)
	Loam, silt loam, silt, sandy clay loam	1 1/3 – 2 2/3 (1 – 2 lbs ai)
	Silty clay loam, clay loam, sandy clay, silty clay, clay	3 – 4 (2.25 – 3 lbs ai)
Red pine	Loamy sand, sandy loam	1 1/3 – 2 2/3 (1lb – 2 lbs ai)
	Loam, silt loam, silt, sandy clay loam	2 2/3 – 4 (2 – 3 lbs ai)
	Silty clay loam, clay loam, sandy clay, silty clay, clay	4 – 5 1/3 (2 – 4 lbs ai)

Established Trees

- 4 years of age from transplanting on coarse-textured soils
- 3 years of age from transplanting on medium-textured soils
- 2 years of age from transplanting for Red Pine

WESTERN US

Application rates by soil type for Tide Hexazinone 75 WDG in the following western conifers: Blue spruce, Douglas fir, Engelman spruce, Grand fir, Jeffrey pine, Lodgepole pine, Noble fir, Ponderosa pine, Sitka spruce, Western hemlock and White fir.

Soil Description	Tide Hexazinone 75 WDG (Lbs./Acre)
Loamy sand, sandy loam	1 1/3 – 3 (1 – 2.25 lbs ai)
Loam, silt loam, sandy clay loam	2 2/3 – 4 (2 – 3 lbs ai)
Silt, silty clay loam, clay loam, sandy clay, silty clay, clay	3 – 4 (2.25 – 3 lbs ai)

For first year plantings utilizing bare root stock, treat only transplant stock that is 2 years old (2-0, 1-1) or more, except (1-0) for Ponderosa and Jeffrey pines. Apply Tide Hexazinone 75 WDG only if rainfall has settled the soil around the base and root systems of the transplants.

BRUSH CONTROLLED

Tide Hexazinone 75 WDG is labeled for the control or suppression of the following species in conifer release sites:

Ash	<i>Fraxinus</i> spp.
Aspen, big tooth	<i>Populus grandidentata</i>
Aspen, trembling	<i>Populus tremuloides</i>
Birch	<i>Betula</i> spp.
Elder, box	<i>Acer negundo</i>
Brambles	<i>Rubus</i> spp.
Cherry, black	<i>Prunus serotina</i>
Cherry, pin	<i>Prunus pensylvanica</i>
Deerbrush	<i>Ceanothus integerrimus</i>
Dogwood, flowering*	<i>Cornus florida</i>
Elm	<i>Ulmus</i> spp.
Hawthorn	<i>Crataegus</i> spp.
Hazel	<i>Corylus</i> spp.
Honeysuckle*	<i>Lonicera</i> spp.
Manzanita, Greenleaf	<i>Arctostaphylos patula</i>
Maple, red*	<i>Acer rubrum</i>
Oaks	<i>Quercus</i> spp.
Poplar, balsam	<i>Populus balsamifera</i>
Snowbrush (varnishleaf)	<i>Ceanothus velutinus</i>
Sourwood*	<i>Oxydendrum arboretum</i>
Sweetgum	<i>Liquidambar</i> spp.
Willows	<i>Salix</i> spp.

* Suppression – a visible reduction in plant population and/or plant vigor as compared to an untreated area and generally not accepted as control.

In addition to brush controlled, herbaceous species listed in **Weeds Controlled** section of **Release-Herbaceous Weed Control** may be controlled with these applications.

SPRAY EQUIPMENT

When applied as a liquid spray using water as the carrier, apply Tide Hexazinone 75 WDG by ground equipment or by air (helicopter only).

For ground applications, use sufficient spray volume for thorough and uniform coverage of the site to be treated, usually a minimum of 25 gallons per acre. For aerial applications (helicopter only), use a minimum of 5 gallons per acre.

Restriction

When using a mechanically-pressurized handgun, do not exceed 1.5 pounds Tide Hexazinone 75 WDG (1.12 lbs a.i.) per gallon.

GRID APPLICATION

Mix 2 2/3 pounds of Tide Hexazinone 75 WDG (2 lbs ai) with sufficient water to make one gallon of suspension and thoroughly agitate. To maintain the Tide Hexazinone 75 WDG in suspension, use intermittent agitation.

Apply the Tide Hexazinone 75 WDG suspension directly to the soil surface in a grid pattern using an exact delivery handgun applicator. This equipment delivers a thin stream of predetermined volume. Tide Hexazinone 75 WDG must be applied during the period from hardwood bud break to early summer.

Application rate and grid pattern will depend on soil texture and woody plant composition. Use the lower rates on coarse textured soils and when the major component of the hardwoods are susceptible species. Use the high rates on fine-textured soils and where weeds identified in this label as "partial control or suppression" predominate.

Application Patterns and Rates for Tide Hexazinone 75 WDG Suspension			
	ML/Spot	Grid (Ft)	Lb/Acre
Coarse	0.5	3x4	1.3* (0.975 lb ai)
	1.2	3x6	2 (1.5 lbs ai)
	2.1	4x6	2.6 (1.95 lbs ai)
Medium/Fine	1.2	3x3	4 (3 lbs ai)
	2.3	3x6	4 (3 lbs ai)
	1.6	3x3	5.3 (3.97 lbs ai)
	3.1	3x6	5.3 (3.97 lbs ai)

* Use on deep sands with pines four years or more of age.

BASAL (SOIL) SINGLE STEM TREATMENT

Mix 2 2/3 pounds of Tide Hexazinone 75 WDG (2 lbs ai) with sufficient water to make one gallon of suspension and thoroughly agitate. Apply the Tide Hexazinone 75 WDG suspension with an exact-delivery handgun applicator. This equipment delivers a thin stream of predetermined volume when triggered. Apply the Tide Hexazinone 75 WDG suspension at the rate of 2 to 4 ml for each inch of stem diameter at breast height. Direct the treatment to the soil within 3 feet of the root collar of woody plants to be controlled.

For multi-stemmed and low-growing brush that have stem diameters that are difficult to determine, apply the Tide Hexazinone 75 WDG suspension at the rate of 2 to 4 ml per 3 feet of canopy width. For tall, slender (columnar) brush types, apply 4 to 8 ml per 3 feet of height. Base the rate on whichever canopy dimension is greater (width or height). Apply the lower volumes for coarse textured soils or soils with low organic matter and the higher volumes for fine textured soils or high organic matter soils.

When treating brush that requires more than a single delivery of the Tide Hexazinone 75 WDG suspension, apply subsequent deliveries equally spaced around the target plant. If treating brush on sloping sites, apply most of the suspension on the uphill side of the stem. If treating resprouts from brush disturbed by cutting or other mechanical methods, the rate of application must be proportional to the original tree size, not just the size of sprout regrowth.

USE PRECAUTIONS AND RESTRICTIONS RELEASE – GRID & SINGLE STEM

- Application of Tide Hexazinone 75 WDG basal soil spot treatments closer than 36 inches to conifer seedlings in their first season or directly up slope from these seedlings may result in injury or mortality.
- Use Tide Hexazinone 75 WDG on seedlings in their first or fourth year and older. Injury may result from use on two and three year old seedlings where root growth is extensive but hardiness is lacking.
-

RELEASE HERBACEOUS WEED CONTROL

Tide Hexazinone 75 WDG is labeled for controlling herbaceous weeds where these pine species are grown:

EASTERN US

Loblolly pine	Slash pine
Longleaf pine	Red pine

WESTERN US

Blue spruce	Noble fir
Douglas fir	Ponderosa pine
Engleman spruce	Sitka spruce
Grand fir	Western hemlock
Jeffrey pine	White fir
Lodgepole pine	

APPLICATION INFORMATION**EASTERN US**

Apply Tide Hexazinone 75 WDG as a broadcast or banded spray in the spring prior to conifer bud break to lessen conifer injury potential.

WESTERN US

Rainbelt (areas of high spring rainfall): For best results, apply as a broadcast or banded spray in the late winter or spring when weeds are actively growing, but prior to conifer budbreak. If application is made after conifer bud break, use directional spray equipment to prevent contact with conifer foliage, as injury may result.

Snowbelt (areas of low spring rainfall): For best results, apply as a broadcast or banded spray in the fall before soil freezes and after the final resting bud has hardened on the conifers. Or, make spring applications after snow cover melts in anticipation of rainfall prior to conifer bud break. Weed control results from spring treatments will be dependent on sufficient rainfall following application to activate Tide Hexazinone 75 WDG.

USE RATES

The rates listed below are for broadcast application. For band application, use proportionately less. For example, use $\frac{1}{2}$ of the broadcast rates when treating a 3-foot band where row spacing is 6 feet. Use the higher rate range for harder to control* (suppression) weeds in the table below.

EASTERN US

Soil Description	Tide Hexazinone 75 WDG (Lbs./Acre)	
	First Year Plantings	Established Trees
Loamy sand, sandy loam (50 – 85% sand)	1 $\frac{1}{3}$ (1 lb ai)	1 $\frac{1}{3}$ – 1 $\frac{2}{3}$ (1 – 1.25 lbs ai)
Loam, silt loam, silt, sandy clay loam	1 $\frac{1}{3}$ – 1 $\frac{1}{2}$ (1 – 1.12 lbs ai)	1 $\frac{2}{3}$ – 2 $\frac{1}{3}$ (1.25 – 1.75 lbs ai)
Silty clay loam, clay loam, sandy clay, silty clay, clay	1 $\frac{1}{2}$ - 1 $\frac{8}{10}$ (1.12 – 1.35 lbs ai)	2 $\frac{1}{3}$ – 2 $\frac{2}{3}$ (1.75 – 2 lbs ai)

Red pine only- Refer to labeled rates in the **FORESTRY RELEASE – Use Rates Eastern US** section of the label.

WESTERN US

Refer to labeled rates in the **FORESTRY RELEASE – Use Rates Western US** section of the label.

WEEDS CONTROLLED

Tide Hexazinone 75 WDG is labeled for the control or suppression of the following species in release sites:

Asters	<i>Aster</i> spp.
Aster, heath*	<i>Aster ericoides</i>
Barnyardgrass	<i>Echinochloa crus-galli</i>
Bentgrass	<i>Agrostis</i> spp.
Bluegrass, annual	<i>Poa annua</i>
Brackenfern	<i>Pteridium aquilinum</i>
Bromegrass	<i>Bromus</i> spp.
Carrot, wild	<i>Daucus carota</i>
Crabgrass*	<i>Digitaria</i> spp.
Daisy, oxeye	<i>Chrysanthemum leucanthemum</i>
Dandelion, common*	<i>Taraxacum officinale</i>
Dandelion, false* (spotted catsear)	<i>Hypochaeris radicata</i>
Dock, curly*	<i>Rumex crispus</i>
Fescue*	<i>Festuca</i> spp.
Fireweed* (willowweed)	<i>Epilobium angustifolium</i>
Fleabane	<i>Conyza</i> spp.
Foxtail	<i>Setaria</i> spp.
Goldenrod*	<i>Solidago</i> spp.
Groundsel, common	<i>Senecio vulgaris</i>
Horseweed/marestail	<i>Conyza canadensis</i>
Orchardgrass*	<i>Dactylis glomerata</i>
Panicums	<i>Panicum</i> spp.
Pinegrass	<i>Calamagrostis rubescens</i>
Ragweed, common	<i>Ambrosia elatior</i>
Ryegrass, Italian (annual)	<i>Lolium multiflorum</i>
Ryegrass, perennial*	<i>Lolium perenne</i>
Smartweed, Pennsylvania	<i>Polygonum pensylvanicum</i>
Squawcarpet	<i>Ceanothus prostrates</i>
Velvetgrass, common	<i>Holcus lanatus</i>

* Suppression – a visible reduction in plant population and/or plant vigor as compared to an untreated area and generally not accepted as control.

FORESTRY

IMPREGNATION ON DRY BULK FERTILIZER

Tide Hexazinone 75 WDG is labeled for impregnating or coating dry bulk fertilizer to be applied on forested sites for the establishment or release of conifer plantations (except longleaf pine) as specified on this label.

PLANTS CONTROLLED

Fertilizer impregnated with Tide Hexazinone 75 WDG is labeled for the control and suppression of the weeds and brush identified for the specific applications on this label. Consult the appropriate segment of this label to determine the appropriate rate of Tide Hexazinone 75 WDG to be applied per acre. Apply this amount of Tide Hexazinone 75 WDG to the volume of fertilizer to be applied per acre.

IMPREGNATION EQUIPMENT

To impregnate or coat the fertilizer use a system consisting of conveyor or closed drum used to blend dry bulk fertilizer.

IMPREGNATION INSTRUCTIONS

To impregnate dry bulk fertilizer with Tide Hexazinone 75 WDG, mix the amount as prescribed above in a sufficient quantity of water to uniformly coat the desired amount of fertilizer. Suspensions of Tide Hexazinone 75 WDG will require thorough agitation.

Direct the spray nozzles of the impregnation equipment to deliver a fine spray of the mixture toward the fertilizer for thorough coverage while avoiding contact with mixing equipment. The use of a spray pattern indicator will help to visually determine the uniformity of impregnation.

Uniform impregnation of dry bulk fertilizer can vary. If absorption of the spray is not adequate, use an absorptive powder or additive, such as "Microcel E" or "HiSil 233", to produce a dry, free flowing mixture.

Apply the fertilizer as soon as possible after impregnation for optimum performance. Impregnated fertilizer may become lumpy and difficult to apply following storage.

Diammonium phosphate, potassium chloride, 16-16-16 and 24-4-4-have been successfully impregnated.

APPLICATION EQUIPMENT

Apply impregnated fertilizer by ground equipment or by air (helicopter only). Accurate calibration and patterning of the equipment is essential for uniform distribution of the impregnated fertilizer on the soil surface.

USE PRECAUTIONS AND RESTRICTIONS FORESTRY – IMPREGNATED FERTILIZER

- If fertilizer materials are excessively dusty, use a suitable additive to reduce dust prior to impregnation. Application of dusty fertilizer which has been impregnated may result in off-target drift and injury to desirable vegetation. Such drift and associated injury may be aggravated by high wind conditions.
- The dry fertilizer must be properly impregnated and uniformly applied to avoid pine injury/mortality and poor weed and brush control.
- Uniform and precise application of the impregnated fertilizer is essential for satisfactory weed and brush control and to minimize pine injury. Overlaps or skips between adjoining swaths or non-uniform distribution of impregnated fertilizer within the swath will deliver poor results and may result in pine injury or mortality.
- Do not impregnate potassium nitrate, sodium nitrate or triple super phosphate fertilizers with Tide Hexazinone 75 WDG as herbicidal action will be lost.

USE PRECAUTIONS AND RESTRICTIONS FORESTRY

- For all forestry uses application by fixed-wing aircraft is prohibited.
- When using a mechanically-pressurized handgun, reduce rate to 1.5 pounds Tide Hexazinone 75 WDG (1.12 lbs a.i.) per gallon.
- Do not use Tide Hexazinone 75 WDG in nurseries, seedbeds, or ornamental plantings.
- On tracts of land where various soil types are present and Tide Hexazinone 75 WDG rate selection is difficult, conifer damage or less-than-expected vegetation suppression may occur due to the different rates required for various soil types.
- Poor weed and brush control may result from the following:
 - Heavy duff or slash present at time of application
 - Use on poorly drained sites
 - Applications made when the soil is saturated with water and rain is imminent within 24 hours
 - Applications to soils high in organic matter (greater than 5%)
- Following harvest, allow stumps and injured trees sufficient time to adequately resprout before applying Tide Hexazinone 75 WDG.
- Where burning is desired, burn vegetation after any brush has completely defoliated, at least twice, allowing for sufficient root uptake of Tide Hexazinone 75 WDG.
- Do not use Tide Hexazinone 75 WDG on frozen soils; use in spring after snow melt.
- Do not add a surfactant in applications over the top of conifers.
- Weed control results from spring applications depend on sufficient moisture to activate Tide Hexazinone 75 WDG.
- When applying Tide Hexazinone 75 WDG after transplanting, wait until rainfall has settled the soil around the base and root systems of the transplants before making the treatment.
- Crop injury may occur when Tide Hexazinone 75 WDG is used:
 - On trees that show poor vigor, insect damage, disease, winter injury, or other stress conditions
 - On any soil containing less than 1% organic matter
 - On loamy sand or sandy loam with less than 2% organic matter, except Jeffrey pine and Ponderosa pine
 - On conifer foliage after conifer bud break
 - On gravelly or rocky soils, exposed subsoils, clay knobs, sand, or sandy soil with 85% or more sand.
 - Livestock may be grazed immediately following a broadcast application of Tide Hexazinone 75 WDG at rates of 1.5 pounds (1.12 lbs ai) per acre or less, and treated vegetation may be cut, dried, and fed after 38 days.
 - Do not cut treated vegetation for feed, or graze livestock on treated areas for 60 days following application of Tide Hexazinone 75 WDG at broadcast rates exceeding 1.5 pounds (1.12 lbs ai) per acre.

YELLOW POPLAR PLANTINGS

Tide Hexazinone 75 WDG is labeled for the control of herbaceous weeds in the establishment of yellow poplar plantations. Make applications over the top of planted seedlings after the soil has settled around the root systems but before the seedlings have broken dormancy (bud break). Make a subsequent application before dormancy break in the Spring of the second year. **USE RATES:** Use the rate range specified in the “**RELEASE – HERBACEOUS WEED CONTROL**” section for pine plantations – **Eastern US**.

For ground application, use sufficient spray volume for uniform and thorough coverage of the site to be sprayed, usually a minimum of 25 gallons per acre. For aerial applications, use a minimum of 5 gallons of water per acre. For broader spectrum control, tank-mix Tide Hexazinone 75 WDG with DuPont™ ESCORT® XP herbicide (or other metsulfuron formulations approved for this use). Add ESCORT® XP at a rate of ½ ounce per acre with the prescribed rate of Tide Hexazinone 75 WDG.

USE PRECAUTIONS AND RESTRICTIONS

YELLOW POPLAR PLANTINGS

- Applications of Tide Hexazinone 75 WDG and tank mixes of Tide Hexazinone 75 WDG and DuPont™ ESCORT® XP (or other metsulfuron formulations approved for this use) made to yellow poplar seedlings that are suffering from loss of vigor caused by insects, disease, drought, winter damage, animal damage, excessive soil moisture, planting shock or other stresses may injure or kill the seedlings.
- Applications of Tide Hexazinone 75 WDG and tank mixes of Tide Hexazinone 75 WDG and ESCORT® XP (or other metsulfuron formulations approved for this use) must only be made after adequate rainfall has closed the planting slit and settled the soil around the roots following transplanting.
- The use of surfactant with Tide Hexazinone 75 WDG is not advised for applications made of the tops of seedlings.
- Careful consideration must be given by an experienced and knowledgeable forester to ensure the specific growth requirements of yellow poplar will be provided by the selected planting site. Treatment of yellow poplar planted on a site inadequate to meet its requirements may injure or kill the seedlings.

PASTURE/RANGELAND

Tide Hexazinone 75 WDG is labeled for control of brush and weeds in pasture.

BERMUDAGRASS/BAHIAGRASS

Tide Hexazinone 75 WDG is labeled for control of smutgrass and other weeds in established stands of bermudagrass and bahiagrass in pastures/rangelands.

APPLICATION INFORMATION

Make a single application of Tide Hexazinone 75 WDG per year when weeds are actively growing.

WEEDS CONTROLLED – USE RATES

Tide Hexazinone 75 WDG effectively controls the following weeds at the rates shown in pastures. Use a lower rate on coarse-textured soils (sand to sandy loam). Use the higher rate on fine-textured soils (clay loam to clay) and on soils high in organic matter.

9/10 – 1 ½ Lbs./Acre (0.67 – 1.12 lbs ai/acre)	
Barley, little	<i>Hordeum pusillum</i>
Barnyardgrass	<i>Echinochloa crus-galli</i>
Dogfennel	<i>Eupatorium capillifolium</i>
Fescue	<i>Festuca</i> spp.
Lespedeza	<i>Lespedeza cuneata</i>
Oxalis	<i>Oxalis</i> spp.
Passionflower, maypop	<i>Passiflora incarnate</i>
Pepperweed, Virginia	<i>Lepidium virginicum</i>
Pigweed	<i>Amaranthus</i> spp.
Smutgrass*	<i>Sporobolus indicus</i>

*Suppression may result with some of the giant (larger) smutgrass species.

Suppression – a visible reduction in plant population and/or plant vigor as compared to an untreated area and generally not accepted as control.

SPRAY EQUIPMENT

Apply Tide Hexazinone 75 WDG uniformly over the desired area using ground equipment only.

For ground application, use enough water for thorough coverage usually a minimum of 25 gallons per acre. The use of a surfactant may increase the potential for bermudagrass or bahiagrass injury.

USE PRECAUTIONS AND RESTRICTIONS

BERMUDAGRASS/BAHIAGRASS

- For bermudagrass grown in the states of ID, OR, UT, or WA, determine the suitability of using Tide Hexazinone 75 WDG by treating a small area at a labeled application rate prior to treating larger areas. The smaller treated area must be observed for any signs of herbicidal injury during 60 days of normal growing conditions to determine if the treatment is safe to bermudagrass.
- Use Tide Hexazinone 75 WDG only in stands of bermudagrass and bahiagrass established for at least one year. Do not treat newly sprigged or sodded areas.
- Some temporary discoloration of the bermudagrass or bahiagrass may occur after application.
- Treatment of mixed pastures containing forage species other than bermudagrass or bahiagrass may result in injury or mortality to the other forage species.
- Injury may result when desirable grasses are under stress from drought, insects, disease, cold temperature, or poor fertility.
- Injury to or loss of desirable trees or other plants may result if Tide Hexazinone 75 WDG is applied or if equipment is drained or flushed on or near desirable trees or other plants, on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Severe crop injury may occur if applications are made on gravelly or rocky soils, thinly covered subsoils, or soils with less than 1% organic matter.
- Livestock may be grazed immediately following a broadcast application of Tide Hexazinone 75 WDG at rates of 1.5 pounds (1.12 lbs ai) per acre or less, and treated vegetation may be cut, dried and fed after 38 days.
- Do not apply to residential areas/lawns or golf courses.

PASTURE/RANGELAND BRUSH CONTROL

Apply Tide Hexazinone 75 WDG either by broadcast or as a basal-soil treatment for the control of undesirable brush in pasture or rangeland.

APPLICATION INFORMATION

Apply Tide Hexazinone 75 WDG from late winter through summer, pre-budbreak until new growth hardens off.

In areas where the soil remains frozen during the winter and spring rains are usually inadequate for soil activation, make a fall or winter treatment before the soil freezes.

For broadcast rates needed to control the species below, see the **Forestry – Release, Use Rates** section.

BRUSH CONTROLLED

Tide Hexazinone 75 WDG is labeled for the control or suppression of the following brush species in pasture and rangeland:

Alder	<i>Alnus</i> spp.
Ash	<i>Fraxinus</i> spp.
Aspen	<i>Populus</i> spp.
Birch	<i>Betula</i> spp.
Blackgum	<i>Nyssa sylvatica</i>
Bay, sweet	<i>Magnolia virginiana</i>
Catclaw acacia	<i>Acacia greggii</i>
Cedar, Eastern red	<i>Juniperus virginiana</i>
Cherry, black	<i>Prunus serotina</i>
Chinaberry*	<i>Melia azedarach</i>
Deerbrush	<i>Ceanothus integerrimus</i>
Dogwood, flowering*	<i>Cornus florida</i>
Elm, American	<i>Ulmus Americana</i>

Elm, Chinese	<i>Ulmus parvifolia</i>
Hackberry, common	<i>Celtis occidentalis</i>
Hawthorn	<i>Crataegus</i> spp.
Hazel	<i>Corylus</i> spp.
Hickory	<i>Carya</i> spp.
Huisache	<i>Acacia farnesiana</i>
Juniper	<i>Juniperus</i> spp.
Locust	<i>Robinia</i> spp.
Lotebush	<i>Ziziphus obtusifolia</i>
Manzanita, Greenleaf	<i>Arctostaphylos patula</i>
Maple, red	<i>Acer rubrum</i>
Mesquite	<i>Prosopis glandulosa</i>
Mulberry	<i>Morus</i> spp.
Oaks	<i>Quercus</i> spp.
Osage-orange	<i>Maclura pomifera</i>
Persimmon	<i>Diospyros</i> spp.
Plum, wild	<i>Prunus munsoniana</i>
Poplar, balsam	<i>Populus balsamifera</i>
Poplar, yellow	<i>Liriodendron tulipifera</i>
Privet	<i>Ligustrum</i> spp.
Rose, multiflora	<i>Rosa multiflora</i>
Sassafras*	<i>Sassafras albidum</i>
Soapweed, small (yucca)	<i>Yucca glauca</i>
Snowbrush (varnishleaf)	<i>Ceanothus velutinus</i>
Sourwood	<i>Oxydendrum arboretum</i>
Sumac	<i>Rhus</i> spp.
Sweetgum	<i>Liquidambar</i> spp.
Tallow, Chinese	<i>Sapium sebiferum</i>
Waxmyrtle	<i>Myrica cerifera</i>
Whitebrush	<i>Aloysia gratissima</i>
Willow	<i>Salix</i> spp.

*Suppression – a visible reduction in plant population and/or plant vigor as compared to an untreated area and generally not accepted as control.

SPRAY EQUIPMENT AND APPLICATION TECHNIQUES

Basal (Soil) – Mix 2 2/3 pounds of Tide Hexazinone 75 WDG (2 lbs ai) with sufficient water to make one gallon of suspension and thoroughly agitate. Apply the Tide Hexazinone 75 WDG suspension with an exact-delivery handgun applicator. This equipment delivers a thin stream of predetermined volume when triggered. Apply the Tide Hexazinone 75 WDG suspension at the rate of 2 to 4 ml for each inch of stem diameter at breast height. Direct the treatment to soil within 3 inches of the root collar of woody plants to be controlled. When treating large stems and when more than one delivery of the Tide Hexazinone 75 WDG suspension is needed per stem, make applications on opposite sides of the stem. Do not apply more than 1/3 gallon of the Tide Hexazinone 75 WDG suspension per acre per year. To maintain the Tide Hexazinone 75 WDG in suspension, use intermittent agitation.

USE PRECAUTIONS AND RESTRICTIONS PASTURE/RANGELAND

- Injury to or loss of desirable trees or other plants may result if Tide Hexazinone 75 WDG is applied or if equipment is drained or flushed on or near desirable trees or other plants, on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Poor weed and brush control may result from the following:
 - Use on poorly drained sites
 - Applications made when the soil is saturated with water and rain is imminent within 24 hours
 - Applications to soils high in organic matter (greater than 5%)

- Following mechanical cutting or clearing, allow stumps and injured trees sufficient time to adequately resprout before applying Tide Hexazinone 75 WDG.
- Do not use Tide Hexazinone 75 WDG on frozen soils
- Leave treated soil undisturbed to reduce the potential for Tide Hexazinone 75 WDG movement by soil erosion due to wind or water.
- Weed and brush control results depend on sufficient moisture to activate Tide Hexazinone 75 WDG.
- When Tide Hexazinone 75 WDG is applied as a basal soil treatment, there is no restriction on grazing by domestic animals nor on cutting surrounding vegetation for forage or hay.
- Livestock may be grazed immediately following a broadcast application of Tide Hexazinone 75 WDG at rates of 1.5 pounds (1.12 lbs ai) per acre or less, and treated vegetation may be cut, dried, and fed after 38 days.
- Do not cut treated vegetation for feed, or graze livestock on treated areas for 60 days following application of Tide Hexazinone 75 WDG at broadcast rates exceeding 1.5 pounds (1.12 lbs ai) per acre.

NON-AGRICULTURAL USES

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Use on non-crop sites including industrial turfgrasses are not within the scope of the Worker Protection Standard.

When applied as a spray do not enter or allow worker entry into treated areas until sprays have dried.

Respirator fit testing, medical qualification, and training

Using a program that conforms to OSHA's requirements (see 29 CFR Part 1910.134), employers must verify that any handler who uses a respirator is:

- Fit-tested and fit-checked,
- Trained, and
- Examined by a qualified medical practitioner to ensure physical ability to safely wear the style of respirator to be worn. A qualified medical practitioner is a physician or other licensed health care professional who will evaluate the ability of a worker to wear a respirator. The initial evaluation consists of a questionnaire that asks about medical conditions (such as heart condition) that would be problematic for respirator use. If concerns are identified, then additional evaluations, such as physical exam, might be necessary. The initial evaluation must be done before respirator use begins. Handlers must be reexamined by a qualified medical practitioner if their health status or respirator style or use-conditions change.
- Upon request by local/state/federal/tribal enforcement personnel, employers must provide documentation demonstrating how they have complied with these requirements.

APPLICATION INFORMATION

Tide Hexazinone 75 WDG is labeled for general weed and brush control as follows: uncultivated nonagricultural areas (such as, airports, highway, railroad and utility right-of way, sewage disposal areas); uncultivated agricultural areas (non-crop producing, which includes: farmyards, fuel storage areas, fence rows, barrier strips); industrial site (outdoor, such as, lumberyards, pipeline and tank farms).

RESTRICTION: Do not apply more than 2.93 lbs. Tide Hexazinone 75 WDG (2.2 lbs. a.i.) per acre via aerial application to uncultivated agricultural areas (non-crop producing), which includes: farmyards, fuel storage areas, fence rows, barrier strips and rights-of-way.

NON-CROP SITES

Tide Hexazinone 75 WDG is labeled for control of many annual, biennial, and perennial weeds in non-crop sites.

APPLICATION INFORMATION

Apply Tide Hexazinone 75 WDG as a preemergence or postemergence spray when weeds are actively germinating or growing.

WEEDS CONTROLLED – USE RATE

Tide Hexazinone 75 WDG effectively controls the following weeds when applied at the use rates shown in industrial sites. When applied at lower rates, Tide Hexazinone 75 WDG provides short-term control of the weeds listed; when applied at higher rates, weed control is increased and extended.

Use lower rate on coarse-textured soils (sand to sandy loam). Use the higher rate on fine-textured soils (clay loam to clay) and on soils high in organic matter.

2 2/3 -6 2/3 Lbs./Acre (2 – 5 lbs ai/acre)	
Barnyardgrass	<i>Echinochloa crus-galli</i>
Bindweed, field*	<i>Convolvulus arvensis</i>
Bouncingbet*	<i>Saponaria officinalis</i>
Bromegrass	<i>Bromus</i> spp.
Buffalograss*	<i>Buchloe dactyloides</i>
Burdock	<i>Arctium</i> spp.
Cocklebur	<i>Xanthium</i> spp.
Crabgrass	<i>Digitaria</i> spp.
Crown vetch	<i>Coronilla varia</i>
Curly dock*	<i>Rumex crispus</i>
Dandelion, common*	<i>Taraxacum officinale</i>
Dandelion, false* (spotted catsear)	<i>Hypochaeris radicata</i>
Dogbane*	<i>Apocynum cannabinum</i>
Fiddleneck, tarweed	<i>Amsinckia lycopsoides</i>
Filaree	<i>Erodium</i> spp.
Fleabane, flax-leaved	<i>Conyza bonariensis</i>
Goatsbeard vine (sweet briar)	<i>Aruncus sylvestris</i>
Goldenrod	<i>Solidago</i> spp.
Horseweed/marestail	<i>Conyza canadensis</i>
Lespedeza	<i>Lespedeza cuneata</i>
Milkweed, common*	<i>Asclepias syriaca</i>
Mustard, wild	<i>Sinapis arvensis</i>
Nutsedge*	<i>Cyperus</i> spp.
Oats, wild*	<i>Avena fatua</i>
Orchardgrass*	<i>Dactylis glomerata</i>
Orchardgrass (seedling)	<i>Dactylis glomerata</i>
Oxalis	<i>Oxalis</i> spp.
Paragrass	<i>Panicum purpurascens</i>
Parsnip, wild	<i>Pastinaca sativa</i>
Pigweed	<i>Amaranthus</i> spp.
Purslane, common	<i>Portulaca oleracea</i>
Quackgrass	<i>Agropyron repens</i>
Ryegrass, Italian (annual)	<i>Lolium multiflorum</i>
Smartweed	<i>Polygonum</i> spp.
Spurge	<i>Euphorbia</i> spp.
Star thistle	<i>Centaurea</i> spp.
Trumpetcreeper*	<i>Campsis radicans</i>

8 - 10 2/3 Lbs./acre (6 – 8 lbs ai/acre)	
Aster, heath	<i>Aster ericoides</i>
Bahiagrass*	<i>Paspalum notatum</i>
Bermudagrass*	<i>Cynodon dactylon</i>
Blackberry	<i>Rubus</i> spp.
Bluegrass	<i>Poa</i> spp.
Broomsedge	<i>Andropogon virginicus</i>
Camphorweed	<i>Heterotheca subaxillaris</i>
Canada thistle*	<i>Cirsium arvense</i>
Carrot, wild	<i>Daucus carota</i>
Chickweed	<i>Stellaria media</i>
Clovers	<i>Trifolium</i> spp.
Dewberry	<i>Rubus trivialis</i>
Dogfennel	<i>Eupatorium capillifolium</i>
Fescue*	<i>Festuca</i> spp.
Fingergrass	<i>Digitaria ciliaris</i>
Foxtail	<i>Setaria</i> spp.
Guineagrass	<i>Panicum maximum</i>
Honeysuckle	<i>Lonicera</i> spp.
Horseweed/marestail	<i>Conyza canadensis</i>
Lantana	<i>Lantana camara</i>
Lettuce, prickly	<i>Lactuca serriola</i>
Natalgrass (red top)	<i>Rhynchelytrum repens</i>
Plantain	<i>Plantago</i> spp.
Ragweed, common	<i>Ambrosia elatior</i>
Smutgrass**	<i>Sporobolus indicus</i>
Spanishneedles	<i>Bidens bipinnata</i>
Vaseygrass	<i>Paspalum urvillei</i>

* Suppression – a visible reduction in plant population and/or plant vigor as compared to an untreated area and generally not accepted as control.

** Suppression may result with some of the giant (larger) smutgrass species.

SPECIFIC WEED PROBLEMS

Control of Canada Thistle in Crown Vetch – Tide Hexazinone 75 WDG is labeled for control of Canada thistle in established stands of crown vetch on noncrop sites. Make a single application of 1 – 1 2/3 lbs. of Tide Hexazinone 75 WDG (0.75 – 1.25 lbs ai) from late spring through mid-summer, when thistle is actively growing prior to flowering. Do not use a surfactant. Some discoloration of the crown vetch foliage may occur after application.

SPRAY EQUIPMENT

Apply Tide Hexazinone 75 WDG uniformly over the desired area using ground equipment or helicopter. Do not apply more than 8 lbs. (6 lbs ai) per acre by air.

Use enough water for thorough coverage. For ground application this is usually a minimum of 25 gallons per acre. Use higher application volumes to obtain uniform application with handgun equipment. For aerial applications (helicopter only) this is usually a minimum of 5 gallons per acre. Use higher volumes of water when water temperatures are cold or the higher rates of Tide Hexazinone 75 WDG are used.

NON-CROP BRUSH CONTROL

Tide Hexazinone 75 WDG is labeled for the control of undesirable brush in non-crop sites.

APPLICATION INFORMATION

Apply Tide Hexazinone 75 WDG from late winter through summer, pre- bud break until new growth hardens off.

In areas where soils remains frozen during the winter and spring rains are usually inadequate for soil activation, make a fall or winter treatment before the soil freezes.

BROADCAST

Apply 5 1/3 to 10 2/3 lbs. of Tide Hexazinone 75 WDG (4 – 8 lbs ai) per acre as a coarse spray by ground equipment or 5 1/3 to 8 lbs. (4 – 6 lbs ai) per acre by air (helicopter only). Use enough water for thorough coverage. For ground, equipment usually a minimum of 25 gallons per acre. For aerial equipment, usually a minimum of 10 gallons per acre. Use higher volumes of water when water temperatures are cold or the higher rates of Tide Hexazinone 75 WDG are used.

BASAL (SOIL) SINGLE STEM TREATMENT

Mix 2 2/3 pounds of Tide Hexazinone 75 WDG (2 lbs ai) with sufficient water to make one gallon of suspension and thoroughly agitate. Apply the Tide Hexazinone 75 WDG suspension with an exact-delivery handgun applicator. This equipment delivers a thin stream of predetermined volume when triggered. Apply the Tide Hexazinone 75 WDG suspension at the rate of 2 to 4 ml for each inch of stem diameter at breast height.

Direct the treatment to the soil within 3 feet of the root collar of woody plants to be controlled.

For multi-stemmed and low-growing brush that have stem diameters that are difficult to determine, apply the Tide Hexazinone 75 WDG suspension at the rate of 2 to 4 ml per 3 feet of canopy width. For tall, slender (columnar) brush types, apply 4 to 8 ml per 3 feet of height. Base the rate on whichever canopy dimension is greater (width or height).

When treating brush that requires more than a single delivery of the Tide Hexazinone 75 WDG suspension, apply subsequent deliveries equally spaced around the target plant. If treating brush on sloping sites, apply most of the suspension on the uphill side of the stem. If treating resprouts from brush disturbed by cutting or other mechanical methods, the rate of application must be proportional to the original tree size, not just the size of sprout regrowth.

LACING/STREAKING – Mix Tide Hexazinone 75 WDG with water to form a concentrated suspension. Apply 5 1/3 to 10 2/3 lbs. of Tide Hexazinone 75 WDG per acre. Adjust the application equipment to deliver a narrow or straight stream spray pattern such that the swath width on soil surface is 6 to 12 inches wide. Direct the spray at the base of the brush. Swaths or treated bands must be 2 to 4 feet apart. Apply the lower volumes for coarse textured soils or soils with low organic matter and the higher volumes for fine textured soils or soils with high organic matter.

USE RATES

Tide Hexazinone 75 WDG is labeled for the control or suppression of the following species in non-crop sites. Use lower rate on coarse-textured soils (sand to sandy loam). Use the higher rate on fine-textured soils (clay loam to clay) and on soils high in organic matter.

5 1/3 – 10 2/3 Lbs./Acre (4 – 8 lbs ai/acre)	
Alder	<i>Alnus</i> spp.
Ash	<i>Fraxinus</i> spp.
Aspen	<i>Populus</i> spp.
Birch	<i>Betula</i> spp.
Blackgum	<i>Nyssa sylvatica</i>
Bay, sweet	<i>Magnolia virginiana</i>
Catclaw acacia	<i>Acacia greggii</i>
Cedar, Eastern red	<i>Juniperus virginiana</i>
Cherry, black	<i>Prunus serotina</i>
Chinaberry*	<i>Melia azedarach</i>
Deerbrush	<i>Ceanothus integerrimus</i>

Dogwood, flowering*	<i>Cornus florida</i>
Elm, American	<i>Ulmus Americana</i>
Elm, Chinese	<i>Ulmus parvifolia</i>
Hackberry, common	<i>Celtis occidentalis</i>
Hawthorn	<i>Crataegus</i> spp.
Hazel	<i>Corylus</i> spp.
Hickory	<i>Carya</i> spp.
Huisache	<i>Acacia farnesiana</i>
Juniper	<i>Juniperus</i> spp.
Locust	<i>Robinia</i> spp.
Lotebush	<i>Ziziphus obtusifolia</i>
Manzanita, Greenleaf	<i>Arctostaphylos patula</i>
Maple, red	<i>Acer rubrum</i>
Mesquite	<i>Prosopis glandulosa</i>
Mulberry	<i>Morus</i> spp.
Oaks	<i>Quercus</i> spp.
Osage-orange	<i>Maclura pomifera</i>
Persimmon	<i>Diospyros</i> spp.
Plum, wild	<i>Prunus munsoniana</i>
Poplar, balsam	<i>Populus balsamifera</i>
Poplar, yellow	<i>Liriodendron tulipifera</i>
Privet	<i>Ligustrum</i> spp.
Rose, multiflora	<i>Rosa multiflora</i>
Sassafras*	<i>Sassafras albidum</i>
Soapweed, small (yucca)	<i>Yucca glauca</i>
Snowbrush (varnishleaf)	<i>Ceanothus velutinus</i>
Sourwood	<i>Oxydendrum arboretum</i>
Sumac	<i>Rhus</i> spp.
Sweetgum	<i>Liquidambar</i> spp.
Tallow, Chinese	<i>Sapium sebiferum</i>
Waxmyrtle	<i>Myrica cerifera</i>
Whitebrush	<i>Aloysia gratissima</i>
Willow	<i>Salix</i> spp.

* Suppression – a visible reduction in plant population and/or plant vigor as compared to an untreated area and generally not accepted as control.

INDUSTRIAL TURFGRASS

Tide Hexazinone 75 WDG is labeled for selective weed control is established stands of bermudagrass and/or bahiagrass in noncrop areas.

APPLICATION TIMING

Make a single application of Tide Hexazinone 75 WDG per year when weeds are actively growing.

WEEDS CONTROLLED – USE RATE

Tide Hexazinone 75 WDG effectively controls the following weeds at the rates shown in industrial turf (unimproved only). Use a lower rate on coarse textured soils (sand to sandy loam). Use the higher rate on fine-textured soils (clay loam to clay) and on soils high in organic matter.

9/10 – 1 ½ Lbs./Acre (0.67 – 1.12 lbs ai/acre)	
Barley, little	<i>Hordeum pusillum</i>
Barnyardgrass	<i>Echinochloa crus-galli</i>
Dogfennel	<i>Eupatorium capillifolium</i>
Fescue	<i>Festuca</i> spp.
Lespedeza	<i>Lespedeza cuneata</i>
Oxalis	<i>Oxalis</i> spp.

Passionflower, maypop	<i>Passiflora incarnate</i>
Pepperweed, Virginia	<i>Lepidium virginicum</i>
Pigweed	<i>Amaranthus</i> spp.
Smutgrass*	<i>Sporobolus indicus</i>

*Suppression may result with some of the giant (larger) smutgrass species.

Suppression – a visible reduction in plant population and/or plant vigor as compared to an untreated area and generally not accepted as control.

SPRAY EQUIPMENT

Apply Tide Hexazinone 75 WDG uniformly over the desired area using ground equipment only.

For ground application, use enough water for thorough coverage usually a minimum of 25 gallons per acre. The use of a surfactant is not advised.

USE PRECAUTIONS AND RESTRICTIONS ALL NON-CROP SITES

- For bermudagrass grown in the states of ID, OR, UT, or WA, determine the suitability of using Tide Hexazinone 75 WDG by treating a small area at a labeled application rate prior to treating larger areas. The smaller treated area must be observed for any signs of herbicidal injury during 60 days of normal growing conditions to determine if the treatment is safe to bermudagrass.
- Injury to or loss of desirable trees or other plants may result if Tide Hexazinone 75 WDG is applied or if equipment is drained or flushed on or near desirable trees or other plants, on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Application spray drift may injure desirable plants.
- Poor weed and brush control may result from the following:
 - Use on poorly drained sites
 - Applications made when the soil is saturated with water and rain is imminent within 24 hours.
 - Applications to soils high in organic matter (greater than 5%)
- Following mechanical cutting or clearing, allow stumps and injured trees sufficient time to adequately resprout before applying Tide Hexazinone 75 WDG.
- Do not use Tide Hexazinone 75 WDG on frozen soils.
- Leave treated soil undisturbed to reduce the potential for Tide Hexazinone 75 WDG movement by soil erosion due to wind or water.
- Do not use Tide Hexazinone 75 WDG on lawns, driveways, tennis courts, or other residential or recreational areas.
- Weed and brush control results from spring applications depend on sufficient moisture to activate Tide Hexazinone 75 WDG.
- Livestock may be grazed immediately following a broadcast application of Tide Hexazinone 75 WDG at rates of 1.5 pounds (1.12 lbs ai) per acre or less, and treated vegetation may be cut, dried and fed after 38 days.
- Do not cut treated vegetation for feed, or graze livestock on treated areas for 60 days following application of Tide Hexazinone 75 WDG at broadcast rates greater than 1.5 pounds (1.12 lbs ai) and up to 8 pounds (6 lbs ai) per acre.
- For Tide Hexazinone 75 WDG rates above 8 pounds (6 lbs ai) per acre, do not cut treated vegetation for forage or hay nor graze domestic animals for 1 year following application.
- There are no grazing or haying restrictions for the directed basal-soil applications of Tide Hexazinone 75 WDG.
- Use Tide Hexazinone 75 WDG only in stands of bermudagrass and bahiagrass turfgrasses established for at least one year. Do not treat newly sprigged or sodded areas.
- Some discoloration of the bermudagrass or bahiagrass may occur after application.
- Injury may result when desirable grasses are under stress from drought, insects, disease, cold temperature, or poor fertility.

- Severe turfgrass injury may occur if applications are made on gravelly or rocky soils, thinly covered subsoils, or soils with less than 1% organic matter.

ADDITIONAL INSTRUCTIONS FOR AGRICULTURAL AND NON-AGRICULTURAL USES

SPRAY TANK CLEAN OUT

Thoroughly clean all traces of Tide Hexazinone 75 WDG from application equipment immediately after use. Flush the tank, pump, hoses, and boom with several changes of water after removing nozzle tips and screens (clean these parts separately). Dispose of the equipment washwater by applying it to a use-site listed on this label.

DRIFT CONTROL ADDITIVES

Using product compatible drift control additives can reduce drift potential. When a drift control additive is used, read and carefully observe cautionary statements and all other information on the additive's label. If using an additive that increases viscosity, ensure that the nozzles and other application equipment will function properly with a viscous spray solution.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment.

Pesticide Storage: Store in original container only in a cool, dry place.

Pesticide Disposal: Pesticide wastes may be hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment by shaking and tapping sides and bottom to loosen clinging particles. If not emptied in this manner, the bag may be considered an acute hazardous waste and must be disposed of in accordance with local, state and federal regulations. When completely empty, offer for recycling if available.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

Tide International, USA, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Tide International, USA, Inc., and Buyer and User assumes the risk of any such use. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, TIDE INTERNATIONAL, USA, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product in the event of ineffectiveness or other unintended consequences that may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Tide International, USA, Inc. or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Tide International, USA, Inc. and Seller harmless for any claims relating to such factors.

To the extent consistent with applicable law, in no event shall Tide International, USA, Inc. or Seller be liable for any incidental, consequential, or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER AND BUYER, AND THE EXCLUSIVE LIABILITY OF TIDE INTERNATIONAL, USA, INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE, AT THE ELECTION OF TIDE INTERNATIONAL, USA, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT, OR COMPENSATION LIMITED TO DAMAGES NOT EXCEEDING THE FAIR MARKET PURCHASE PRICE, AND SHALL NOT INCLUDE INCIDENTAL OR CONSEQUENTIAL DAMAGES.**

Tide International, USA, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of Tide International, USA, Inc.

Escort is a registered trademark of E.I. duPont de Nemours and Company.

Microcel E is a trademark of Celite Corporation.

HiSil 233 is a trademark of PPG Industries Ohio, Inc.

[Hexar is a trademark of Tide International, USA, Inc.]

[EPA approval date]