



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

May 27, 2021

Brien O'Loughlin
Regulatory Consultant
Tide International USA, Inc.
c/o Pyxis Regulatory Consulting Inc.
4110 136th St. Ct. NW
Gig Harbor, WA 98332

Subject: Notification per PRN 98-10 – Change in Primary Brand Name
Product Name: Tide Tebuconazole 45WDG Fungicide
EPA Registration Number: 84229-43
Application Date: 12/23/2020
Decision Number: 571540

Dear Mr. O'Loughlin:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped "Notification" and will be placed in our records.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If you have any questions, you may contact Senedu Alemu 703 305-6178 or by email at alemu.senedu@epa.gov.

Page 2 of 2
EPA Reg. No. 84229-43
Decision No. 571540

A handwritten signature in cursive script, appearing to read "Anne Miller".

Acting Product Manager 21
Fungicide Branch
Registration Division (7505P)
Office of Pesticide Programs

AmTideTide Tebuconazole 45WDG Fungicide**ACTIVE INGREDIENT:**

Tebuconazole,

alpha-[2-(4-chlorophenyl)ethyl]alpha-(1,1-dimethylethyl)-1*H*-1,2,4-triazole-1-ethanol 45%**OTHER INGREDIENTS:** 55%**TOTAL** 100%**STOP - Read the label before use****KEEP OUT OF REACH OF CHILDREN
CAUTION**

TO THE USER: If you do not understand the label, find someone to explain it to you in detail. Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

Refer to inside Label Booklet for Precautionary Statements, Storage and Disposal, and Directions for Use.

FIRST AID

If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
Have a product container or label with you when calling a poison control center or doctor, or going for treatment. For MEDICAL Emergencies Call telephone no. 800-858-7378 For PRODUCT USE Information Call telephone no. 949-753-4729	
Note To Physician: No specific antidote. Treat symptomatically. Symptoms of Poisoning: The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.	

EPA Reg. No. [83851-484229-43](#)**EPA Est. No.****Manufactured for:**

[AmTide-LLCTide International USA, Inc.](#)
[P.O.Box 200821 Hubble](#)
[Laguna Hills, CA 92654Irvine, CA 92618](#)

Net Weight: 2 Pounds**NOTIFICATION**

84229-43

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

05/27/2021

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Causes moderate eye irritation. Harmful if swallowed. Avoid contact with eyes or clothing. Avoid breathing dust or spray mist.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA Chemical Resistant Category Selection Chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside.
- Wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to estuarine and marine invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory:

Tebuconazole is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water Advisory:

This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted-entry interval and notification to workers. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI). The REI for each crop is listed in the application directions associated with each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Chemigation: Do not apply this product through any type of irrigation system.

Spray Volume: AmTideTide Tebuconazole 45 WDG Fungicide may be applied in 50 gallons or more of spray solution per acre by ground spray equipment or in 15 gallons or more per acre by aircraft equipment except as noted under "Product Use Directions" or "Remarks" for each crop. Increase spray volume as crop growth increases to ensure thorough coverage of the foliage and fruit. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied.

Aerial application is prohibited in New York State.

Mixing: Add specified amount of AmTideTide Tebuconazole 45 WDG Fungicide into the spray tank while filling with water to the desired level. Operate the agitator while mixing. If other materials are added to the spray tank, the AmTideTide Tebuconazole 45 WDG Fungicide should be thoroughly dispersed prior to the addition of other materials.

Compatibility: To determine the compatibility of AmTideTide Tebuconazole 45 WDG Fungicide with other products, the following procedure should be followed:

Pour the specified proportions of the products into a suitable container of water, mix thoroughly and allow to stand at least five (5) minutes. If the combination remains mixed or can be re-mixed readily, the mixture is considered physically compatible.

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES; RESERVOIRS; RIVERS; PERMANENT STREAMS, MARSHES OR NATURAL PONDS; ESTUARIES AND COMMERCIAL FISH FARM PONDS.

Spray Drift Management: The interaction of many equipment and weather related factors determines the potential for spray drift. The applicator is responsible for considering all of these factors when making application decisions. Avoiding spray drift is the responsibility of the applicator.

Buffer Zone Requirements: For soil or foliar applications, do not apply by ground within 25 feet, or by air within 150 feet of lakes; reservoirs; rivers; permanent streams, marshes or natural ponds; estuaries and commercial fish farm ponds.

Recommendations for Aerial Applications: The spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter.

Importance of Droplet Size: An important factor influencing drift is droplet size. Small droplets (<150 - 200 microns) drift to a greater extent than large droplets. Within typical equipment specifications, applications should be made to deliver the largest droplet spectrum that provides sufficient control and coverage. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure.

Spray should be released at the lowest possible height consistent with good pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

Wind Speed Restrictions: Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size, canopy and equipment specifications determine drift potential at any given wind speed. Do not apply when winds are greater than 15 mph and avoid gusty and windless conditions. Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Restrictions During Temperature Inversions: Do not make aerial or ground applications during temperature inversions. Drift potential is high during temperature inversions. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however if fog is not present, inversions can also be identified by the movement of smoke from a ground source. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Airblast (Air Assist) Specific Recommendations for Tree Crops and Vineyards: Airblast sprayers carry droplets into the canopy of trees/vines via a radially, or laterally directed air stream. The following specific drift management practices should be followed:

- Adjust deflectors and aiming devices so that spray is only directed into the canopy;
- Block off upward pointed nozzles when there is no overhanging canopy;
- Use only enough air volume to penetrate the canopy and provide good coverage;
- Do not allow the spray to go beyond the edge of the cultivated area (i.e., turn off sprayer when turning at end rows);
- Only spray inward, toward the orchard or vineyard, for applications to the outside rows.

Runoff Management: Do not cultivate within 10 feet of the aquatic areas to allow growth of a vegetative filter strip. When used on erodible soils, best management practices for minimizing runoff should be employed. Consult your local Soil Conservation Service for recommendations in your use area.

ROTATIONAL CROPS

Any crop not specified on this label may be planted into treated areas 120 days after last application.

RESISTANCE MANAGEMENT STRATEGY:

AmTideTide Tebuconazole 45WDG Fungicide is a demethylation inhibitor (DMI) fungicide. It may be applied in a tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy. Tank-mixing **AmTideTide** Tebuconazole 45WDG Fungicide with other DMI fungicides is not recommended.

APPLICATION DIRECTIONS

CROP	DISEASE TREATED	PRODUCT APPLICATION RATE
Grape	Black Rot (<i>Guignardis bidwellii</i>) Powdery Mildew (<i>Uncinula necator</i>)	4 oz. per acre
<p>Powdery Mildew: Apply in a preventative spray schedule. Make the first application before bloom and continue applications using spray intervals of up to 21 days in low to moderate disease pressure. Use a 14-day spray interval when disease pressure is severe.</p> <p>Black Rot: <u>Preventive Schedule:</u> Make the first application when new shoots are 1 to 3 inches and continue every 7 to 14 days through 5° BRIX stage or until veraison (berry coloring) is complete. Apply when new shoots are 1 inch and every 7 to 10 days on highly susceptible varieties or under severe disease conditions. <u>Post-Infection Schedule:</u> Follow a post-infection schedule beginning when new shoots are 1 inch until 5° Brix stage. Apply within 72 hours after the beginning of an infection period. Do not apply AmTideTide Tebuconazole 45WDG Fungicide more frequently than every 7 days. Continue AmTideTide Tebuconazole 45WDG Fungicide applications using the preventive schedule if the post-infection schedule is discontinued, then follow the preventive schedule.</p> <p>Botrytis: If AmTideTide Tebuconazole 45WDG Fungicide is applied in a powdery mildew spray schedule, it will complement the activity of registered fungicides used for control of Botrytis bunch rot. Apply every 14 days for suppression of Botrytis.</p> <p>COMMENTS: Preharvest interval (PHI) is 14 days. Do not apply more than 2 pounds of AmTideTide Tebuconazole 45WDG Fungicide per acre per crop season. Do not apply more frequently than every 7 days.</p> <p>Use sufficient water to ensure thorough coverage of vines and fruit. Increase spray volume as vine growth increases.</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		
Banana Plantain	Black Leaf Streak (<i>Mycosphaerella fijiensis</i>) Yellow Sigatoka (<i>Mycosphaerella musicola</i>)	3.2 oz per acre
<p>Apply specified labeled rate as a foliar spray in a minimum of 20 gallons or more of water per acre to ensure thorough coverage. A spray oil recommended for use on banana or plantain should be utilized at 4.3 pints per acre in tank-mix with AmTideTide Tebuconazole 45WDG Fungicide. Thoroughly mix AmTideTide Tebuconazole 45WDG Fungicide in the spray solution before the addition of a spray oil. Make the first application before disease becomes established and repeat applications every 14 days as necessary to maintain disease control.</p> <p>Do not apply AmTideTide Tebuconazole 45WDG Fungicide to non-bagged bananas. Do not apply more than 16 oz of AmTideTide Tebuconazole 45WDG Fungicide per acre per harvest cycle. Preharvest interval (PHI) is 0 days.</p> <p>Restricted entry interval (REI) = 12 hours.</p>		

CROP	DISEASE TREATED	PRODUCT APPLICATION RATE
Tree Nuts: Almonds (see specific use directions) Beechnut Brazil Nut Butternut Cashew Chestnut Chinquapin Filbert (see specific use directions) Hickory Nut Macadamia Nut Pistachios (see specific use directions) Walnut	Brown rot blossom blight <i>(Monilinia laxa, M. fructicola)</i> Botryosphaeria panicle and shoot blight <i>(Botryosphaeria dothidea)</i> Eastern filbert blight <i>(Anisogramma anomala)</i>	8 oz. per acre
<p>Begin applications when conditions are favorable for disease but before infection. Apply on a 7- to 14-day spray schedule.</p> <p>A maximum of 32 oz of product may be applied per acre per crop season. Do not cut cover crops in treated areas for feed or allow livestock to graze treated areas. AmTideTide Tebuconazole 45WDG Fungicide may be applied up to 35 days before harvest.</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		
Almond	Brown rot blossom blight (<i>Monilinia laxa, M. fructicola</i>)	8 oz per acre
<p>Blossom blight: Begin application at pink bud. If the bloom period is extended and/or severe disease conditions exist, make a second application at full bloom. If conditions remain favorable for disease, make another application at petal fall.</p> <p>COMMENTS: Apply AmTideTide Tebuconazole 45WDG Fungicide in a minimum spray volume of 15 gallons per acre by air or 50 gallons per acre by ground. Reduce the application interval for varieties that are highly susceptible to the indicated diseases or when severe disease conditions exist. The use of ground application after petal fall is preferred because of difficulty in penetrating the canopy and obtaining thorough coverage of the foliage and fruit by air. A maximum of 32 oz of product may be applied per acre per crop season. Do not cut cover crops in treated areas for feed or allow livestock to graze treated areas. AmTideTide Tebuconazole 45WDG Fungicide may be applied up to 35 days before harvest.</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		

CROP	DISEASE TREATED	PRODUCT APPLICATION RATE
Pistachio	Botryosphaeria panicle and shoot blight (<i>Botryosphaeria dothidea</i>)	8 oz per acre
<p>Botryosphaeria: Begin application at 25 to 50% bloom and repeat again 10 to 14 days later to protect young flower clusters and fruit. Make two additional applications of AmTideTide Tebuconazole 45WDG Fungicide 14 days apart beginning 49 days before harvest.</p> <p>COMMENTS: Apply AmTideTide Tebuconazole 45WDG Fungicide in a minimum spray volume of 15 gallons per acre by air or 50 gallons per acre by ground. Reduce the application interval for varieties that are highly susceptible to the indicated diseases or when severe disease conditions exist. The use of ground application after flowering is preferred because of difficulty in penetrating the canopy and obtaining thorough coverage of the foliage and fruit by air.</p> <p>A maximum of 32 oz of product may be applied per acre per crop season. Do not cut cover crops in treated areas for feed or allow livestock to graze treated areas. AmTideTide Tebuconazole 45WDG Fungicide may be applied up to 35 days before harvest</p> <p>Control of insect vectors and a thorough pruning to remove plant tissue infected by <i>Botryosphaeria</i> are critical for optimum control of this disease.</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		
Filbert (Hazelnut)	Eastern filbert blight (<i>Anisogramma anomala</i>)	8 oz. per acre

Apply [AmTideTide](#) Tebuconazole 45WDG Fungicide in a preventative spray schedule when conditions are favorable for disease (such as budbreak when the first green tissue is viable). Continue applications of [AmTideTide](#) Tebuconazole 45WDG Fungicide at 7- to 14-day intervals if weather conditions are conducive to disease development. Use shorter spray intervals during budbreak and rapid shoot elongation. The expanding shoot tip must be protected. Reduce the application interval for varieties that are highly susceptible to the indicated disease or when severe disease conditions exist.

Apply [AmTideTide](#) Tebuconazole 45WDG Fungicide in sufficient spray volume for thorough coverage. Tank-mixing the lowest labeled rate of a spray surfactant with [AmTideTide](#) Tebuconazole 45WDG Fungicide may improve spray coverage and penetration of the active ingredient into plant tissue. The use of ground application is preferred because of the difficulty in penetrating the canopy and obtaining thorough coverage of the foliage and stems by air. A maximum of 32 oz of [AmTideTide](#) Tebuconazole 45WDG Fungicide may be applied per acre per crop season. Do not cut cover crops in treated areas for feed or allow livestock to graze treated areas. [AmTideTide](#) Tebuconazole 45WDG Fungicide may be applied up to 35 days before harvest.

Restricted-entry interval (REI) = 12 hours.

CROP	DISEASE	PRODUCT APPLICATION RATE	
Pome Fruits: Apple Crabapple Loquat Mayhaw Pear Pear, Oriental Quince	Scab (<i>Venturia</i> spp.) Cedar-apple rust (<i>Gymino-sporangium juniperi-virginianae</i>) Powdery mildew (<i>Podosphaera leucotricha</i>)	4 to 8 oz. per acre*	2 oz per 100 gallons**

Scab:

Preventative Schedule: Apply at 7- to 10-day intervals from green tip through the second cover spray.

Post-Infection Schedule: Apply within 72 hours after the beginning of a scab infection period. [AmTideTide](#) Tebuconazole 45WDG Fungicide applications should not be closer than 7 days apart. Continue [AmTideTide](#) Tebuconazole 45WDG Fungicide applications using the preventative schedule if the post-infection schedule is discontinued.

Cedar-apple rust: Apply at 7- to 10-day intervals from pink stage through the second cover spray.

Powdery mildew: Apply at 7- to 10-day intervals from tight cluster through the second cover spray.

COMMENTS:

* The amount of [AmTideTide](#) Tebuconazole 45WDG Fungicide required per acre will depend on tree size and volume of foliage present. The rate per acre is based on a standard of 400 gallons of dilute spray solution per acre for large trees. For smaller trees, multiply 2 ounces times the number of 100 gallons of spray solution required to thoroughly wet, to the point of runoff, one acre of the trees being treated.

** For concentrate sprays, apply the same amount of product per acre as would be applied in a dilute spray based on tree size and foliage volume.

[AmTideTide](#) Tebuconazole 45WDG Fungicide may be applied up to 75 days before harvest. A total of 3 pounds of [AmTideTide](#) Tebuconazole 45WDG Fungicide may be applied per acre per crop season.

Restricted-entry interval (REI) = 5 days.

CROP	DISEASE TREATED	PRODUCT APPLICATION RATE	
Stone Fruits: Apricot Cherry Nectarine Peach Plum Plumcot Prune (fresh)	Brown rot (blossom blight, fruit rot) (<i>Monilinia laxa</i> , <i>M. fructicola</i>) Cherry leaf spot (<i>Blumeriella jaapii</i>) Powdery mildew (<i>Podosphaera</i> spp., <i>Sphaerotheca pannosa</i>)	4 to 8 oz. per acre**	2 oz per 100 gallons*
Plum Plumcot Prune (fresh) Peach	Rust (<i>Tranzschelia discolor</i>)	5 to 8 oz per acre**	

*When using the 2 oz per 100 gallon rate, use at least 4 oz of **AmTideTide** Tebuconazole 45WDG Fungicide per acre.
 The amount of **AmTideTide Tebuconazole 45WDG Fungicide required per acre will depend on tree size and volume of foliage present. The rate per acre is based on a standard of 400 gallons of dilute spray solution per acre for large trees. For smaller trees, multiply 2 ounces times the number of 100 gallons of spray solution required to thoroughly wet, to the point of runoff, one acre of the trees being treated. For concentrate sprays, apply the same amount of product per acre as would be applied in a dilute spray based on tree size and foliage volume, but not less than 4 oz of **AmTideTide** Tebuconazole 45WDG Fungicide per acre. Apply 6 to 8 oz per acre of **AmTideTide** Tebuconazole 45WDG Fungicide when severe disease conditions exist.

Blossom blight: Begin application at white bud on cherry, pink bud on peach and nectarine, and green tip on plums and prunes. Apply again at 50% bloom and at petal fall if conditions continue to be favorable for disease development.

Fruit rot: Begin application at two to three weeks before harvest and continue on 7-day intervals through the day of harvest. The blossom and fruit stages must be protected for optimum control of brown rot. If **AmTideTide** Tebuconazole 45WDG Fungicide is applied during only one of these stages, another registered fungicide should be applied to the other stage to provide optimum protection. Additional cover sprays during the early post-bloom period are also important for preventing quiescent fruit infections on sweet cherry and peach.

Leaf spot: Begin application at petal fall or when first leaves unfold and continue applications at 7- to 14-day intervals. Applications should be made at 7-day intervals early in the growing season when terminal growth is rapid and/or under severe disease conditions. A post-harvest application may be made to maintain control and reduce overwintering inoculum.

Powdery mildew: Follow leaf spot schedule until terminal growth ceases.

Rust: Begin applications after canker emergence and continue applications at 14-day intervals under severe disease conditions.

COMMENTS:

AmTideTide Tebuconazole 45WDG Fungicide may be applied up to and including the day of harvest (0-day PHI). Do not apply more than 3 lbs of product per acre per crop season.

Restricted-entry interval (REI) = 5 days.

CROP	DISEASE TREATED	PRODUCT APPLICATION RATE
Cherry (Post-Harvest)	Brown rot (<i>Anisogramma anomala</i>) Gray mold (<i>Botrytis cinerea</i>) Rhizopus rot (<i>Rhizopus</i> spp.)	8 to 16 oz per 25,000 lb of fruit

Mix the specified rate of **AmTideTide** Tebuconazole 45WDG Fungicide in 100 gallons of water or wax/oil emulsion (or aqueous dilution of a wax/oil emulsion). Apply through a spray or flood system on a commercial packing line after stem cutting, sizing, sorting, or washing. For spray treatments, use conventional spray equipment such as T-Jet, flat-fan, or flood jet spray nozzles. Use enough fruit coating material to fully coat fruit. Workers handling treated fruit need to wear waterproof gloves. Do not apply more than once prior to boxing and do not rinse.

Plums (Post-Harvest) Including: Chickasaw, Damson, Japanese, Plumcot, Prune (fresh)	Brown rot (<i>Anisogramma anomala</i>) Gray mold (<i>Botrytis cinerea</i>) Rhizopus rot (<i>Rhizopus</i> spp.)	8 oz per 200,000 lb of fruit
---	---	------------------------------

Mix the specified rate of **AmTideTide** Tebuconazole 45WDG Fungicide in 100 gallons of water, with or without wax. Maintain **AmTideTide** Tebuconazole 45WDG Fungicide suspension by direct or by-pass agitation sufficient to prevent settling of product in spray mixture. Apply as spray to fruit as they travel along the packing line. After sizing, sorting, or washing, **AmTideTide** Tebuconazole 45WDG Fungicide application may be incorporated into a waxing system using conventional equipment that provides for even and complete coverage of fruit. Do not apply more than once prior to boxing and do not rinse.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Do not store below freezing. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If container is leaking or material is spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Dispose of pesticide as directed below. In spill or leak incidents, keep unauthorized people away.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment by shaking and tapping sides and bottom to loosen clinging particles. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. If not emptied in this manner, the bag may be considered an acute hazardous waste and must be disposed in accordance with local, state and federal regulations. When completely empty, offer for recycling if available or dispose of bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of **AmTide-LLCTide International USA, Inc.** To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: **AMTIDE-LLCTIDE INTERNATIONAL USA, INC.** MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. NO AGENT OF **AMTIDE-LLCTIDE INTERNATIONAL USA, INC.** IS AUTHORIZED TO MAKE ANY WARRANTIES BEYOND THOSE CONTAINED HEREIN OR TO MODIFY THE WARRANTIES CONTAINED HEREIN. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, **AMTIDE-LLCTIDE INTERNATIONAL USA, INC.** DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT,

NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT AMTIDE-LLC/TIDE INTERNATIONAL USA, INC.'S ELECTION, THE REPLACEMENT OF PRODUCT.

EPA 20201223