

84467-2

09/26/2008

1/18



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (H7505C)  
1200 Pennsylvania Avenue, N.W.  
Washington, D.C. 20460

EPA Reg. Number:  
84467-2

Date of Issuance:  
SEP 26 2008

Term of Issuance:  
Unconditional

Name of Pesticide Product:  
Antimite-6.5EC

NOTICE OF PESTICIDE:

Registration  
 Reregistration

(Under FIFRA as amended)

Name and Address of Registrant (include ZIP Code):

UPI-ACETO, LLC  
630 Freedom Business Center, Suite 402  
King of Prussia, PA 19406

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided:

1. You will submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. You make the following label changes before releasing the product for shipment:
  - a) Revise the EPA Registration Number to read "EPA Reg. No. 84467-2".

Continued on the next page

Signature of Approving Official:

Date:

9-26-2008

George LaRocca, Product Manager (13), RD/IB (7505P)

b) Direction for Use: Use Restrictions.

As required by the recent amendment to the Propargite RED, replace the three crop rotation statements with:

- “Do not plant any root crop in rotation within 6 months after last application of propargite to labeled crops.”
- “Do not plant other crops in rotation within 2 months after last application of propargite.”

c) Crop tables

If the crop table exceeds one page in length you must retain the headings ‘Crop’, ‘Mites Controlled’, and ‘Directions’, for each of the respective columns on each continuing page

d) Field Corn & Popcorn.

Change “At least 3 pints per acre should be used under conditions of severe mite pressure.” to “Three pints per acre should be used under conditions of severe mite pressure.”


e) Sorghum.

In the crop column add the Sorghum (CA only) and retain the associated directions for use.

2. You are required to generate a one year storage stability study (830.6317) for the subject product under warehouse conditions. It is recommended that observations be made at 0, 3, 6, 9, and 12 months. The corrosion characteristics study (830.6320) may be conducted concurrently. An accelerated study will not substitute for a one year study under warehouse conditions.
3. The basic Confidential Statement of Formula (CSF) dated May 2, 2008 is acceptable and is the basic CSF of record.
4. Please submit three (3) copies of your final printed labeling before releasing the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing amended labeling constitutes acceptance of these conditions.

A stamped copy of the label and the acute toxicology and product chemistry reviews are enclosed for your records. If you have any questions regarding this registration action, please contact Kevin Sweeney at (703) 305-5063. Thank you for your cooperation in this matter.

Sincerely,



for

George LaRocca  
Product Manager 13  
Insecticide Branch  
Registration Division (7505P)

Enclosure

4/18

**Restricted Use Pesticide due to eye and skin effects. For retail sale to use by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.**

# Antimite™-6.5EC

## AGRICULTURAL MITICIDE

**ACCEPTED  
with COMMENTS  
in EPA Letter Dated**

**Active Ingredient:**

Propargite* .....	<del>Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 84467-2</del>	73.6%
Other Ingredients: .....		26.4%
Total: .....		100.0%

\*Contains 6.55 lbs. active ingredient per gallon.

EPA Reg. No. 84467-xxx

EPA Est. No. xxx

**KEEP OUT OF REACH OF CHILDREN**

**DANGER • PELIGRO**

**Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)**

### FIRST AID

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15- 20 minutes. Call a poison control center or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

**IF SWALLOWED:** Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

5/18

**EMERGENCY TELEPHONE NUMBERS:**  
**FOR MEDICAL EMERGENCIES: (866) 673-6671 (Rocky Mountain Poison Control Center)**  
**FOR SPILLS OR TRANSPORTATION EMERGENCIES: (800) 424-9300 (Chemtrec)**

Net contents: \_\_\_\_\_

Sold by:  
UPI-Aceto, LLC  
630 Freedom Business Center, Suite 402  
King of Prussia, PA 19406  
1 800-438-6071

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**DANGER**

Corrosive. Causes irreversible eye damage. Causes skin burns. Harmful if swallowed. Harmful absorbed through skin. Harmful if inhaled. Do not get in eyes or on clothing. Wear protective eyewear such as goggles, face shield or safety glasses. Wear coveralls over long-sleeved shirt and long pants, chemical resistant footwear and socks, chemical resistant gloves (such as barrier laminate, butyl rubber, nitrile rubber, viton, section category C). Avoid breathing spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Remove contaminated clothing and wash clothing before reuse.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical-resistant to this product are listed below. For more options, follow the instructions for Category C on an EPA chemical-resistance category selection chart.

**All Mixers, Loaders, and Applicators and Handlers must wear:**

- Coveralls over long-sleeved shirt and long pants,
- Chemical resistant footwear and socks,
- Goggles, face shield or safety glasses,
- Chemical resistant gloves such as nitrile rubber, butyl rubber, viton and/or barrier laminate, expect for flaggers and applicators using closed cabs,
- Chemical resistant apron for mixers and loaders and persons exposed to the concentrate.
- For overhead exposure wear chemical-resistant headgear.

See Engineering controls for further requirements

**USER SAFETY REQUIREMENTS**

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS**

Users should:

- Remove clothing/PPE immediately if pesticide gets inside.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, change into clean clothing and follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exists, use detergent and hot water.
- Keep and wash PPE separately from the laundry.

**ENGINEERING CONTROLS**

Mixers and loaders supporting aerial applications to corn and cotton must: use a closed system that meets the requirements listed in the Worker Protection Standards (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4)], and must:

- Wear the personal protective equipment required above for mixers/loaders,
- Wear protective eyewear if the system operates under pressure, and
- Be provided and must have it immediately available for use in an emergency, such as a broken package, spill, or equipment breakdown: coveralls, chemical-resistant footwear, and a NIOSH approved respirator with an organic vapor removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval prefix TC-14G), or an organic vapor cartridge or canister with N,R,P or HE prefilter , and chemical resistant headgear, if overhead exposure.

**Applicators using airblast spray equipment must:**

- Wear the personal protective equipment required above for mixers and loaders,
- Be provided and must have immediately available when they must enter the treated area: coveralls, chemical-resistant gloves, chemical-resistant footwear, chemical-resistant headgear, if overhead exposure,
- Take off any PPE that was worn in the treated area before reentering the cab if the applicators are operating in a closed cab, and
- Store all such PPE in a chemical-resistant container, such as plastic bag, to prevent contamination of the inside of the cab or other enclosed area.

Direct flaggers need to wear the personal protective equipment required above for mixers/loaders and applicators/handlers, unless they are in an enclosed vehicle which need not be pressurized. When handlers and/or applicators use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standards(WPS) for agricultural pesticides[40 CFR 170.240(d)(4-6)], the PPE requirements may be reduced or modified as specified in the WPS.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish and aquatic invertebrates and may cause reproductive effects in wildlife. Do not apply directly to water or areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in the neighboring areas. Do not contaminate water when disposing of equipment washwaters.

**PHYSICAL OR CHEMICAL HAZARDS**

Combustible. Do not use or store the product near heat or open flame.

**DIRECTIONS FOR USE**

**RESTRICTED USE PESTICIDE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.



**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker to enter into treated areas during the restricted entry interval (REI) listed except as provided for by the WPS.

The REI and Exceptions are listed in the Directions for Use associated with the crop.

Notify workers of these exceptions (including when entry is permitted for each of the tasks named in the exception).

For early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls over long-sleeved shirt and long pants,
- chemical-resistant gloves, made of any waterproof material,
- chemical-resistant footwear plus socks, and
- protective eyewear.

**Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.**

**SPRAY DRIFT PRECAUTIONS**

**AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR AND GROWER.**

**WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS:**

- Do not apply by ground within 50 feet or by air within 75 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds; estuaries and commercial fish farm ponds.
- Risk of exposure to sensitive aquatic areas can be reduced by making applications when the wind direction is away from the aquatic area.

**Wind direction and speed**

- Only apply this product if the wind direction favors on target deposition.
- Do not apply when the wind velocity exceeds 15 mph.

**Temperature inversion**

- Do not make aerial or ground applications into temperature inversions.
- Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

**Droplet size**

- For ground and aerial applications apply the coarsest droplet size spectrum that provides sufficient coverage and mite control.

**Additional Requirements for Ground Applications**

- For ground boom applications, apply using a nozzle height of no more than 4 feet above the ground or crop canopy.
- For airblast applications, turn off outward pointing nozzles at row ends and when spraying the outer two rows. To minimize spray loss over the top in orchard applications, spray must be directed into the canopy.

**Additional Requirements for Aerial Applications**

- The spray boom should be mounted on the aircraft as to minimize drift caused by wingtip or rotor vortices. The minimum practical boom length should be used and must not exceed 75% of the wing span or 80% rotor diameter.
- Spray must be released at the lowest height consistent with pest control and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety.
- When applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

**RUNOFF PRECAUTIONS**

Under some conditions, propargite may have a high potential for runoff into surface water for several days after application. Do not apply in the following areas:

- frequently flooded areas (excluding artificially flooded areas)
- areas where intense or sustained rainfall is forecasted to occur within 48 hours.

Use best management practices for minimizing surface runoff in the following areas:

- poorly draining or wet soils with readily visible slopes towards adjacent surface water.
- areas with in-field canals or ditches that drain to surface water.
- areas not separated from adjacent surface waters with vegetated filter.

Consult your soil conservation service for recommendations in your use area.

### USE RESTRICTIONS

- Do not use in spray solution above pH 10.
- It is best to use ANTIMITE-6.5EC at pH 7 or lower.
- Beans- Do not feed or forage treated bean vines or bean trash after harvesting.
- Cotton- Apply only before bolls open. Do not feed treated foliage or cotton trash to livestock.
- Peanuts- Do not graze or feed livestock on treated areas or cut treated forage for hay.

Unless the rotated crop is a registered use for propargite, the following applies:

- Do not plant leafy vegetables in rotation within 2 months after the last application of propargite to cotton or corn.
- Do not plant small grains in rotation within the 82 days after the last application of propargite to cotton and corn.
- Do not plant any other feed or food crop in rotation within 6 months after the last application of propargite unless the crop is a registered use for propargite.

#### Registered-labeled field and vegetable uses for propargite are:

Beans, dry	Potatoes
Corn (sweet, field and popcorn)	Sorghum
Cotton	Seed crops only:
Hops	Alfalfa
Jojoba	Carrots
Mint	Clover
Peanuts	Sugar beets

### GENERAL INSTRUCTIONS

Antimite-6.5EC is a liquid emulsifiable concentrate formulation. Antimite-6.5EC needs to be diluted with water prior to spray applications for control of the mites listed in the Use Instructions. For effective control, complete coverage of the fruit and both the upper and lower leaf surfaces is necessary since Antimite-6.5EC is not systemic in action. For best results, add the recommended dosage from the Use Instructions of Antimite-6.5EC to the spray tank, agitate, and spray thoroughly to cover fruit, nuts and foliage. Rate of application is variable according to pest pressure, timing of sprays, and field scouting. Product performance is best when day temperatures average above 70° F.

12/18

**DOSAGE INSTRUCTIONS FOR USE ON BEARING CROPS**

CROP	MITES CONTROLLED	DIRECTIONS
<p>General Information – All labeled crops below</p>		<p>Apply high rate to areas with high mite infestation and areas which historically have had high mite infestations. See Agricultural Use Requirements For general REI exception. There is no exception allowed for hand weeding or thinning.</p>
<p>Beans dry (including dry lima beans) (West of the Rocky Mts.)</p>	<p>Two-spotted Spider Mite</p>	<p>Apply Antimite-6.5EC at 2-3 pints per acre. Apply a minimum spray solution of 20 gallons per acre by ground. For aerial application apply in a minimum spray solution of 5 gallons per acre. Do not apply more than 2 applications per year. The minimum spray interval is 21 days. Restricted Entry Interval (REI) of 9 days. The pre harvest interval (PHI) is 14 days. Do not apply more than 4.5 pints Antimite-6.5EC (3.7 lbs. propargite a.i. per acre) per year.</p>
<p>Oranges &amp; Grapefruits (Bearing) (Florida &amp; Texas Only)</p>	<p>Citrus Red Mite Texas Citrus Mite Six Spotted Mite Citrus Rust Mite False Spider Mite Broad Spider Mite</p>	<p>Apply Antimite-6.5EC at 2-3 pints per acre. Apply a minimum spray solution of 25 gallons per acre by ground. For aerial application apply in a minimum spray solution of 10 gallons per acre. Do not apply more than 2 applications per year. The minimum spray interval is 28 days. Restricted Entry Interval (REI) of 20 days. The pre-harvest interval (PHI) is 21 days. The grazing or feeding of livestock on treated areas or cutting treated forage for hay is prohibited. Apply in sufficient water for thorough coverage. Apply no more than 5 pints per year. Do not apply petroleum oils 30 days prior to or 40 days after applying Antimite-6.5EC as phytotoxicity may occur on leaves and/or fruit. Do not apply Antimite-6.5EC mixed with petroleum oils or any materials containing petroleum based solvents as phytotoxicity may occur on leaves and/or fruit. Do not mix Antimite-6.5EC with or apply to, or after, alkaline materials such as lime, Bordeaux mixture or lime sulfur. Avoid Antimite-6.5EC applications to immature leaves and fruit, and prevent spray solution from puddling, dripping, or running off leaves and/or fruit.</p>
<p>General Information – for Field Corn &amp; Popcorn listed below:</p>		<p>At least 3 pints per acre should be used under conditions of severe mite pressure. Treat early. Best results are achieved when mite populations are low, before the mites have damaged the plants. Treatment is recommended when mite populations are just beginning to build, and the corn is still small enough (usually 3 to 4 feet) to allow good coverage. Ensure applications are made only when corn leaves are</p>

		dry. This is especially true for white corn varieties which are the most susceptible to plant injury. Additional adjuvants are not recommended for use on corn in California.
Field Corn & Popcorn	Two-spotted Spider Mite Banks grass Mite	Apply Antimite-6.5EC at 2-3 pints per acre. Apply a minimum spray solution of 20 gallons per acre by ground. For aerial application apply in a minimum spray solution of 2 gallons per acre. Do not apply more than 1 application per year. Restricted Entry Interval (REI) of 13 days. The pre- harvest interval (PHI) is 30 days.
Sweet Corn (California only)	Two-spotted Spider Mite Banks grass Mite	Apply Antimite-6.5EC at 2-3 pints per acre. Apply a minimum spray solution of 20 gallons per acre by ground. For aerial application apply in a spray solution of 10-20 gallons per acre. Do not apply more than 1 application per year. Restricted Entry Interval (REI) of 13 days. The pre-harvest interval (PHI) is 30 days.
General Information – for All Cotton listed below:		Commercially acceptable mite control may be difficult to obtain where synthetic pyrethroids are or have been used. Do not apply more than 4.0 pints Antimite-6.5EC (3.3 lbs. propargite a.i.) per acre per year. Do not apply more than 2 applications per year. The minimum spray interval is 21 days. Restricted Entry Interval (REI) of 6 days. The pre- harvest interval (PHI) is 50 days.
Cotton(California & Arizona only)	Two-spotted Spider Mite Pacific Spider Mite Strawberry Spider Mite	<b>Early:</b> Apply Antimite-6.5EC at 1 pint per acre. Apply a spray solution of 15-30 gallons per acre by ground. Aerial application not recommended. Antimite-6.5EC may be phytotoxic to young cotton under certain conditions. Age and climatic conditions result in physiological hardening when plants are generally between 6 and 12 inches tall. Applications can begin after plants are hardened. Antimite-6.5EC may be used in concentrate sprayers designed for concentrate row crop applications. Maintain Antimite-6.5EC dosage rate per acre. <b>Mid Season to Layby:</b> Apply Antimite-6.5EC at 1 1/2-2 pints per acre for ground applications for ground applications. Apply a spray solution of 25-40 gallons per acre by ground. Antimite-6.5EC may be used in concentrate sprayers designed for concentrate row crop applications. Maintain Antimite-6.5EC dosage rate per acre. Apply Antimite-6.5EC at 2 pints per acre for aerial applications. For aerial application apply in a spray solution of 5-15 gallons per acre Antimite-6.5EC may be applied in

		<p>vegetable oils by air with ULV sprayers.</p> <p><b>Layby:</b> Apply Antimite-6.5EC at 2 pints per acre.</p> <p>Apply a spray solution of 25-50 gallons per acre by ground. Antimite-6.5EC may be used in concentrate sprayers designed for concentrate row crop applications. Maintain Antimite-6.5EC dosage rate per acre.</p> <p>For aerial application apply in a spray solution of 5-15 gallons per acre. Antimite-6.5EC may be applied in vegetable oils by air with ULV sprayers.</p> <p><b>Layby to Boll Opening:</b> Apply Antimite-6.5EC at 2 pints per acre for aerial applications.</p> <p>For aerial application apply in a spray solution of 5-15 gallons per acre. Antimite-6.5EC may be applied in vegetable oils by air with ULV sprayers.</p>
<p>Cotton (East of the Rocky Mts.)</p>	<p>Two-spotted Spider Mite Pacific Spider Mite Strawberry Spider Mite</p>	<p><b>Early:</b> Apply Antimite-6.5EC at 1 pint per acre.</p> <p>Apply a spray solution of 15-30 gallons per acre by ground.</p> <p>Aerial application not recommended.</p> <p>The feeding of treated foliage or cotton trash to livestock and applications after bolls have opened are prohibited.</p> <p>Antimite-6.5EC may be phytotoxic to young cotton under certain conditions. Age and climatic conditions result in physiological hardening when plants are generally between 6 and 12 inches tall. Applications can begin after plants are hardened.</p> <p>Antimite-6.5EC may be used in concentrate sprayers designed for concentrate row crop applications. Maintain Antimite-6.5EC dosage rate per acre.</p> <p><b>Mid Season to Layby:</b> Apply Antimite-6.5EC at 1-2 pints per acre.</p> <p>Apply a spray solution of 25-40 gallons per acre by ground. Antimite-6.5EC may be used in concentrate sprayers designed for concentrate row crop applications. Maintain Antimite-6.5EC dosage rate per acre.</p> <p>For aerial application apply in a spray solution of 5-15 gallons per acre. Antimite-6.5EC may be applied in vegetable oils by air with ULV sprayers.</p> <p><b>Layby to Boll Opening:</b> Apply Antimite-6.5EC at 2 pints per acre.</p> <p>Apply a spray solution of 25-50 gallons per acre by ground. Antimite-6.5EC may be used in concentrate sprayers designed for concentrate row crop applications. Maintain Antimite-6.5EC dosage rate per acre.</p> <p>For aerial application apply in a spray solution of 5-15 gallons per acre. Antimite-6.5EC may be applied in vegetable oils by air with ULV sprayers.</p>
<p>Jojoba</p>	<p>Tetranychus Spider Mite</p>	<p>Apply Antimite-6.5EC at 2 pints per acre.</p> <p>Apply a spray solution of 20-40 gallons per acre by ground.</p> <p>For aerial application apply in a minimum spray solution of 5 gallons per acre.</p> <p>Do not apply more than 2 applications per year.</p> <p>The minimum spray interval is 10 days.</p> <p>Restricted Entry Interval (REI) of 2 days.</p> <p>Apply to foliage when mite populations reach damaging</p>

<p>Mint (except California)</p>	<p>Two-spotted Spider Mite Mint Bud (squirelly mint effect)</p>	<p>numbers. Apply Antimite-6.5EC at 2-21/2 pints per acre. Apply a minimum spray solution of 20 gallons per acre by ground. For aerial application apply in a minimum spray solution of 8 gallons per acre. Do not apply more than 2 applications per year. The minimum spray interval is 14 days. Restricted Entry Interval (REI) of 7 days. The earliest harvest days after last applications (last sprays) is 14 days. Do not apply more than 5.0 pints Antimite-6.5EC (4.1 lbs. propargite a.i. per acre) per year.</p>
<p>Peanuts (except California)</p>	<p>Two-spotted Spider Mite</p>	<p>Apply Antimite-6.5EC at 2 pints per acre. Apply a minimum spray solution of 20 gallons per acre by ground. For aerial application apply in a minimum spray solution of 5 gallons per acre. Do not apply more than 2 applications per year. The minimum spray interval is 14 days. Restricted Entry Interval (REI) of 2 days. The pre- harvest interval (PHI) is 14 days. When temperatures are greater than 90°F with high humidity some leaf phytotoxicity may occur. The grazing or feeding if livestock in treated areas or cutting treated forage of hay is prohibited.</p>
<p>Potatoes (Pacific Northwest Only)</p>	<p>Two-spotted Spider Mite</p>	<p>Apply Antimite-6.5EC at 2-21/2 pints per acre. Apply no more than 5 pints (4.1 lbs. propargite a.i.) per acre per year. The 21/2 pint per acre rate should be used when the crop is under extreme pressure. Apply a spray solution of 20-50 gallons per acre by ground: For aerial application apply in a minimum spray solution of 10 gallons per acre. Do not apply more than 2 applications per year. The minimum spray interval is 21 days. Minimum spray interval is 14 days in Washington State only. Restricted Entry Interval (REI) of 2 days. The pre- harvest interval (PHI) is 14 days. The grazing or feeding of livestock on cover crops is prohibited.</p>
<p>Sorghum (East of the Rocky Mts.)</p>	<p>Two-spotted Spider Mite Banks grass Mite</p>	<p><b>East of the Rocky Mountains:</b> Apply Antimite-6.5EC at 1 1/2-2 pints per acre. Apply a minimum spray solution of 20 gallons per acre by ground. For aerial application apply in a minimum spray solution of 5 gallons per acre. Due to the large number of varieties of sorghum, it is recommended that a small test plot be treated to check the phytotoxicity before using Antimite-6.5EC on a large area. Do not apply more than 1 application per year. Restricted Entry Interval (REI) of 2 days.</p>

		<p>The post harvest interval for the last spray for silage is 30 days and grain is 60 days.</p> <p><b>CA Only:</b> Apply Antimite-6.5EC at 1 1/2-2 pints per acre. For aerial application apply in a minimum spray solution of 10 gallons per acre.</p> <p>Do not apply more than 1 application per year.</p> <p>Restricted Entry Interval (REI) of 2 days.</p> <p>The pre-harvest interval (PHI) for silage is 30 days and grain is 45 days.</p>
--	--	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

### COMPATIBILITIES

#### TANK MIX COMPATIBILITIES

ANTIMTIE-6.5EC is compatible with most pesticides. Observe all cautions and limitations on labeling of all products used in mixtures.

NOTE:

A. Leaf injury may occur if the following conditions are not observed.

1. ANTIMTIE-6.5EC is not compatible with petroleum based foliar spray oils. (Further information available from supplier.)

2. Do not use penetrating surfactants. Under certain conditions a non-penetrating surfactant can be added. Rely on your previous use experience or consult your local surfactant supplier for specific recommendations.

B. ANTIMTIE-6.5EC is not compatible with alkaline materials such as lime, lime sulfur or Bordeaux. The effectiveness of ANTIMTIE-6.5EC will be reduced.



### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a dry and secure location.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once triple rinsed or pressure rinsed (or equivalent), recycle if available. Some agricultural pesticide containers can be taken to a container collection site or pick up for recycling. To find the nearest site, contact your chemical dealer or manufacturer. If recycling is not available, dispose of in a sanitary landfill or by incineration if allowed by state and local ordinances.

### IMPORTANT INFORMATION

#### READ BEFORE USING PRODUCT

#### CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product reflect the opinion of experts based on field use and tests and must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of UPI-Aceto, LLC or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of UPI-Aceto, LLC and Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold UPI-Aceto, LLC and Seller harmless for any claims relating to such factors.

To the extent consistent with applicable law, UPI-Aceto, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes

18/18

stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or UPI-Aceto, LLC, and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, UPI-ACETO, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, UPI-Aceto, LLC or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF UPI-ACETO, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICES OF THE PRODUCT OR, AT THE ELECTION OF UPI-ACETO, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

UPI-Aceto, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of UPI-Aceto, LLC.

© 2008 UPI-Aceto, LLC. All rights reserved.

ANTIMITE-6.5EC is a trademark of United Phosphorus, Inc.