



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

85724-10

Date of Issuance:

1/26/21

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

AKOFOS 48 EC

Name and Address of Registrant (include ZIP Code):

Bert Volger
Agent for Aako B.V.
c/o Ceres International, LLC
1087 Heartsease Drive
West Chester, PA 19382

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Michael Walsh, Product Manager 11
Invertebrate and Vertebrate Branch 2
Registration Division (7505P)

Date:

1/26/21

2. You are required to comply with the data requirements described in the DCI and EDSP Orders identified below:
 - a. Chlorpyrifos GDCI-059101-967
 - b. Chlorpyrifos EDSP-059101

You must comply with all of the data requirements within the established deadlines. If you have questions about the Generic DCI or EDSP Orders listed above, you may contact the Chemical Review Manager in the Pesticide Reevaluation Division:

<http://iaspub.epa.gov/apex/pesticides/f?p=chemicalsearch:1>

3. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 85724-10.”
4. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 08/12/2020

If you have any questions, please contact Briana Hanlon by phone at (703) 347-8337, or via email at hanlon.briana@epa.gov.

Attachment

RESTRICTED USE PESTICIDE
FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

CHLORPYRIFOS | **GROUP 1B** | **INSECTICIDE**

AKOFOS 48 EC

For control of listed insects infesting certain field, fruit, nut, and vegetable crops.

ACTIVE INGREDIENT	% BY WT.
Chlorpyrifos: 0,0-diethyl-0-(3,5,6-trichloro-2-pyridinyl) phosphorothioate	42.5%
INERT INGREDIENTS*:	57.5%
TOTAL	100.0%

Contains 4 pounds of chlorpyrifos per gallon

* Contains petroleum distillates

Contains chlorpyrifos, the active ingredient used in Lorsban® 4E.

AKOFOS 48 EC is not manufactured or distributed by Dow AgroSciences LLC.

EPA Reg. No. 85724-xx

EPA Est. No. XXXX


Contents: 2.5 gal.

KEEP OUT OF REACH OF CHILDREN WARNING-AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID Organophosphate	
IF SWALLOWED:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Do not give any liquid to a person.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
IF IN EYES:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For medical emergencies involving this product, call the Poison Control Center at 1-800-222-1222.	
Note to Physician: This product contains an organophosphate that inhibits cholinesterase. Treat symptomatically. If exposed, plasma and red blood cell cholinesterase tests may indicate significance of exposure (baseline data are useful). Atropine, only by injection, is the preferable antidote. Oximes, such as 2-PAM/protopam, may be therapeutic if used early; however, use only in conjunction with atropine. In case of severe acute poisoning, use antidote immediately after establishing an open airway and respiration. Contains petroleum distillate. Vomiting may induce aspiration pneumonia.	

Manufactured for:
Aako B.V.
Arhemseweg 87, P.O. Box 205
3830 AE Leusden, The Netherlands



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING**

May be fatal if swallowed. Causes substantial but temporary eye injury. Causes skin irritation. Harmful if absorbed through skin. Do not get in eyes, on skin, or on clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All mixers, loaders, other applicators, and other handlers must wear:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate or viton \geq 14 mils
- Chemical-resistant apron when mixing or loading or exposed to the concentrate
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure
- Wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any R or P filter; OR a NIOSH-approved powered airpurifying respirator with an HE filter.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

When handlers use closed systems or closed cab motorized ground application equipment in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [(40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates, small mammals, and birds. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Protective information may be obtained from your cooperative agricultural extension service.

PHYSICAL OR CHEMICAL HAZARDS

Combustible. Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the Agency responsible for pesticide regulation. Read all Directions for Use carefully before applying.

AGRICULTURAL USE REQUIREMENTS

Use this product in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

The REI for each crop is listed in the direction for use associated with each crop. Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of:

- 4 days for fruit trees
- 5 days for citrus trees
- 3 days for cauliflower
- 24 hours for all other crops not listed above

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves made of barrier laminate or viton \geq 14 mils
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical resistant headgear for overhead exposure

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

NON AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Keep children, pets, and other unprotected persons out of treated area until sprays have dried.

USE INFORMATION

AKOFOS 48 EC insecticide forms an emulsion when diluted with water and is suitable for use in all conventional spray equipment. Consult your State Experiment Station or State Extension Service for proper timing of applications.

When an adjuvant is to be used with this product, Aako B.V. suggests the use of a Chemical Producers and Distributors Association certified adjuvant.

USE PRECAUTIONS

Do not formulate this product into other end use products. **Attention:** Do not cut or weld container.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, AKOFOS 48 EC contains a Group 1B insecticide. Any (insect/mite) population may contain individuals naturally resistant to AKOFOS 48 EC and other Group 1B insecticides. The resistant individuals may dominate the insect/mite population if this group of insecticides/acaricides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed:

Rotate the use of AKOFOS 48 EC or other Group 1B insecticides/acaricides within a growing season, or among growing seasons, with different groups that control the same pests.

Use tank mixtures with insecticides/acaricides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):

- Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
- Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
- When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
- Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
- The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide/acaricides use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact an Aako representative at 484-459-9203.

SPRAY DRIFT

Do not allow spray to drift from the application site and contact people, structures people occupy at any time and the associated property, parks and recreation areas, nontarget crops, aquatic and wetland areas, woodlands, pastures, rangelands, or animals.

- For ground boom applications, do not apply within 25 feet of rivers, natural ponds, lakes, streams, reservoirs, marshes, estuaries, and commercial fish ponds. Apply with nozzle height no more than 4 feet above the ground or crop canopy and when wind speed is 10 mph or less at the application site as measured by an anemometer. Use fine or coarser spray according to ASAE 572 definition for standard nozzles or VMD for spinning atomizer nozzles.
- For orchard/vineyard airblast applications, do not apply within 50 feet of rivers, natural ponds, lakes, streams, reservoirs, marshes, estuaries, and commercial fish ponds. Direct spray above trees/vines and turn off outward pointing nozzles at row ends and outer rows. Apply only when wind speed is 3-10 mph at the application site as measured by an anemometer outside of the orchard/vineyard on the upwind side.
- For aerial applications, do not apply within 150 feet of rivers, natural ponds, lakes, streams, reservoirs, marshes, estuaries, and commercial fish ponds. The boom width must not exceed 75% of the wingspan or 90% of the rotary blade. Use upwind swath displacement and apply only when wind speed is 3-10 mph as measured by an anemometer. Use fine or coarser spray according to ASAE 572 definition for standard nozzles or VMD for spinning atomizer nozzles. If application includes a no-spray zone, do not release spray at a height greater than 10 feet above the ground or the crop canopy.
- For overhead chemigation, do not apply within 25 feet of rivers, natural ponds, lakes, streams, reservoirs, marshes, estuaries, and commercial fish ponds. Apply only when wind speed is 10 mph or less.

The applicator also must use all other measures necessary to control drift.

SPRAY DRIFT MANAGEMENT

The buffer distances specified in the below table are the distances in feet that must exist to separate sensitive sites from the targeted application site. Buffers are measured from the edge of the sensitive site to the edge of the application site.

Sensitive sites are areas frequented by non-occupational bystanders (especially children). These include residential lawns, pedestrian sidewalks, outdoor recreational areas such as school grounds, athletic fields, parks and all property associated with buildings occupied by

humans for residential or commercial purposes. Sensitive sites include homes, farmworker housing, or other residential buildings, schools, daycare centers, nursing homes, and hospitals. Non-residential agricultural buildings, including barns, livestock facilities, sheds, and outhouses are not included in this prohibition.

Application rate (lb ai/A)	Nozzle Droplet Type	Required Setback (Buffer Zones) (feet)		
		Aerial	Airblast	Ground
> 0.5 - 1	Coarse or very coarse	10	10	10
> 0.5 - 1	Medium	25	10	10
> 1 - 2	Coarse or very coarse	50	10	10
> 1 - 2	Medium	80	10	10
> 2 - 3	Coarse or very coarse	80 ¹	10	10
> 2 - 3	Medium	100 ¹	10	10
> 3 -4	Medium or coarse	NA ²	25	10
> 4	Medium or coarse	NA	50	10

¹Aerial application of greater than 2 lbs. ai/A is only permitted for Asian Citrus Psylla control, up to 2.3 lbs. ai/A.

²NA is not allowed.

Only pesticide handlers are permitted in the setback area during application of this product. Do not apply this product if anyone other than a mixer, loader, or applicator, is in the setback area. Exception: Vehicles and persons riding bicycles that are passing through the setback area on public or private roadways are permitted.

SPRAY MIX DIRECTIONS

To prepare the spray, add a portion of the required amount of water to the spray tank and with agitation, add the AKOFOS 48 EC. Complete filling the tank with the balance of water needed. Maintain sufficient agitation during both mixing and application to ensure uniformity of the spray mixture.

AKOFOS 48 EC may also be used in tank mixtures with certain herbicides and/or with non-pressure fertilizer solutions as recommended under specific crop use directions. Prepare tank mixtures in the same manner as recommended above for use of AKOFOS 48 EC alone. When tank mixtures of AKOFOS 48 EC and herbicides are involved, add wettable powders first, flowables second, and emulsifiable concentrates last. Maintain constant agitation during both mixing and application to ensure uniformity of the spray mixture. Do not allow spray mixtures to stand overnight.

Note: Test compatibility of the intended tank mixture before adding AKOFOS 48 EC to the spray or tank mix. Add proportionate amounts of each ingredient to a pint or quart jar, cap, shake, and let set 15 minutes. Formation of precipitates that do not readily redisperse indicates an incompatible mixture that should not be used.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

SPRINKLER IRRIGATION

AKOFOS 48 EC may be applied by sprinkler irrigation for the following crop uses: alfalfa, citrus orchard floors, corn (field and sweet), cotton, cranberry, mint (peppermint and spearmint), sorghum, soybeans, and wheat.

See the use sections for the individual crops for further application information. Do not apply this product to the above listed crops through any other type of irrigation system. Do not apply this product by chemigation to any other crop.

SPRINKLER USE DIRECTIONS

The following use directions are to be followed when AKOFOS 48 EC is applied through sprinkler irrigation systems. Thoroughly clean the injection system and tank of any fertilizer or chemical residues, and dispose of the residues according to state and federal laws. Flush the injector with soap and water. Determine the amount of insecticide needed to cover the desired acreage. Pump the required AKOFOS 48 EC into a steel tank, start mechanical or hydraulic agitation, and add in order the non-emulsifiable oil and/or water. Continually agitate the mixture containing AKOFOS 48 EC. Set the sprinkler system to deliver the desired inches of water per acre. Start the water pump and sprinkler, and let the system achieve the desired pressure and speed before starting the injector. Start the injector and calibrate the injector system according to number 14 in **SPECIAL USE PRECAUTIONS** section. The mixture containing AKOFOS 48 EC must be injected continuously and uniformly into the irrigation water line as the sprinkler is moving. This procedure is necessary to deliver the desired rate per acre in a uniform manner. When the application is finished, allow the entire irrigation and injector system to be thoroughly flushed clean before stopping the system.

SPRINKLER USE PRECAUTIONS

The following use precautions will result in a safe and successful application of mixture containing AKOFOS 48 EC.

1. Apply this product only through sprinkler irrigation systems including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, micro sprinkler, or hand move. Do not apply this product through any other type of irrigation system.
2. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
3. If you have questions about calibration, you should contact State Extension Service Specialist, equipment manufacturers, or other experts.
4. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system.
5. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

6. The system must contain a functional check valve, vacuum relief valve, and a low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. Refer to the American Society of Agricultural Engineer's Engineering Practice 409 for more information.
7. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
8. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
9. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
10. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
11. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. The metering pump must provide a greater pressure than that of the irrigation system at the point of injection. The pump must meet Section 675 for "Electrically Driven or Controlled Irrigation Machines" NEC 70 and must contain Viton or Teflon seals.
12. To insure uniform mixing of the insecticide into the water line, inject the mixture through a nozzle placed in the fertilizer injection port or just ahead of an elbow or tee in the irrigation line so that the turbulence created at those points will assist in mixing. It is suggested that the injection point be higher than the insecticide tank to prevent siphoning.
13. The steel tank holding the insecticide mixture should be large enough to allow the system to complete a revolution with one filling. It should be free of rust, fertilizer sediment, and foreign material, and equipped with an in-line strainer situated between the tank and the injector pump.
14. In order to calibrate the irrigation system and injector to apply the mixture containing AKOFOS 48 EC, determine the following: 1) Calculate the number of acres irrigated by the system; 2) Set the irrigation rate and determine the number of minutes for the system to cover the intended treatment area; 3) Calculate the total gallons of insecticide mixture needed to cover the desired acreage. Divide the total gallons of insecticide mixture needed by the number of minutes to cover the treatment area. This value equals the gallons per minute output that the injector must deliver. Convert the gallons per minute to milliliters or ounces per minute. Calibrate the injector pump with the system in operation at the desired irrigation rate. It is suggested that the injector pump be calibrated at least twice before operation, and monitor the system.
15. Do not apply when wind speed favors drift beyond the area intended for treatment. End guns must be turned off during the application if they irrigate nontarget areas.
16. Do not allow irrigation water to collect or runoff and pose a hazard to livestock, wells, or adjoining crops.
17. Allow foliage to dry before reentering the field.
18. Do not apply through sprinkler systems that deliver a low coefficient of uniformity such as certain water drive units.

CROPS

ALFALFA

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Use AKOFOS 48 EC to control the following pests at the dosages indicated by application as a broadcast foliar spray:

PEST	AKOFOS 48 EC
Aphids (suppression)	½ pint/acre (0.25 lb. a.i./acre)
Corn rootworm adults (spotted cucumber beetle), Grasshoppers, Leafhoppers	½ - 1 pint/acre (0.25 -0.5 lb. a.i./acre)
Alfalfa blotch leafminer, Alfalfa looper, Alfalfa weevil larvae and adults, Armyworms, Cutworms, Egyptian alfalfa weevil larvae and adults, Plant bugs, Leafhoppers, Spittlebugs	1-2 pints/acre (0.5-1 lb. a.i./acre)

NOTE: Use higher rates to control spotted alfalfa aphid in California and Nevada. Stubble spray may be applied to control leafhopper in the Northeast.

Mix the required dosage with enough water to ensure thorough coverage of crop foliage and apply using aerial (fixed-wing or helicopter) or power-operated ground spray equipment. For aerial application, use 2 to 5 gallons of water per acre. For best coverage when using ground application, a minimum of 20 gallons of water per acre with hollow cone nozzles is recommended. Control may be reduced at low spray volumes under high temperature and wind conditions. Treat when field counts or crop injury indicates that damaging pest populations are developing or present; however, do not apply more than once per crop cutting. Some reduction in insect control may be evident under excessively cool conditions. For Egyptian alfalfa weevil control in California, apply the specified dosage in a minimum of 5 gallons of water per acre when larvae are actively feeding and populations reach 15 to 20 larvae per 180° sweep with a 15-inch diameter net.

AKOFOS 48 EC may also be applied through sprinkler irrigation systems as a postemergence broadcast application to control the above listed foliar pests. For best results, use the listed rate of AKOFOS 48 EC per acre. Maintain vigorous tank agitation to assure uniformity of the application throughout the injection period. See **SPRINKLER IRRIGATION** section for further information.

Do not tank mix AKOFOS 48 EC with pesticides, surfactants, or fertilizer formulations unless prior use has shown the combination noninjurious under your current conditions of use. Some phytotoxic symptoms may be observed on young, tender, rapidly growing alfalfa when treated with AKOFOS 48 EC. Alfalfa will outgrow the symptoms and no yield loss should be expected.

This product is highly toxic to bees exposed to direct treatment on alfalfa. Do not apply if nearby bees are clustered outside of hives and bees are foraging. Protective information may be obtained from your agricultural extension service.

RESTRICTIONS: Do not cut or graze treated alfalfa within 7 days after application of ½ pint (0.25 lb. a.i.) of AKOFOS 48 EC per acre, within 14 days after application of 1 pint (0.5 lb. a.i.) per acre, or within 21 days after application of rates above 1 pint (0.5 lb. a.i.) per acre. Do not make more than 4 applications per year or apply more than once per crop cutting. Do not make a second application of AKOFOS 48 EC within 10 days of the first application. To avoid contamination of irrigation floodwaters, do not flood irrigate within 24 hours following an application of AKOFOS 48 EC. The maximum single application rate is 1.0 lb. a.i. chlorpyrifos per acre.

APPLE TREE TRUNK

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 days unless PPE required for early entry is worn.

Use AKOFOS 48 EC to control American plum borer, apple bark borer, broad necked root borer, dogwood borer, flatheaded appletree borer, roundheaded apple tree borer. Mix 1½ qts of AKOFOS 48 EC with 100 gallons of water and using a low-volume handgun or shielded spray apply directly to the trunk from a distance no further than 4 ft. Do not allow spray to contact foliage or fruit.

RESTRICTIONS: For use only in states East of the Rockies (except Mississippi). Do not apply within 28 days before harvest. Make no more than one application to the tree trunk per year as either a pre-bloom or post-bloom application. Do not apply if a pre-bloom application has been made during the year using any other chlorpyrifos product. Do not allow meat or dairy animals to graze in treated orchards. Treat the lower 4 feet of the trunk only. If wind speed is greater than 10 mph, do not apply.

ASPARAGUS

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Use AKOFOS 48 EC to control armyworms, asparagus aphids, asparagus beetles, cutworms, and grasshoppers by application at the rate of 2 pints per acre (1 lb. a.i.). Mix the specified dosage in sufficient water to ensure thorough coverage of treated plants and apply a broadcast foliar spray. For cutworms, it is preferable to apply AKOFOS 48 EC when the soil is moist and worms are active on or near the soil surface. Applications may be made during the fern stage for control of asparagus beetles and asparagus aphids when field counts or crop injury indicates that damaging pest populations are developing or present.

RESTRICTIONS: Do not make more than one preharvest application per season or apply within one day of harvest. Do not make more than two postharvest applications during the fern stage. Do not make a second application of AKOFOS 48 EC within 10 days of the first application. Based on available residue data, the use of AKOFOS 48 EC on asparagus is limited to the Midwest and Pacific Northwest (AZ, CA, ID, IL, IA, KS, MI, MN, MO, NE, ND, OH, OR, SD, WA, WI). The maximum single application rate preharvest or postharvest is 1.0 lb. a.i. chlorpyrifos per acre.

CHERRIES (Trunk Spray)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 days unless PPE required for early entry is worn.

Use AKOFOS 48 EC for the control of lesser peach tree borer, greater peach tree borer, and American plum borer by application as a trunk spray. Mix 1½ to 3 quarts (1.5 to 3 lbs. a.i.) of AKOFOS 48 EC with 100 gallons of water and apply as a coarse, low-pressure spray to give uniform coverage of tree trunks and lower limbs. Make a second application two weeks after the first one and a third application after harvest. Avoid contact with foliage in sweet cherries as premature leaf drop may result. Consult your State Agricultural Experiment Station or Extension Service Specialist for proper time to treat in your area.

In addition, one of the three allowable applications per year may be applied as a dormant spray for control of San Jose scale, peach twig borer, and climbing cutworm. For control of these pests, tank mix ½ to 1 pint (0.25 to 0.5 lb. a.i.) of AKOFOS 48 EC with 1 to 2 gallons of a petroleum oil recommended for dormant use in 100 gallons of water and spray the entire tree by application to runoff using ground spray equipment. For low volume (concentrate) sprays (40 to 100 gallons of spray mixture per acre), use the same amounts of AKOFOS 48 EC and spray oil per acre required for application as a dilute spray and apply in a manner that will ensure thorough coverage of the trees. Use the higher dosage of AKOFOS 48 EC for severe infestations. Use oil as recommended by your State Agricultural Experiment Station or Extension Service Specialist.

RESTRICTIONS: Make only three applications per year. Do not apply within 21 days before harvest. Do not allow meat or dairy animals to graze in treated orchards.

CHRISTMAS TREE PLANTATIONS

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours days unless PPE required for early entry is worn.

Use AKOFOS 48 EC at the rate indicated to control the following insects on the tree varieties listed.

TREE VARIETY	PEST	DOSAGE AKOFOS 48 EC	REMARKS

Balsam fir Concolor fir Douglas fir Eastern white pine Fraser fir Grand fir Noble fir Scotch pine White spruce	Ants (except fire ants, carpenter ants, harvester ants, and pharaoh ants), Aphids, Adelgids (Cooley, Eastern spruce gall), European pine shoot moth, European pine sawfly, Grasshopper, Gypsy moth, Mites (European red spider, Two spotted spider)[except in WA & OR], Pales weevil (adult), Pine needle midge, Pine spittlebug, Plant bugs, Spittlebugs, Spruce budworm, Spruce needleminer, Scale (Pine needle, Pine tortoise, Spruce bud, Black pine, Striped pine)	1 quart/acre (1 lb. a.i./acre)	Do not treat plants under extreme heat and drought stress. Apply as a foliar spray using ground equipment. Thorough coverage of foliage is essential. Use a minimum 10 gpa of finished spray. For effective control of adult spider mites if large numbers of eggs are present, apply a second spray 7 -10 days after initial treatment to control newly hatched nymphs. For scale control, apply when scale crawlers are active. Before treating other conifer species, make application and observe for 7 to 10 days for symptoms of phytotoxicity. The user is responsible for determining if it is safe to treat other conifer species under commercial growing conditions.
	Pales weevil	3 quarts/100 gal. (3 lbs. a.i./100 gal.)	Apply as a cut stump drench.

RESTRICTIONS: Use AKOFOS 48 EC on tree plantations only in Connecticut, Maine, Maryland, Massachusetts, Michigan, New Hampshire, New Jersey, New York, North Carolina, Ohio, Oregon, Pennsylvania, South Carolina, Tennessee, Vermont, Virginia, Washington, and Wisconsin. Do not allow livestock to graze in treated areas. Do not make more than 3 applications per season. Do not make a second application of AKOFOS 48 EC within 7 days of the first application. Do not apply by aerial application.

CITRUS FRUITS

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 5 days unless PPE required for early entry is worn.

Use AKOFOS 48 EC at the rates indicated according to the designated geographic area to control the following pests. Use the lower rates for light infestations and increase the dosage for heavier infestations. A petroleum spray oil recommended for use on citrus trees may be added to dilute spray mixtures only at a rate of up to 1.8 gallons per 100 gallons of water to improve control of aphids, mealybugs, scale insects, and thrips. Treat when insects become a problem or in accordance with the local spray schedule recommended by your State Extension Service Specialist.

AKOFOS 48 EC may be applied in tank mixtures with ethion, dicofol. See **MIXING DIRECTIONS** for further instructions. Read and carefully follow all applicable directions, restrictions, and precautions on labeling for the other products used in combination with AKOFOS 48 EC.

PRECAUTIONS: Observe local use directions for tank-mix combinations especially in regard to applications of AKOFOS 48 EC plus spray oil. Consult with a county farm advisor, county agency, extension service personnel, agricultural commissioner, or pest control advisor for such information regarding a given locality. Do not apply when trees are stressed by drought or high temperatures. Do not tank mix AKOFOS 48 EC with Difolatan 80 Sprills as crop injury may occur. AKOFOS 48 EC is highly toxic to bees exposed to direct treatment.

RESTRICTIONS:

Do not apply when bees are actively visiting the area. During the bloom period in California, apply from one hour after sunset until two hours before sunrise. Do not apply more than 2 applications (does not include citrus orchard floor) or more than 15 pints (7.5 lbs. a.i.) of AKOFOS 48 EC per acre per year. Do not make second foliar application within 30 days of the first application. The preharvest interval restriction is 21 days for citrus, at a dose up to 7 pints (3.5 lbs. a.i.) of AKOFOS 48 EC per acre nor within 35 days for application of rates above 7 pints (3.5 lbs. a.i.) per acre. Do not allow livestock to graze in treated areas. The maximum application rate for a single application is 4 lbs. ai/A in citrus. The use of application rates greater than 8 pints (4 lbs. a.i.) per acre are allowed only in the following California counties: Fresno, Tulare, Kern, Kings, Madera, where it may be applied at 6 lbs. ai/A for control of red scale by ground application. Do not allow meat or dairy animals to graze in treated areas.

Preharvest interval restrictions:

Citrus: 21 days.

CITRUS FRUITS

CROP	GEOGRAPHIC LOCATION	PEST	DOSAGE PT./ACRE	SPRAY VOLUME GAL./ACRE	REMARKS
Calamondin, Chironja, Citrus Citron, Citrus Hybrids, Grapefruit, Kumquat, Lemons, Limes, Mandarin (tangerine), Oranges, Pummelo, Satsuma mandarin, Tangelo, Tangor and Other Citrus Fruit	California- Fresno, Tulare, Kern, Kings, and Madera Counties Only	Red scale	8-12 (4-6 lbs. a.i.)	Ground: 100-2400	Do not use a spray concentration of AKOFOS 48 EC of less than ½ pt./100 gal. of total volume. Additional Precautions for California and Arizona: Do not use AKOFOS 48 EC in combination with spray oil when temperatures are expected to exceed 95°F the day of application or for several consecutive days thereafter. Do not apply during the months of December, January, or February.
Calamondin, Chironja, Citrus Citron, Citrus Hybrids, Grapefruit, Kumquat, Lemons, Limes, Mandarin (tangerine), Oranges, Pummelo, Satsuma mandarin, Tangelo, Tangor and Other Citrus Fruit	California, Arizona	Aphids, Katydid, Lepidopterous larvae, Avocado leafroller, Cutworms, Fruit-tree leafroller, Orange tortrix, Western tussock moth	2-7 (1-3.5 lbs. a.i.)	Ground: 100-750 Aerial: min. 15	Do not use a spray concentration of AKOFOS 48 EC of less than ½ pt./100 gal. of total volume. Additional Precautions for California and Arizona: Do not use AKOFOS 48 EC in combination with spray oil when temperatures are expected to exceed 95°F the day of application or for several consecutive days thereafter. Do not apply during the months of December, January, or February.
		Scale insects (Black scale, Brown soft scale, California red scale)	8 (4 lbs. a.i.)	100-2400	
		Thrips (suppression), Mealybugs	6-12 (3-6 lbs. a.i.)	100-750	
CROP	GEOGRAPHIC LOCATION	PEST	DOSAGE PT./ACRE	SPRAY VOLUME GAL./ACRE	REMARKS
Calamondin, Chironja, Citrus Citron, Citrus Hybrids, Grapefruit, Kumquat, Lemons, Limes, Mandarin (tangerine), Oranges, Pummelo, Satsuma mandarin, Tangelo, Tangor and Other Citrus Fruit	Florida	Aphids, Grasshoppers, Orangedogs, Mealybugs, Scale insects (Snow scale, Florida red scale, Purple scale, Long scale, Chaff scale, Black scale, Brown soft scale)	2-4 (1-2 lbs. a.i.)	Ground: 100-1400 Aerial: min. 20	Do not use a spray concentrate of AKOFOS 48 EC of less than ½ pt./100 gal. of water per acre. Lubber grasshoppers must be controlled when they are small (less than 1 inch in length) by direct contact with spray.
		Citrus rust mites	4-7 (2-3.5 lbs. a.i.)	100-700	
	Texas	Aphids, Cutworms, Katydid, Mealybugs, Scale insects (Brown soft scale, California red scale, Chaff scale)	4-7 (2-3.5 lbs. a.i.)	200-700	Do not use less than ½ pt. of AKOFOS 48 EC per 100 gallons of water in dilute applications.
		Citrus rust mites (suppression)	4-7 (2-3.5 lbs. a.i.)	200-700	
Small Transplanted Grapefruit, Orange, and other Citrus Trees	Texas	Aphids, Cutworms, Katydid, Mealybugs, Scale insects (Brown soft scale, California red scale, Chaff scale)	Max of 7 (3.5 lbs. a.i.)		Apply AKOFOS 48 EC at a rate of 1 fl. oz./1 gal. of water with a backpack sprayer. Apply to runoff.

CITRUS ORCHARD FLOORS

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 5 days unless PPE required for early entry is worn.

Use AKOFOS 48 EC to control ant species (except fire ants, carpenter ants, harvester ants, and pharaoh ants), by applying the specified dose in 25 or more gallons of water with ground application equipment that will uniformly apply the spray to the orchard floor. Do not allow spray to contact foliage or fruit. To control foraging ants, apply AKOFOS 48 EC to the orchard floor at the rate of ¾ to 1 quart (0.75 to 1 lb. a.i.) per acre. Repeat as needed. For best insect control, uniform coverage of the orchard floor is necessary. Do not apply where weed growth or other obstructions would impede uniform coverage of the orchard floor. Foliar applications of AKOFOS 48 EC may be made in addition to the orchard floor, but must comply with the 10-day retreatment interval.

AKOFOS 48 EC may also be applied to citrus orchard floors through sprinkler irrigation systems only if the system uniformly covers the soil surface at the base of the tree. Use the listed amount of AKOFOS 48 EC per acre. See **SPRINKLER IRRIGATION** section for further information.

RESTRICTIONS: Do not apply last treatment within 28 days before harvest. Do not allow livestock or dairy animals to graze in treated areas. Do not apply more than 3 quarts (3 lbs. a.i.) per acre per year. Do not make more than 3 applications per year (does not include foliar applications to citrus trees). Do not make a second application within 10 days of the first application. The maximum single application rate is 1 lb. a.i. chlorpyrifos per acre.

CORN (FIELD), CORN (SWEET) (INCLUDING CORN GROWN FOR SEED)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Conservation Tillage: Preplant, At-Plant, or Preemergence Applications

Apply as a broadcast spray to surface trash and exposed soil using power-operated ground spray equipment. Use a total spray volume of 20 gpa or more. Use a higher rate in the rate range to extend residual control.

Maximum application rates for a single application: Corn: 1.0 lb ai/A.

Maximum number of applications per season: Corn (field, sweet, seed): 3 lbs. ai/A.

Maximum amount a.i. to be applied per acre per season: Corn (field, sweet, seed): 3 lbs. ai/A.

Tank-Mixing: AKOFOS 48 EC may also be applied in tank mixtures with paraquat or glyphosate and/or liquid fertilizer solutions. See **MIXING DIRECTIONS** section in this label for tank mixing instructions. Read and carefully follow all applicable directions, restrictions, and precautions on labeling for each product used in combination with AKOFOS 48 EC.

TARGET PESTS	AKOFOS 48 EC
Armyworms, Cutworms	1-2 pints/acre (0.5 - 1 lb. a.i./acre)

Postemergence Application

Apply as a postemergence broadcast spray using sufficient spray volume to ensure thorough coverage of treated plants, but no less than 15 gallons per acre (gpa) for ground spray equipment or 2 to 5 gpa for aircraft equipment. Control may be reduced at low spray volumes under high temperature and wind conditions.

RESTRICTION: Do not apply aerially in Mississippi.

Chemigation: AKOFOS 48 EC may be broadcast applied postemergence through sprinkler irrigation systems at listed application rates to control listed foliar pests. For best results, tank mix AKOFOS 48 EC with 2 pints of non-emulsifiable oil. See **SPRINKLER IRRIGATION** Section of this product label for application instructions.

TARGET PESTS	AKOFOS 48 EC
Grasshoppers	½-1 pint/acre (0.25 lb.-0.5 lb. a.i./acre)
Aphids, Armyworms, Chinch bugs (1), Corn rootworm adults (2), Cutworms (3), European corn borer (5), Flea beetle adults (1), Southern corn leaf beetle, Webworms (4), Western bean cutworm	1-2 pints/acre (0.5 - 1 lb. a.i./acre)
Corn earworm, Southwestern corn borer (6)	1½-2 pints/acre (0.75-1 lb. a.i./acre)
Billbugs (1), Common stalk borer (9), Corn rootworm larvae (7),(8), Lesser cornstalk borer	2 pints/acre (1 lb. a.i./acre)

Pest Specific Use Directions:

1. For best billbug, chinch bug, or flea beetle control, ground apply in a minimum spray volume of 20 to 40 gpa at 40 psi. If corn is less than 6 inches tall, apply in a 9- to 12- inch wide band over the row. For corn greater than 6 inches tall, apply using drop nozzles directed to the base of the plant. Do not reduce the application rate for banded or directed applications. Concentrate the full labeled dosage rate in the treated zone. When chinch bugs continue to immigrate to corn over a prolonged period or under extreme pest pressure, a second application may be needed.
2. The listed dosage will control silk clipping by corn rootworm adults.
3. For cutworms, it is preferable to apply AKOFOS 48 EC when soil is moist and worms are active on or near the soil surface. If ground is dry, cloddy, or crusted at time of treatment, worms may be protected from the spray and effectiveness will be reduced. Shallow incorporation using a rotary hoe or other suitable equipment immediately before or soon after treatment may improve control. A second application may be required if damage or density levels exceed economic thresholds established for your area.
4. For webworm control, shallow incorporation using a rotary hoe or other suitable equipment immediately before or soon after treatment is necessary.

5. For European corn borer control, use 1 ½ to 2 pints per acre when application is made with power-operated ground or aerial equipment or 1 to 2 pints per acre when application is made through a sprinkler irrigation system. University research indicates that achieving greater than 50% control of first-generation European borer with a single liquid insecticide treatment is highly dependent upon timing, insecticide placement, and weather conditions.
6. For southwestern corn borer, a second application may be applied 21 days later if needed due to reinfestation.
7. For postemergence control of corn rootworm larvae, apply at cultivation. Direct the spray to both sides of the row at the base of the plants just ahead of the cultivator shovels. Cover the insecticide with soil around the brace roots. A cultivation application of AKOFOS 48 EC may be made in addition to an at-planting application of Chlorpyrifos 15G.
8. AKOFOS 48 EC may also be applied through sprinkler irrigation systems at the rate of 2 pints per acre to control corn rootworm larvae. Time the application to coincide with the appearance of the second instar larvae. Apply with enough water to wet the root zone to the depth control needed. If soils are wet, allow enough soil drying to occur such that an application using a minimum amount of water will not produce surface runoff. See Chemigation (**SPRINKLER IRRIGATION**) section of this label for application instructions.
9. Do not use AKOFOS 48 EC in combination with a burndown herbicide for control of common stalk borer. For common stalk borer control, treat approximately 11 days after application of glyphosate or after burndown with paraquat herbicide is complete (3 to 5 days).

RESTRICTIONS (CORN (FIELD), CORN (SWEET) (INCLUDING CORN GROWN FOR SEED):

Do not apply within 21 days before harvest of grain or ears. Do not apply more than 3 lbs. a.i. chlorpyrifos (6 pints of AKOFOS 48 EC) per acre per season. Do not make more than 3 applications per season of any product containing chlorpyrifos including the maximum allowed of 2 granular applications at the 1 lb. a.i. chlorpyrifos rate. Do not make a second application of AKOFOS 48 EC or other product containing chlorpyrifos within 10 days of the first application. The maximum single application rate is 1 lb. a.i. chlorpyrifos (2 pints of AKOFOS 48 EC) per acre. If more than 1 lb. a.i. granular chlorpyrifos per acre is applied at-plant (for a maximum of 1.3 lbs. a.i. per acre per season), only 1 additional application of a liquid product containing chlorpyrifos at 1 lb. a.i. per acre is allowed per season, for a total of 2.3 lbs. a.i. chlorpyrifos per acre per season.

COTTON

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Use AKOFOS 48 EC for control of the following pests in all states except Arizona and California at the dosages indicated:

PEST	AKOFOS 48 EC
Cotton fleahopper, Plant bugs (Lygus, Mirids)	3/8-1 pint/acre (0.18 -0.5 lb. a.i./acre)
Fall armyworm, Grasshopper, Thrips, Yellowstriped armyworm	½ - 1 pint/acre (0.25 -0.5 lb. a.i./acre)
Cotton aphid	½ -2 pints/acre (0.25-1 lb. a.i./acre)
Spider mites	1 pint/acre (0.5 lb. a.i./acre)
Beet armyworm, Cotton budworm, Tobacco budworm, Cutworms, Pink bollworm, Salt marsh caterpillar	1½ - 2 pints/acre (0.75 - 1 lb. a.i./acre)

NOTE: The listed dosage rate of 3/8 pint per acre will not achieve the high degree of control of the highest label rate but will minimize the damage done by plant bugs and cotton fleahopper and allow the beneficial insects to survive, build up, and be available to aid in the control of bollworms infesting cotton. For infestations of cotton aphids that are difficult to control, use a higher dosage within the indicated rate range.

Use AKOFOS 48 EC for control of the following pests in Arizona and California at the dosages indicated:

PEST	AKOFOS 48 EC
Armyworms, Cotton Aphid, Cotton fleahopper, Lygus, Saltmarsh caterpillar, Thrips	1-2 pints/acre (0.5 - 1 lb. a.i./acre)
Cotton bollworm, Tobacco budworm, Boll weevil, Cutworms, Pink bollworm	2 pints/acre (1 lb. a.i./acre)

Note: The 2-pint rate will aid in the suppression of cotton leafperforator and spider mites.

Mix the required dosage with sufficient water to ensure thorough coverage of plants and apply using aerial or power-operated ground spray equipment. For aerial application, use at least 2 gallons of spray per acre. For ground application, use sufficient spray volume to ensure thorough coverage of treated plants but not less than 10 gallons of spray per acre. Increase spray volume when foliage is dense and/or when pests populations are high, and/or under high temperature and wind conditions. Treat when field counts indicate damaging insect populations are developing or present. Retreat as necessary to maintain control.

AKOFOS 48 EC may also be applied through sprinkler irrigation systems as a postemergence broadcast application to control the above listed foliar pests. For best results, use the listed rate of AKOFOS 48 EC per acre. Maintain vigorous tank agitation to assure uniformity of the application throughout the injection period. See **SPRINKLER IRRIGATION** section for further information. For effective control of spider mites when large numbers of eggs are present, apply a second spray 3 to 5 days after initial treatment to control newly hatched nymphs.

For best results on bollworms and budworms, it is suggested that fields be scouted twice per week and treatments made when worms are 1/4 inch or less in length.

RESTRICTIONS: Do not apply within 14 days before harvest or make more than 3 applications per season. Do not apply more than 6 pints (3 lbs. a.i. chlorpyrifos) per acre per season. Do not make a second application within 10 days of the first application. Do not allow meat or dairy animals to graze in treated areas. Do not feed gin trash or treated forage to meat or dairy animals. The maximum single application rate is 1.0 lb. a.i. chlorpyrifos per acre.

CRANBERRIES

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Use AKOFOS 48 EC by application as a broadcast, foliar spray to control brown spanworm, cranberry fruitworm, cranberry weevil, cutworms, fireworms, and *Sparganothis* fruitworm at the rate of 3 pints (1.5 lbs. a.i.) per acre. Mix the specified dosage in enough water to ensure thorough coverage and apply no less than 5 gallons of spray per acre when using aerial equipment or no less than 15 gallons of spray per acre when using ground equipment. For weevil control, apply once at flower bud development (late May, early June) and, if weevils are present, once after 100% bloom (early to mid-July). For other insects, treat when field counts indicate damaging insect populations are developing or present. Apply only after the winter flood has been removed. To avoid pesticide contamination of flood water, make no applications while bogs are flooded.

AKOFOS 48 EC may also be applied through sprinkler irrigation systems to control the above listed pests. For best results, use the listed rate of AKOFOS 48 EC per acre. Maintain vigorous tank agitation to assure uniformity of the application throughout the injection period. See **SPRINKLER IRRIGATION** section for further information.

RESTRICTIONS: Do not make more than two applications per year or apply within 60 days before harvest. Do not make a second application within 10 days of the first application. The maximum single application rate is 1.5 lbs. a.i. chlorpyrifos per acre.

FIGS

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 days unless PPE required for early entry is worn.

Use AKOFOS 48 EC at the rate of 2 quarts (2 lbs. a.i.) per acre for control of dried fruit beetle by application in sufficient water to the soil surface followed by incorporation into the top 3 inches of soil. Apply to fig orchard soil as a dormant application in late winter prior to beetle emergence and prior to leaf formation. Use a spray volume of 10 gallons per acre or more and apply as a broadcast spray to the soil surface using ground equipment. On the day of treatment incorporate into the top 3 inches of soil using suitable equipment.

RESTRICTIONS: Make only one application per year. Do not apply within 7 months (217 days) of harvest. Based on available residue data, use of AKOFOS 48 EC on figs is restricted to California. The maximum single application rate is 2.0 lbs. a.i. chlorpyrifos per acre.

GRAPES

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Use AKOFOS 48 EC as a soil surface application for control of grape root borer by application just before the pest emerges from the soil. Mix 4½ pints (2.25 lbs. a.i.) of AKOFOS 48 EC with 100 gallons of water and apply 2 quarts of the diluted spray mixture to the soil surface on a 15 square foot area around the base of each vine. Do not allow spray to contact fruit or foliage. AKOFOS 48 EC may also be used as a prebloom application to control climbing cutworm. Apply one quart per acre as a spray drench ground application using a minimum spray volume of 25 gallons per acre. Do not use in conjunction with soil surface application for grape borer control.

RESTRICTIONS: Do not make more than one application per season or apply within 35 days before harvest. Based upon available residue data, the use of AKOFOS 48 EC in grapes is restricted to states east of the Rocky Mountains/Continental Divide. Do not make more than one application per season of AKOFOS 48 EC or other product containing chlorpyrifos. The maximum single application rate for soil surface application is 2.12 lbs. a.i. chlorpyrifos per 100 gallons. The maximum single application rate for prebloom application is 1.0 lb. a.i. chlorpyrifos per acre. Not for use in Mississippi.

LEGUME VEGETABLES (Succulent or Dried) (except soybean)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

CROP	SPECIFIC DIRECTIONS
Legume Vegetables including adzuki bean, asparagus bean, bean, blackeyed pea, broad bean (dry and succulent), catjang, chickpea, Chinese longbean, cowpea, crowder pea, dwarf pea, edible pod pea, English pea, fava bean, field bean, field pea, garbanzo bean, garden pea, grain lupin, green pea, guar, gyacinth bean, jackbean, kidney bean, lablab bean, lentil, lima bean, moth bean, mung bean, navy bean, pea, pigeon pea, pinto bean, rice bean, runner bean, snap bean snow pea,	For control of seed maggots, apply 2 pints/acre as a preplant broadcast application. Apply AKOFOS 48 EC in a minimum of 10 gpa of spray to the soil surface using suitable ground equipment. AKOFOS 48 EC must be incorporated in the top 1 to 3 inches of soil to improve the activity against seed maggots. To apply an at-plant T-Band application, apply 1.8 fl. oz. of AKOFOS 48 EC per 1000 feet of row at 30-inch row spacing. To achieve shallow incorporation, apply the spray in a 3 to 5 inch wide band over the row behind the planter shoe and in front of the press wheel to achieve shallow incorporation. Mix the specified dosage in the table below in a minimum of 10 gpa of spray and apply to the soil surface as a ground spray. Equivalent rates of insecticide spray required per 100 feet of row for listed row spacing are given in the table below. Incorporate AKOFOS 48 EC into the top ½ to 1 inch of soil to improve activity

southern pea, sugar snap pea, sweet lupin, sword bean, tepary bean, urd bean, wax bean, white lupin, white sweet lupin, yardlong bean	against seed maggots.
---	-----------------------

Spray Volume Per Acre (Gallons)	Fl. oz. of Spray Volume per 100 Feet of Row			
	30-inch	28-inch	24-inch	22-inch
10	7.3	6.9	5.9	5.4
15	11	10.3	8.8	8.1
20	14.7	13.7	11.8	10.8

RESTRICTIONS: Do not make more than one application of AKOFOS 48 EC per year. Do not apply AKOFOS 48 EC at-plant if the field was treated with a preplant incorporated treatment of AKOFOS 48 EC. Do not apply more than 2 pints of AKOFOS 48 EC (1 lb. a.i. chlorpyrifos) per acre. Not for use in Mississippi.

MINT (Peppermint and Spearmint)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Use AKOFOS 48 EC by application as a broadcast foliar spray to control cutworms at the rate of 2 to 4 pints (1 to 2 lbs. a.i.) per acre and mint root borer at the rate of 4 pints per acre. Mix the specified dosage in water to give no less than 10 gallons of spray per acre and apply using ground spray equipment. For cutworm control, treat during May and June when field counts indicate damaging insect populations are developing or present. When larvae are less than 3/4 inch in length, use the 2 pint rate. When larvae are 3/4 inch or more in length, use the higher rate. For mint root borer control, apply postharvest when field counts indicate damaging insect populations are developing or present. Follow treatment with approximately 1-acre inch of sprinkler irrigation immediately after application to incorporate the insecticide into the soil.

AKOFOS 48 EC may also be applied through sprinkler irrigation systems as a postemergence broadcast application to control the above listed pests. For best results, use the listed rate of AKOFOS 48 EC per acre. Maintain vigorous tank agitation to assure uniformity of the application throughout the injection period. See **SPRINKLER IRRIGATION** for further information.

RESTRICTIONS: Do not apply within 90 days before harvest. Make only one application during the growing season. Do not make more than one preplant incorporated application in the spring. Do not use in conjunction with a broadcast foliar application for cutworm control. Make only one postharvest application per season. The maximum single application rate is 2.0 lbs. a.i. chlorpyrifos per acre. Not for use in Mississippi.

NECTARINES, PEACHES, ALMONDS (Trunk Spray or Preplant Dip)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 days unless PPE required for early entry is worn.

Use AKOFOS 48 EC for the control of peach tree borer in almonds, nectarines, and peaches by application as a trunk spray before newly hatched borers enter the trees. Mix 3 quarts (3 lbs. a.i.) of AKOFOS 48 EC with 100 gallons of water and apply as a coarse, low-pressure spray to give uniform coverage of tree trunks. Thoroughly wet all bark areas from ground level to scaffold limbs. Do not allow spray to contact fruit. Consult your State Agricultural Experiment Station or Extension Service Specialist's written recommendations for proper time to treat in your area.

AKOFOS 48 EC may also be used as a preplant dip application for non-bearing peach and nectarine trees (only) at the equivalent application rate of 3 quarts (3 lbs. a.i.) per 100 gallons of water for control of peach tree borer. Dip trees several inches above the grafting bud scar and plant immediately or allow to dry before returning to storage. Do not allow peach trees to remain in contact with the dip solution.

RESTRICTIONS: Make only one application per season. The preharvest interval restriction is 21 days. Do not allow meat or dairy animals to graze in treated orchards. Not for use in Mississippi.

ONIONS (DRY BULB)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

At-Plant Soil Drench Application: Use AKOFOS 48 EC to control onion maggot by application as an in-furrow drench. Apply AKOFOS 48 EC at the rate of 1.1 fluid ounce per 1000 linear feet of row at an 18-inch row spacing. Use a minimum of 40 gallons of total drench per acre. Incorporate to a depth of 1 to 2 inches. Do not make more than one application per year following these instructions.

Post-Plant Soil Drench Application: Apply one quart (1 lb. a.i.) per acre as an early season directed spray to the base of onion seedlings or transplants during peak onion maggot egg laying. Use a minimum of 100 gallons per acre for thorough wetting.

RESTRICTIONS: Do not make more than two applications per year (at plant plus post plant). Do not harvest within 60 days of application. The maximum single application rate is 0.03 lb. a.i. chlorpyrifos per 1000 feet of row. Do not aerially apply this product in Mississippi.

PEANUTS

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

For suppression of wireworms, apply AKOFOS 48 EC at a rate of 4 pints (2 lbs. a.i.) per acre as a preplant broadcast spray to the soil surface followed by immediate soil incorporation to a depth of 3 to 4 inches. Use a minimum of 10 gallons of total spray per acre.

RESTRICTIONS: Do not make more than one application per season. The maximum single application rate is 2.0 lbs. a.i. chlorpyrifos per acre. Do not harvest within 21 days after treatment. Do not feed treated peanut forage or hay to meat or dairy animals. Aerial application to peanuts is prohibited.

**PEARS
(California, Oregon and Washington)**

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 days unless PPE required for early entry is worn.

Apply 4 pints (2 lbs. a.i.) per acre in 100 to 400 gallons of spray per acre to control codling moth. Apply using an airblast speed sprayer or other suitable ground equipment.

RESTRICTIONS: For this use, do not make more than one post harvest application prior to dormancy per year, do not harvest or use treated fruit for food or feed, do not allow meat or dairy animals to graze in treated orchards. If unauthorized entry into a treated orchard cannot be prevented, then the orchard shall be posted with appropriate signs according to the Worker Protection Standard while treated, unharvested fruit remains on the tree.

SORGHUM-GRAIN SORGHUM (MILO)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Use AKOFOS 48 EC Insecticide for control of the following pests at the dosage indicated:

PEST	AKOFOS 48 EC	SPECIFIC DIRECTIONS
Sorghum midge	½ pint/acre (0.25 lb. a.i./acre)	Apply when 30 to 50% of the seed heads are in bloom. Repeat at 3-day intervals if necessary.
Grasshoppers, Yellow sugar cane aphid and other aphids	½-1 pint/acre (0.25 - 0.5 lb. a.i./acre)	
Greenbug	½-2 pints/acre (0.25 - 1 lb. a.i./acre)	For infestations of greenbug that are difficult to control, use a higher dosage within the indicated rate range.
Chinch bugs, Lesser cornstalk borer	1-2 pints/acre (0.5 - 1 lb. a.i./acre)	Apply as a directed spray toward the base of the plant using power-operated ground spray equipment with sufficient water to ensure coverage of an 8-12 inch band centered on the row. On plants less than 6 inches high, apply an 8- to 12-inch band over the row. Do not reduce the dosage for banded or directed applications. Concentrate the full labeled dosage rate in the treated zone.
Webworms	1 pint/acre (0.5 lb. a.i./acre)	
Armyworms, Corn earworm, Cutworms	1-2 pints/acre (0.5 - 1 lb. a.i./acre)	
European and Southwestern corn borer	1½-2 pints/acre (0.75 - 1 lb. a.i./acre)	

Mix the specified dosage in enough water to ensure thorough coverage and apply using suitable aerial or ground spray equipment. To minimize chemical injury, do not apply AKOFOS 48 EC to drought stressed grain sorghum within 3 days following irrigation or rain except where the product is applied in irrigation water.

AKOFOS 48 EC may also be applied through sprinkler irrigation systems as a postemergence broadcast application to control the above listed foliar pests. For best results, use the listed rate of AKOFOS 48 EC per acre. Maintain vigorous tank agitation to assure uniformity of the application throughout the injection period. See **SPRINKLER IRRIGATION** section for further information.

RESTRICTIONS: The treated crop is not to be used for grain, forage, fodder, hay, or silage within 30 days after application of 1 pint of AKOFOS 48 EC per acre or within 60 days after application of rates above 1 pint (0.5 lb. a.i.) per acre. Do not treat sweet varieties of sorghum. Do not apply more than 3 pints (1.5 lbs. a.i.) of AKOFOS 48 EC per acre per season. Do not make more than 3 applications per season. Do not make a second application within 10 days of the first application. Do not apply by air in Mississippi.

SOYBEANS

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

For use to control armyworms, bean leaf beetle, corn earworm, cutworms, European corn borer, grasshoppers, green cloverworm, lesser cornstalk borer, Mexican bean beetle, saltmarsh caterpillar and other woollybears, southern green stink bug, spider mites, and velvetbean caterpillar.

Soil Treatment

Use AKOFOS 48 EC at the rate of 1 to 2 pints (0.5 to 1 lb. a.i.) per acre to control cutworms and lesser cornstalk borer. Mix the specified dosage in a minimum of 10 gallons of spray per acre and apply to the soil surface using suitable ground spray equipment. Equivalent rates of insecticide spray required per 100 feet of row for various row spacing are given in the accompanying table. For at-plant treatments, apply the insecticide over the row in a 4- to 6-inch band in front of the planter shoe or press wheel or after the press wheel followed by a drag chain for light incorporation. Do not apply as an in-furrow treatment. For postemergence rescue treatments, apply as a directed spray in a 9- to 12-inch band at the base of the plant. To plants under 6 inches high, apply over-the-top in a 6- to 12-inch band. Treat when field counts or conditions indicate that pests are or may become a problem.

Fluid Ounces of Spray Required Per 100 Feet of Row for Various Row Spacing

Volume of Spray Per Acre	36"	32"	28"	24"
10 gallons	8.8	7.9	6.9	5.9
15 gallons	13.2	11.8	10.3	8.8
20 gallons	17.6	15.7	13.7	11.8

Foliar Treatment

Use AKOFOS 48 EC at the following rate by application in sufficient water to ensure thorough coverage of treated plants:

PEST	AKOFOS 48 EC
European corn borer, Southern green stink bug	2 pints/acre (1 lb. a.i./acre)
Bean leaf beetle, Cutworms, Corn earworm, Saltmarsh caterpillar and other woollybears, Soybean aphid	1-2 pints/acre (0.5 -1 lb. a.i./acre)
Mexican bean beetle, Armyworms	1-1½ pints/acre (0.5-0.75 lb. a.i./acre)
Velvetbean caterpillar, Grasshoppers, Green cloverworm, Spider mites	½ -1 pint/acre (0.25 - 0.5 lb. a.i./acre)

Apply as a broadcast spray using either aerial or ground equipment when field counts indicate damaging insect populations are developing or present; retreat as necessary to maintain control. Mix the specified dosage in a minimum of 15 gallons of spray per acre for ground spray equipment or 5 gallons of spray per acre for aircraft equipment. For effective control of spider mites when large numbers of eggs are present, apply a second spray 3 to 5 days after initial treatment to control newly-hatched nymphs. On determinate soybeans, do not apply more than one application after pod set.

AKOFOS 48 EC insecticide may also be applied through sprinkler irrigation systems as a postemergence broadcast application to control the above listed foliar pests. For best results, use the listed rate of AKOFOS 48 EC per acre. Maintain vigorous tank agitation to assure uniformity of the application throughout the injection period. See **SPRINKLER IRRIGATION** section for further information.

RESTRICTIONS: Do not apply more than 6 pints (3 lbs. a.i.) of AKOFOS 48 EC per acre per season. Do not make more than 3 applications per season. Do not apply last treatment within 28 days before harvest nor apply last two treatments closer than 10 days apart. Do not allow livestock to graze in treated areas or otherwise feed treated soybean forage, hay, and straw to meat or dairy animals. Not for use in Mississippi.

STRAWBERRIES

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Preplant Incorporation Treatment: Use AKOFOS 48 EC insecticide at the following rates by application in sufficient water to the soil surface and incorporate into the soil in the spring for protection of strawberries during the following year:

PEST	AKOFOS 48 EC
Garden symphylans, grub	4 pints/acre (2 lbs. a.i./acre)

Foliar Application: Use AKOFOS 48 EC by application as a broadcast foliar spray to control strawberry bud weevil (clipper) at the rate of 1 quart per acre. Apply in a minimum of 40 gallons of spray per acre when buds first appear and 10 to 14 days later. Do not apply after berries start to form or when berries are present. Do not tank mix AKOFOS 48 EC with pesticides, surfactants, or fertilizer formulations unless prior use has shown the combination noninjurious under your current conditions of use. Phytotoxicity may occur when AKOFOS 48 EC is applied to strawberries experiencing high temperature and drought stress.

Post Harvest Application: For strawberry crown moth, apply 1 quart (1 lb. a.i.) per acre as a directed spray to crown of strawberry plants immediately after harvest and after plants are topped. Use a minimum spray volume of 100 gallons per acre and repeat application if required 14 to 18 days later. Do not sprinkle irrigate for one week following application.

RESTRICTIONS: For prebloom use only, do not apply after berries start to form or when berries are present. Do not apply within 21 days before harvest. For preplant application, do not make more than one application per year. For foliar and post harvest applications, do not make more than two applications per year. Do not make a second application within 10 days of the first foliar application and within 14 days for post harvest application. The maximum single application rate is 2.0 lbs. a.i. chlorpyrifos per acre and 1.0 lb. a.i. chlorpyrifos per acre for foliar and postharvest application. Not for use in Mississippi.

SUGAR BEETS

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Soil Treatment (At Planting or Preplant Incorporated): To reduce feeding damage from early season insects such as cutworms, apply at planting or as a preplant treatment and incorporate to a depth of 1 to 2 inches. Do not apply as an in-furrow treatment. Apply 1 pint of AKOFOS 48 EC per planted acre to a 10-inch wide band centered over the row for furrows 30 inches apart. (For rows 30 inches apart, this is equivalent to 9.2 fl. oz. of AKOFOS 48 EC per 10,000 feet of row). For other row widths, adjust the spray volume per planted acre in proportion to the length of row actually treated.

Postemergence Treatment: Apply specified rate as a broadcast or banded foliar spray. Treat when field counts indicate that damaging insect populations are developing or present.

Broadcast Application: Apply the specified dosage in water using 2 to 5 gallons per acre of finished spray when using aerial spray equipment or 10 to 30 gallons per acre when using ground spray equipment.

Banded Foliar Spray: Apply the specified rate within the band using a minimum of 7 gallons of spray volume in a 5- to 7-inch wide band centered over the row. Do not reduce the rate for band applications. Concentrate the full labeled dosage rate (see band rates in table below) in the treated zone. For best results, band-applied treatments should be lightly incorporated, either mechanically or with irrigation.

Target Pests	AKOFOS 48 EC	
	Broadcast (pint/acre)	Band (pint/acre)
Grasshoppers	½ -1 (0.25-0.5 lb. a.i.)	-
Spider mites	1 (0.5 lb. a.i.)	¾ (0.33 lb. a.i.)
Tarnished plant (Lygus)	1 (0.5 lb. a.i.)	-
Fall armyworms, Yellowstriped armyworm, Webworms	1-2 (0.5 – 1 lb. a.i.)	¾-1 1/3 (0.33 - 0.66 lb. a.i.)
Beet armyworm	1 ½ -2 (0.75 – 1 lb. a.i.)	1 – 1 1/3 (0.5-0.66 lb. a.i.)
Cutworms, Flea beetle adults	2 (1 lb. a.i.)	1 1/3 (0.66 lb. a.i.)
Sugarbeet root maggot adults	½ -1 (0.25-0.5 lb. a.i.)	-
Sugarbeet root maggot larvae	-	1 1/3- 2 (0.66 – 1 lb. a.i.)
Sugarbeet root maggot larvae	2 (1 lb. a.i.)	1 1/3 – 2 (0.66 – 1 lb. a.i.)
Aphids*	1-2 (0.5 - 1 lb. a.i.)	-

*Not for use in California on aphids.

For grasshopper control, the low rate will control small nymphs (1st through 3rd instar). For sugarbeet root maggot adults, apply anytime from 7 days before until 3 days after peak adult emergence in order to target adults present at time of application based on local field trap monitoring. For sugarbeet root maggot larvae, use as primary treatment to control root maggot larvae. Base application timing on local field trap monitoring. Apply anytime from 7 days before until 3 days after peak adult emergence. Use as a supplemental postemergence treatment following an at-plant insecticide application for control of root maggot larvae. Base application timing on local field trap monitoring. Apply anytime from 7 days before until 3 days after peak adult emergence. To prevent potential development of insecticide resistance in sugarbeet root maggot, producers are encouraged to take the following steps: (1) avoid making more than 2 applications of AKOFOS 48 EC per season when adults are active; (2) If an organophosphate insecticide was applied at planting, make no more than one postemergence application of AKOFOS 48 EC when adults are active.

RESTRICTIONS: Do not apply within 30 days before harvest of beet roots and tops. Do not make more than 3 applications per season. Do not apply more than a total of 6 pints (3 lbs. a.i.) per acre per season. The maximum single application rate is 1.0 lb. a.i. chlorpyrifos per acre. Do not make a second application within 10 days of the first application. Do not allow livestock to graze in treated areas nor harvest treated beet tops as feed for meat or dairy animals within 30 days after last treatment. Not for use in Mississippi.

SUNFLOWERS

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

For use to control cutworms, sunflower beetle larvae and adults, stem weevil, sunflower moth, banded sunflower moth, woollybears, seed weevil, and grasshoppers.

Preplant Incorporation Treatment

Use AKOFOS 48 EC insecticide at the following rates by application in sufficient water to the soil surface and incorporate into the soil:

PEST	AKOFOS 48 EC
Cutworms	2-4 pints/acre (1 - 2 lbs. a.i./acre)

Use listed rate in not less than 10 gallons of water per acre and apply as a broadcast spray to the soil surface using suitable power-operated ground spray equipment. On the same day of treatment, incorporate the insecticide into the top 2 to 4 inches of soil using a disc, field cultivator, or equivalent equipment.

Postemergence Treatment

Use AKOFOS 48 EC for control of the following pests at the dosage indicated by application in sufficient water to ensure thorough coverage of treated plants:

PEST	AKOFOS 48 EC
Cutworms	2-3 pints/acre (1-1.5 lbs. a.i./acre)
Sunflower beetle larvae and adults, Stem weevil, Sunflower moth, Banded sunflower moth, Woollybears, Seed weevil	1-1½ pints/acre (0.5-0.75 lb. a.i./acre)
Grasshoppers	1 pint/acre (0.5 lb. a.i./acre)

Apply as a broadcast spray using either aerial (fixed-wing or helicopter) or power-operated ground spray equipment when field counts indicate that pests are or may become a problem. For cutworm control, a second treatment may be made 7 to 10 days later if needed. For stem weevil control, optimal treatment time is within 5 to 7 days after adult weevils begin to appear. For sunflower moth control, make first application during early 1 to 5 percent bloom stage. A second treatment may be made 10 days later if needed. For seed weevil control, treat when field counts indicate there are 10 to 12 adults per plant for oil crops and 1 to 3 adults per plant on confectionery crops.

Additional treatments should be made at successive 10-day intervals if field counts indicate need to retreat. For sunflower beetle larvae or adult control, treat when field counts indicate there are 10 larvae or 1 to 2 adults per seedling. Additional treatments may be made at successive 10-day intervals if field counts indicate need to retreat.

RESTRICTIONS: Do not apply more than 6 pints (3 lbs. a.i.) of AKOFOS 48 EC per acre per season. Do not make more than 3 applications per season. Do not apply within 42 days before harvest. Do not make a second application within 10 days of the first application. Do not allow livestock to graze in treated areas. The maximum single application rate is 2.0 lbs. a.i. chlorpyrifos per acre for preplant incorporation and 1.5 lbs. a.i. chlorpyrifos per acre for postemergence broadcast treatment. Not for use in Mississippi.

SWEET POTATOES

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Use AKOFOS 48 EC to reduce the feeding damage caused by populations of *Conderus* wireworm, *Systema* flea beetle, and the sweet potato flea beetle. Apply at the rate of 4 pints (2 lbs. a.i.) per acre as a broadcast (overall) spray to the soil surface followed by incorporation. Mix the specified dosage with enough water to obtain uniform coverage and apply as a coarse spray using suitable ground spray equipment. Incorporate the insecticide to a depth of 4 to 6 inches as soon as possible after application by using a rotary hoe, disc cultivator, or other suitable incorporation equipment. Plant the crop in the usual manner no later than 14 days after treatment (any delay in planting will reduce the length of time that AKOFOS 48 EC will protect against feeding damage). AKOFOS 48 EC will not control false wireworms or whitefringed beetle or other grubs that attack sweet potatoes.

RESTRICTIONS: Do not make more than one application per season. Do not harvest within 125 days of treatment. The maximum single application rate is 2.0 lbs. a.i. chlorpyrifos per acre. Do not aerially apply this product in Mississippi.

TOBACCO

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Use AKOFOS 48 EC for preplant treatment to control larvae of cutworms, flea beetles, mole crickets, root maggots, and wireworms. Apply 2 pints (1 lb. a.i.) of AKOFOS 48 EC per acre in not less than 10 gallons of water as a broadcast (overall) spray to the soil surface one week before transplanting. Immediately following application, incorporate the insecticide into the soil to a depth of 2 to 4 inches using suitable equipment.

To control the above insects and low to moderate populations of rootknot nematodes in North Carolina, South Carolina, and Virginia, use AKOFOS 48 EC at the rate of 2 pints (1 lb. a.i.) per acre.

Apply the specified dosage in not less than 10 gallons of water as a broadcast (overall) spray to the soil surface 24 to 48 hours before bedding and transplanting. Immediately following application, incorporate into the soil surface 24 to 48 hours before bedding and transplanting. Immediately following application, incorporate into the soil to a depth of at least 4 inches using suitable equipment.

RESTRICTIONS: Do not apply more than 2 pints (1 lb. a.i.) per application. Do not make more than one application per season. Do not apply more than 2 pints (1 lb. a.i.) per acre per season. Do not aerially apply this product in Mississippi.

TREE FRUITS (Dormant/Delayed Dormant Sprays)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 days unless PPE required for early entry is worn.

Use AKOFOS 48 EC as a dormant or delayed dormant spray at the rates indicated to control the following insects on the crops listed. While AKOFOS 48 EC may be used without oil, oil is recommended to control additional pests such as European red mite and brown almond mite. Applications may be made on pears after harvest. See specific use directions below.

CROP	PEST	AKOFOS 48 EC PER 100 GALLONS OF SPRAY*
Apples (1)	Rosy apple aphid, San Jose scale, Lygus, <i>Pandemis</i> leafroller, Climbing cutworms, Obliquebanded leafroller	$\frac{1}{2}$ -1 pint (0.25 – 0.5 lb. a.i.) (Use a minimum of 1.5 pints/acre)
Pears	San Jose scale, Climbing cutworms, Pear psylla adults	
Plums, Prunes	San Jose scale, Mealy plum aphid, Climbing cutworms, Peach twig borer	
Almonds, Peaches, Nectarines	San Jose scale, Peach twig borer, Climbing cutworms	

* Based on 200 to 600 gallons per acre as a dilute spray.

(1) Only one application of any chlorpyrifos containing product may be made per year. The application can either be a prebloom dormant/dormant delayed spray to the canopy or the trunk, OR a post-bloom application to the lower 4 feet of the trunk. For postbloom application instructions and restrictions for apple, see the Apple (Tree Trunk) section.

For dilute spray, tank mix the specified dosage with 1 to 2 gallons of a petroleum spray oil recommended for dormant use in 100 gallons of water and spray the entire tree by application to run off using suitable ground spray equipment. (See **Additional Precautions Specific to California** section for use in California.)

For low volume (concentrate) sprays (less than 200 gallons of spray mixture per acre), use the same amount of AKOFOS 48 EC as for a dilute spray and apply in a manner that will ensure thorough coverage of the trees. Use the higher dosage of AKOFOS 48 EC for severe infestations. Use oil as recommended by your State Agricultural Experiment Station or Extension Service Specialist.

PRECAUTIONS: Because cold or dry conditions may cause AKOFOS 48 EC plus oil sprays to infuse trees resulting in bud damage or drop, do not apply until winter rains or irrigation has replenished soil moisture such that bark and twigs are not desiccated.

RESTRICTIONS: Do not use more than 4 pints of AKOFOS 48 EC per acre.

Additional Precautions Specific to California: Use a minimum of 100 gallons of total spray volume per acre. Do not use more than 4 gallons of spray oil per acre on almonds, and 6 gallons per acre on peaches and nectarines. Do not use any adjuvants or surfactants in addition to or as a substitute for a petroleum spray oil in a tank mix with AKOFOS 48 EC. Do not apply on almonds in the following counties in California: Butte, Colusa, Glenn, Solano, Sutter, Tehama, Yolo, and Yuba. Do not use more than 1% dormant oil in almond orchards less than 4 years old.

RESTRICTIONS: Make only one application during the dormant/delayed dormant season, applying no more than 4 pints (2 lbs. a.i.) per acre per season. Do not allow meat or dairy animals to graze in treated orchards. The preharvest interval restriction is 21 days. Only one application of any chlorpyrifos containing product may be made per year. The application can either be a prebloom dormant/dormant delayed spray to the canopy or the trunk, OR a post-bloom application to the lower 4 feet of the trunk. For postbloom application instructions and restrictions for apple, see the Apple (Tree Trunk) section.

Do not make a soil or foliar application within 10 days of a dormant/delayed dormant application of chlorpyrifos to the orchard.

TREE NUTS

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Foliar Spray: Use AKOFOS 48 EC at the dosage indicated by application as a foliar spray to control pests listed in the following table. Mix the required dosage in sufficient water to ensure thorough and complete coverage of the foliage and crop, and apply as a concentrate or dilute spray using conventional, power-operated spray equipment. For dilute sprays applied to tree nut crops, mix the required dosage in sufficient water to allow for spray to runoff. For concentrate sprays, apply an equivalent amount of AKOFOS 48 EC per acre. Treat when pests appear or in accordance with local conditions. Insect control by aerial application may be less than control by ground application because of less coverage. Consult your state agricultural experiment station, certified pest control advisor, or extension service specialist for specific use information in your area. To avoid contamination of irrigation floodwaters, do not flood irrigate within 24 hours following an application of AKOFOS 48 EC. AKOFOS 48 EC is highly toxic to bees exposed to direct treatment. Do not apply when bees are actively foraging in the treated area.

ALMONDS, FILBERTS, PECANS, WALNUTS

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Use AKOFOS 48 EC at the rates indicated to control the listed pests.

CROP	PEST	DOSAGE
Almonds	Navel orangeworm, Peach twig borer, San Jose scale	4 pints/acre (2 lbs. a.i./acre)
Filberts	Eye-spotted bud moth, Filbert aphid, Filbert leafroller, Filbert worm, Obliquebanded leafroller, Omnivorous leaf-tier, Winter moth	3-4 pints/acre (1.5 –2 lbs. a.i./acre)
Walnuts	Codling moth, Walnut scale, Walnut husk fly	4 pints/acre (2 lbs. a.i./acre)
Pecans	Pecan nut casebearer, Fall webworm	$1\frac{1}{2}$ - 4 pints/acre (0.75 –2 lbs. a.i./acre)

	Phylloxera spp.1, Black pecan aphid, Hickory shuckworm 2, Pecan leaf scorch mite (suppression) 3, Ant species (except fire ants, carpenter ants, harvester ants, and pharaoh ants) 4	2-4 pints/acre (1- 2 lbs. a.i./acre)
	Yellow pecan aphid5, Black margined aphid5, Spittlebugs	1-4 pints/acre (0.5-2 lbs. a.i./acre)

1. For best *Phylloxera* spp. control, make 2 applications at 10 day intervals using a minimum of 1.0 pint of AKOFOS 48 EC per acre starting at bud swell.
2. For best results, make 2 applications, 10-14 days apart.
3. To suppress pecan leaf scorch mite, use a preventative program.
4. For ant control, apply as an orchard floor spray. Do not apply where weed growth or other obstructions prevent uniform coverage of the orchard floor.
5. For control of yellow pecan aphid and black margined aphid, apply in tank mix combination with the listed rate of a pyrethroid insecticide labeled for control or suppression of these aphids.

Dormant or Delayed Dormant Spray: For control of additional pest such as European red mite, it is recommended to mix AKOFOS 48 EC with oil, although it can be used without oil. Using conventional, power-operated spray equipment, apply as a concentrate or dilute spray. For dilute sprays (greater than 200 gpa), use sufficient spray volume to saturate tree foliage, but not to point of runoff. For concentrate sprays (less than 200 gpa), evenly distribute an equivalent amount of product per acre.

Precautions: Cold or dry conditions may cause this product and oil mixtures to infuse into trees, resulting in bud damage or bud drop. Do not apply until winter rains or irrigation has replenished soil moisture so that bark and twigs are not desiccated. Do not flood irrigate within 24 hours of application of this product to avoid contamination of irrigation tail waters.

RESTRICTIONS: Make no more than 3 foliar applications per season on almonds, filberts, and pecans and no more than 2 applications per season on walnuts. Make no more than one application of chlorpyrifos during the dormant season. Do not make a soil or foliar application of AKOFOS 48 EC or products containing chlorpyrifos within 10 days of a dormant/delayed dormant application. Do not apply within 14 days of harvest of almonds, filberts, and walnuts, or 28 days of harvest of pecans. Do not allow livestock to graze in treated orchards. Do not apply more than 8 pints (4 lbs. a.i.) per acre per season as a foliar spray. Do not make a second application within 10 days of the first application. Do not apply more than 4 pints of AKOFOS 48 EC (2.0 lbs. a.i. chlorpyrifos) per acre per season as a dormant/delayed dormant application. Do not aerially apply this product in Mississippi.

CALIFORNIA RESTRICTIONS:

Do not use more than 1% dormant oil and/or penetrating surfactants in almond orchards less than 4 years old. Use a minimum of 100 gpa of total spray volume. Use up to 2% supreme oil with no more than 4 gpa on almonds. In orchards with high overwintering populations of European red mite or brown almond mite, use the higher spray volumes that allow for the use of higher per acre rates of oil. Do not use any adjuvants or surfactants in addition to, or as a substitute for petroleum spray oil in a tank mix with this product. Do not apply on almonds during dormant and delayed dormant in the following counties in California: Butte, Colusa, Glenn, Solano, Sutter, Tehama, Yolo, and Yuba.

ALMOND, PECAN, WALNUT ORCHARD FLOORS

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Use AKOFOS 48 EC to control ants (except fire ants, carpenter ants, harvester ants, and pharaoh ants), and pavement ant by applying the specified dose with ground application equipment that will uniformly apply the spray to the orchard floor. Use when ant activity becomes evident within the orchard.

Since worker ants cease most of their foraging activity at temperatures above 90°F, best results will be achieved with applications made at temperatures below 90°F at the time of application. Dosage of AKOFOS 48 EC and spray volume may vary depending on the irrigation method employed in the orchard as follows:

Ant Control in Sprinkler- or Drip-irrigated Orchards

Apply AKOFOS 48 EC as a broadcast spray to the entire orchard floor using ground spray equipment at 4 to 8 pints (2 -4 lbs. a.i.) per acre in 25 or more gallons of water. Use the high rate for heavy infestations and the low rate for light infestations. In orchards where ant activity is concentrated around the irrigation emitters, apply the high rate to a 6- to 8-foot band along the drip-irrigation line and the low rate to the rest of the orchard.

Ant Control in Flood-irrigated Orchards

Apply AKOFOS 48 EC at 4 to 8 pints (2-4 lbs. a.i.) per acre in 25 or more gallons of water to the entire orchard floor using ground spray equipment. Apply the high rate to heavily infested areas and the low rate to lightly infested areas. Where ant colonies are abundant only in the berm areas, apply AKOFOS 48 EC at 8 pints (4 lbs. a.i.) per treated acre in 50 or more gallons of water to a 6- to 10-foot band along the treeline (berm).

Do not apply where weed growth or other obstructions would impede uniform coverage of the orchard floor. Mow or chemically control weeds before the application of AKOFOS 48 EC. Foliar applications of AKOFOS 48 EC may be made in addition to the orchard floor treatment.

RESTRICTIONS: Do not make more than 2 applications per season. Do not apply more than 8 pints (4 lbs. a.i.) per acre per season to the orchard floor. If the 8 pint rate is used, then a second application is not permitted. Do not apply the last treatment within 14 days of harvest. Do not allow livestock to graze in treated orchards. Do not allow spray to contact fruit or foliage. Do not make a second application within 10 days of the first application.

TURF GRASS GROWN FOR COMMERCIAL SOD

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Use AKOFOS 48 EC to control the pests listed in the following table by application at the listed dosages. Dilute AKOFOS 48 EC in water and apply using suitable application equipment. For best results, turf should be moist at time of treatment.

PESTS	AMOUNT OF AKOFOS 48 EC	
	1000 sq. ft.	Acre
Ants (except fire ants, carpenter ants, harvester ants, and pharaoh ants), armyworms (such as: beet, fall, yellowstriped), chinch bugs, crickets, cutworms, earwigs, European crane fly larvae, fiery skipper, gnats, grasshoppers, greenbug aphids, green June beetle grubs, leafhoppers, Lucerne moth, millipedes, mites (such as: clover, Bermudagrass stunt, winter grain), pillbugs, springtails, sod webworms (lawn moths) (1), sowbugs	¾ fl oz	1 qt. (1 lb. a.i.)
Billbug adults (such as: bluegrass, Denver, hunting) (2)	¾ - 1 ½ fl. oz.	1-2 qt. (1-2 lbs. a.i.)
Annual bluegrass weevil (<i>Hyperodes</i>) (3), black turfgrass ateniens adults (4), mole crickets (5)	1 ½ fl. oz.	2 qt. (2 lbs. a.i.)
White grubs (such as: black turfgrass ateniens, European chafer, Japanese beetle larvae, and northern and southern masked chafers)(6)	1 ½ - 3 fl. oz.	2-4 qt. (2-4 lbs. a.i.)

Numbers in parentheses (-) refer to Specific Use Directions below.

SPECIFIC USE DIRECTIONS:

1. For sod webworms, watering or mowing of the treated area should be delayed for 12 to 24 hours after treatment.
2. For billbugs, spray early in the season just prior to or coinciding with first appearance of adults as recommended by your local Agricultural Extension Service Specialist.
3. To control annual bluegrass weevil, spray suspected problem areas in mid-April and again in mid May, or as recommended by your local Agricultural Extension Service Specialist.
4. For black turfgrass ateniens adults, spray early in the season as recommended by your local Agricultural Extension Service Specialist. A repeat application may be needed 1 to 2 weeks later.
5. To control mole crickets in turfgrass, apply AKOFOS 48 EC through high-pressure injection or other suitable subsurface placement application equipment. Depending on the application equipment used, follow the manufacturer's recommendation for calibration and the volume of spray per acre needed to provide control or as recommended by your local Agricultural Extension Service Specialist. For best results, apply when young nymphs are active.
6. For white grubs, spray when grubs are young and actively feeding near the soil surface, usually during late July and August or as recommended by your local Agricultural Extension Service Specialist. For best results, soil should be moist prior to treatment. For best results, immediately after spraying, irrigate the treated area with ½ to 1 inch of water to wash the insecticide into the thatch and underlying soil.

VEGETABLES (Brassica, Cole Crops)

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 3 days for cauliflower and 24 hours for all other vegetables unless PPE required for early entry is worn.

Use AKOFOS 48 EC at the dosages indicated to control the pests listed in the following table. To avoid phytotoxicity in vegetables except Brussels sprouts, do not mix with other pesticide products or treat plants that are under extreme heat and drought stress.

CROP	PEST	DOSAGE AKOFOS 48 EC	USE DIRECTIONS	RESTRICTIONS
Cauliflower-soil application	Root maggot	1.6-2.4 fl. oz./1000 linear ft. of row	For direct-seeded crops, apply the specified dosage in a water-based spray as a 4-inch wide band over the row at planting time. Shallow incorporation is necessary. Placement behind the planter shoe and in front of the press wheel is recommended. For transplanted crops, apply AKOFOS 48 EC as a water-based spray directed to the base of the plants immediately after setting. Use a minimum of 40 gallons of total spray per acre. Do not add any additional adjuvants, surfactants, or spreader stickers. Do not apply as a foliage application.	Cauliflower-soil Preharvest Interval Restriction is 21-30 days. Do not apply more than 2 pints of AKOFOS 48 EC to cauliflower planted in 40-inch rows. Use proportional amounts for other row spacings not to exceed 4 pints of AKOFOS 48 EC per acre. Do not apply more than 2.6 pints (1.3 lbs. a.i.) of AKOFOS 48 EC per acre to broccoli, Brussels sprouts, cabbage, Chinese cabbage, collards, kale, kohlrabi, and turnips planted in 40-inch rows. Do not apply more than 4.5 pints (2.25 lbs. a.i.) of AKOFOS 48 EC per acre to these crops in 20-inch rows (or two rows per bed). Use proportional amounts for other row
Broccoli, Brussels Sprouts, Cabbage, Chinese Cabbage, Collards, Kale, Kohlrabi, Turnips-soil application		1.6-2.75 fl. oz./1000 linear ft. of row		

Broccoli, Cabbage-soil application	Root aphid	1.2 fl. oz./1000 ft. of row for single row plantings, and 2.4 fl. oz./1000 linear ft. of row for double row plantings	Apply AKOFOS 48 EC in a water emulsion or with liquid fertilizer injected as a sidedress on each side of the row after plants are established. Avoid mechanical damage to crop roots. Use a minimum of 15 gallons of total spray volume per acre. See MIXING DIRECTIONS section of the label for mixing instructions with liquid fertilizers.	spacings not to exceed 4½ pints (2.25 lbs. a.i.) of AKOFOS 48 EC per acre. Do not make more than one application per season within 30 days before harvest on broccoli, Brussels sprouts, cabbage, Chinese cabbage, collards, kale, kohlrabi, and turnips. Do not make more than one soil application per crop. Do not make a foliar application within 10 days of a soil application.
Brussels sprouts-foliar application	Armyworms, Cabbage aphid, Cutworms, Imported cabbage worm, Striped flea beetle (adult)	1-2 pints/acre (0.5-1 lb. a.i./acre)	Apply AKOFOS 48 EC with conventional power-operated equipment in 20 to 150 gallons of water per acre. Apply when insects appear on foliage and at 10 to 14 day intervals or thereafter as needed. Consult your State Agricultural Station, Extension Service Specialist, or Integrated Pest Control Advisor for proper time to treat in your area.	Do not make more than 3 applications per season. Do not apply within 21 days before harvest. Do not make a second application within 10 days of the first application. Do not make a foliar application within 10 days of a soil application.

CROP	PEST	DOSAGE AKOFOS 48 EC	USE DIRECTIONS	RESTRICTIONS
Radishes-soil application	Root maggot	1.0 fl. oz./1000 linear ft. of row	Apply the specified dosage as a water-based drench in the seed furrows with the seed at planting time. Use a minimum of 40 gallons of total drench per acre.	Do not apply more than 5½ pints (2.75 lbs. a.i.) of AKOFOS 48 EC per acre or make more than one soil application per season. Do not make a foliar application within 10 days of a soil application. Do not apply within 30 days of harvest.
Rutabagas-soil application	Root maggot	1.6-3.2 fl. oz./1000 linear ft. of row	Apply the specified dosage in a water-based spray as a 4-inch wide band over the row at planting time, behind the planter shoe and in front of the press wheel to achieve shallow incorporation. Use a minimum of 40 gallons of total spray volume per acre.	Do not apply more than 4½ pints (2.25 lbs. a.i.) of AKOFOS 48 EC per acre or make more than one soil application per season. Do not use rutabaga tops for food or feed purposes. Do not make a foliar application within 10 days of a soil application. Do not apply within 30 days of harvest.

WHEAT

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

For use only in Arizona, California, Colorado, Idaho, Kansas, Minnesota, Montana, Nebraska, New Mexico, Nevada, North Dakota, Oklahoma, Oregon, South Dakota, Texas, Utah, Washington, and Wyoming.

For use to control aphids (including Russian wheat aphid), wheat midge, brown wheat mite, grasshoppers, army cutworms and to provide suppression of other cutworm species.

PEST	DOSAGE	TIMING/SPECIAL DIRECTIONS
Aphids (including Russian wheat aphid, Greenbug, English grain aphid), Brown wheat mite,	½ to 1 pint/acre (0.25-0.5 lb. a.i./acre)	From emergence to flowering, treat when 15-20% of tillers are infested. From flowering to early milk stage, treat when 20% or more of tillers are infested.

Grasshoppers		
Wheat midge (orange wheat blossom midge)	1 pint/acre (0.5 lb. a.i./acre)	Treatment is recommended when 75% of the wheat heads have emerged from the boot and when midge adults are found in the crop (1 midge per 4-5 heads). Application timing is critical to ensure good control. If possible, apply in the late afternoon or early evening when temperatures exceed 50° F and wind speed is less than 7 mph.
Army cutworms, other cutworm species (suppression only)	1 pint/acre (0.5 lb. a.i./acre)	Control may be reduced under high temperature conditions (greater than 80°F), under dry soil conditions, or if larvae are more than ½ inch long. Treat when field counts or crop injury indicates that damaging pest populations are developing or present. A second application of 1 pint/acre may be made for additional control.

Mix the required dosage with water and apply in a minimum of 2 gallons per acre finished spray volume. Apply using aerial (fixed wing or helicopter) or power-operated ground spray equipment.

RESTRICTIONS: Do not make more than 2 applications per season. Do not apply within 14 days of harvest for forage and hay and within 28 days of harvest for grain and straw. Do not allow livestock to graze or otherwise feed on treated forage within 14 days of application. Do not feed straw from treated wheat within 28 days of application.

TURF AND OTHER NONRESIDENTIAL OUTDOOR USES

AKOFOS 48 EC is an emulsifiable concentrate for control of pests located around industrial buildings (turf and ornamental), road medians (turf and ornamentals), and golf course turf only. The maximum single application rate for golf course is 1 lb. ai/A. Pests controlled by AKOFOS 48 EC are listed in the following tables. AKOFOS 48 EC is compatible with fungicides, insecticides, and miticides commonly recommended except for alkaline materials such as Bordeaux mixtures and lime. A small jar compatibility test should always be conducted using proper proportions of chemicals and water to check for physical compatibility prior to tank mixing.

RESTRICTIONS: Do not treat vegetable gardens. Do not allow livestock to graze in treated areas. Do not feed treated grass cuttings (hay) or seed screenings to livestock or use hay for livestock bedding. Do not use AKOFOS 48 EC in poultry houses. Keep out of fish pools and other bodies of water.

ORNAMENTALS AROUND INDUSTRIAL BUILDINGS AND ROAD MEDIANS

AKOFOS 48 EC may be used to treat evergreens, vines, flowers, shrubs, shade and flowering trees, non-bearing fruit, nut and citrus trees found around industrial buildings and road medians infested with pests listed in the following table. Dilute AKOFOS 48 EC with water according to the directions given in the table and apply using suitable hand- or power-operated spray equipment. Ensure complete and uniform coverage. Uniform coverage is critical for effective insect and mite control. Apply a coarse spray to thoroughly wet both upper and lower leaf surfaces and infested limb and trunk areas. Attempt to penetrate dense foliage but avoid over-spraying to the point of excessive runoff. Treat when pests appear and repeat at 7 to 10 day intervals, if needed. For application timing and other specific use information, consult your State Agricultural Experiment Station or Extension Service Specialist.

Note: Environmental factors have significant effects on phytotoxic expression. AKOFOS 48 EC has been tested on numerous ornamental plants without causing serious phytotoxicity at listed use rates. Some varieties of azaleas, camellias, poinsettias, rose bushes, or variegated ivy have shown varying degrees of phytotoxicity following treatment with AKOFOS 48 EC. Before treating large numbers of plants (especially those previously listed), treat a small block of plants and observe for 7 to 10 days to determine phytotoxic potential.

PEST*	Amount of AKOFOS 48 EC in Water to Make		Specific Directions
	Per Acre	100 Gallons	
Adelgids: (Cooley, Eastern spruce galls, Pine bark), Ants (except fire ants, carpenter ants, harvester ants, and pharaoh ants), Aphids: (Apple, Chrysanthemum, Cottonwood, Elm leaf, Peach, Rose, Spirea, Woolly), Armyworms: (Fall, Yellowstriped), Bagworms ¹ , Boxelder bugs, Cankerworms, Catalpa sphinx, Chiggers (for control of chiggers in golf courses, road medians, and industrial sites only), Mealy bugs, Elm spanworms, Fall webworms ² , Grasshoppers, Green fruitworms, Hornworms, Jackpine budworms, Juniper webworms, Katydid, Lace bugs, Leafhoppers, Leafrollers ³ , Maple leafcutters ⁴ , Mites ⁵ : (Clover, Red spider, Southern red, Spruce spider, Twospotted spider), Oleander caterpillars, Orange tortrix, Periodical cicada, Plant bugs, Poplar tentmaker, Psyllids, Puss caterpillars, Rose chafers, Sawflies, exposed: (Pin oak, Pine, Redheaded), Sowbugs, Spittlebugs, Spring elm caterpillars, Springtails, Spruce budworms: (Eastern, Western), Tent caterpillars: (Eastern, Western, Forest), Thornbug, Walnut caterpillars, Whiteflies, Yellownecked caterpillars	1 pt. – 1 qt. (0.5 - 1 lb. a.i.)	8-16 fl. oz.	¹ Treat when bagworm larvae are small and actively feeding. ² Direct spray into web and immediately adjacent foliage for control of fall webworms. ³ For effective control of leafrollers, spray before leaves are tightly rolled. ⁴ Apply spray to maple leafcutter larvae as cases are being formed for effective control. Do not treat sugar maple trees intended for maple syrup production. ⁵ For effective control of spider mites when large numbers of eggs are present, apply a 2 nd spray 3-5 days in the South or 7-10 days in the North after initial treatment to control newly hatched nymphs.
Armyworms: (Beet), Beetles: (Fuller rose, Native elm bark ¹), Browntail moth, Cutworms, Leafhoppers, Mahogany webworms, Mealbugs, Mimosa webworms, Moths: (Browntail, Cypress tip, Douglas fir tussock, European pine shoot, Gypsy ² , Holly bud, Nantucket pine tip, Pandora, Pitch pine tip, Subtropical pine tip, Tussock, Oakworms: (California, Orangestriped, Redhumped), Redhumped caterpillars, Thrips: (Exposed), Weevils: (Blackvine ³ , Pine production, Yellow poplar)	1 qt. (1 lb. a.i.)	1 pt. (0.5 lb a.i.)	¹ Make applications in the spring or early summer to reduce twig and branch feeding by bark beetles. ² To kill migrating and invading gypsy moth larvae, treat trunks and foliage. ³ Blackvine weevils are night feeders. Late afternoon spraying will maximize control.
Foliar feeding beetles: (Blister leaf, Cottonwood leaf ¹ , Elm leaf, Flea, Fuller rose, Japanese, June, Willow leaf)	1 qt. (1 lb. a.i.)	1 pt. (0.5 lb. a.i.)	¹ Use AKOFOS 48 EC in water to control cottonwood leaf beetle larvae and adults infesting cottonwoods.

			Make the treatment when field counts indicate damaging beetle populations are developing or present.
Borers ¹ , Clearwing moths: (Ash, Dogwood, Lesser peachtree, Lilac, Oak, Rhododendron), Metallic wood: (Bronze birch, Flathead apple tree, Twolined chestnut), Longhorned beetles: (Locust, Red oak), Cranberry girdler larvae ² , Leafminers, Needleminers: (Jeffrey pine, Lodgepole pine, Spruce), Scale insects ³ : (Cottonycushion, Cottony maple, Euonymus, Fletcher, Florida wax, Golden oak, Hemispherical, Lecanium, Magnolia, Oak kermes, Oak lacanium, Oystershell, Pine needle, San Jose, Tea, White birch, White peach)	1 qt. (1 lb. a.i.)	1 qt. (1 lb. a.i.)	¹ For borers, apply AKOFOS 48 EC to the trunks and lower limbs of trees and shrubs when the adults begin to emerge. Consult your State Agricultural Experiment Station or Extension Service Specialist for proper time to treat. Apply uniformly a coarse, low-pressure spray. Pheromone traps may aid in detection of adult clearwing moths. ² Apply 1 qt. of AKOFOS 48 EC for cranberry girdler larvae. Direct spray at the base of tree using 50 gallons of water per acre. Irrigate immediately after applications for soil penetration of 1-2 inches. Treat after egg laying during the summer. ³ Time applications for control of scale insects when crawlers or first two stages of settled nymphs are present.
Borers: (Cottonwood, Peactree ¹)	1 qt. (1 lb. a.i.)	3 qts. ** (3 lbs. a.i.)	¹ For peachtree borers, apply AKOFOS 48 EC in water to flowering trees and shrubs of the genus <i>Prunus</i> as a trunk spray before newly-hatched larvae enter the trees. Apply as a coarse, low-pressure spray. Thoroughly wet all bark areas from ground level to scaffold limbs. ** When using the 3 qt. per 100 gallon dilution, do not exceed 1qt. of AKOFOS 48 EC per acre.

PEST*	Amount of AKOFOS 48 EC in Water to Make		PEST*
	Per Acre	100 gallons	
Beetles ¹ : (Includes wood infesting, Ambrosia, Anobiidae, Black turpentine, Cottonwood leaf, Elm leaf, European elm bark, Flea, Fuller rose, Japanese, June, Native elm bark ² , Southern pine, Willow leaf)	1 qt. (1 lb. a.i.)	2 gals. ***	¹ For preventative treatment, apply the spray to the main trunk of trees in the early spring or when threat of attack exists from nearby infested trees. For remedial treatment, apply the spray to the main trunk of infested trees when damage occurs but before adult beetles begin to emerge. ² To prevent native elm bark beetles from overwintering in uninfested trees, apply AKOFOS 48 EC in water to the bottom 9 ft. of the trunk. Wet the trunk thoroughly but do not spray to runoff. Care should be taken to apply the spray right to the base of the root flare. Application can be made with either a backpack mistblower or a hydraulic pressure sprayer from spring through early fall. *** When using the 2-gallon per 100-gallon dilution, do not exceed 1 qt. of AKOFOS 48 EC per acre.

FOR COMMERCIAL ORNAMENTALS IN NURSERIES

FOR USDA QUARANTINE USE ONLY AND FOR DIRECT MOUND AND/OR DRENCH APPLICATION ONLY

AKOFOS 48 EC may be used for USDA quarantine use only to treat containerized, potted, or balled and burlapped nursery stock to control the insects in the soil attached to the roots of these plants. Completely submerge the container with drain holes or root ball stabilized by burlap in a tank containing diluted AKOFOS 48 EC. Do not remove burlap wrap or plastic containers with drain holes prior to submerging. Keep the container or root ball submerged until complete soil saturation has occurred, normally about 30 seconds.

Precautions: During all operations (submerging, drenching, injecting), wear a chemical-resistant apron in addition to other PPE listed for applicators and other handlers. Make applications in a well-ventilated area.

Note: Environmental factors have significant effects on phytotoxic expression. AKOFOS 48 EC has been tested on numerous ornamental plants without causing serious phytotoxicity at listed use rates. However, because of the numerous varieties grown, treat a small group of plants at the listed rate under the anticipated growing conditions and observe for at least 7 days to determine phytotoxic potential before treating a larger number of plants.

PEST*	Amount of AKOFOS 48 EC in Water to Make		Specific Directions
	1 Gallon	100 Gallons	
Fire Ants ¹	1/25 fl. oz.	4 fl. oz.	¹ As an alternative to submerging potted plants, dilute 4 oz. of AKOFOS 48 EC in 100 gallons of water. Apply this dilution to the point of runoff twice daily for 3 consecutive days. Do not remove burlap wrap or container from plants prior to treatment.
White Grubs ² Weevils ³ (such as Blackvine) Coffee root mealybug ²	3/8 fl. oz. 1/6 fl. oz.	2 qts. ** (2 lbs. a.i.) 1 pt. (0.5 lb. a.i.)	² An alternative treatment to submerging containerized plants is to drench the container with the diluted insecticide solution applying approximately 10 to 12 fl. oz. of diluted insecticide solution per gallon of container size (4-5 fl. oz./100 cubic inches of container). The container media should be pre-moistened by irrigation or rainfall before drenching. Do not remove container from plants prior to treatment. ³ An alternate treatment to submerging balled and burlapped plants is to inject AKOFOS 48 EC into the root ball. Equally distribute 1 to 3 quarts of the dilute AKOFOS 48 EC solution per cubic foot of soil volume through an injection rod inserted into the soil ball surrounding the plant roots. Uniform distribution of the insecticide throughout the soil of the root ball is critical for effective control. It is recommended that the injection rod be inserted in at least 4 equally spaced locations around the stem of the plant at a 30-45 degree angle from the plant between the stem and the upper, outer perimeter of the ball. This technique has been shown to be most effective with small root balls (up to 1.5 ft in diameter). Larger root balls may require more injection points to ensure thorough soil distribution of the insecticide. The injection rod should be coupled to a flow meter to monitor the correct volume applied per root ball using an injection pressure of at least 30 psi. The application should be made such that splash-back and runoff are minimized. **Do not exceed more than 1 quart (1 lb. a.i.) of AKOFOS 48 EC per acre.

*Superscripts refer to specific directions.

IN INDUSTRIAL PLANT SITES AND ROAD MEDIANS (DORMANT SPRAY OF TREE PESTS) and ORNAMENTALS

AKOFOS 48 EC may be used as a dormant or delayed dormant spray at the rates indicated to control the listed insects. AKOFOS 48 EC may be used without oil; however, oil is recommended to control additional pests such as the European red mite.

For high volume (dilute) sprays (200 to 600 gallons of spray mixture per acre), tank mix the specified dosage with 1 to 2 gallons of a petroleum spray oil recommended for dormant use in 100 gallons of water. Spray the entire tree to runoff using suitable ground spray equipment.

For low volume (concentrate) sprays (less than 200 gallons of spray mixture per acre), use the same amount of AKOFOS 48 EC as for a dilute spray and apply in a manner that will ensure thorough coverage of the trees. Use oil as recommended by your State Agricultural Experiment Station or Extension Service specialist.

Precautions: Do not apply until rain or irrigation have replenished soil moisture such that bark and twigs are not desiccated since cold dry conditions may cause AKOFOS 48 EC plus oil to infuse trees resulting in bud damage or drop.

Restrictions: Make only one application during the dormant season except for the control of the apple ermine moth. Do not allow meat or dairy animals to graze in treated areas.

PEST*	Amount of AKOFOS 48 EC in Water to Make			Specific Directions
	1 Gallon	3 Gallons	100 Gallons	
Aphids: (Mealy plum, Rosy Apple, Woolly apple), Borers: (Peach twig), Cutworms: (Climbing), Leafrollers: (Pandemis), Pear psylla adults, Plant bugs, Scale: (San Jose)	1/12 – 1/6 fl. oz.	1/4 - 1/2 fl. oz.	1/2 - 1 pt.	Tank mix with 1-2 gallons of a petroleum spray oil recommended for dormant use in 100 gallons of water.
Apple ermine moth	1/12 fl. oz.	1/4 fl. oz.	1/2 pt.	For control on <i>Malus</i> species make 2 applications at a 7-14 day interval in combination with a petroleum spray oil at the rate of 2-4% (v/v) in a spray to wet application to ensure thorough coverage of all stems and branches. When using tank mixtures, follow all label directions for the mixing partner (oil). Use appropriate application equipment and spray volumes to ensure complete coverage of the plant(s) or control will be compromised.

TURF, INDUSTRIAL PLANT, AND ROAD MEDIAN OUTDOOR USES

Apply AKOFOS 48 EC to control the pests listed in the following table at the listed dosages and in accordance with the directions given below or as recommended by your local Agricultural Extension Service specialist. Dilute AKOFOS 48 EC in water and apply as a coarse, low-pressure spray using suitable application equipment. Except as noted, thoroughly water immediately after treatment to wash the insecticide into the turf. The area to be treated should be moist at the time of treatment. Spray when pests first appear; retreat when needed. The maximum single application rate of golf course turf is 1 lb. ai/A.

PEST*	Amount of AKOFOS 48 EC per		Specific Directions
	1,000 sq. feet	Acre	
Ticks ¹ : (American dog, Cattle fever, Gulf coast, Lone star)(for control of ticks in golf courses, road medians, and industrial plant sites only)	¼ fl. oz.	1 ½ pts.	¹ For control of ticks, treat soil and other areas likely to serve as harborage sites for ticks that have removed themselves from their host. Spray surfaces to be treated until wet but do not create excessive runoff. Note: This application is intended as a premise spray only. Do not use as a direct spray on livestock or any sites that may come in contact with livestock.
Ants (except fire ants, carpenter ants, harvester ants, and pharaoh ants), Armyworms: (Beet, Fall, Yellowstriped), Chiggers ¹ (for control of chiggers in golf courses, road medians and industrial plant sites only), Chinch bugs, Clover mites, Cutworms, Crickets, Deer ticks ² (for control of ticks in golf courses, road medians, and industrial sites only), Earwigs, Fiery skipper, Gnats, Grasshoppers, Greenbug aphids, June beetles, Leafhoppers, Lucerne moths, Millipedes, Mites: (Clover, Bermudagrass stunt, Formula grass, Winter grain), Pillbugs, Sod webworms ³ : (Lawn moths), Sowbugs, Ticks ¹ (for control of ticks in golf courses, road medians and industrial plant sites only)	¾ fl. oz.	1 qt.	¹ Apply AKOFOS 48 EC for area control of ticks and chiggers infesting golf course turf, turf in road medians, and industrial plant sites where these pests are present and create a nuisance or a possible public health problem. Do not allow public use of treated areas during application or until spray has dried. Apply AKOFOS 48 EC in water at the rate of ½ pint/acre (equivalent to 1/6 fl. oz. per 1,000 sq. ft.) using a hydraulic sprayer, mist applicator, knap sack sprayer, or other suitable hand or power-operated spray equipment. Treat low underbrush, grassy areas, weeds, and ground surface and debris using enough spray volume to obtain thorough coverage, usually 40-100 gals./acre. ² Apply AKOFOS 48 EC in water at the rate of 1 quart per acre or ¾ fl. oz. per 1,000 sq. ft. for control of deer ticks. Treat low underbrush, turf, grassy areas, weeds, and ground surface and debris, using enough spray volume to obtain thorough coverage. ³ For sod webworms, watering or mowing of the treated area should be delayed for 24 hours after treatment.
European crane fly	1 fl. oz.	1 qt.	
Turfgrass weevil (<i>Hyperodes</i>) ¹	1 ½ fl. oz.	1 qt.	¹ Make application to problem areas in mid-April and again in mid-May or as recommended by your local Agricultural Extension Service specialist.
White grubs ¹ : (Black turfgrass atatenius, European chafer, Japanese beetle larvae, Southern and Northernmasked chafer)	1 ½ - 3 fl. oz.	1 qt.	¹ Spray when white grubs are young and actively feeding near the soil surface, usually during late July and August or as recommended by your local Agricultural Extension Service specialist. Immediately after spraying, irrigate the treated area with ½ to 1 inch of water to wash the insecticide deep into the thatch or into the underlying soil.
Billbug adults, such as: (Bluegrass, Denver, Hunting)	¾ - 3 fl. oz.	1 qt.	Spray early in the season when adult billbugs first appear.
Mole Crickets	1 ½ fl. oz.	1 qt.	For mole crickets in golf course turf, turf in road medians, and industrial plant site turfgrass, apply through high-pressure injection or other suitable subsurface placement application equipment. Depending on the application equipment used, follow the manufacturer's recommendation for calibration and the volume of spray per acre needed to provide control or as recommended by your local Agricultural Extension Service specialist. Apply when young nymphs are active.

*Superscripts refer to Specific Directions.

OUTSIDE SURFACES AND AROUND INDUSTRIAL PLANT SITES (SUCH AS AROUND WAREHOUSES, FOOD PROCESSING AND FOOD MANUFACTURING SITES)

AKOFOS 48 EC may be applied as a residual spray to and around outside surfaces of nonresidential buildings and structures. Permitted areas of use include fences, pre-construction foundations, refuse dumps, outside of walls, and other areas where pests congregate or have

been seen. Do not allow adults, children, or pets to contact treated surfaces until sprays have dried. Keep out of fish pools and other bodies of water. Do not feed treated grass cuttings (hay) or seed screenings to livestock, or use treated hay for livestock bedding. Do not treat vegetable gardens. Repeat treatment as needed to maintain effectiveness. Unless prohibited by a products' label, users, at their own discretion, can tank mix pesticides currently labeled for similar use patterns. Always perform a small jar compatibility test using proper proportions to check for physical compatibility prior to tank mixing. Do not tank mix this product with products containing dichlorvos (DDVP).

PEST*	Amount of AKOFOS 48 EC in Water to Make			Specific Directions
	1 Gallon	10 Gallons	50 Gallons	
	For Band Treatment ¹			
Ants (except fire ants, carpenter ants, harvester ants, and pharaoh ants), Beetles, Boxelder bugs (or other true bugs), Clover mites, Crickets, Earwigs, Elf leaf beetles (adults), Firebrats, Millipedes, Pillbugs, Silverfish, Sowbugs, Spiders (excluding black widow and brown recluse spiders), Springtails, Ticks (for control of ticks in golf courses, road medians and industrial plant sites only.	¼ tsp.	¾ fl. oz.	4 fl. oz.	¹ To help prevent infestation of non-residential buildings, treat a band of soil 6-10 ft. wide around and adjacent to buildings including the building foundation to a height of 2-3 ft. where pests are active and may find entrance. Use 4 fl. oz. of AKOFOS 48 EC per 50 gals. of water and apply as a coarse spray at the rate of about 10 gals. of spray mixture per 1,000 sq. ft. to thoroughly and uniformly wet the band area. Maximum application rates for a single application: Golf course turf: 1 lb. ai/A
	For Outside Surfaces			
	1 ½ fl. oz.	13 ½ fl. oz.	2 qts.**	Do not exceed 1 qt. of AKOFOS 48 EC per acre.

* Superscripts refer to specific directions

ANTS (EXCEPT FIRE ANTS, CARPENTER ANTS, HARVESTER ANTS, AND PHAROAH ANTS) AND MISCELLANEOUS PESTS

PEST	Amount of AKOFOS 48 EC in Water to Make			Specific Directions
	1 Gallon	3 Gallons	100 Gallons	
Ants (except fire ants, carpenter ants, harvester ants, and pharaoh ants), Cockroaches: (American, Asian, Brownbanded, German, Oriental, Smokey brown, Wood), Sowbugs, Springtails	1/6 fl. oz.	½ fl. oz.	1 pt.	

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in secured dry storage area. Prevent cross-contamination with other pesticides and fertilizers. Do not store above 100°F for extended periods of time. Storage below 20°F may result in formation of crystals. If product crystallizes, store at 50°F to 70°F and agitate to redissolve crystals. If container is damaged or spill occurs, use product immediately or dispose of product and damaged container as indicated below.

PESTICIDE DISPOSAL: Open dumping is prohibited. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the hazardous waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

Nonrefillable Container (five gallons or less): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

LIMITATION OF WARRANTY AND LIABILITY

Read the entire directions for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following **CONDITIONS, DISCLAIMER OF WARRANTIES** and **LIMITATIONS OF LIABILITY**.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Aako B.V. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, Aako B.V. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Aako B.V. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Aako B.V. disclaims any liability whatsoever for special, incidental or consequential damages resulting from

the use or handling of this product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Aako B.V.'s election, the replacement of product.