



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Antimicrobials Division (7510P)
 1200 Pennsylvania Ave., N.W.
 Washington, D.C. 20460

EPA Reg. Number:

85837-6

Date of Issuance:

4/27/21

NOTICE OF PESTICIDE:

Registration
 Reregistration
 (under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Peraoxy Pro

Name and Address of Registrant (include ZIP Code):

Innovasource
 11515 Vanstory Drive, Suite 110
 Huntersville, NC 28078

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Antimicrobials Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Steven Snyderman, Product Manager 33
 Regulatory Management Branch 2
 Antimicrobials Division (7510P)
 Office of Pesticide Programs

Date:

4/27/21

EPA Form 8570-6

2. You are required to comply with the data requirements described in the DCI or EDSP Order identified below:

- a. Hydrogen Peroxide: GDCI-000595-1127; EPA-HQ-OPP-2009-0546

- b. Peroxyacetic Acid: GDCI-000595-1127, EPA-HQ-OPP-2009-0546-0013; GDCI-063201-1125, EPA-HQ-OPP-2009-0546-0012; GDCI-063604-1126, EPA-HQ-OPP-2009-0546-0011

You must comply with all of the data requirements within the established deadlines. If you have questions about the Generic DCI listed above, you may contact the Reevaluation Team Leader (Team 36): <http://www2.epa.gov/pesticide-contacts/contacts-office-pesticide-programs-antimicrobial-division>

3. Make the following label changes before you release the product for shipment:
- Revise the EPA Registration Number to read, “EPA Reg. No. 85837-6.”
4. Submit one copy of the final printed label for the record before you release the product for shipment.


Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. See FIFRA section 2(p)(2). If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process, FIFRA section 12(a)(1)(B). Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Assurance.

If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated November 13, 2020

If you have any questions, please contact Zebora Johnson by phone at (703) 308-7080, or via email at <mailto:johnson.zebora@epa.gov>.

Sincerely,



Steven Snyderman, Product Manager 33
Regulatory Management Branch II
Antimicrobials Division (7510P)

Page 3 of 3
Admin. No. 85837-6
Action Case No./Decision No. 00217108

Office of Pesticide Programs

Enclosure: Approved Label

Bactericidal, Virucidal Δ, Fungicidal, Tuberculocidal

For use on hard, non-porous surfaces.

Application:

PERAOXY PRO disinfects as it cleans in one operation. PERAOXY PRO can be used to disinfect floors, walls and other hard nonporous surfaces such as tables, chairs, countertops, bathroom fixtures, sinks, bed frames, doors, shelves, racks, carts, refrigerators, coolers, glazed tile, linoleum, vinyl, nonporous glazed porcelain, glazed ceramic, plastic (such as polypropylene and polyethylene), stainless steel, glass, aluminum, non-porous baked enamel, chrome, laminated or painted surfaces or sealed stone. PERAOXY PRO should not be used on marble or brass surfaces.

For health care, institutional, and industrial use.

Areas of use in hospitals: PERAOXY PRO may be used for surgical and obstetrical suites; housekeeping surfaces; physical therapy departments; nursing services; dental facilities; autopsy facilities; intensive care units; pharmacies; and clinical laboratories. Also, use PERAOXY PRO in nursing homes, other health-care facilities, schools, colleges, veterinary clinics, animal life science laboratories, industrial facilities, office buildings, recreational facilities, industrial facilities; hotels; retail facilities; office buildings; retail and wholesale establishments.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

Dilute PERAOXY PRO with the appropriate amount of water to an effective concentration of 1135ppm Peroxalic acid and 1665ppm hydrogen peroxide 0.65 fl. oz. per 1gallon of water. Apply solution with a cloth, mop, sponge, auto-scrubber, or hand pumped trigger sprayer such that all surfaces remain wet for 1 minute to kill bacteria, viruses, and fungi as cited on the label. Use a 3-minute contact time for Streptococcus pneumoniae, and Vancomycin-Resistant Enterococcus faecalis. Use a 10-minute contact time for Poliovirus Type 1 and Tuberculocidal activity. PERAOXY PRO is effective against Mycobacterium bovis (TB surrogate) at ambient temperature (22°C). Allow surface to air dry. For heavily soiled areas, a preliminary cleaning is required. Prepare a fresh solution daily or more often if the use solution becomes visibly soiled clouded or diluted.

This product is not for use on medical device surfaces.

(OR)

Kills HIV-1 (AIDS Virus), HBV, and HCV:

On pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids in health care settings or other settings in which there is an expected likelihood of soiling of surfaces/objects with blood or body fluids, and in which the surfaces / objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency virus Type 1 (HIV-1) (associated with AIDS), Hepatitis B Virus, or Hepatitis C Virus.

Special Instructions for cleaning and decontamination against HIV-1 (AIDS Virus), HBV, and HCV of surfaces/ objects soiled with blood/body fluids. Personal Protection: Disposable protective gloves, gowns, face masks, or eye coverings as appropriate must be worn during all cleaning of body fluids, blood and decontamination procedures. Cleaning Procedure: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application. Contact Time: HIV-1, HBV, and HCV are inactivated after 1 minute of contact.

Infectious material: Blood and other bodily fluids should be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal.

PERAOXY PRO is effective against the following organisms in 1 minute with 5% organic soil load and 400ppm hard water: For use on hard, non-porous surfaces.

* Contact time is increased to 10 minutes to be effective against Poliovirus Type 1 and Mycobacterium bovi (TB surrogate).

§ Contact time is increased to 3 minutes to be effective against Streptococcus pneumoniae and Vancomycin resistant Enterococcus faecalis (VRE).

Bacteria	Human virusesΔ
<i>Acinetobacter baumannii</i> ATCC 19606	Herpes simplex virus Type 1 ATCC VR-260
<i>Clostridium perfringens</i> ATCC 13124	Herpes simplex virus Type 2 ATCC VR-734
Community-associated MRSA, USA300 strain (CA-MRSA) CI 08001	Human immunodeficiency virus type 1
(HIV-1) (Zeptomatrix Corporation)	
<i>Escherichia coli</i> O157:H7 ATCC 35150	Human Influenza A virus (A/Hong Kong/8/68-H3N2) SPAFAS
<i>Klebsiella pneumoniae</i> ATCC 4352	Human Rotavirus ATCC VR-2018
<i>Listeria monocytogenes</i> ATCC 19111	Poliovirus Type 1* ATCC 1562
Methicillin-resistant <i>Staphylococcus aureus</i> (MRSA) ATCC 33592	Respiratory Syncytial Virus ATCC VR-26
Methicillin-resistant <i>Staphylococcus epidermidis</i> (MRSE) ATCC 51625	Rhinovirus Type 37 ATCC 1147
<i>Mycobacterium bovis</i> (TB surrogate)*	
(Organon Teknika Corp)	Vaccinia virus ATCC VR-156
<i>Pseudomonas aeruginosa</i> ATCC 15442	Human Coronavirus strain 229e
ATCC VR-740	
<i>Salmonella enterica</i> ATCC 10708	Bovine viral diarrhea virus (Human Hepatitis C surrogate) (American Bioresearch Laboratories)
<i>Salmonella enterica</i> serovar Typhimurium ATCC 13311	Duck hepatitis B virus (Human Hepatitis B surrogate) (Hepanda Virus Testing)
<i>Serratia marcescens</i> ATCC 13880	Feline calicivirus (Norovirus surrogate) (University of Ottawa)
<i>Shigella dysenteriae</i> serotype 1 ATCC 29026	
<i>Staphylococcus aureus</i> ATCC 6538	Animal viruses Δ
<i>Streptococcus pneumoniae</i> § ATCC 6304	Avian influenza A (Turkey/Wis/66-H9N2) SPAFAS
<i>Streptococcus pyogenes</i> ATCC 19615	
Vancomycin-intermediate <i>Staphylococcus aureus</i> (VISA) ATCC 700787	Fungi
Vancomycin-resistant Enterococcus faecalis (VRE) § ATCC 51575	<i>Candida albicans</i> ATCC 10231
<i>Vibrio cholerae</i> ATCC 14035	Trichophyton mentagrophytes ATCC 9533
<i>Yersinia enterocolitica</i> ATCC 35669	

FOR ANTIMICROBIAL USE WITH AQUEOUS TREATMENT FLUIDS IN SUBTERRANEAN OILFIELD AND GAS-FIELD WELL OPERATIONS SUCH AS WELL DRILLING, FORMATION

FRACTURING.

PRODUCTIVITY ENHANCEMENT AND SECONDARY RECOVERY.

PERAOXY PRO can be for control of slime forming

and spoilage bacteria, yeast and fungi and anaerobic sulfate reducing bacteria, that lead to reservoir souring and metal corrosion. This product must be introduced through a closed mixed/loading and delivery transfer system equipped with a metering device that is appropriate for its intended uses.

DRILLING MUDS, FRACTURING FLUIDS, WELL SQUEEZED

FLUIDS

For the preservation of drilling muds, workover and completion fluids and other product susceptible to contamination, pre-mix with the fluid or add directly at the point of use at 3.75 fluid ounces per 1000 gallons of water (5 ppm peroxyacetic acid) to 75 fluid ounces per 1000 gallons of water (100 ppm peroxyacetic acid) as required. Depending on the severity of the contamination, initial application may be added up to 749 fluid ounces per 1000 gallons of water (1000 ppm).

FLOODING, INJECTION AND PRODUCED WATER

For Water Flooding operations, add initially at 3.75 fluid ounces per 1000 gallons of water (5 ppm peroxyacetic acid) to 75 fluid ounces per 1000 gallons of water (100 ppm peroxyacetic acid) and repeat until control is achieved. Subsequent treatment may be continued on a weekly basis or as required.

Injection wells associated with gas storage systems may be treated up to 100 ppm when diluted in the formation water. Any additional top-up water should be treated as required.

For hydrostatic systems, apply 3.75 fluid ounces per 1000 gallons of water (5 ppm peroxyacetic acid) to 75 fluid ounces per 1000 gallons of water (100 ppm peroxyacetic acid) depending on the water quality and the duration of the shut-in.

PIPELINE AND TANK MAINTENANCE

For microbial control in water-bottoms in crude and refined hydrocarbon storage tanks, piping and transportation systems. Apply 3.75 fluid ounces per 1000 gallons of water (5 ppm peroxyacetic acid) to 75 fluid ounces per 1000 gallons of water (100 ppm peroxyacetic acid) in the aqueous phase, directly injected into the water-bottom, pipeline or may be added to the hydrocarbon phase.

Treatment may be applied daily or monthly for both storage and transportation systems as needed.

FOR DISINFECTION OF SEWAGE AND WASTEWATER EFFLUENTS IN TREATMENT PLANTS

Use PERAOXY PRO to treat sewage and wastewater effluent related to public and private wastewater treatment plants. PERAOXY PRO can be applied directly to the effluent or may be used with an appropriate activator such as hydrogen peroxide or other technology. PERAOXY PRO may be applied to effluent water discharged from trickle bed or percolating fluidized bed filters. The application rate for individual facilities will depend on the degree of bioloading of the effluent stream to be discharged and the local microbial discharge limit. Adjust application rate to meet the need of the individual facility.

1. Add PERAOXY PRO to effluent water at a concentration of 0.5 ppm to 15 ppm. Allow contact time of approximately 15 to 60 minutes.

2.The maximum amount of peroxy acid that can be discharged from the treatment facility is 1 ppm. Use an appropriate peroxy acid test kit analyzer to ensure that this level is not exceeded. Contact Innovasource, LLC or assistance establishing treatment regimes.

INFLUENT WATER SYSTEMS:

PERAOXY PRO should be fed continuously to incoming fresh water streams (nonpotable use only) at dosages ranging from 10 to 975 ppm peroxyacetic acid (65 to 6500 ppm PERAOXY PRO).

MILL PROCESS WATERS:

*Continuous Feed – PERAOXY PRO should be fed continuously at dosages ranging from 10 to 975 ppm peroxyacetic acid (65 to 6500 ppm PERAOXY PRO). This range is equivalent to 0.13 to 13 lbs. PERAOXY PRO per ton (dry basis) of pulp or paper produced.

*Intermittent Feed – PERAOXY PRO should be fed intermittently (6 to 8 times per day) at dosages ranging from 10 to 975 ppm peroxyacetic acid (65 to 6500 ppm PERAOXY PRO). This range is equivalent to 0.13 to 13 lbs. PERAOXY PRO per ton (dry basis) of pulp or paper produced.

*Shock Dose – PERAOXY PRO should be shock dosed at dosages ranging from 98 to 2048 ppm peroxyacetic acid (648 to 13,638 ppm PERAOXY PRO). This range is equivalent to 1.3 to 27.3 lbs. PERAOXY PRO per ton (dry basis) of pulp or paper produced.

CONTROL OF SLIME FORMING BACTERIA AND BIOFOULING IN ONCE-THROUGH AND RECIRCULATING COOLING WATER (COOLING TOWERS, EVAPORATIVE CONDENSERS, AIR WASHERS) AND ORNAMENTAL OR RECREATIONAL WATER FEATURES

Severely fouled systems must be cleaned before adding this product. This product must be added in the water system directly, and not mixed with any other chemicals or additives. Never add this product into any feeding device, such as sht feeders, filter housings, by-pass feeders, or miscellaneous piping of any kind, because dangerous acute decomposition can occur. Discontinue the use of chlorine or bromine products prior to using this product. Contamination with other chemicals could result in product decomposition. Add this product to only water at a point in the system where uniform mixing and even distribution will occur.

For shock (slug) treatment for moderately to severely fouled systems add 5-20 fl. oz. of this product per 1000 gallons of process water (7-27 ppm peroxyacetic acid). Repeat as necessary until microbiological control is evident.

Thereafter, to maintain control use 1.5-7.5 fl. oz. of this product per 1000 gallons of process water (2-10 ppm of peroxyacetic acid) as a continuous treatment method. Continuous dosing methods usually require 1.5-5 fl. oz. per 1000 gallons of water (2-7 ppm peroxyacetic acid) to achieve adequate results.

Intermittent dosing treatment usually require dose cycles of a minimum once per every other day, up to 6 times per 24 hours. Recommended rates for intermittent dose cycles are 5-10 fl. oz. of this product per 1000 gallons of process water (7-14 ppm peroxyacetic acid).

STORAGE AND DISPOSAL

Storage and Disposal
Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original containers in a cool, well-ventilated area, away from direct sunlight. Do not allow product to become overheated in storage. This may cause increased degradation of the product, which will decrease product effectiveness. In case of spill, flood area with large quantities of water. Do not store in a manner where cross-contamination with other pesticides or fertilizers could occur.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Open dumping is prohibited. If wastes cannot be disposed of according to label directions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container unless the directions for use allow a different (concentrated) product to be diluted in the container.*

Containers less than 5 gallons:

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Containers greater than 5 gallons:

Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip the container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand container on its end and tip back and for the several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use of disposal. Repeat the procedure two more times. Then offer for recycling or dispose in a sanitary landfill, or by incineration, if allowed by state and local authorities by burning.