



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

APR 9 2010

Mr. Michael Kellogg Pyxis Regulatory Consulting, Inc. Agent for United Turf Alliance 4110 13th. St. NW Gig Harbor, WA 98332

SUBJECT:

Application for Pesticide Notification (PRN 98-10)

Request Primary Brand Name "ArmorTech® THREESOMETM"

EPA Reg. No. 86064-5

Application Dated March 23, 2010

Dear Mr. Kellogg:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 03/23/10 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

Please read instructions on	reverse before comple.	<u>તાલ form.</u>	·**	Form Approv	/ed:,MB No. 2€)70-006C	O. Approval expires 2-28-95
\$EPA	Environmental	United States Environmental Protection Ager Washington, DC 20460			Registrat Amendm Other		OPP Identifier Number
		Application fo	or Pestici	de - Sectio	n I		
1. Company/Product Numbe 86064-5				Product Manage	·	\ r1	oposed Classification
4. Company/Product (Name) United Turf Alliance / UTA 00			PM#	PM# 23			None
5. Name and Address of Applicant (Include ZIP Code) United Turf Alliance c/o Pyxis Regulatory Consulting, Inc. 4110 136th St. NW Giq Harbor. WA 98332 Check if this is a new address			6. Expedited Reveiw. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. Product Name				
		S	Section - I				
Amendment - Explain Resubmission in resp Notification - Explain Explanation: Use addition	ponse to Agency letter	dated			lication.	NOT	TIFICATION PR 9 - 2010
Submission of change in prin consistent with the provision confidential statement of for understand that if this notific be subject to enforcement an	ns of PR Notice 98-10 and mula of this product. I un cation is not consistent with	id EPA regulations at inderstand that it is a virth the terms of PR N	t 40 CFR 152.4 violation of 18 lotice 98-10 an	16, and no other o U.S.C. Sec. 1001	changes have beer 1 to willfully make	n made to any false	o the labeling or the statement to EPA. I further
		S	ection - I	11			
1. Material This Product Wil	Il Be Packaged In:						
Child-Resistant Packaging Yes No	Unit Packaging Yes No		Yes √ No			Metal Plastic Glass	
* Certification must be submitted	If "Yes" Unit Packaging wgt.		"Yes" ickage wgt			Paper Other (S	pecify)
3. Location of Net Contents ✓ Label ✓ C	Information Container	4. Size(s) Retail Co 2.5, 30,	ontainer 55, 250 gallo	l r	Location of Label On Label On Labeling a		•
6. Manner in Which Lebel is	Affixed to Product	Lithograph Paper glued Stenciled		Other _			
		Sc	ection - I	<u> </u>			
1. Contact Point (Complete	items directly below fo	or identification of in	ndividual to be	e contacted, if n	ecessary, to proc	ess this	application.)
Name Michael Kellogg		Title Age				•	e No. (Include Area Code) 53-7369 ^{°, °}
I certify that the state I acknowledge that ar both under applicable	ements I have mede on ny knowlinglly false or r I law.	Certification this form and all att misleading statemen	tachments the	ereto are true, a	ccurate and comport or imprisonment of	pletc. o	8. Dute Application Received
2. Signature Munda Day		3, Titl	ent				(((((((((((((((((((
4. туреd Name Michael Kellogg		5. Dat	ite 8/23/16	0			

304/

PYXIS REGULATORY CONSULTING, INC

4110 136th St. NW Gig Harbor, WA 98332

Phone: 253-853-7369 Fax: 253-853-5516 www.PyxisRC.com

March 23, 2010

COURIER DELIVERY

Kathryn Montague (PM 23)
Document Processing Desk (NOTIF)
Office of Pesticide Programs
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 S. Crystal Drive
Arlington, VA 22202-4501

RE: United Turf Alliance – UTA 001 (EPA Reg. No. 86064-5) Notification of a Change in Primary Brand Name per PRN 98-10

Dear Ms. Montague,

On behalf of United Turf Alliance please find the enclosed notification of a change in primary brand name for UTA 001 (EPA Reg. No. 86064-5). The new primary brand name is "ArmorTech® THREESOMETM".

In support of this primary brand name notification, please find the following:

- 1. Application for Pesticide Registration (EPA Form 8570-1)
- 2. One (1) copy of the ArmorTech THREESOME label with changes tracked
- 3. One (1) copy of the ArmorTech THREESOME label with changes incorporated
- 4. Certification with Respect to Label Integrity
- 5. One (1) copy of the ArmorTech THREESOME label on CD

Please note: Pursuant to the Agency Notice of Registration dated December 16, 2009, United Turf Alliance will submit the child-resistant packaging certification to the Agency prior to the first shipment of product being made.

Please feel free to contact me if you have any questions or need any additional information.

Regards

Michael Kellogg

Enclosures

cc: T. Zech; United Turf Alliance

ArmorTech[®] THREESOME™

SELECTIVE BROADLEAF WEED CONTROL FOR TURFGRASS INCLUDING USE ON SOD FARMS. TO CONTROL CLOVER, DANDELION, HENBIT, PLANTAINS, WILD ONION, AND MANY OTHER BROADLEAF WEEDS. ALSO FOR HIGHWAYS, RIGHTS-OF-WAY AND OTHER SIMILAR NON-CROP AREAS.

CONTAINS 2,4-D, MECOPROP-p AND DICAMBA	APR 9 - 2010
ACTIVE INGREDIENTS:	
Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	30.56%
Dimethylamine Salt of (+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid**‡	8.17%
Dimethylamine Salt of Dicamba (3,6-Dichloro-o-anisic Acid)***	2.77%
OTHER INGREDIENTS:	<u>58.50%</u>
TOTAL:	100.00%
Isomer Specific Method, Equivalent to:	
*2,4-Dichlorophenoxyacetic Acid	
**(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid	6.75%, 0.63 lbs./gal.
***3,6-Dichloro-o-anisic Acid	2.30%, 0.22 lbs./gal.
‡CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p	_

INTENDED FOR USE BY TURF MAINTENANCE PERSONNEL, LANDSCAPING OR COMMERCIAL APPLICATORS ONLY.

KEEP OUT OF REACH OF CHILDREN DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

	FIRST AID
If in eyes:	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed:	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing:	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
	HOT LINE NUMBER
treatment. You	ict container or label with you when calling a poison control center or doctor, or going for may also contact 1-800-222-1222 for emergency medical treatment information. SICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

EPA Reg. No. 86064-5 Manufactured for: United Turf Alliance 12480 Ford Drive Fishers, IN 46038

Net Contents:

EPA Est. No.



50911

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER - PELIGRO

Corrosive. Causes irreversible eye damage. Do not get in eyes, or on skin or clothing. Harmful if swallowed.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are made of any waterproof material. If you want more options, follow the instructions for Category A on an EPA chemical-resistance category selection chart.

All mixers, loaders, applicators and other handlers must wear:

- · Goggles or face shield
- · Long-sleeved shirt and long pants
- · Shoes plus socks, and
- Chemical-resistant gloves
- Chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate

See engineering controls for additional requirements.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands, face, and arms with soap and water before eating, smoking, drinking or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing.
 As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

This product is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees and Roughs) and similar turf areas. Also for use on Sod Farms. For turf use, the maximum number of broadcast applications per treatment site is 2 per year. Product should not be used in or near greenhouses.

.....**2**

c č o

(0001)

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Aerial application is prohibited.

Use of this product in certain portions of California, Oregon and Washington is subject to the January 22, 2004 Order for injunctive relief in <u>Washington Toxics Coalition</u>, et al. v. EPA, C01-0132C, (W.D. WA). For further information, please refer to http://www.epa.gov/espp.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls worn over short-sleeved shirt and short pants
- Waterproof gloves
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of ground application can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

Use only Medium or coarser spray nozzles according to ASAE (S572) definition for standard nozzles or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 10 mph. Only apply this product if the wind direction favors an on target deposition and there are not sensitive areas (including, but not limited to residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at downwind edge of treated area.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: 1) conditions of temperature inversion exist, or 2) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybears

(vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates. Do not apply with a nozzle height greater than 4 feet above the crop canopy.

USE PRECAUTIONS

Avoid mist to vegetables, flowers, ornamentals, shrubs, trees and other desirable plants. Do not pour spray solutions near these plants. Do not spray on carpetgrass, dichondra nor on lawns or turf where desirable clovers are present. Avoid fine mists. Except as noted, use only lawn-type sprayers. Coarse sprays are less likely to "wind-drift". Use coarse spray droplets. Do not spray roots of ornamentals and trees. Do not exceed specified dosages for any area; be particularly careful within the dripline of trees and other species. Do not apply to newly seeded grasses until well established. Avoid broadcast applications when air temperature exceeds 90°F. When using small, spot treatment applications in temperature over 90°F, turf injury may occur.

When treating Carpetgrass, avoid broadcast applications when air temperature exceeds 80°F. When air temperatures exceed 80°F, limit application to spot treatment only.


Apply only to dormant St. Augustine (Common, Raleigh and Seville varieties grown in Texas, Louisiana, Mississippi only) and dormant Centipede grasses (Texas, Louisiana, Mississippi only).

Avoid applying during excessively dry or hot periods unless irrigation is used. For optimum results: (1) turf should not be mowed for 1 to 2 days before and after application; (2) do not apply if rain is expected within 4 hours after the application, delay irrigation cycle for 24 hours. Reseed no sooner than three to four weeks after application of this product. Failure to observe all precautions may result in injury to turf and/or susceptible plants.

This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizer and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test is recommended prior to mixing in application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed.

WEEDS CONTROLLED

Bedstraw Black medic	Dandelion Dock	Lespedeza Mallow	Poison oak Purslane	Wild carrot Wild garlic
Buckhorn	Ground ivy	Morningglory	Ragweed	Wild lettuce
	•		•	
Burdock	Healall	Peppergrass	Sheep sorrel	Wild onion
Chicory	Henbit	Pigweed	Shepherdspurse	Yarrow
Chickweed	Knotweed	Plantain	Speedwell	and other
Clover	Lambsquarter	Poison ivy	Spurge	broadleaf weeds



8091

TURF, ORNAMENTAL (golf courses, cemeteries, parks, sports fields, turfgrass, lawns and other grass areas) AND

TURF (grown for seed or sod)

APPLICATION RATES

SITE	USE RATE (Fluid	SPRAY VOLUME	USE RATE (Pints)	SPRAY VOLUME	USE DIRECTIONS
	Ounces)	(Gallons)	()	(Gallons)	
	PER 1,000	SQ. FT.	PER A		
Bahiagrass,					Normal Applications
Bluegrass, Common Bermudagrass, Fescue, Ryegrass, Zoysiagrass	1.1 to 1.5	0.5 to 5	3 to 4	5 to 220	For Professional Lawn Maintenance - Higher water volumes may be used when tank-mixed with a turf fertilizer. Follow fertilizer labels for proper amounts to add. Use reduced rates if grass is stressed from heat or drought. Exercise care when applying during growth stages from dormancy to green-up and from green-up to dormancy. Some temporary discoloration may occur on warm season grasses. If Bermudagrass is dormant, up to 4 pints per acre may be used. However, some hybrid Bermudagrasses may be sensitive to this product. Contact your local extension service weed control
Dormant St. Augustine (Common, Raleigh and Seville varieties grown in Texas, Louisiana, Mississippi only). Dormant Centipede grasses (Texas, Louisiana,	0.75 to 0.90 Maximum	1 to 5	2 to 2.5 Maximum	40 to 200	specialist. Use reduced rates if grass is stressed from heat or drought. Exercise care when applying during growth stages from dormancy to green-up and from green-up to dormancy. Some temporary discoloration may occur on warm season grasses.
Mississippi					
only). Bentgrass (Putting and Bowling Greens)	0.67 Maximum	3.3	1.8 Maximum	145	Apply on closely mowed Bentgrass, preferably in May or mid-August through September. Slight turf yellowing will disappear after about one week. On closely mowed golf course fairways and tees or bowling greens – apply when weeds are actively growing and temperatures are low. Exercise care to avoid over application to prevent injury. Temporary discoloration may occur.

NOTE: Care should be taken to avoid overdosing Bentgrass or injury may result. Large volumes of spray water (i.e. one fluid ounce in 5 gallons of water per 1,500 square feet) will aid in obtaining uniform coverage. If hand-type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom as sideways application with a boom where the spray from more than one nozzle is

allowed to fall on the same area will result in heavy local over-application and subsequent turf discoloration or injury.

Ornamental Turf: Make no more than 2 applications per year.

<u>Turf Grown for Seed/Sod:</u> Make no more than 2 applications per year. Minimum interval of 30 days between applications.

LOW VOLUME SPRAY APPLICATION EQUIPMENT

EQUIPMENT	JIPMENT APPLICATION USE RATE		TE	USE DIRECTIONS	
		Fluid Ounces per 1,000 Square Feet	Pints per Acre		
Controlled Droplet Applicators (CDA),	Cool season grasses	1.1 to 1.5	3 to 4	Use in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment). Do not overlap spray patterns.	
Atomizers and Spinning Disk Applicators	Listed warm season grasses	0.75 to 0.9	2 to 2.5	Use in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment); and follow CDA spray instructions for cool season grasses. Use reduced rates if grass is stressed from heat, drought, etc.	
Lower Volume Equipment	Cool Season and Listed Warm Season Grasses	1.1 to 1.5	3 to 4	Use as little as 5 gallons of water per acre. Use only application equipment that is capable of spreading a uniform droplet, wetting each weed surface.	

NON-CROPLAND

(fencerows, hedgerows, roadsides, ditches, rights-of-way, utility, powerlines, railroads, airports, and industrial sites)

Applications to non-cropland areas are not applicable to treatment of commercial timber or other plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

Roadsides (including aprons and guardrails), rights-of-way and other similar non-crop areas: For control of broadleaf weeds, mix at a rate of 1.5 to 3.5 quarts of this product per acre in enough water to wet all parts of the foliage. This may require 50 to 300 gallons of water per acre. This mixture will cover 43,560 square feet. Thoroughly saturate all weeds with spray mixture. Apply any time between the time when plants come into full leaf (spring) to when the plants begin to go dormant. Best results are obtained when weeds are young and actively growing. Do not cut weeds until herbicide has translocated throughout the plant causing root death. For small broadleaf weeds, use the lower rate. Heavy, dense stands require the higher rate with the high water volume. For small (spot) applications with small tank sprayers, apply at the rate of 4 ounces of this product per gallon of water and spray to thoroughly wet all foliage.

Annual/Perennial Control: Make no more than 2 applications per year.

For control of woody plants, apply to both stems and foliage any time from the time foliage is completely matured until the time plants start to go dormant. All leaves, stems and suckers must be completely wet to the ground line for effective control. Regrowth may be anticipated on the more resistant species.

Woody Brush: Add 1to 1.5 gallons of this product per acre to wet all parts of the brush foliage, stem and bark. This may require up to 200 to 600 gallons of water per 43,560 square feet depending upon the height and thickness of the brush. Mix thoroughly before spraying.

Woody Plant Control: Make no more than 1 application per year.

This product will either kill, control or suppress the weeds listed in the label bookles for this product. Some of these species may require repeat spot applications even under ideal conditions for application.

Maximum Seasonal Application Rate to non-crop sites is 4 pounds 2,4-D Acid equivalent per accept per application site.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter this product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed, labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying.

(Nonrefillable container ≤ 5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(Nonrefillable > 5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of United Turf Alliance or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold United Turf Alliance and Seller harmless for any claims relating to such factors.

United Turf Alliance warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or United Turf Alliance, and Buyer and User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UNITED TURF ALLIANCE MAKES NO WARRANTIES OF MERCHANTABILITY OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither United Turf Alliance nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. To the extent consistent with applicable law, the exclusive remedy of the user or

BUYER, AND THE EXCLUSIVE LIABILITY OF UNITED TURF ALLIANCE AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF UNITED TURF ALLIANCE OR SELLER, THE REPLACEMENT OF THE PRODUCT.

United Turf Alliance and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of United Turf Alliance.

ArmorTech® is a registered trademark and THREESOME™ is a trademark of United Turf Alliance.

EPA 20100323

