



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

86064-10

Date of Issuance:

9/23/21

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

ArmorTech TEBU 3.6F

Name and Address of Registrant (include ZIP Code):

Michael Kellogg
c/o Pyxis Regulatory Consulting Inc.
United Turf Alliance, LLC.
4110 136th St. Ct. NW
Gig Harbor, WA 98332

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Nathan Mellor, Acting Product Manager 21
Fungicide Branch, Registration Division (7505P)

Date:

9/23/21

2. You are required to comply with the data requirements described in the DCI identified below:

- Tebuconazole GDCI-128991-1598

You must comply with all of the data requirements within the established deadlines. If you have questions about the Generic DCI listed above, you may contact the Chemical Review Manager in the Pesticide Reevaluation Division: <http://iaspub.epa.gov/apex/pesticides/f?p=chemicalsearch:1>

3. Make the following label changes before you release the product for shipment:

- Revise the EPA Registration Number to read, "EPA Reg. No. 86064-10."

4. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 04/27/2021

If you have any questions, please contact Marcel Howard by phone at (703)305-6784, or via email at howard.marcel@epa.gov

Enclosure

{Note to reviewer: [Text] in brackets denotes optional or explanatory language}
{Note to reviewer: {Text} in braces denotes where in the final label text will appear}
{BOOKLET FRONT PANEL LANGUAGE}

TEBUCONAZOLE	GROUP 3	FUNGICIDE
--------------	---------	-----------

ArmorTech® TEBU 3.6 F

Contains tebuconazole, the active ingredient used in Torque.

For use of listed diseases on ornamentals and golf courses turf.

ACTIVE INGREDIENT:	(% by weight)
Tebuconazole, a-[2-(4-chlorophenyl)ethyl]-a-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol	38.7%
OTHER INGREDIENTS:	61.3%
TOTAL	100.0%

Contains 3.6 pounds of Tebuconazole per gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See [below] [inside label booklet] for [additional] [First Aid,] [Precautionary Statements] and [Directions for Use].


ArmorTech TEBU 3.6 F is not manufactured or distributed by Cleary Chemical LLC seller of Torque.

EPA Reg. No.: 86064-xx

EPA Est. No.:

Net Contents:

Manufactured for:
United Turf Alliance, LLC
8014 Cumming Highway, Suite 403-282
Canton, GA 30115



FIRST AID	
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by the poison control center or doctor. • DO NOT give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
Note to Physician: No specific antidote. Treat symptomatically. The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.	
HOT LINE NUMBER	
For information on this pesticide (including general health concerns or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378, Monday through Friday, 8:00 AM to 12:00 PM Pacific Standard Time. In the event of a medical emergency, call your poison control center at 1-800-222-1222.	

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves: barrier laminate, or butyl rubber ≥ 14 mils or nitrile rubber ≥ 14 mils or neoprene rubber ≥ 14 mils or polyvinyl chloride ≥ 14 mils or Viton ≥ 14 mils
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

<p>User Safety Recommendations</p> <p>Users should:</p> <ul style="list-style-type: none"> • Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Drift

and runoff may be hazardous to aquatic organisms in neighboring areas. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory: Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT mix or allow coming in contact with oxidizing agent. Hazardous Chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated including plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves: barrier laminate or butyl rubber ≥ 14 mils or nitrile rubber ≥ 14 mils or neoprene rubber ≥ 14 mils or polyvinyl chloride ≥ 14 mils or Viton ≥ 14 mils
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box only apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Golf Course Turf and Landscape Uses: Keep children and pets out of treated areas until sprays have dried.

PRODUCT INFORMATION

Read the entire Directions for Use and Limitation of Warranty and Liability before using this product.

Chemigation: DO NOT apply this product through any type of irrigation system.

OBSERVE THE FOLLOWING RESTRICTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS INCLUDING LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES

- DO NOT apply within 100 feet of aquatic areas listed above.
- DO NOT cultivate within 10 feet of an aquatic area to allow growth of a vegetation filter strip.
- See Spray Drift Management section for further information.

SPRAY DRIFT MANAGEMENT

Make ground application when wind velocity favors on-target product deposition (approximately 3 to 10 mph). DO NOT apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperatures.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.

Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.

Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

BOOM HEIGHT – Ground Boom

For ground equipment, the boom should remain level with the target area and have minimal bounce.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Boomless Ground Applications:

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications:

Take precautions to minimize spray drift.

Spray Volume: For best results ArmorTech® TEBU 3.6 F may be applied in 66-132 gallons of water per acre for turf using ground-based equipment. For ornamentals, 50-300 gallons of finished spray per acre are advised depending upon equipment, plant species and plant growth stage at time of application. For the most effective results, check calibration equipment regularly. When using lower spray volumes, be sure to maintain uniform application and full crop coverage so as to ensure effective control. Increase spray volume to ensure proper application, if required.

Compatibility Test for Mix Components:

Before mixing components, always perform a compatibility jar test. For 66 gallons per acre spray volume, use 5 cups of water in a clear, clean mixing jar. For other spray volumes adjust accordingly. Only use water from the intended source at the source temperature. Add components in the sequence indicated below in Mixing Order using 3 teaspoons for each pound of dry product or 1% teaspoon for each pint of liquid product of directed label rate per acre. Always cap the jar and invert 10 cycles between component additions. When the components have all been added to the jar and fully mixed, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution will not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture is spray solution is compatible. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent and use the compatibility agent as directed on its label.

Mixing: Continuous agitation is required during mixing. When, mixing this product and water, use the specified application rates as listed for each crop on this label. Before combining any other substances with the mixture, ensure that the ArmorTech TEBU 3.6 F is completely dispersed in the mixture. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Mixing Procedure:

1. Water. Add three-quarters of the required volume to a thoroughly clean sprayer tank.
2. Agitation. Start agitation and maintain constant agitation throughout mixing and application.
3. Inductor. If an inductor is used, rinse it thoroughly after each component has been added.
4. Products in PVA Bags. Place any product contained in water soluble PVA bags into the mixing tank. Wait until all water soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5. Water Dispersible Products. Including dry flowables (OF), wettable powders (WP), suspension concentrates (SC) or suspo-emulsions (SE).
6. Water-soluble products.
7. Emulsifiable concentrates (for example oil concentrates when applicable).
8. Water soluble additives (for example AMS or UAN when applicable).
9. Remaining quantity of water.

RESISTANCE MANAGEMENT

For resistance management, ArmorTech TEBU 3.6 F contains a group 3 fungicide. Any fungal population may contain individuals naturally resistant to ArmorTech TEBU 3.6 F and other Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Follow appropriate resistance management strategies.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of ArmorTech TEBU 3.6 F or other Group fungicides within a growing season sequence with different groups the control the same pathogens.
- Use tank mixtures with fungicide/bactericides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide/bactericide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide/bactericide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal/bacterial populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact United Turf Alliance, LLC at 614-284-4442 or at info@utaarmortech.com. You can also contact your pesticide distributor or university extension specialist to report resistance.

DISEASE CONTROL IN GOLF COURSE TURF

TURF USE RESTRICTIONS

For use on golf course turf only.

DO NOT use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (i.e., elementary, middle and high school), campgrounds, churches, and theme parks.

Not for residential use.

Not for use on turf being grown for sale or commercial use as sod.

DO NOT use clippings for animal feed.

DO NOT exceed 3.6 fl. oz. (4.4 lb a.i./A) of ArmorTech TEBU 3.6 F per 1,000 sq ft per year.

DO NOT apply more than 1.1 fl. oz. (1.3 lb a.i./A) of ArmorTech TEBU 3.6 F per 1,000 sq ft per application.

DO NOT apply more than 6 applications per year.

The minimum retreatment interval is 21 days.

In the State of New York, **DO NOT** exceed 1.8 fl. oz. (2.2 lb a.i./A) of ArmorTech TEBU 3.6 F per 1,000 sq ft per year.

Application Information

For use on all Golf turf applications of cool season and warm season grasses (including Bentgrasses, Bluegrasses, Fescues, Ryegrasses, St. Augustine grasses, and Zoysia) or their mixtures. ArmorTech

TEBU 3.6 F is not phytotoxic to any of the above-mentioned grasses when used in accordance with the label.

Note: Bermudagrass can be sensitive to ArmorTech TEBU 3.6 F under certain conditions. **DO NOT** apply consecutive applications during or just after dormancy break.

Note: For all grasses, avoid consecutive applications of ArmorTech TEBU 3.6 F or other DMI containing fungicides when temperatures are expected to exceed 85 degrees F.

ArmorTech TEBU 3.6 F can be used for the prevention and control of the diseases mentioned in table below. Begin applications when conditions favor disease development and repeat applications as long as these conditions persist. Preventative treatments can be applied using 21-day intervals as indicated. When treating golf greens, always treat aprons and approaches. Spray uniformly over the area to be treated with properly calibrated equipment.

Apply the specified amount of ArmorTech TEBU 3.6 F in sufficient water for thorough coverage. Use a volume of 66 - 132 gallons per acre (1.5 - 3.0 gallons per 1,000 sq ft). Apply using properly calibrated low volume, handheld, mechanical or motorized ground broadcast equipment. Application to small areas may be made with low-pressure handwand or backpack equipment. Maintain constant agitation during application.

Depending on the disease, water ArmorTech TEBU 3.6 F into the crown and active root zone for best results. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. For best results use spray mixture the same day it is prepared.

Golf Course Turf Disease Control

DISEASE	RATE of ArmorTech TEBU 3.6 F (fl. oz./1000 SQ Ft)	APPLICATION INSTRUCTIONS
Dollar Spot (<i>Sclerotinia homoeocarpa</i>) Copper Spot (<i>Gloeocercospora sorghi</i>) Powdery Mildew (<i>Erysiphe graminis</i>) Corticium Red Thread (<i>Laetisaria fuciformis</i>) Rusts (<i>Puccinia</i> spp.) Red Thread (<i>Laetisaria fuciformis</i>) Pink Patch (<i>Limonomyces rosipellis</i>)	0.6-1.1 (0.7 – 1.3 lb a.i./A)	For prevention, begin applications when conditions are favorable for disease development using the low to medium rate. For curative treatments, use the medium to high rate. ArmorTech TEBU 3.6 F can be tank mixed with chlorothalonil or Spectro® (chlorothalonil/thiophanate-methyl; EPA Reg. No. 1001-72) at labeled rates. DO NOT make two consecutive applications of ArmorTech TEBU 3.6 F or other DMI containing fungicides. Alternate with another fungicide with a different mode of action. Apply subsequent application after 21 days.
Brown Ring Patch (<i>Waitea circinata</i>)	0.6-1.1 (0.7 – 1.3 lb a.i./A)	For prevention, begin applications when conditions are favorable for disease development. Make two applications using the low to medium rate tank mixed with polyoxin D zinc salt at labeled rates. Repeat application at 21-day interval.

DISEASE	RATE of ArmorTech TEBU 3.6 F (fl. oz./1000 SQ Ft)	APPLICATION INSTRUCTIONS
Brown Patch/Rhizoctonia Blight, Large Patch (<i>Rhizoctonia sofan</i> ,) Anthracnose Basal and Foliar (<i>Colletotrichum cereale</i>)	0.6-1.1 (0.7 – 1.3 lb a.i./A)	For prevention, begin applications when conditions are favorable for disease development using the low to medium rate. For curative treatments, use the medium to high rate. ArmorTech TEBU 3.6 F can be tank mixed with chlorothalonil, Spectro® (chlorothalonil/thiophanate-methyl; EPA Reg. No. 1001-72), thiophanate-methyl or polyoxin D zinc salt at labeled rates. DO NOT make two consecutive applications of ArmorTech TEBU 3.6 F or other DMI containing fungicides. Alternate with another fungicide with a different mode of action. Apply subsequent application after 21 days.
Bermudagrass decline (<i>Gaeumannomyces graminis</i> var. <i>graminis</i>)	0.6-1.1 (0.7 – 1.3 lb a.i./A)	For prevention, begin applications two or four weeks prior to the historical appearance of disease symptoms. Make two applications at the low to medium rate, 21 days apart or one application at the high rate. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and root zone of the turf. The amount of water is dependent on the depth of the root zone. Alternate with another fungicide with a different mode of action as needed. Initiate cultural control practices at the same time the fungicide is applied. Refer to your local County Extension Service for this information.
Take All Patch (<i>Gaeumannomyces graminis</i> var. <i>avenae</i>)	0.6-1.1 (0.7 – 1.3 lb a.i./A)	For prevention, make two applications at low rate, 21 days apart or one application at high rate in the fall when root zone soil temperatures reach 75° F and again in the spring at 55-60° F root zone soil temperatures. Applications in both fall and spring may be necessary. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.

DISEASE	RATE of ArmorTech TEBU 3.6 F (fl. oz./1000 SQ Ft)	APPLICATION INSTRUCTIONS
Gray Leaf Spot (<i>Pyricularia grisea</i>)	0.6-1.1 (0.7 – 1.3 lb a.i./A)	For prevention, begin applications when conditions are favorable for disease development (July-Aug) using the low to medium rate. For curative treatments, use the medium to high rate and tank mix with chlorothalonil or Spectro® (chlorothalonil/thiophanate-methyl; EPA Reg. No. 1001-72) at labeled rates. DO NOT make two consecutive applications of ArmorTech TEBU 3.6 F or other DMI containing fungicides. Alternate with another fungicide with a different mode of action, for example thiophanate-methyl. Apply subsequent application after 21 days.
Stripe Smut (<i>Ustilago striiformis</i>)	0.6-1.1 (0.7 – 1.3 lb a.i./A)	Make a single application to historical disease areas in spring as grass growth begins. For curative treatments, use the medium to high rate.
Spring Dead Spot (<i>Leptosphaeria korrea</i> , <i>L. narmari</i> , <i>Ophiosphaerel/a herpotricha</i> , <i>Gaeumannomyces graminis</i>) Necrotic Ring Spot (<i>Leptosphaeria korrea</i>)	0.6-1.1 (0.7 – 1.3 lb a.i./A)	For prevention, make two applications at the low to medium rate, 21 days apart or one application at the high rate in the fall when root zone soil temperatures reach 75° F. ArmorTech TEBU 3.6 F can be combined with thiophanate-methyl at labeled rates. A spring application may be necessary in areas where disease pressure is known to be heavy. Apply using low to medium rate when root zone soil temperature reaches 55-60° F. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
Fusarium Patch (<i>Fusarium roseum</i>)	0.6-1.1 (0.7 – 1.3 lb a.i./A)	Apply first application in mid-June or 2-4 weeks prior to time this blight normally becomes evident. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone. Alternate with another fungicide with a different mode of action, for example thiophanate-methyl. Apply subsequent application after 21 days.

DISEASE	RATE of ArmorTech TEBU 3.6 F (fl. oz./1000 SQ Ft)	APPLICATION INSTRUCTIONS
Summer Patch (<i>Magnaporthe poae</i>)	0.6-1.1 (0.7 – 1.3 lb a.i./A)	For prevention, apply in the spring when root zone soil temperatures reach 55-60°F. DO NOT make two consecutive applications of ArmorTech TEBU 3.6 F or other DMI containing fungicides. Alternate with another fungicide with a different mode of action, for example thiophanate-methyl. Make additional applications at 21-day intervals. See local university guidelines for additional information. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.

DISEASE	RATE of ArmorTech TEBU 3.6 F (fl. oz./1000 Sq Ft)	APPLICATION INSTRUCTIONS
Zoysia Patch, Large Patch of zoysia (<i>Rhizoctonia solani</i>)	0.6-1.1 (0.7 – 1.3 lb a.i./A)	Make first application in early fall (mid-September to mid-October) prior to development of disease symptoms. Alternate with another fungicide with a different mode of action, for example polyoxin D zinc salt. Second and third applications may be made at 21-day intervals. See local university guidelines for additional information. An additional application in early spring may be necessary in areas where disease pressure is known to be heavy.
Gray Snow Mold/ Typhula Blight (<i>Typhula incarnate</i>) Pink Snow Mold/Microdochium Patch (<i>Microdochium nivale</i>)	0.6-1.1 (0.7 – 1.3 lb a.i./A)	Apply in the fall, before anticipated turf dormancy and before first snow cover. If turf breaks dormancy during winter months a second application may be made. DO NOT apply over snow cover, or when turf is dormant. It is advised that ArmorTech TEBU 3.6 F be tank-mixed with other registered snow mold products, including 26/36® (thiophanate-methyl/iprodione; EPA Reg. No. 228-630), polyoxin D zinc salt, chlorfenapyr or chlorothalonil for best season-long results.
Fairy Ring <i>Chlorophyllum</i> (<i>Lepiota</i>), <i>Lycoperdon</i> , <i>Marasmius</i>	0.6-1.1 (0.7 – 1.3 lb a.i./A)	For prevention in Cool season turf, make two applications at the low-medium rate in the spring when root zone soil temperatures reach 55-60° F. Make a second application using a 21-day interval. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone. DO NOT use a wetting agent unless hydrophobic soil conditions exist. For curative treatment, use the medium to high rate. For warm season turf breaking dormancy, DO NOT make two consecutive applications of ArmorTech TEBU 3.6 F or other DMI containing fungicides. Alternate with another fungicide with a different mode of action, including polyoxin D zinc salt, Heritage (acibenzolar-s-methyl/azoxystrobin; EPA Reg. No. 100-1550) or flutolanil. For hydrophobic areas, use an appropriate wetting agent to effectively penetrate the hydrophobic zone commonly created with this disease.
<p>APPLICATION DIRECTIONS: Apply the specified amount of ArmorTech TEBU 3.6 F in 1.0 to 3.0 gallons of water per 1000 sq. ft. Make all applications after mowing and allow foliage to dry thoroughly before irrigation.</p> <p>RESTRICTIONS: DO NOT use clippings for animal feed. DO NOT exceed 3.6 fl.oz. (4.4 lb a.i./A) of ArmorTech TEBU 3.6 F per 1000 sq. ft. per year. DO NOT apply more than 1.1 fl. oz. (1.3 lb a.i./A) of ArmorTech TEBU 3.6 F per 1,000 sq ft per application. DO NOT exceed 6 applications per year. The minimum retreatment interval is 21 days.</p>		

DISEASE CONTROL IN FIELD, NURSERY AND CONTAINER ORNAMENTALS AND ORNAMENTALS IN COMMERCIAL and RESIDENTIAL LANDSCAPES

ORNAMENTAL USE RESTRICTIONS

For use on ornamental plants only; not for woodlands or forest management. **DO NOT** use in, on, or around any structure, vehicle, article, surface, or area associated with the household, including but not limited to areas including non-agricultural outbuildings, non-commercial greenhouses, pleasure boats and recreational vehicles, or in any preschool or day care. Not for residential use.

Intended for use by professional applicators.

DO NOT apply more than 10 fl. oz. per acre (12.2 lb a.i./A) is a single application.

DO NOT apply more than 0.31 gallons (40 fl. oz.) of this product (equal to 1.13 lbs. of tebuconazole) per acre per year.

DO NOT make more than 4 applications per year.

The minimum retreatment interval is 14 days.

DO NOT apply to bearing fruit trees or vegetables.

Application Information

ArmorTech TEBU 3.6 F can be used in a preventative and curative disease control program for the listed plant types and disease in the table below. Optimum disease management is obtained when ArmorTech TEBU 3.6 F is used in conjunction with sound disease management practices.

Apply material with properly calibrated handheld, mechanical or motorized spray equipment. Begin applications when disease first appears and repeat at 14-21-day intervals during the growing season. Use the shortest interval when conditions are unusually favorable for the development of disease. For handheld, mechanical, or motorized applications, mix as directed below and apply as a full coverage spray to drip for the prevention and control of the diseases listed below. Choose a finished spray volume appropriate for the size of the plants and amount of foliage, which will provide thorough coverage throughout the canopy. Allow sprays to dry before overhead irrigation is applied.

Apply ArmorTech TEBU 3.6 F at rates of 4-10 fl oz per acre in 100 gallons of water (4.9 - 12.2 lb a.i./A). Spray volume may range from 50 up to 300 gallons of finished spray per acre depending upon equipment, plant species and plant growth stage at time of application.

Note: The "Directions for Use" of this product reflect the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied to any plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results. This product is not directed for use on African Violets, Begonias, Boston Fern, and Geraniums.

Ornamentals Disease Control

PLANTS	DISEASE	APPLICATION	
		To Prevent Diseases	To Treat Existing Disease
Roses	Black Spot, Powdery Mildew, Rust	Apply every 14-21 days during the growing season, starting when leaves first appear.	Apply every 14 days for a total of 3 applications beginning at the first sign of disease.

Flowers	Leaf Spot, Powdery Mildew, Rust, Southern Blight	Apply at least 3 times per year, 14-21 days apart, beginning with Spring bud break. Rotation or Tank mixing with barrier protectant fungicides is advised for resistance management.	
Crabapples (Ornamental), Dogwoods and Other Landscape (Ornamental) Trees	Anthrachnose, Leaf Spot, Powdery Mildew, Rust, Scab		
Azaleas, Camellias, Rhododendrons and Other Landscape (Ornamental) Shrubs	Anthrachnose, Black Spot, Leaf Spot, Petal Blight, Powdery Mildew, Rust, Southern Blight		
Ground Covers and Vines			
HOW MUCH TO USE FOR SMALL PLANTINGS: ADD 1 TEASPOON TO 2.5 GALLONS OF WATER.			

Pump Style Sprayers

1. Add the appropriate amounts of concentrate and water to the sprayer tank.
2. Close the sprayer, shake well and pressurize.
3. Adjust nozzle to a coarse spray pattern and apply.
4. Occasionally re-pressurize the sprayer if needed to maintain a good spray pattern.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

[For Residential uses]

[Nonrefillable container. DO NOT reuse or refill this container. If empty: Offer for recycling if available or discard in a sanitary landfill. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.]

[For Commercial Uses]

[For plastic containers ≤ 5 gallons: Nonrefillable Container: DO NOT reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.]

[For plastic containers > 5 gallons: Nonrefillable container. DO NOT reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.]

LIMITATION OF WARRANTY AND LIABILITY

IMPORTANT: READ BEFORE USE. Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of United Turf Alliance, LLC. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, United Turf Alliance, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither United Turf Alliance, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

ArmorTech® is a registered trademark of United Turf Alliance, LLC
[Torque®] is a registered trademark of Cleary Chemical LLC

{LANGUAGE ON LABEL AFFIXED TO CONTAINER}

TEBUCONAZOLE	GROUP	3	FUNGICIDE
--------------	-------	----------	-----------

ArmorTech® TEBU 3.6 F

[Contains tebuconazole, the active ingredient used in Torque.]
For use of listed diseases on ornamentals and golf courses turf.

ACTIVE INGREDIENT: (% by weight)
Tebuconazole, a-[2-(4-chlorophenyl)ethyl]-a-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol.....38.7%

OTHER INGREDIENTS:.....61.3%

TOTAL 100.0%

Contains 3.6 of Tebuconazole per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If swallowed:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. DO NOT induce vomiting unless told to do so by the poison control center or doctor. DO NOT give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
Note to Physician: No specific antidote. Treat symptomatically. The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.	
HOT LINE NUMBER	
For information on this pesticide (including general health concerns or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378, Monday through Friday, 8:00 AM to 12:00 PM Pacific Standard Time. In the event of a medical emergency, call your poison control center at 1-800-222-1222.	

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to mammals, fish and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where

surface water is present or to intertidal areas below the mean high-water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory: Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.
PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

[For Residential uses]

[Nonrefillable container. DO NOT reuse or refill this container. If empty: Offer for recycling if available or discard in a sanitary landfill. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.]

[For Commercial Uses]

[For plastic containers ≤ 5 gallons: Nonrefillable Container: **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.]

[For plastic containers > 5 gallons: Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.]

See inside label booklet for additional Precautionary Statements and Directions for Use.

[ArmorTech TEBU 3.6 F is not manufactured, or distributed by Competitor Company, seller of Torque.]

Manufactured for:
United Turf Alliance, LLC
8014 Cumming Highway, Suite 403-282
Canton, GA 30115

EPA Reg. No.: 86064-xx
EPA Est. No.: _____
Net Contents: _____
Batch No.