

86363-21

01-31-2012

1/28



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (7505P)  
Ariel Rios Building  
1200 Pennsylvania Ave., NW  
Washington, D.C. 20460

EPA Reg. Number:

Date of Issuance:

86363-21

JAN 31 2012

NOTICE OF PESTICIDE:

Registration  
 Reregistration  
(under FIFRA, as amended)

Term of Issuance: Conditional

Name of Pesticide Product:

KT CTL 720 Fungicide

Name and Address of Registrant (include ZIP Code)

Kaizen Technologies  
1555 Main Street  
Suite A3-206  
Windsor, CO 80550

Mailed to: Scott Baker, Agent  
Lighthouse Product Services  
3937 Cedarwood Lane  
Johnstown, CO 80534

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

You must make the following labeling revisions:

- 1. The EPA Reg. No. must be changed to 86363-21.

Signature of Approving Official:

Tony Kish  
Product Manager, Team 22  
Fungicide Branch  
Registration Division (7505P)

Date:

2  
28

2. On page 2, add a space not a line between the last sentence in the "WPS Uses" section and the Non WPS Uses section.

By 2/1/2013 you must generate a one year GLP storage stability (830.6317) and corrosion characteristics (830.6320) data o the proposed product. The results must be submitted to the Agency in electronic and hard copy format.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release of the shipment of your product constitutes acceptance of these conditions.

A copy of the label stamped "Accepted with Comments" is enclosed for your records.

Enclosures

3  
28

# KT CTL 720 FUNGICIDE

ACTIVE INGREDIENT:	% BY WT.
Chlorothalonil (tetrachloroisophthalonitrile) .....	54.0%
INERT INGREDIENTS: .....	46.0%
TOTAL .....	100.0%

Contains 6.0 Pounds of Active Ingredient Per Gallon (720 Grams Per Liter)

## KEEP OUT OF REACH OF CHILDREN WARNING-AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail).

FIRST AID
<b>IF IN EYES:</b> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
<b>IF SWALLOWED:</b> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious or convulsing person.
<b>IF INHALED:</b> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for treatment advice.
<b>IF ON SKIN OR CLOTHING:</b> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor or going for treatment.
<b>NOTE TO PHYSICIAN:</b> Persons having temporary irritation may respond to treatment with antihistamines or steroid creams and/or systemic steroids.

See inside booklet for complete Precautionary Statements, Directions for Use, and Conditions of Sale and Warranty.

EPA Reg. No. 86363-

EPA Est. No. 53883-TX-002

Net Contents: 1, 2.5, 15, 55, \_\_\_\_\_ Gallons

**Manufactured for:**  
Kaizen Technologies  
1555 Main Street  
Suite A3-206  
Windsor, CO 80550

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated  
JAN 31 2012**

**Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.  
86363-21**

012612

4  
28

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
WARNING**

Causes eye irritation. May cause skin irritation. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Do not get into eyes, on skin, or on clothing. Avoid prolonged contact with skin. Avoid breathing spray mist. DO NOT take internally. Note to user: This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation, and redness or rash on exposed skin areas. Persons having allergic reactions should contact a physician.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical resistance category selection chart.

For WPS or non-WPS applications made in enclosed areas such as greenhouses, applicators and other handlers must wear a NIOSH approved respirator with any N, P, R, or HE filter.

**WPS Uses (commercial production on farms, nurseries, sodfarms, and in greenhouses):**

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as nitrile rubber, natural rubber, or butyl rubber
- Shoes plus socks
- Protective eyewear such as goggles, safety glasses, or face shield

---

**Non-WPS Uses (such as applications to non-residential turf, golf courses, etc.):**

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as nitrile rubber, natural rubber, or butyl rubber
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**ENGINEERING CONTROLS STATEMENT**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR Part 170.240(d)(46)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**USER SAFETY RECOMMENDATIONS**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove PPE immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**ENVIRONMENTAL HAZARDS**

This product is toxic to aquatic invertebrates and wildlife. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high-water mark.

5  
28

Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

This chemical is known to leach through soil into groundwater under certain conditions as a result of labeled use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

This chemical can contaminate surface water through spray drift. Under some conditions, it may also have a high potential for runoff into surface waters for several days to weeks after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlaying tile drainage systems that drain to surface water.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, (WPS) 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the REI of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as nitrile rubber, natural rubber, or butyl rubber
- Shoes plus socks
- Protective eyewear such as goggles, safety glasses, or face shield

Special Eye Irritation Provisions: This product is a severe eye irritant. Although the restricted-entry interval expires after 12 hours, for the next 6.5 days, entry is permitted only when the following safety measures are provided:

- (1) At least one container designed specifically for flushing eyes must be available in operating condition at the WPS-required decontamination site intended for workers entering the treated area.
- (2) Workers must be informed, in a manner they can understand:
  - that residues in the treated area may be highly irritating to their eyes,
  - that they should take precautions, such as refraining from rubbing their eyes, to keep residues out of their eyes,
  - that if they do get residues in their eyes, they should immediately flush their eyes using the eye flush container that is located at the decontamination site or using other readily available clean water, and
  - how to operate the eye flush container.

6  
28

### **NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides, 40 CFR Part 170. The WPS applies when the product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Do not enter or allow others to enter the treated area until sprays have dried.

### **APPLICATION INSTRUCTIONS**

KT CTL 720 FUNGICIDE, a flowable product containing chlorothalonil, is for use as a spray for the control of many important plant diseases.

### **RESISTANCE MANAGEMENT**

To avoid the development of tolerant or resistant strains of fungi, KT CTL 720 FUNGICIDE should always be tank mixed with a fungicide of different chemistry, and/or a fungicide of different chemistry should be alternated with KT CTL 720 FUNGICIDE at each application. If after using KT CTL 720 FUNGICIDE as recommended and the treatment is not effective, a tolerant or resistant strain of fungi may be present. Discontinue the use of KT CTL 720 FUNGICIDE for at least one season. KT CTL 720 FUNGICIDE is effective for use in programs that attempt to minimize disease resistance to fungicides. KT CTL 720 FUNGICIDE has a multi-site mode of action and may be used to delay or prevent the development of resistance to single-site fungicides. Consult with your Federal or State Cooperative Extension Service representatives for guidance on the proper use of KT CTL 720 FUNGICIDE in programs that seek to minimize the occurrence of disease resistance to other fungicides.

### **PRODUCT PRECAUTIONS**

KT CTL 720 FUNGICIDE can be used effectively in dilute or concentrate sprays. Thorough, uniform coverage is essential for disease control. Do not combine KT CTL 720 FUNGICIDE in a spray tank with pesticides, surfactants, or fertilizers, unless prior use has shown the combination to be physically compatible, effective, and non-injurious under your conditions of use. Do not combine KT CTL 720 FUNGICIDE with DiPel 4L, Foil, Triton AG-98, Triton B-1956 as phytotoxicity may result from the combination when applied to crops listed on this label.

**Note:** Prior to pouring, slowly invert container several times to assure uniform mixture.

The required amount of KT CTL 720 FUNGICIDE should be added slowly into the spray tank during filling. With concentrate sprays, premix the required amount of KT CTL 720 FUNGICIDE in a clean container and add to the spray tank as it is being filled. Keep agitator running when filling spray tank and during spray operations. Dosage rates on this label indicate pints of KT CTL 720 FUNGICIDE per acre, unless specified otherwise. Under conditions favoring disease development, the high rate specified and shortest application interval should be used.

### **APPLICATION PRECAUTIONS AND REQUIREMENTS**

This product must not be applied within 150 feet for aerial and airblast applications, or 25 feet for ground applications of marine/estuarine water bodies unless there is an untreated buffer area of that width between the area to be treated and the water body.

### **SPRAY DRIFT MANAGEMENT**

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator and grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to public health uses, or to applications using dry formulations.

1. The distance of the outermost nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Excluding helicopters, nozzles must always point backward parallel with the airstream and never be pointed downward more than 45 degrees.

Where states have more stringent regulations, they must be observed. The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information.

**Aerial Drift Reduction Advisory Information:**

**INFORMATION ON DROPLET SIZE**

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly or under unfavorable conditions (see Wind, Temperature).

**CONTROLLING DROPLET SIZE — General Techniques**

- Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

**CONTROLLING DROPLET SIZE — Aircraft**

- Number of nozzles - Use the minimum number of nozzles that provide uniform coverage.
- Nozzle orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- Nozzle type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift potential.

**BOOM HEIGHT**

Setting the boom at the lowest labeled height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.

**BOOM LENGTH**

For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

**APPLICATION HEIGHT**

Application should not be made at a height greater than 10 ft. above the top of the largest plants, unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

**SWATH ADJUSTMENT**

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the upwind and downwind edges of the field, the application must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.).

**WIND**

Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential.

8  
28

**NOTE:**

Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

**TEMPERATURE AND HUMIDITY**

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

**TEMPERATURE INVERSIONS**

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

**SENSITIVE AREAS**

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, nontarget crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

**SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

**AIR ASSISTED (AIR BLAST) FIELD CROP SPRAYERS**

Air assisted field crop sprayers carry droplets to the target via a downward directed air stream. Some may reduce the potential for drift, but if a sprayer is unsuitable for the application and/or set up improperly, high drift potential can result. It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application, is configured properly, and that drift is not occurring. NOTE: Air assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration.

**AIR ASSISTED (AIR BLAST) TREE AND VINE SPRAYERS**

Air assisted tree and vine sprayers carry droplets into the canopy of trees and vines via a radically or laterally directed air stream. In addition to the general drift management principles already described, the following specific practices will further reduce the potential for drift:

- Adjust deflectors and aiming devices so that spray is only directed into the canopy.
- Block off upward pointed nozzles when there is no overhanging canopy.
- Use only enough air volume to penetrate the canopy and provide good coverage.
- Do not allow spray to go beyond the edge of the cultivated area. Spray the outside row only from outside the planting.

**APPLICATION AND CALIBRATION TECHNIQUES FOR CHEMIGATION**

Apply this product only through center pivot, motorized lateral move, traveling gun, solid set, and portable (wheel move, side roll, end tow, or hand move) irrigations system(s). Do not apply this product through any other type of irrigation system. Use only on crops specifically



9  
28

designated in the DIRECTIONS FOR USE.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers, or other experts. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

**Specific Instructions for Public Water Systems:**

1. Public water system means a system for the provision to the public of piped water from human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Always inject KT CTL 720 FUNGICIDE into irrigation water after it discharges from the irrigation pump and after it passes through the check valve. Never inject pesticides on the intake line on the suction side of the pump.
8. Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur.
9. Do not apply when wind speed favors drift beyond the area intended for treatment.

**Specific Instructions for Sprinkler Irrigation Systems:**

KT CTL 720 FUNGICIDE may be used through two basic types of sprinkler irrigation systems as outlined in Sections A and B. Determine which type of system is in place, then refer to the appropriate directions provided for each type.

A. Center Pivot, Motorized Lateral Move, and Traveling Gun Irrigation Equipment For injection of pesticides, these continuously moving systems must use a positive displacement injection pump, of either diaphragm or piston type, constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock and capable of injection at pressures approximately two to three times those encountered within the irrigation water line. Venturi application units cannot be used on these systems.

Fill chemical supply tank of injection equipment with water. Operate system for one complete revolution or run across the field, measuring time required, amount of water injected, and acreage covered. Thoroughly mix recommended amount of KT CTL 720 FUNGICIDE for acreage to be covered into same amount of water used during calibration and inject into system continuously for one revolution or run. Mixture in the chemical supply tank must be continuously agitated during the injection run. Shut off injection equipment after one revolution or run, but

19  
28

continue to operate irrigation system until KT CTL 720 FUNGICIDE has been cleared from last sprinkler head.

B. Solid Set and Portable (Wheel Move, Side Roll, End Tow, or Hand Move) Irrigation Equipment With stationary systems, an effectively designed in-line Venturi applicator unit is preferred which is constructed of materials that are compatible with pesticides; however, a positive-displacement pump can also be used. Determine acreage covered by sprinkler. Fill tank of injection equipment with water and adjust flow to use contents over a 30 to 45 minute period. Mix desired amount of KT CTL 720 FUNGICIDE for acreage to be covered with water so that the total mixture of KT CTL 720 FUNGICIDE plus water in the injection tank is equal to the quantity of water used during calibration, and operate entire system at normal pressures recommended by the manufacturer of injection equipment used for amount of time established during calibration. No agitation should be required.

KT CTL 720 FUNGICIDE can be injected at the beginning or end of the irrigation cycle or as a separate application. Stop injection equipment after treatment is completed and continue to operate irrigation system until KT CTL 720 FUNGICIDE has been cleared from last sprinkler head. Do not use on greenhouse grown crops.

**CROP DIRECTIONS-FIELD CROPS AS A SPRAY (Ground or Aerial Equipment)** - Apply KT CTL 720 FUNGICIDE at the rate shown; use sufficient water to provide thorough coverage. Gallonage will vary with crop and amount of plant growth. Spray volume usually will range between 20 to 150 gallons per acre (200 to 1,400 liters per hectare) for dilute sprays and 5 to 10 gallons per acre (50 to 100 liters per hectare) for concentrate ground sprays and aircraft applications. Either ground or aircraft methods of application are recommended unless specific directions are given for a crop. Application through sprinkler irrigation systems is not recommended unless specific directions are given for a crop. See the following instructions for application and calibration.

**FIELD CROPS**

CROP	DISEASES CONTROLLED	RATE OF KT CTL 720 FUNGICIDE PER APPLICATION PINTS/ACRE	SEASONAL LIMITS (PINTS/ ACRE/ YEAR)	APPLICATION DIRECTIONS
<b>ASPARAGUS</b>	Rust Purple Spot Cercospora Leaf Blight	2.0-4.0	12.0	Begin application after harvest of spears, when conditions favor disease development on ferns, generally when leaf wetness occurs. Repeat applications at 2 to 4 week intervals until ferns are no longer productive. Use high rate and shortest application interval when conditions favor disease development. Do not apply within 190 days (120 days in CA and AZ) before harvest.
<b>BEANS, DRY</b> Including but not limited to: Navy Bean Pinto Bean Kidney Bean Lima Bean	Rust (Phakopsora spp.) Anthracnose Downy Mildew Cercospora Leaf Spot (for	1.37-2.0	8.0	Use in sufficient water to obtain adequate coverage. Begin applications at first onset of disease which may occur as early as 2 to 4 weeks before flowering. Repeat applications at 7 to 10 day intervals. For use only on beans to be

11  
28

CROP	DISEASES CONTROLLED	RATE OF KT CTL 720 FUNGICIDE PER APPLICATION PINTS/ACRE	SEASONAL LIMITS (PINTS/ ACRE/ YEAR)	APPLICATION DIRECTIONS
Broad Bean Pink Bean Jack Bean Cow Pea Chick Pea (Garbanzo) Blackeyed Pea Southern Pea, etc.	Blackeyed Pea only) Ascochyta Blight			harvested dry with pods removed. Do not apply within 14 days of harvest. KT CTL 720 FUNGICIDE may be applied through sprinkler irrigation equipment. See calibration directions which appear on the product label.
<b>BEANS, SNAP</b>	Rust (Phakopsora spp.)	1.37-3.0	12.0	Use in sufficient water to obtain adequate coverage. Begin applications during early bloom stage or when disease first threatens and repeat at 7 day intervals. For resistance management of rust, alternate with another fungicide registered for bean rust control. Do not apply within 7 days of harvest.
	Botrytis Blight (Gray Mold)	3.0		
<b>BLUEBERRY</b>	Mummy Berry (suppression)Ant hracnose	3.0 - 4.0	12.0	Apply in sufficient water to obtain adequate coverage, normally 20 to 100 gallons per acre. Begin applications at bud break (green tip). Repeat applications at 10-day intervals until early bloom. Do not apply after full bloom, otherwise phytotoxicity may occur to developing fruit. Do not apply within a week before or after an oil application or a tank-mix containing oil-based pesticides. Do not apply within 42 days before harvest.
	Septoria Leaf Spot Rust	3.0 - 4.0	12.0	After all berries are harvested, a foliar application may be made to maintain healthy leaves for the following season. Apply in sufficient water (normally 20 to 100 gallons per acre) and repeat at 10 to 14 day intervals.
<b>CABBAGE BROCCOLI CAULIFLOWER CHINESE BROCCOLI CHINESE CABBAGE (only tight-</b>	Alternaria Leaf Spot Downy Mildew	1.5	11.7	Use in sufficient water to obtain adequate coverage. Begin applications after transplants are set in field, or shortly after emergence of field-seeded crop, or when conditions favor disease development. Repeat at 7 to 10 day intervals. Do not apply within 7 days

12  
28

CROP	DISEASES CONTROLLED	RATE OF KT CTL 720 FUNGICIDE PER APPLICATION PINTS/ACRE	SEASONAL LIMITS (PINTS/ ACRE/ YEAR)	APPLICATION DIRECTIONS
headed varieties) BRUSSELS SPROUTS				of harvest.
BRUSSELS SPROUTS (CA only)	Ring Spot	2.0	16.0	For field-seeded Brussels sprouts begin application at time of early sprout development or when conditions favor disease development. Repeat at 7 to 10 day intervals. Do not apply within 7 days of harvest.
CARROT	Cercospora (Early) Blight Alternaria (Late) Blight	1.5-2.0	20.0	Use in sufficient water to obtain adequate coverage. Start applications when disease threatens and repeat at 7 to 10 day intervals. KT CTL 720 FUNGICIDE may be applied the day of harvest. KT CTL 720 FUNGICIDE may be applied through sprinkler irrigation equipment (solid set, portable wheel move, motorized lateral move, or center pivot systems only). See calibration directions preceding this section.
CELERY	Cercospora (Early) Blight Septoria Late Blight Basal Stalk Rot (Rhizoctonia solani)	2.0 - 3.0	24.0	Start applications when transplants are set in the field. Apply in sufficient water to obtain adequate coverage. KT CTL 720 FUNGICIDE may be applied through sprinkler irrigation equipment (solid set, portable wheel move, motorized lateral move, or center pivot systems only). See calibration directions preceding this section. Do not apply within 7 days of harvest.
	Pink Rot (suppression)	3.0		
	Early Blight Late Blight	1.5-2.0 pints per 100 gallons	24.0	
CORN (Sweet) CORN (Grown for seed)	Helminthosporium Leaf Blight Rust	0.75-2.0	12.0	Use in sufficient water to obtain adequate coverage. Begin applications when conditions favor disease development and repeat at

13  
28

CROP	DISEASES CONTROLLED	RATE OF KT CTL 720 FUNGICIDE PER APPLICATION PINTS/ACRE	SEASONAL LIMITS (PINTS/ ACRE/ YEAR)	APPLICATION DIRECTIONS
				7 day intervals. Under severe disease conditions, use 1.5 to 2 pints per acre. Do not apply within 14 days of harvest. Do not apply to sweet corn to be processed. Do not ensile treated corn or use as livestock forage. Do not allow livestock to graze in treated fields.
<b>CRANBERRY</b>	Fruit Rot Lophodermium Leaf/Twig Blight	4.0-6.5	20.0	Apply at early bloom and repeat at 10 to 14 day intervals. Under severe disease conditions, use the 6.5 Pints per acre rate on a 10 day schedule. Do not apply within 50 days of harvest. Do not apply to bogs when flooded or allow release of irrigation water from bogs for at least 3 days following application. KT CTL 720 FUNGICIDE may be applied through sprinkler irrigation equipment. Use 300 gallons of water per acre through solid set systems only. See calibration directions preceding this section.
	Upright Dieback	4.0-6.5	20.0	Apply in sufficient water to uprights and runners making the first application before bloom when shoots begin growth in the spring. Apply at 10 to 14 day intervals. Do not apply within 50 days of harvest. Do not apply to bogs when flooded or allow release of irrigation water from bogs for at least 3 days following application. KT CTL 720 FUNGICIDE may be applied through sprinkler irrigation equipment. Use 300 gallons of water per acre through solid set systems only. See calibration directions preceding this section.
<b>CUCURBITS</b> Cantaloupe Cucumbers Honeydew Muskmelon Pumpkin Squash Watermelon	Anthracnose Downy Mildew Target Spot	1.5 - 2.0	21.0	Use in sufficient water to obtain adequate coverage. Begin applications when plants are in first true leaf stage or when conditions are favorable for disease development. Repeat applications at 7 day intervals. KT CTL 720 FUNGICIDE may be

14  
28

CROP	DISEASES CONTROLLED	RATE OF KT CTL 720 FUNGICIDE PER APPLICATION PINTS/ACRE	SEASONAL LIMITS (PINTS/ ACRE/ YEAR)	APPLICATION DIRECTIONS
				<p>applied the day of harvest. KT CTL 720 FUNGICIDE may be applied through sprinkler irrigation equipment (solid set, portable wheel move, or center pivot systems only). See Calibration directions preceding this section.</p> <p>Note: Spraying mature watermelons may result in sunburn of the upper surface of the fruit. Do not apply KT CTL 720 FUNGICIDE to watermelons when any of the following conditions are present:</p> <ul style="list-style-type: none"> <li>• Intense heat and sunlight,</li> <li>• Drought conditions,</li> <li>• Poor vine canopy,</li> <li>• Other crop and environmental conditions which may be conducive to increased natural sunburn.</li> </ul> <p>Do not combine KT CTL 720 FUNGICIDE with anything except water for application to watermelons unless your prior use has shown the combination to be non-injurious to watermelons under your conditions of use.</p>
<b>GRASSES GROWN FOR SEED</b>	Stem Rust Leaf Rust Stripe Rust Septoria Leaf Spot Glume Blotch Bipolaris Leaf Spot Drechslera Leaf Spot	1.0-1.5	6.0	Use in sufficient water to obtain adequate coverage. Begin applications during stem elongation when conditions favor disease development. Re-apply at flag (top) leaf emergence and repeat applications at 14 day intervals. Do not apply within 14 days of harvest. Do not allow livestock to graze on treated areas or feed hay Produced before harvest. Feeding of treated plant parts after harvest of seed is allowed. KT CTL 720 FUNGICIDE may be applied through sprinkler irrigation equipment (solid set, portable wheel move, or center pivot systems only). See calibration directions preceding this section.
	Selenophoma (Eyespot)	1.0-2.0		

15  
28

CROP	DISEASES CONTROLLED	RATE OF KT CTL 720 FUNGICIDE PER APPLICATION PINTS/ACRE	SEASONAL LIMITS (PINTS/ ACRE/ YEAR)	APPLICATION DIRECTIONS								
MANGO	Anthracnose	2.0-3.5	32.0	Use a water volume of 20 to 300 gallons per acre. Begin applications at early bloom and repeat on a 7-14 day interval until early fruit development. Begin the season with the 2 pint rate on a 14-day interval. If disease pressure is severe, use the higher rate and shorter interval. Do not apply within 21 days of harvest.								
MINT (IN, MI, ND, OR, WI only)	Rust Septoria Leaf Spot	1.37	4.0	Use in sufficient water to obtain adequate coverage, normally 20 to 150 gallons per acre for dilute sprays and 5 to 10 gallons per acre for concentrate ground and aircraft applications. Begin applications when emerging plants are 4 to 8 inches high. Repeat applications at 7 to 10 day intervals. Do not apply within 80 days of harvest. Do not feed fresh or extracted mint hay from treated fields to livestock.								
ONION (Dry bulb) GARLIC	Botrytis Leaf Blight/Blast Purple Blotch Suppression: Botrytis Neck Rot Downy Mildew	1.0-3.0	9.0	Apply in sufficient water to obtain thorough coverage of tops. KT CTL 720 FUNGICIDE is recommended for use with disease monitoring systems which adjust fungicide rates and frequency of application according to disease hazard. Apply as follows: <table border="0" style="margin-left: 40px;"> <tr> <td style="text-align: right;">Rate/Acre</td> <td style="text-align: left;">Frequency</td> </tr> <tr> <td style="text-align: right;"><b>Low Disease Hazard, prior to Infection</b></td> <td>1 pint      10 days</td> </tr> <tr> <td style="text-align: right;"><b>Low Disease Hazard, some disease present</b></td> <td>1.37 pints      7-10 days</td> </tr> <tr> <td style="text-align: right;"><b>High Disease Hazard</b></td> <td>3 pints      7 days</td> </tr> </table> For suppression of neck rot (Botrytis spp.) during storage, a minimum of 3 weekly applications prior to lifting, using 1 3/8 to 3 pints of KT CTL 720 FUNGICIDE per acre is recommended. Do not apply within 7 days of harvest.	Rate/Acre	Frequency	<b>Low Disease Hazard, prior to Infection</b>	1 pint      10 days	<b>Low Disease Hazard, some disease present</b>	1.37 pints      7-10 days	<b>High Disease Hazard</b>	3 pints      7 days
Rate/Acre	Frequency											
<b>Low Disease Hazard, prior to Infection</b>	1 pint      10 days											
<b>Low Disease Hazard, some disease present</b>	1.37 pints      7-10 days											
<b>High Disease Hazard</b>	3 pints      7 days											

16  
28

CROP	DISEASES CONTROLLED	RATE OF KT CTL 720 FUNGICIDE PER APPLICATION PINTS/ACRE	SEASONAL LIMITS (PINTS/ ACRE/ YEAR)	APPLICATION DIRECTIONS
<b>ONION (Green bunching)</b> <b>LEEK SHALLOT</b> <b>ONION AND GARLIC GROWN FOR SEED</b>	Botrytis Leaf Blight/Blast Purple Blotch Downy Mildew (suppression)	1.5-3.0	9.0	Use in sufficient water to obtain thorough coverage of tops. Begin applications prior to favorable infection periods and repeat at 7 to 10 day intervals for as long as conditions favor disease. Use the high rate and a 7 day schedule of applications when heavy dew or rain persists. Do not apply within 7 days of harvest on garlic. Do not apply within 14 days of harvest on green bunching onions, leeks, or shallots. KT CTL 720 FUNGICIDE may be applied through sprinkler irrigation equipment (solid set, portable wheel move, or center pivot systems only). See Calibration directions preceding this section.
<b>PAPAYA</b>	Alternaria Fruit Spot Anthracnose Stem End Rot	1.5-3.0	9.0	Apply with ground equipment only. Use sufficient water to obtain adequate coverage of fruit and leaves. Begin treatment when conditions favor development of disease and continue treatments at 14 day intervals until weather conditions no longer favor disease development. KT CTL 720 FUNGICIDE may be applied the day of harvest.
<b>PARSNIP</b>	Alternaria Leaf Spot Downy Mildew Anthracnose Botrytis Blight (Gray Mold) Bottom Rot (Rhizoctonia)	1.5-2.0	8.0	Apply in sufficient water to obtain adequate coverage. Make the first application at the first sign of disease or when conditions are favorable for infection. Continue applications on a 7 to 10 day schedule. Do not apply within 10 days of harvest. KT CTL 720 FUNGICIDE may be applied through sprinkler irrigation equipment (solid set, portable wheel move, or center pivot systems only). See calibration directions preceding this section.
<b>PASSION FRUIT (HI only)</b>	Alternaria Fruit and Leaf Spot (Passion Fruit Brown Spot)	2.0	10.0	Apply with ground equipment in sufficient water to obtain adequate coverage of fruit and leaves. Begin treatment when fruit spots appear



17  
28

CROP	DISEASES CONTROLLED	RATE OF KT CTL 720 FUNGICIDE PER APPLICATION PINTS/ACRE	SEASONAL LIMITS (PINTS/ ACRE/ YEAR)	APPLICATION DIRECTIONS
	Anthracnose Cercospora Fruit Spot			(April to July) and continue treatments at 14 day intervals until weather conditions no longer favor disease development. Do not apply within 7 days of harvest.
PEANUT	Early Leaf Spot (Cercospora) Late Leaf Spot (Cercosporidium) Pepper Spot	1.0-1.5	12.0	Apply in sufficient water for coverage when leaf wetness first occurs or 30 to 40 days after planting. Repeat at 14 day intervals. When conditions favor late leaf spot or when rust or web blotch occur, apply 1.5 pints per acre at 14 day intervals for the remainder of the season. Do not apply within 14 days of harvest. KT CTL 720 FUNGICIDE may be applied through sprinkler irrigation equipment. Use 1.5 pints per acre in solid set, portable wheel move, center pivot, motorized lateral move, or traveling gun sprinkler irrigation equipment. See calibration directions preceding this section. It is recommended to alternate chemigation applications With ground or aerial applications. Do not allow livestock to graze in treated areas. Do not feed hay or threshings from treated fields to livestock.
	Rust Web Blotch	1.5		
POTATO	Late Blight Early Blight Botrytis Vine Rot Black Dot	0.75 then 1.0-1.5	15.0	Begin applications at the low rate when vines are first exposed and leaf wetness occurs. Repeat applications at 5 to 10 day intervals. Begin applying the higher label rates at 5 to 10 day intervals when any one of the following events occur: <ul style="list-style-type: none"> <li>• Vines close within the rows;</li> <li>• Late blight forecasting measures 18 disease severity values (DSV);</li> <li>• The crop reaches 300 P-days</li> </ul> Increase water spray volume as canopy density increases. Use the highest rate and shortest interval

18  
28

CROP	DISEASES CONTROLLED	RATE OF KT CTL 720 FUNGICIDE PER APPLICATION PINTS/ACRE	SEASONAL LIMITS (PINTS/ ACRE/ YEAR)	APPLICATION DIRECTIONS
				when plants are rapidly growing and disease conditions are severe. Do not apply within 7 days of harvest. KT CTL 720 FUNGICIDE may be applied through sprinkler irrigation equipment (solid set, portable wheel move, center pivot, or motorized lateral move systems only). Do not exceed a 10 day interval between applications when using this technique. See calibration directions preceding this section.
SOYBEAN	Anthracnose Diaporthe Pod and Stem Blight Frogeye Leaf Spot ( <i>Cercospora sojina</i> ) Purple Seed Stain Cercospora Leaf Blight ( <i>Cercospora kikuchii</i> ) Septoria Brown Spot Rust (Suppression)	See Below	See Below	Apply in sufficient water to obtain complete coverage, using at least 5 gallons of water per acre for aerial application. Use the three application program in areas having a history of moderate to severe disease intensity. The minimum retreatment interval is 14 days. KT CTL 720 FUNGICIDE may be applied through sprinkler irrigation equipment. Follow application and calibration direction preceding this section. Do not apply within 6 weeks of harvest. Do not feed hay or threshings from treated fields to livestock.
		1.5-2.5	6.0	Two application program: For determinate varieties, make the first application at early pod set (R3 stage, when the majority of pods are 1/8 to 3/8 inches in length) and the second at beginning of seed formation (R5). For indeterminate varieties, make the first application when largest pods are 1 to 1.25 inches in length. Make the second application 14 days later.
		1.0-2.0	6.0	Three application program: For determinate varieties, make the first application at the beginning of flowering (R1), the second at early pod set (R3), and the third at beginning of seed formation (R5). For the indeterminate varieties, make the first application one week

19  
28

CROP	DISEASES CONTROLLED	RATE OF KT CTL 720 FUNGICIDE PER APPLICATION PINTS/ACRE	SEASONAL LIMITS (PINTS/ ACRE/ YEAR)	APPLICATION DIRECTIONS
	Stem Canker ( <i>Diaporthe phaseolorum var. caulivora</i> )	1.0	6.0	after first flowering and continue applications at 14 day intervals. Apply in 10 to 20 gallons of water per acre as a band treatment directing spray to provide coverage of entire plant. Make the first application at time of emergence of the second trifoliolate leaves (V2). If conditions favor stem canker disease, make a second and third application. Make all applications at 14 day intervals.
TOMATO	Foliage: Early Blight Late Blight Gray Leaf Spot Gray Leaf Mold Septoria Leaf Spot Target Spot	1.37-2.0	20.0	Apply in sufficient water to obtain adequate coverage. Begin applications when dew or rain occurs and disease threatens. Apply every 7 to 10 days for foliage diseases. For fruit diseases, begin at fruit set and apply every 7 to 14 days. Use the highest rate and shortest interval when disease is severe. KT CTL 720 FUNGICIDE may be applied the day of harvest. KT CTL 720 FUNGICIDE may be combined in the spray tank with EPA-registered pesticide products that claim copper as the active ingredient and are labeled for control of bacterial diseases in tomatoes. Check the copper manufacturer's label for specific instructions, precautions, and limitations prior to mixing with KT CTL 720 FUNGICIDE. Do not use with Copper Count N in concentrated spray suspensions. KT CTL 720 FUNGICIDE may be applied through sprinkler irrigation equipment (solid set or portable wheel move systems only). See calibration directions preceding this section.
	Fruit: Anthracnose Alternaria Fruit Rot (Black Mold) Botrytis Gray Mold Late Blight Fruit Rot Rhizoctonia Fruit Rot	2.0-2.75		

20  
28

CROP	DISEASES CONTROLLED	RATE OF KT CTL 720 FUNGICIDE PER APPLICATION PINTS/ACRE	SEASONAL LIMITS (PINTS/ ACRE/ YEAR)	APPLICATION DIRECTIONS
<b>STRAWBERRY (non-bearing nurseries)</b>	Ramularia leaf spot ( <i>Ramularia tulasnei</i> )	1.5	20	Apply in sufficient water to obtain adequate coverage. Begin application when conditions favor leaf spot development, usually following rainy weather or sprinkler irrigation. Repeat applications at 10 to 14 day intervals. Use the shortest interval when disease conditions are severe. Continue applications until runners are dug. KT CTL 720 FUNGICIDE may be applied to strawberry plants in nurseries through sprinkler irrigation equipment. Refer to the KT CTL 720 FUNGICIDE label for chemigation instructions. Do not use KT CTL 720 FUNGICIDE on strawberry plants in commercial fruit production.
<b>STRAWBERRY TRANSPLANTS (preplant dip)</b>	Ramularia leaf spot ( <i>Ramularia tulasnei</i> )	1.5 per 100 gallons of water.	20	Mix KT CTL 720 FUNGICIDE in water and stir the suspension thoroughly. Stir periodically to assure a uniform mixture. Dip strawberry transplants into the suspension for 5 to 10 minutes until plant surfaces are completely wetted. Transplant treated plant stock into nursery beds without rinsing. Wear chemical-resistant gloves of any waterproof material when mixing and applying KT CTL 720 FUNGICIDE as a transplant dip treatment and while handling treated stock. Do not use KT CTL 720 FUNGICIDE on strawberry plants in commercial fruit production.

21  
28

CROP	DISEASES CONTROLLED	RATE OF KT CTL 720 FUNGICIDE PER APPLICATION PINTS/ACRE	SEASONAL LIMITS (PINTS/ ACRE/ YEAR)	APPLICATION DIRECTIONS
------	---------------------	---	-------------------------------------	------------------------

**TREE AND ORCHARD CROPS—APPLICATION INSTRUCTIONS**

Apply KT CTL 720 FUNGICIDE in sufficient water and with proper calibration to obtain uniform coverage of tree canopy. Application with ground equipment is preferable to aerial application because ground applications generally give better coverage of the tree canopy. If application with ground equipment is not feasible, KT CTL 720 FUNGICIDE may be applied with aircraft using the spray volume in the table below. When concentrate sprays are used or when treating non-bearing or immature trees, the lower rate of KT CTL 720 FUNGICIDE listed may be used. Both ground and aircraft methods of application are recommended unless specific directions are given for a crop. Application through sprinkler irrigation systems is not recommended unless specific directions are given for a crop. See the following instructions for application and calibration. DO NOT allow livestock to graze treated areas. The following spray volumes are recommended as gallons of spray per acre:

CROP	SPRAY VOLUME (Gallons per Acre)	
Almonds	20 (concentrate) to 300 (full dilute)	
Filberts (Hazelnuts) (Oregon only)	20 (concentrate) to 300 (full dilute)	
Peach, Nectarine, Apricot, Tart Cherry, Plum, Prune	20 (concentrate) to 300 (full dilute)	
<b>Conifers:</b>	<b>Dilute</b>	<b>Concentrate</b>
Christmas Trees	100	10 to 50 (aircraft or ground equipment)
Nursery Beds	100	5 to 10 (ground equipment only)

22  
28

CROP	DISEASES CONTROLLED	KT CTL 720 FUNGICIDE PINTS/ACRE	KT CTL 720 FUNGICIDE PINTS/100 GAL*	SEASONAL LIMITS PINTS/ACRE	APPLICATION DIRECTIONS
Almonds	Anthracnose Blossom Blight/ Brown Rot Shothole Scab	4	1.33	25	For blossom blight, begin application at popcorn (pink bud) and follow with an application at full bloom. If weather is still conducive for disease development, another application may be made at petal fall. For control of shothole, make an application in the autumn at leaf fall. In the spring, make the first application at bud break, followed by an application at shuck split to control nut infections and to control scab. Do not apply within 150 days of harvest.
Filberts (Hazelnuts)	Eastern Filbert Blight	4	1.33	12.0	Begin applications at leaf bud break and repeat applications at 2 to 4 week intervals. Do not apply within a week before or after an oil application or a tank-mix containing oilbased pesticides. Do not apply within 120 days before harvest.

23  
28

CROP	DISEASES CONTROLLED	KT CTL 720 FUNGICIDE PINTS/ACRE	KT CTL 720 FUNGICIDE PINTS/100 GAL*	SEASONAL LIMITS PINTS/ACRE	APPLICATION DIRECTIONS
<b>FRUIT TREES</b> Apricot Cherry (Sweet) Cherry (Tart) Nectarine Peach Plum Prune	Leaf Curl Coryneum Blight (Shothole)	3.1-4.1	1.0-1.35	20.5	For best control of both diseases, apply at leaf fall in late autumn, using sufficient water and proper sprayer calibration to obtain uniform coverage. When conditions favor high disease levels, use the high rate of application and apply once or twice more in mid-to late winter before bud swell. If the leaf fall application is not practical, application of Equus 720 SST for control of leaf curl may be made at any time prior to bud swell the following spring. Where Coryneum blight (shothole) occurs, also apply at bud break to protect newly emerging leaves and at shuck split to prevent fruit infections. Make applications at a minimum of 10 day intervals. Equus 720 SST may be applied the day of harvest.
	Brown Rot Blossom Blight Lacy Russet Scab (Plum/Prune)	3.1-4.1	1.0-1.275		Make one application at popcorn (pink, red, or early white bud) and a second application at full bloom. If weather conditions favor disease development, make an additional application at petal fall. Make applications at a minimum of 10 day intervals. KT CTL 720 FUNGICIDE may be applied the day of harvest.

24  
28

	Cherry Leaf Spot Scab Black Knot (Cherry, Plum)	3.1-4.1	1.0-1.275		In addition to the bloom application listed above, make one application at shuck split. Do not apply KT CTL 720 FUNGICIDE after shuck split and before harvest. If additional disease control is needed before harvest, use another registered fungicide. For control of cherry leaf spot after harvest, make one application to foliage within 7 days after fruit is removed. In orchards with a history of high leaf spot incidence, make a second application 10 to 14 days later. Make applications at a minimum of 10 day intervals. Equus 720 SST may be applied the day of harvest.
*Volumetric rates to be used only with full dilute spray volume specified on this label for tree and orchard crops.					



25  
28

CROP	DISEASES CONTROLLED	KT CTL 720 FUNGICIDE PINTS/ACRE	SEASONAL LIMITS PINTS/ACRE	APPLICATION DIRECTIONS
CONIFERS Pines, Spruces	See Below	See Below	22.0	The minimum retreatment interval for established trees is 21 days. The minimum retreatment in nursery beds is 7 days.
	Swiss Needlecast	2.75-5.5		Single-application technique: In Christmas tree plantations make one application in the spring when new shoot growth is 1/2 to 2 inches in length.
	Scleroderris Canker (Pines), Swiss Needlecast	1.5-2.75		Make the first application in spring when new shoot growth is 1/2 to 2 inches in length. Make additional applications at 4 week intervals until conditions no longer favor disease development. For use in nursery beds, apply the highest rate specified on a 4 week schedule.
	Sirococcus Tip Blight	2.0-3.5		Apply in early spring prior to bud break. Repeat applications at approximately 6 to 8 week intervals, until spore release ceases in late fall. Apply monthly during periods of frequent rainfall and where Lophodermium infections occur during dormancy (Pacific Northwest). During drought periods, applications may be suspended, then resumed upon next occurrence of needle wetness.
	Rhizosphaera Needlecast (Spruces), Scirrhia Brown Spot (Pines)	5.5		Apply at bud break and repeat at 3 to 4 week intervals until needles are fully elongated and conditions no longer favor disease development. In plantations of mixed provenance, or when irregular bud break occurs, apply weekly until all trees have broken bud, then every 3 to 4 weeks as specified above. In nursery beds, use the high rate on a 3 week schedule.
	Cyclaneusma and Lophodermium Needlecasts (Pines)	2.75-5.5		Begin applications in nursery beds when seedlings are 4 inches tall and when cool, moist conditions favor disease development. Make additional applications at 7 to 14 day intervals as long as disease favorable conditions persist.
	Rhabdocline Needlecast (Douglas fir)	1.5-2.75		Begin applications when 10% of buds have broken and twice thereafter at 7 to 10 day intervals.
	Botrytis Seedling Blight Phoma Twig Blight	1.5-2.75		
Autoecious Needle Rust (Weir's Cushion) (Spruces)	5.5			

26  
28

**MUSHROOMS:** Verticillium Brown Spot and Dry Bubble – Apply 2.75 to 5.5 fl. oz. of KT CTL 720 FUNGICIDE per 1,000 sq. ft. of mushroom bed. Apply as a drench to the mushroom bed surface in at least 12.5 gallons of water per 1,000 sq. ft. of mushroom bed. Make two applications. Apply the high rate (5.5 fl.oz.) of KT CTL 720 FUNGICIDE in the first application and the low rate (2.75 fl. oz.) of KT CTL 720 FUNGICIDE in the second application. The first application should be made within two days of topdressing the spawn-colonized mushroom compost with a casing layer. The second application should be made at pinning. Do not apply within 5 days of first harvest. Make no more than two applications per cropping cycle. Do not apply more than 8.25 fl. oz. of KT CTL 720 FUNGICIDE per cropping cycle.

**GRASS: SODFARMS**

Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields located on or next to schools (i.e., elementary, middle, and high schools), campgrounds, churches, and theme parks.

Apply KT CTL 720 FUNGICIDE in 30 to 40 gallons of water per acre. Begin applications when conditions favor disease development and repeat applications as long as these conditions persist using the rates recommended in the following table.

Under severe disease conditions, a single application of 15 pints per acre may be made with a 7 day retreatment interval. Subsequent applications must follow the rates and retreatment intervals outlined in the following table for the remainder of the year.

Do not mow or water after treatment until spray deposited on grass is thoroughly dry. KT CTL 720 FUNGICIDE should always be used in conjunction with good turf management practices.

Sodfarm turf treated with chlorothalonil prior to harvest must be mechanically cut, rolled, and harvested. Follow all provisions outlined in the Agricultural Use Requirements box.

DISEASES CONTROLLED	LOW DISEASE PRESSURE TREATMENT REGIME		EXTREME DISEASE CONDITION		Application Limit Per Year for Sod farms (Pints/Acre)*
	Retreatment Interval (Days)	Application Rate (Pints/Acre)	Maximum Single Application Allowed in a Year (Pints/Acre)	Minimum Retreatment Interval for the Maximum Single Application (Days)	
Dollar Spot	7-10	2.75 <sup>a</sup> -5.5	15	7	17
	14-21	5.5-9.66			
Leaf Spot, Melting Out, Brown Blight	7-10	5.5			
	14-21	5.5-9.66			
Brown Patch	7-14	5.5-9.66			
Gray Leaf Spot	7-10	5.5-9.66			
Red Thread	7-10	5.5-9.66			
Anthracnose	7-14	8.12-9.66			

<sup>a</sup> Low rate is not effective on intensively mowed grasses.

Diseases are caused by some of the following fungi:

Dollar Spot: *Sclerotinia homeocarpa*, *Lanzia* or *Moellerodiscus* spp.

Leaf Spot, Melting Out and Brown Blight: *Drechslera* spp., *Bipolaris* spp., *Curvularia* spp.

Brown Patch: *Rhizoctonia* spp.

Anthracnose: *Collectotrichum*

\* Do not use for sodfarms at application rates greater than 13 lbs. a.i. (17 pints of KT CTL 720 FUNGICIDE) per acre per year.

27  
28

#### **STORAGE AND DISPOSAL**

Do not contaminate water, foodstuffs, feed, or seed by storage or disposal.

**PESTICIDE STORAGE:** Store in a cool place. Protect from excessive heat. Store product in original container only away from water, food, or feed. Keep container closed to prevent spills and contamination. Carefully open containers. After partial use, replace lid and close tightly.

Do not put concentrate or diluted product into food or drink containers.

**PESTICIDE DISPOSAL:** Do not contaminate water, food, or feed by disposal. Improper disposal of excess pesticide, pesticide spray, or rinsate is a violation of Federal law. Wastes resulting from the use of this product that cannot be used according to the label instructions or chemically reprocessed may be disposed of on site or at a landfill or waste disposal facility approved for pesticide disposal, or in accordance with all applicable Federal, state, or local regulations. For further guidance, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER HANDLING:** Empty containers retain vapor and product residues.

**Nonrefillable Container (five gallons or less):** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Nonrefillable Container (greater than five gallons):** Non refillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Refillable Container:** Refillable container. Refill this container with chlorothalonil only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

#### **CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials, or other influencing factors in the use of the product, which are beyond the control of Kaizen

28  
28

Technologies or Seller. All such risks shall be assumed by the Buyer and User, and Buyer and User agree to hold Kaizen Technologies and Seller harmless for any claims relating to such factors. Kaizen Technologies warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Kaizen Technologies, and Buyer and User assume the risk for such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

To the extent consistent with applicable law, neither Kaizen Technologies nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF KAIZEN TECHNOLOGIES OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, STRICT LIABILITY, TORT OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR, AT THE ELECTION OF KAIZEN TECHNOLOGIES OR SELLER, THE REPLACEMENT OF THE PRODUCT.

Kaizen Technologies and Seller offer this product, and Buyer and User accept it, subject to the forgoing Conditions of Sale and Limitation of Warranty and Liability which may not be modified except by written agreement signed by a duly authorized representative of Kaizen Technologies.

Copper-Count N is a registered trademark of Mineral Research and Development Corporation.  
DiPel is a registered trademark of Abbott Laboratories.  
Foil is a registered trademark of Ecogen, Inc.  
Triton is a registered trademark of Union Carbide Corp.