



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Biopesticides and Pollution Prevention Division (7511P)
 1200 Pennsylvania Ave., N.W.
 Washington, D.C. 20460

EPA Reg. Number:

86583-5

Date of Issuance:

4/6/2017

NOTICE OF PESTICIDE:

Registration
 Reregistration
 (under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

EcoBird 14.5

Name and Address of Registrant:

Roth Chemical Company Ltd.
 6600 College Blvd.
 Suite 220
 Overland Park, KS 66211

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Biopesticides and Pollution Prevention Division prior to use of the label in commerce. In any correspondence on this product, always refer to the above EPA Registration Number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA or the Act).

Registration is in no way to be construed as an endorsement or recommendation of this product by the U.S. Environmental Protection Agency (EPA). In order to protect health and the environment, the Administrator, on his or her motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under the Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration or registration review of your product when the EPA requires all registrants of similar products to submit such data.

Signature of Approving Official:

Andrew Bryceland, Team Leader
 Biochemical Pesticides Branch
 Biopesticides and Pollution Prevention Division (7511P)
 Office of Pesticide Programs

Date:

4/6/2017

2. Make the following labeling change before you release this product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 86583-5.”
3. Submit one (1) copy of the final printed labeling for the record before you release this product for shipment.

Should you wish to add/retain a reference to your company’s website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA-approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6. A stamped copy of the labeling is enclosed for your records. Please also note that the record for this product currently contains the following acceptable Confidential Statement of Formula (CSF):

- Basic CSF dated 03/17/2017

Any CSFs other than those listed above are superseded.

If you have any questions, please contact Cody Kendrick of my branch by phone at (703) 347-0468 or via email at kendrick.cody@epa.gov.

Sincerely,




Andrew Bryceland, Team Leader
Biochemical Pesticides Branch
Biopesticides and Pollution
Prevention Division (7511P)
Office of Pesticide Programs

Enclosure

MASTER LABEL

EcoBird 14.5

Goose and Bird Repellent



For Commercial, Agricultural and Home Use
To be used on Agricultural Commodities (Berries, Apples, Stone Fruits, Cereal Grains, and Sunflowers) and Non-Agricultural Sites including Turf, Hydroseeding, Garbage, Food Stuffs, Waste & Trash, and Spot Repelling

ACTIVE INGREDIENT:
Methyl Anthranilate (MA), methyl 2-aminobenzoate 14.5%
OTHER INGREDIENTS: 85.5%
TOTAL100.0%

This product contains 1.23 pounds/0.31 pounds/0.15 pounds of Methyl Anthranilate per gallon/quart/pint of product.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID	
If in eyes	- Hold eye open and rinse slowly and gently with water for 15-30 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. - Call a poison control center or doctor for treatment advice.
If on skin or clothing	- Take off contaminated clothing. - Rinse skin immediately with plenty of water for 15-20 minutes. - Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.	

Manufactured for:

Roth Chemical Company, Ltd.
941 West 141st Terrace, Suite B
Kansas City, Missouri 64145

EPA Reg. No. 86583-

EPA Est. No.

NET CONTENTS: 1 pint, 1 quart, 1 gallon, 2.5 gallons

Sub-label #1: Commercial and Agricultural Market

EcoBird 14.5

Goose and Bird Repellent
For Commercial and Agricultural Use

To be used on Agricultural Commodities (Berries, Apples, Stone Fruits, Cereal Grains, and Sunflowers)
and Non-Agricultural Sites including Turf, Hydroseeding,
Garbage, Food Stuffs, Waste & Trash, and Spot Repelling

ACTIVE INGREDIENT:
Methyl Anthranilate (**MA**), methyl 2-aminobenzoate 14.5%
OTHER INGREDIENTS: 85.5%
TOTAL 100.0%

This product contains 1.23 pounds/0.31 pounds/0.15 pounds of Methyl Anthranilate per gallon/quart/pint of product.

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID	
If in eyes	- Hold eye open and rinse slowly and gently with water for 15-30 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. - Call a poison control center or doctor for treatment advice.
If on skin or clothing	- Take off contaminated clothing. - Rinse skin immediately with plenty of water for 15-20 minutes. - Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.	

Manufactured for:

Roth Chemical Company, Ltd.
941 West 141st Terrace, Suite B
Kansas City, Missouri 64145

EPA Reg. No. 86583-

EPA Est. No.

NET CONTENTS: 1 pint, 1 quart, 1gallon, 2.5 gallons

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/ maintaining. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users Should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements, specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance **with** its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this labeling about PPE, restricted-entry intervals, and notification to workers. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS).

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of zero (0) hours.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried.

AGRICULTURAL CROPS

USE RESTRICTIONS:

Use this product to limit damage caused by birds on ripening crops. This product must be applied using appropriate spray equipment. Do not apply when crop is wet or rain is expected. Withhold treatment until fruit is susceptible to bird damage. Do not apply through any type of irrigation system.

COMPATIBILITY WITH OTHER PESTICIDES:

Compatibility with other adjuvants has not been tested. If adding other adjuvants or products, test for compatibility first or ask your representative for advice. Do not mix with dry formulations such as wettable powder formulations which may result in residue on the crop.

DILUTION DIRECTIONS:

Shake or stir container well prior to diluting. For better coverage, add a super wetter (such as a non-ionic, organofunctional silicone surfactant, or sticker spreader), as directed on the label. Mix product outside or in a ventilated area. For ground application, mix product with water at a ratio of 1 part product to 10-20 parts water. For aerial application, mix product with water at a ratio of 1 part product to 1-3 parts water.

APPLICATION DIRECTIONS

- Apply spray mixture so that crop receives thorough coverage.
- Repeat treatment if significant bird damage begins to occur.
- This product may be applied up to the day of harvest.

USE SITE	APPLICATION DIRECTIONS	NOTES
Apples	Apply by ground application at a rate of 2.5 - 5 gallons (3.1-6.2 lbs. a.i.) per acre or 7.5-15 fl. oz. per 1000 sq. ft. Use a 1: 10 or 1 :20 dilution with water and apply at rates of 27.5-55 gallons or 52.5-100 gallons, respectively, of spray mixture per acre.	California only: Mixers/handlers and air blast applicators must wear coveralls over normal work clothes and gloves. Harvesters must wear gloves and long- sleeved shirt. For mixers/loaders, daily usage of this product is limited to a maximum of 40 acres.
Stone Fruits Apricots Cherries Peaches Plums	Apply by ground application at a rate of 2.5 - 5 gallons (3.1-6.2 lbs. a.i.) per acre or 7.5-15 fl. oz. per 1000 sq. ft. Use a 1: 10 or 1 :20 dilution with water and apply at rates of 27.5-55 gallons or 52.5-100 gallons, respectively, of spray mixture per acre.	California only: Mixers/handlers and air blast applicators must wear gloves and coveralls over normal work clothes and not prepare spray solutions using more than 50 gallons of concentrate per day. Harvesters must wear gloves and long-sleeved shirt. For mixers/loaders, daily usage of this product is limited to a maximum of 40 acres.
Blueberries	Apply by ground application at a rate of 2.5 - 5 gallons of (3.1-6.2 lbs. a.i.) per acre or 7.5-15 fl. oz. per 1000 sq. ft. Use a 1: 10 or 1 :20 dilution with water and apply at rates of 27.5-55 gallons or 52.5-100 gallons, respectively, of spray mixture per acre.	California only: Mixers/handlers and air blast applicators must wear gloves and coveralls over normal work clothes and not prepare spray solutions using more than 50 gallons of concentrate per day. Harvesters must wear gloves and long-sleeved shirt. For mixers/loaders, daily usage of this product is limited to a maximum of 40 acres.

USE SITE	APPLICATION DIRECTIONS	NOTES
Grapes (Table)	Apply by ground application at a rate of 2.5 - 5 gallons (3.1-6.2 lbs. a.i.) per acre or 7.5-15 fl. oz. per 1000 sq. ft. Use a 1:10 or 1:20 dilution with water and apply at rates of 27.5-55 gallons or 52.5-100 gallons, respectively, of spray mixture per acre.	California only: Mixers/handlers and air blast applicators must wear gloves and coveralls over normal work clothes and not prepare spray solutions using more than 50 gallons of concentrate per day. Harvesters must wear gloves and long-sleeved shirt. For mixers/loaders, daily usage of this product is limited to a maximum of 40 acres. New York only: Aerial application is prohibited. This product is not to be used on grapes within Suffolk County.
Cereal grains Corn (including sweet and pop) Millet Oats Sorghum	Apply by aerial application at a rate of 2.5 - 5 gallons (3.1-6.2 lbs. a.i.) per acre or 7.5-15 fl. oz. per 1000 sq. ft. Use a 1:3 dilution with water and apply at rates of 10-20 gallons of spray mixture per acre.	Begin applications when birds begin feeding or crop begins to ripen. California only: Mixers/handlers and air blast applicators must wear coveralls over normal work clothes and gloves. Harvesters must wear gloves and long-sleeved shirt. For mixers/loaders, daily usage of this product is limited to a maximum of 40 acres. New York only: Aerial application is prohibited.
Sunflowers	Apply by aerial application at a rate of 2.5 - 5 gallons (3.1-6.2 lbs. a.i.) per acre or 7.5-15 fl. oz. per 1000 sq. ft. Use a 1:3 dilution with water and apply at rates of 10-20 gallons of spray mixture per acre.	Reapply: 22 days after application. California only: Mixers/handlers and air blast applicators must wear coveralls over normal work clothes and gloves. Harvesters must wear gloves and long-sleeved shirt. For mixers/loaders, daily usage of this product is limited to a maximum of 40 acres. New York only: Aerial application is prohibited.

TURF AND HYDROSEEDING

USE RESTRICTIONS:

Use this product to repel birds from golf courses, other turf areas (including around agricultural buildings and farmsteads), and hydroseeded areas. This product must be applied using appropriate spray equipment. Do not apply when grass is wet or rain is expected.

DILUTION DIRECTIONS:

Shake or stir container well prior to diluting. For better coverage, add a super wetter (such as a non-ionic, organofunctional silicone surfactant, or sticker spreader), as directed on the label. Mix product outside or in a ventilated area. Mix product with water at a ratio of 1 part product to 20 parts water. For example, mix 2 pints of product with 5 gallons of water to make 5.25 gallons of spray mixture.

APPLICATION DIRECTIONS:

Apply at a rate of 1.2 gallons of spray mixture (0.07 lbs. a.i.) per 1000 sq. ft. of turf. Spray evenly on area to be protected to provide thorough coverage and allow material to dry before permitting human activity on the treated area. Repeat application in 4 days or as warranted by Canada goose activity. Do not mow treated area for several days after application.

GARBAGE, FOOD STUFFS, WASTE, AND TRASH**USE RESTRICTIONS:**

Use this product to repel birds from garbage, spilled grain, and other waste food stuffs. This product must be applied using appropriate spray equipment. Do not apply if rain is expected.

DILUTION DIRECTIONS:

Shake or stir container well prior to diluting. For better coverage, add a super wetter (such as a non-ionic, organofunctional silicone surfactant, or sticker spreader), as directed on the label. Mix product outside or in a ventilated area. Mix product with water at a ratio of 10-50% v/v with water. To make a 20% v/v mixture, mix 1 gallon of product with 5 gallons of water. To make a 50% v/v mixture, mix 1 gallon of product with 1 gallon of water.

APPLICATION DIRECTIONS:

Spray mixture evenly on area to be protected to provide thorough coverage and allow material to dry before permitting human activity on the treated area. Repeat application as warranted by bird activity.

SPOT REPELLING**USE RESTRICTIONS:**

Use this product to immediately repel birds from in or on structures, turf areas, landfills, and other outdoor areas (including around agricultural buildings and farmsteads). This product must be applied using appropriate spray equipment.

DILUTION DIRECTIONS:

Shake or stir container well prior to diluting. Mix product outside or in a ventilated area. Mix product with water at a ratio of 10-50% v/v with water. To make a 20% v/v mixture, mix 1 gallon of product with 5 gallons of water. To make a 50% v/v mixture, mix 1 gallon of product with 1 gallon of water.

APPLICATION DIRECTIONS:

Spray mixture evenly on area to be protected to provide thorough coverage and allow material to dry before permitting human activity on the treated area. Repeat application as warranted by bird activity.

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Product should be stored in its original container in a cool, dry locked place out of reach of children and out of direct sunlight. Do not use or store near heat or open flame.

PESTICIDE DISPOSAL: To avoid wastes, use all material in this container by application according to label directions. If waste cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Triple rinse (or equivalent) promptly after emptying. Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds, after the flow begins to drip. Fill the container 1 /4 full with water and soap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. If burned, stay out of smoke.

WARRANTY STATEMENT

Notice: The statements on this label are believed to be true and accurate, but because conditions are beyond our control, Roth Chemical Company Ltd. does not make, nor does it authorize any agent or representative to make any oral, or written warranty, guarantee, or representation, express

or implied, concerning this material or the use thereof beyond the statements of the label. Roth Chemical Company Ltd. d. and the seller disclaim liability for, and expect the buyer to assume all risk for any claim or personnel injury or property damage or loss resulting from handling, storage, or use of this material not in accordance with the written directions. No other warranty exists.

Sub-label #2: Residential Market

EcoBird 14.5

Goose and Bird Repellent

For Home Use

To be used on Agricultural Commodities (Berries, Apples, Stone Fruits, Cereal Grains, and Sunflowers) and Non-Agricultural Sites including Turf, Hydroseeding, Garbage, Food Stuffs, Waste & Trash, and Spot Repelling

ACTIVE INGREDIENT:

Methyl Anthranilate (MA), methyl 2-aminobenzoate14.5%

OTHER INGREDIENTS:85.5%

TOTAL100.0%

This product contains 1.23 pounds/0.31 pounds/0.15 pounds of Methyl Anthranilate per gallon/quart/pint of product.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID	
If in eyes	- Hold eye open and rinse slowly and gently with water for 15-30 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. - Call a poison control center or doctor for treatment advice.
If on skin or clothing	- Take off contaminated clothing. - Rinse skin immediately with plenty of water for 15-20 minutes. - Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.	

Manufactured for:

Roth Chemical Company, Ltd.
941 West 141st Terrace, Suite B
Kansas City, Missouri 64145

EPA Reg. No. 86583- .

EPA Est. No.

NET CONTENTS: 1 pint, 1 quart, 1 gallon

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

CAUTION: Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wear gloves when handling and applying this product. Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area **will** help avoid run off to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

CROPS

USE RESTRICTIONS:

Use this product to limit damage caused by birds on ripening crops. This product must be applied using appropriate spray equipment. Do not apply when crop is wet or rain is expected. Withhold treatment until fruit is susceptible to bird damage.

COMPATIBILITY WITH OTHER PESTICIDES:

Compatibility with other adjuvants has not been tested. If adding other adjuvants or products, test for compatibility first or ask your representative for advice. Do not mix with dry formulations such as wettable powder formulations which may result in residue on the crop.

DILUTION DIRECTIONS:

Shake or stir container well prior to diluting. For better coverage, add a super wetter (such as a non-ionic, organofunctional silicone surfactant, or sticker spreader), as directed on the label. Mix product outside or in a ventilated area. For ground application, mix product with water at a ratio of 1 part product to 10-20 parts water. For example, mix 1 fl. oz. of product with 2 cups of water to make a spray mixture of 1 part product and 16 parts water.

APPLICATION DIRECTIONS:

- Apply spray mixture so that crop receives thorough coverage.
- Repeat treatment if significant bird damage begins to occur.
- This product may be applied up to the day of harvest
- To cover 100 square feet, 1 fl. oz. of product with the appropriate dilution of water is required (1 :10 or 1 :20).

USE SITE	APPLICATION DIRECTIONS	NOTES
Apples	Apply by ground application at a rate of 7.5-15 fl. oz. per 1000 sq. ft. Use a 1:10 or 1 :20 dilution with water.	

USE SITE	APPLICATION DIRECTIONS	NOTES
Stone Fruits Apricots Cherries Peaches Plums	Apply by ground application at a rate of 7.5-15 fl. oz. per 1000 sq. ft. Use a 1:10 or 1:20 dilution with water.	
Blueberries	Apply by ground application at a rate of 7.5-15 fl. oz. per 1000 sq. ft. Use a 1:10 or 1:20 dilution with water.	
Grapes (Table)	Apply by ground application at a rate of 7.5-15 fl. oz. per 1000 sq. ft. Use a 1:10 or 1:20 dilution with water.	New York only: This product is not to be used on grapes within Suffolk County.

TURF

USE RESTRICTIONS:

Use this product to repel Canada Geese from golf courses and other turf areas (including around agricultural buildings and farmsteads). This product must be applied using appropriate spray equipment, such as a hand, backpack, or pump sprayer. Do not apply when grass is wet or rain is expected.

DILUTION DIRECTIONS:

Shake or stir container well prior to diluting. Mix product outside or in a ventilated area. Mix with water at a ratio of 1 part product to 12 parts water. For example, mix 8 fl. oz. of product with 96 fl. oz. (3 qts.) of water to make 104 fl. oz. of spray mixture.

APPLICATION DIRECTIONS:

Apply at a rate of 1.2 gallons spray mixture (0.07 lbs. a.i.) per 1000 square feet of turf. Spray evenly on area to be protected to provide thorough coverage and allow material to dry before permitting human activity on the treated area. Repeat application in 4 days or as warranted by Canada goose activity. Do not mow treated area for several days after application.

GARBAGE, FOOD STUFFS, WASTE, AND TRASH

USE RESTRICTIONS:

Use this product to repel birds from garbage, trash, spilled grain, and other waste food stuffs. This product must be applied using appropriate spray equipment. Do not apply rain if expected.

DILUTION DIRECTIONS:

Shake or stir container well prior to diluting. For better coverage, add a super wetter (such as a non-ionic, organofunctional silicone surfactant, or sticker spreader), as directed on the label. Mix product outside or in a ventilated area. Mix product with water at a ratio of 10-50% v/v with water. To make a 20% v/v mixture, mix 1 pint of product with 4 pints of water. To make a 50% v/v mixture, mix 1 pint of product with 1 pint of water.

APPLICATION DIRECTIONS:

Spray mixture evenly on area to be protected to provide thorough coverage and allow material to dry before permitting human activity on the treated area. Repeat application as warranted by bird activity.

SPOT REPELLING

USE RESTRICTIONS:

Use this product to immediately repel birds from in or on structures, turf areas, and other outdoor areas. This product must be applied using appropriate spray equipment.

DILUTION DIRECTIONS:

Shake or stir container well prior to diluting. Mix product outside or in a ventilated area. Mix with water at a ratio of 10-50% v/v with water. To make a 20% v/v mixture, mix 1 pint of product with 4 pints of water. To make a 50% v/v mixture, mix 1 pint of product with 1 pint of water.

APPLICATION DIRECTIONS:

Spray mixture evenly on area to be protected to provide thorough coverage and allow material to dry before permitting human activity on the treated area. Repeat application as warranted by bird activity.

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Product should be stored in its original container in a cool, dry locked place out of reach of children and out of direct sunlight. Do not use or store near heat or open flame.

PESTICIDE DISPOSAL: To avoid wastes, use all material in this container by application according to label directions. If waste cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

WARRANTY STATEMENT

Notice: The statements on this label are believed to be true and accurate, but because conditions are beyond our control, Roth Chemical Company, Ltd. does not make, nor does it authorize any agent or representative to make any oral, or written warranty, guarantee, or representation, express or implied, concerning this material or the use thereof beyond the statements of the label. Roth Chemical Company, Ltd. and the seller disclaim liability for, and expect the buyer to assume all risk for any claim or personnel injury or property damage or loss resulting from handling, storage, or use of this material not in accordance with the written directions. No other warranty exists.