



U.S. ENVIRONMENTAL PROTECTION AGENCY  
 Office of Pesticide Programs  
 Registration Division (7505P)  
 1200 Pennsylvania Ave., N.W.  
 Washington, D.C. 20460

**EPA Reg. Number:**  
 87235-4

**Date of Issuance:**  
 2/16/20

**NOTICE OF PESTICIDE:**

Registration  
 Reregistration  
 (under FIFRA, as amended)

**Term of Issuance:**  
 Conditional

**Name of Pesticide Product:**  
 Monark SB

**Name and Address of Registrant (include ZIP Code):**

Andrej Branc  
 North American Business Manager  
 PelGar USA/PelGar International Ltd.  
 215 Evergreen Court  
 Saylorsburg, PA 18353

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

**Signature of Approving Official:**

Gene Benbow, Product Manager 07  
 Invertebrate and Vertebrate Branch 3 Registration Division (7505P)

**Date:**

2/16/20

2. You are required to comply with the data requirements described in the DCI identified below:

a. Difenacoum GDCI-119901

You must comply with all of the data requirements within the established deadlines. If you have questions about the Generic DCI listed above, you may contact the Chemical Review Manager in the Pesticide Reevaluation Division: <http://iaspub.epa.gov/apex/pesticides/f?p=chemicalsearch:1>

3. Make the following label changes before you release the product for shipment:

- Revise the EPA Registration Number to read, “EPA Reg. No. 87235-4.”

4. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.


If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 2/6/2020

If you have any questions, please contact Marianne Lewis by phone at (703) 308-8043, or via email at [lewis.marianne@epa.gov](mailto:lewis.marianne@epa.gov).

Enclosure

[Large Transport Container Label – Front Panel]



**MONARK SB**

[Soft Bait Rodenticide] [Difenacoum Soft Bait]

ACTIVE INGREDIENT:  
 Difenacoum (CAS #56073-07-5):..... 0.005%  
 OTHER INGREDIENTS^:..... 99.995%  
 Total 100.000%

^Contains Denatonium Benzoate

**KEEP OUT OF REACH OF CHILDREN  
 CAUTION**

FIRST AID	
If swallowed	<ul style="list-style-type: none"> <li>•Call a poison control center or doctor immediately for treatment advice.</li> <li>•Have person sip a glass of water if able to swallow.</li> <li>•Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>•Do not give anything by mouth to an unconscious person.</li> </ul>
If on skin or clothing	<ul style="list-style-type: none"> <li>•Take off contaminated clothing.</li> <li>•Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>•Call a poison control center or doctor immediately for treatment advice.</li> </ul>
If in the eyes	<ul style="list-style-type: none"> <li>•Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>•Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>•Call a poison control center or doctor immediately for treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor, or when going for treatment. You may also contact 1-800-222-1222 for emergency medical treatment information.	
<b>Note to Physician</b>	
If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K1 intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.	
<b>Treatment for Pet Poisoning</b>	
If animal eats bait, call veterinarian at once.	
<b>Note to Veterinarian</b>	
Anticoagulant Difenacoum: For animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K1. If needed, check prothrombin times every 3 days until values return to normal (up to 30 days). In severe cases, blood transfusions may be needed.	

**PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if swallowed. Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Keep away from children, domestic animals and pets.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

All handlers (including applicators) must wear: shoes plus socks, long pants and waterproof gloves. Any person who retrieves carcasses or unused bait following application of this product must wear gloves.

**USER SAFETY REQUIREMENTS:** Follow manufacturer's instruction for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS:**

Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This product is extremely toxic to fish, birds and other wildlife. Predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten bait. Do not apply this product directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

**DISCLAIMER:** To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions

SEE PRODUCT LABEL FOR PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE

Manufactured by:  
PelGar International Ltd.  
Unit 13 Newman Lane  
Alton, Hampshire GU34 2QR, UK  
EPA Reg. No. 87235-XXX  
EPA Est. No. 87235-GBR-001

Imported by:  
PelGar USA  
Saylorsburg, PA  
1-610-849-1723

*NOT REGISTERED FOR SALE OR USE IN CALIFORNIA*

DoE:  
DoM:  
Batch #:

[PCO Outer Container Label – Front Panel]

# Monark SB

[Soft Bait Rodenticide] [Difenacoum Soft Bait]

**KILLS RATS, MICE AND MEADOW VOLES  
KILLS WARFARIN RESISTANT NORWAY RATS**

ACTIVE INGREDIENT:	
Difenacoum (CAS #56073-07-5):.....	0.005%
OTHER INGREDIENTS^:.....	<u>99.995%</u>
Total	100.000%

^Contains Denatonium Benzoate

## KEEP OUT OF REACH OF CHILDREN CAUTION

See Back Panel for additional precautionary labeling.

**Net Weight:** 16 lbs, 18 lbs, 24 lbs as (4 pails X 6 lbs each), 24 lbs (as bags of ½ lb to 8 lb)

*Statement required on label of inner packages holding less than 16 pounds of bait:*

For packages holding 4 x 6-lb pails, do not sell these pails individually.

For packages holding ½ lb to 8 lb bags, INDIVIDUAL SALE IS PROHIBITED BY LAW.

Manufactured by: Imported by:

PelGar International Ltd.  
Unit 13 Newman Lane  
Alton, Hampshire GU34 2QR, UK  
EPA Reg. No. 87235-  
EPA Est. No. 87235-GBR-001

PelGar USA  
Saylorsburg, PA  
1-610-849-1723

**NOT REGISTERED FOR SALE OR USE IN CALIFORNIA**

DoE:
DoM:
Batch #:

## [PCO Outer Container Label – Side/Back Panel]

<b>FIRST AID</b>	
If swallowed	<ul style="list-style-type: none"> <li>•Call a poison control center or doctor immediately for treatment advice.</li> <li>•Have person sip a glass of water if able to swallow.</li> <li>•Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>•Do not give anything by mouth to an unconscious person.</li> </ul>
If on skin or clothing	<ul style="list-style-type: none"> <li>•Take off contaminated clothing.</li> <li>•Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>•Call a poison control center or doctor immediately for treatment advice.</li> </ul>
If in the eyes	<ul style="list-style-type: none"> <li>•Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>•Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>•Call a poison control center or doctor immediately for treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor, or when going for treatment. You may also contact 1-800-222-1222 for emergency medical treatment information.	
Note to Physician	
If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K1 intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.	
Treatment for Pet Poisoning	
If animal eats bait, call veterinarian at once.	
Note to Veterinarian	
Anticoagulant Difenacoum: For animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K1. If needed, check prothrombin times every 3 days until values return to normal (up to 30 days). In severe cases, blood transfusions may be needed.	

**PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if swallowed. Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Keep away from children, domestic animals and pets.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

All handlers (including applicators) must wear: shoes plus socks, long pants and waterproof gloves. Any person who retrieves carcasses or unused bait following application of this product must wear gloves.

**USER SAFETY REQUIREMENTS:** Follow manufacturer's instruction for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS:**

Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This product is extremely toxic to fish, birds and other wildlife. Predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten bait. Do not apply this product directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ THIS LABEL:** Read this entire label and follow all use directions and use precautions. Use only for the sites, pests, and application methods described on this label.

**RESTRICTIONS:**

1. Do not expose children, pets, or non-target animals to rodenticides.
2. Store unused product out of reach of children, pets and non-target animals.
3. Bait stations are mandatory for outdoor, above ground use. Tamper resistant bait stations must be used if children, pets, domestic animals, non-target animals or birds may access the bait.
4. Apply bait indoors in locations out of reach of children, pets, domestic animals and non-target animals, or in tamper-resistant bait stations. These bait stations must be resistant to destruction by dogs and by children under six years of age and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooved livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
5. Do not apply bait directly to food of feed crops
6. Do not broadcast bait.
7. Do not apply to burrows
8. Do not sell this product in individual containers holding less than 16 pounds of bait.
9. Fence and perimeter baiting, beyond 100 feet from a structure is prohibited.
10. Not registered for sale or use in California.

This product can only be used to control Norway rats (*Rattus norvegicus*), Roof rats (*Rattus rattus*), House mice (*Mus musculus*), Cotton rat (*Sigmodon hispidus*), Eastern harvest mouse (*Reithrodontomys humuli*), Golden mouse (*Ochrotomys nuttalli*), Meadow vole (*Microtus pennsylvanicus*), Mexican woodrat (*Neotoma Mexicana*), Polynesian rat (*Rattus exulans*), Southern plains woodrat (*Neotoma micropus*) and White throated woodrat (*Neotoma albugula*) in and within 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Examples of such structures include homes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains, aircraft), docks and port or terminal buildings and related structures around and associated with these sites.

**SELECTION OF TREATMENT AREAS:** Determine areas where rats, mice or meadow voles will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, beside burrows, in corners and concealed places, between floors and walls, or in locations where signs of rats, mice or meadow voles have been seen. Protect bait from rain and snow. Remove as much alternative food as possible.

**APPLICATION DIRECTIONS:** Each bag of **Monark Soft Bait** contains approximately 0.54 ounces (15.00 grams) of bait.

**RATS:** Apply 7 to 26 bait bags per placement for controlling rats. Space placements 15 to 30 feet apart at appropriate locations within the infested area. Maintain a continuous supply of fresh bait for at least 10 days or until signs of new feeding by rats on bait cease to appear.

**MICE & MEADOW VOLES:** Apply 2 bait bags per placement usually spaced 8 to 12 feet apart, reducing to every 4 to 6 feet at points of very high mouse or meadow vole activity. Maintain a constant supply of fresh bait for 15 days or until signs of mouse or meadow vole activity cease.

**FOLLOW-UP:** Replace contaminated or spoiled bait immediately. Wearing waterproof gloves, collect and properly dispose of dead animals and leftover bait and bait bags, as well as any bait that has become accessible to non-target animals. To discourage re-infestation, limit sources of rodent food, water, and harborage as much as possible. If re-infestation does occur, repeat treatment. For continuous sources of infestation, set up permanent bait stations and replenish them as needed.

#### **STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store only in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

**Pesticide Disposal:** Wastes resulting from the use of this product may be placed in trash or delivered to an approved waste disposal facility.

**Container Handling:** Non-refillable container. Do not reuse or refill this container. Triple rinse (or equivalent) then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or incineration.

**DISCLAIMER:** To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.



[PCO Inner container – Complete Label – package weight 1/2 LB. TO 8 LB. – Front Panel]

# Monark SB

[Soft Bait Rodenticide] [Difenacoum Soft Bait]

**KILLS RATS, MICE AND MEADOW VOLES  
KILLS WARFARIN RESISTANT NORWAY RATS**

<b>ACTIVE INGREDIENT:</b> Difenacoum (CAS #56073-07-5):.....	0.005%
ACTIVE INGREDIENT: OTHER INGREDIENTS^:.....	99.995%
^Contains Denatonium Benzoate	Total 100.000%

## KEEP OUT OF REACH OF CHILDREN CAUTION

Manufactured by:  
PelGar International Ltd.  
Unit 13 Newman Lane  
Alton, Hampshire GU34 2QR, UK  
EPA Reg. No. 87235-  
EPA Est. No. 87235-GBR-001

Imported by:  
PelGar USA  
Saylorsburg, PA  
1-610-849-1723

*NOT REGISTERED FOR SALE OR USE IN CALIFORNIA*

*[Additional statement on packaging:]*

INDIVIDUAL SALE IS PROHIBITED BY LAW

Net Weight: 8 oz (1/2 lb) to 8 lb

See Back Panel for additional precautionary labeling.

DoE: DoM: Batch #:
--------------------------

[PCO Inner container – Complete Label – package weight 1/2 LB. TO 8 LB.– Back Panel]

<b>FIRST AID</b>	
If swallowed	<ul style="list-style-type: none"> <li>•Call a poison control center or doctor immediately for treatment advice.</li> <li>•Have person sip a glass of water if able to swallow.</li> <li>•Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>•Do not give anything by mouth to an unconscious person.</li> </ul>
If on skin or clothing	<ul style="list-style-type: none"> <li>•Take off contaminated clothing.</li> <li>•Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>•Call a poison control center or doctor immediately for treatment advice.</li> </ul>
If in the eyes	<ul style="list-style-type: none"> <li>•Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>•Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>•Call a poison control center or doctor immediately for treatment advice.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor, or when going for treatment. You may also contact 1-800-222-1222 for emergency medical treatment information.</p>	
<p><b>Note to Physician</b></p>	
<p>If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K1 intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.</p>	
<p><b>Treatment for Pet Poisoning</b></p>	
<p>If animal eats bait, call veterinarian at once.</p>	
<p><b>Note to Veterinarian</b></p>	
<p>Anticoagulant Difenacoum: For animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K1. If needed, check prothrombin times every 3 days until values return to normal (up to 30 days). In severe cases, blood transfusions may be needed.</p>	

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if swallowed. Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Keep away from children, domestic animals and pets.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

All handlers (including applicators) must wear: shoes plus socks, long pants and waterproof gloves. Any person who retrieves carcasses or unused bait following application of this product must wear gloves.

**USER SAFETY REQUIREMENTS:** Follow manufacturer’s instruction for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS:**

Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This product is extremely toxic to fish, birds and other wildlife. Predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten bait. Do not apply this product directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ THIS LABEL:** Read this entire label and follow all use directions and use precautions. Use only for the sites, pests, and application methods described on this label.

**RESTRICTIONS:**

1. Do not expose children, pets, or non-target animals to rodenticides.
2. Store unused product out of reach of children, pets and non-target animals.
3. Bait stations are mandatory for outdoor, above ground use. Tamper resistant bait stations must be used if children, pets, domestic animals, non-target animals or birds may access the bait.
4. Apply bait indoors in locations out of reach of children, pets, domestic animals and non-target animals, or in tamper-resistant bait stations. These bait stations must be resistant to destruction by dogs and by children under six years of age and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooved livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
5. Do not apply bait directly to food or feed crops
6. Do not broadcast bait.
7. Do not apply to burrows
8. Do not sell this product in individual containers holding less than 16 pounds of bait.
9. Fence and perimeter baiting, beyond 100 feet from a structure is prohibited.
10. Not registered for sale or use in California.

This product can only be used to control Norway rats (*Rattus norvegicus*), Roof rats (*Rattus rattus*), House mice (*Mus musculus*), Cotton rat (*Sigmodon hispidus*), Eastern harvest mouse (*Reithrodontomys humuli*), Golden mouse (*Ochrotomys nuttalli*), Meadow vole (*Microtus pennsylvanicus*), Mexican woodrat (*Neotoma Mexicana*), Polynesian rat (*Rattus exulans*), Southern plains woodrat (*Neotoma micropus*) and White throated woodrat (*Neotoma albugula*) in and within 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Examples of such structures include homes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains, aircraft), docks and port or terminal buildings and related structures around and associated with these sites.

**SELECTION OF TREATMENT AREAS:** Determine areas where rats, mice or meadow voles will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, beside burrows, in corners and concealed places, between floors and walls, or in locations where signs of rats, mice or meadow voles have been seen. Protect bait from rain and snow. Remove as much alternative food as possible.

**APPLICATION DIRECTIONS:** Each bag of **Monark Soft Bait** contains approximately 0.54 ounces (15.00 grams) of bait.

**RATS:** Apply 7 to 26 bait bags per placement for controlling rats. Space placements 15 to 30 feet apart at appropriate locations within the infested area. Maintain a continuous supply of fresh bait for at least 10 days or until signs of new feeding by rats on bait cease to appear.

**MICE & MEADOW VOLES:** Apply 2 bait bags per placement usually spaced 8 to 12 feet apart, reducing to every 4 to 6 feet at points of very high mouse or meadow vole activity. Maintain a constant supply of fresh bait for 15 days or until signs of mouse or meadow vole activity cease.

**FOLLOW-UP:** Replace contaminated or spoiled bait immediately. Wearing waterproof gloves, collect and properly dispose of dead animals and leftover bait and bait bags, as well as any bait that has become accessible to non-target animals. To discourage re-infestation, limit sources of rodent food, water, and harborage as much as possible. If re-infestation does occur, repeat treatment. For continuous sources of infestation, set up permanent bait stations and replenish them as needed.

#### **STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store only in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

**Pesticide Disposal:** Wastes resulting from the use of this product may be placed in trash or delivered to an approved waste disposal facility.

**Container Handling:** Non-refillable container. Do not reuse or refill this container. Triple rinse (or equivalent) then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or incineration.

**DISCLAIMER:** To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

# Monark SB – Agricultural Building Use

[Soft Bait Rodenticide] [Difenacoum Soft Bait]

**For Use In and Around Agricultural Buildings Only**

## FOR INDOOR AND OUTDOOR USE

**KILLS RATS, MICE AND MEADOW VOLES  
KILLS WARFARIN RESISTANT NORWAY RATS**

ACTIVE INGREDIENT:		
Difenacoum (CAS #56073-07-5):.....	0.005%	
OTHER INGREDIENTS^:.....	<u>99.995%</u>	
	Total	100.000%

^Contains Denatonium Benzoate

## KEEP OUT OF REACH OF CHILDREN CAUTION

See Back Panel for additional precautionary labeling.

**Net Weight:** 24 lbs (4 pails X 6 lbs each) or 8 pounds up to 22 pounds  
*Statement required on label of inner packages holding less than 8 pounds of bait:*  
For packages holding 4 x 6-lb pails, do not sell these pails individually.

Manufactured by:  
PelGar International Ltd.  
Unit 13 Newman Lane  
Alton, Hampshire GU34 2QR, UK  
EPA Reg. No. 87235-  
EPA Est. No. 87235-GBR-001

Imported by:  
PelGar USA  
Saylorsburg, PA  
1-610-849-1723

**NOT REGISTERED FOR SALE OR USE IN CALIFORNIA**

DoE: DoM: Batch #:
--------------------------

[Monark SB - Agricultural Building Use Outer Container Label – Side/Back Panel]

<b>FIRST AID</b>	
If swallowed	<ul style="list-style-type: none"> <li>•Call a poison control center or doctor immediately for treatment advice.</li> <li>•Have person sip a glass of water if able to swallow.</li> <li>•Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>•Do not give anything by mouth to an unconscious person.</li> </ul>
If on skin or clothing	<ul style="list-style-type: none"> <li>•Take off contaminated clothing.</li> <li>•Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>•Call a poison control center or doctor immediately for treatment advice.</li> </ul>
If in the eyes	<ul style="list-style-type: none"> <li>•Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>•Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>•Call a poison control center or doctor immediately for treatment advice.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor, or when going for treatment. You may also contact 1-800-222-1222 for emergency medical treatment information.</p>	
<p><b>Note to Physician</b></p>	
<p>If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K1 intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.</p>	
<p><b>Treatment for Pet Poisoning</b></p>	
<p>If animal eats bait, call veterinarian at once.</p>	
<p><b>Note to Veterinarian</b></p>	
<p>Anticoagulant Difenacoum: For animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K1. If needed, check prothrombin times every 3 days until values return to normal (up to 30 days). In severe cases, blood transfusions may be needed.</p>	

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if swallowed. Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Keep away from children, domestic animals and pets.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

All handlers (including applicators) must wear: shoes plus socks, long pants and waterproof gloves. Any person who retrieves carcasses or unused bait following application of this product must wear gloves.

**USER SAFETY REQUIREMENTS:** Follow manufacturer’s instruction for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS:**

Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish, birds and other wildlife. Dogs and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten bait. Do not apply this product directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ THIS LABEL:** Read this entire label and follow all use directions and use precautions. Use only for the sites, pests, and application methods described on this label.

**RESTRICTIONS:**

1. Do not expose children, pets, or non-target animals to rodenticides.
2. Store unused product out of reach of children, pets and non-target animals.
3. Bait stations are mandatory for outdoor, above ground use. Tamper resistant bait stations must be used if children, pets, domestic animals, non-target animals or birds may access the bait.
4. Apply bait indoors in locations out of reach of children, pets, domestic animals and non-target animals, or in tamper-resistant bait stations. These bait stations must be resistant to destruction by dogs and by children under six years of age and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooved livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
5. Do not apply bait directly to food or feed crops
6. Do not broadcast bait.
7. Do not apply to burrows
8. Do not sell this product in individual containers holding less than 8 pounds of bait.
9. Fence and perimeter baiting, beyond 100 feet from a structure is prohibited.
10. Not registered for sale or use in California.

This product can only be used to control Norway rats (*Rattus norvegicus*), Roof rats (*Rattus rattus*), House mice (*Mus musculus*), Cotton rat (*Sigmodon hispidus*), Eastern harvest mouse (*Reithrodontomys humuli*), Golden mouse (*Ochrotomys nuttalli*), Meadow vole (*Microtus pennsylvanicus*), Mexican woodrat (*Neotoma Mexicana*), Polynesian rat (*Rattus exulans*), Southern plains woodrat (*Neotoma micropus*) and White throated woodrat (*Neotoma albugula*) in and within 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Examples of such structures include homes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains, aircraft), docks and port or terminal buildings and related structures around and associated with these sites.

**SELECTION OF TREATMENT AREAS:** Determine areas where rats, mice or meadow voles will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, beside burrows, in corners and concealed places, between floors and walls, or in locations where signs of rats, mice or meadow voles have been seen. Protect bait from rain and snow. Remove as much alternative food as possible.

**APPLICATION DIRECTIONS:** Each bag of **Monark Soft Bait** contains approximately 0.54 ounces (15.00 grams) of bait.

**RATS:** Apply 7 to 26 bait bags per placement for controlling rats. Space placements 15 to 30 feet apart at appropriate locations within the infested area. Maintain a continuous supply of fresh bait for at least 10 days or until signs of new feeding by rats on bait cease to appear.

**MICE & MEADOW VOLES:** Apply 2 bait bags per placement usually spaced 8 to 12 feet apart, reducing to every 4 to 6 feet at points of very high mouse or meadow vole activity. Maintain a constant supply of fresh bait for 15 days or until signs of mouse or meadow vole activity cease.

**FOLLOW-UP:** Replace contaminated or spoiled bait immediately. Wearing waterproof gloves, collect and properly dispose of dead animals and leftover bait and bait bags, as well as any bait that has become accessible to non-target animals. To discourage re-infestation, limit sources of rodent food, water, and harborage as much as possible. If re-infestation does occur, repeat treatment. For continuous sources of infestation, set up permanent bait stations and replenish them as needed.

#### **STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store only in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

**Pesticide Disposal:** Wastes resulting from the use of this product may be placed in trash or delivered to an approved waste disposal facility.

**Container Handling:** Non-refillable container. Do not reuse or refill this container. Triple rinse (or equivalent) then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or incineration.

**DISCLAIMER:** To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.



[Bait Bag Label]
<b>MONARK SB</b>
ACTIVE INGREDIENT: Difenacoum....0.005%
<b>CAUTION: KEEP OUT OF REACH OF CHILDREN</b> See outer container for complete directions for use and precautionary statements.
<b>IT IS ILLEGAL TO SELL OR DISTRIBUTE THESE BAIT BAGS INDIVIDUALLY.</b>
Net Weight: 0.54 ounces (15.00 grams)
EPA Reg. No. 87235- EPA Est. No. 87235-GBR-001

**OPTIONAL MARKETING STATEMENTS** (For Pest Control operator and Agricultural use):

Made in the UK

Made in Great Britain

Easy to secure, puncture through center of sachet

Monark SB

Monark Soft Bait(s)

Soft bait

Soft bait sachet[s]

Sachet[s]

Bait contains ingredients including (optional listing to include): wheat flour, chopped/milled wheat, lard

Contains synthetic peanut flavoring

Lard based (soft) bait

Palatable, lard based formula(tion)

Palatable to rats and mice (rodents)

Bait requires no further mixing prior to use

For use in and around homes and buildings

For Use In and Around Industrial and Commercial Buildings

For indoor and outdoor use

For indoor and outdoor rat, house mouse, and meadow vole control

Can be used indoors or outdoors

Kills rats, mice and meadow voles

Kills Norway rats, roof rats, house mice, meadow voles and warfarin resistant Norway rats

Kills house mice

Kills rats [house mice] [meadow voles]

With grains and seeds

Palatable formulation

Helps Provide (effective) control

Helps Control rats, house mice, and meadow voles

Helps provide effective control of rats, house mice, and meadow voles

Easily secured [inside station]

Flexible bait

Flex[ible] bait sachet[s]

Can be used in any bait station(s)

