



U.S. ENVIRONMENTAL PROTECTION AGENCY  
 Office of Chemical Safety  
 and Pollution Prevention  
 Registration Division (H7505P)  
 1200 Pennsylvania Avenue NW  
 Washington, D.C. 20460

EPA Reg. Number:  
 87276-6

Date of Issuance:

Term of Issuance: **Unconditional**

Name of Pesticide Product:  
**Equil Imi 2F Insecticide**

**NOTICE OF PESTICIDE:**

Registration  
 Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Mr. Michael Kellogg  
 Envincio LLC  
 c/o Pyxis Regulatory Consulting, Inc.  
 4110 135<sup>th</sup> St. NW  
 Gig Harbor, WA 98332

*Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.*

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)5. Once a pesticide is registered, however, it is not regarded as permanently acceptable. Registration does not eliminate the need for continual reassessment of pesticides. If the Agency determines that, at any time, additional data are required to maintain in effect an existing registration, the Agency will require submission of such data under FIFRA section (3)(c)(2)(B).

1. Revise the EPA Registration Number to read, EPA Reg. No. "87276-6".

Signature of Approving Official:

*Venus Eagle*

Venus Eagle  
 Product Manager 01  
 Insecticide-Rodenticide Branch  
 Registration Division (7504P)

Date:

**MAY 18 2010**

Page Two  
87276-6

2. Submit two copies of your final printed label before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitute acceptance of these conditions.

A stamped copy of the label is enclosed for your records. If you have any questions regarding this notice contact Dani Daniel at 703 305-5409.

3  
26

# Equil Imi 2 F Insecticide

FOLIAR AND SYSTEMIC INSECT CONTROL

FOR USE ON TURFGRASS, LANDSCAPE ORNAMENTALS, RESIDENTIAL FRUIT AND NUT TREES AND INTERIOR PLANTSCAPES.

PREVENTS AND CONTROLS SUBTERRANEAN TERMITES, DRYWOOD TERMITES, DAMPWOOD TERMITES, CARPENTER ANTS, AND OTHER LISTED WOOD-INFESTING INSECTS

For use by individuals/firms licensed or registered by the state to apply termiticide products. States may have more restrictive requirements regarding qualifications of persons using this product. Consult the structural pest control regulatory agency of your state prior to use of this product.

Intended for use by professional applicators to apply turf maintenance, nursery/greenhouse, and/or landscaping/ornamental products.

GROUP	4	INSECTICIDE
-------	---	-------------

Contains imidacloprid, the active ingredient used in Premise® and Merit®.

**ACTIVE INGREDIENT:**

Imidacloprid: 1-(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine .....21.4%

**OTHER INGREDIENTS:** .....78.6%

**TOTAL:** .....100.0%

Contains 2 pounds of imidacloprid per gallon.  
Shake well before using.

## KEEP OUT OF REACH OF CHILDREN CAUTION


PRECAUCION AL USUARIO: Si usted no puede leer o entender ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.  
(TO THE USER: If you cannot read or understand English, do not use this product until the label has been fully explained to you.)

FIRST AID	
<b>If Swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If Inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If on Skin or Clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing</li> <li>• Rinse skin immediately with plenty of soap and water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If in Eyes</b>	<ul style="list-style-type: none"> <li>• Hold eyelids open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
HOT LINE NUMBER	
Have the product container or label with you when calling a poison central center or doctor, or going for treatment. You may also contact 1-866-257-4118 for emergency medical treatment information.	
<b>NOTE TO PHYSICIAN:</b> No specific antidote is available. Treat patient symptomatically.	

See inside label booklet for additional PRECAUTIONARY STATEMENTS.  
For Chemical Spill, Leak, Fire, or Exposure, Call: CHEMTREC 1-800-424-9300  
For Medical Emergencies Only, Call: 866-257-4118

EPA Reg. No. 87276-

Manufactured for:  
Envincio LLC  
200 Cascade Pointe Lane, Suite 100  
Cary, NC 27513



Net Contents:

EPA Est. No.

Equil Imi 2 F Insecticide is not manufactured or distributed by Bayer Environmental Science, seller of Premise® and Merit®.

u  
-  
26

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Keep children or pets away from treated area until dry.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

**Applicators and other handlers (mixers and loaders) must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyvinylchloride (PVC) or viton.
- Shoes plus socks

In addition: All pesticide handlers must wear protective eyewear when working in a non-ventilated space or when applying termiticide by rodding or subslab injection.

Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS**

**User should:**

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.
- Remove and wash contaminated clothing before reuse.

**ENVIRONMENTAL HAZARDS**

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Apply this product only as specified on this label. Extreme care must be taken to avoid runoff. Apply only to soil or other fill substrate that will accept the solution at the specified rate. Do not treat soil that is water-saturated or frozen or in any conditions where run-off or movement from the treatment area (site) is likely to occur.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not use this product on plants being grown for sale or other commercial use or for commercial seed production or for research purposes.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**APPLICATION AS A TERMITICIDE**

Use this product in and along outside perimeter of structures and building construction to prevent and control termite infestations.

**USE INSTRUCTIONS**

For subterranean termite control, specific treatment recommendations may differ due to regulations, treatment procedures, soil types, construction practices and other factors. The purpose of chemical soil treatment for termite control is to establish a continuous chemical treated zone (horizontal and/or

vertical) between the wood and other cellulose material in the structure and the termite colonies in the soil. Follow all federal, state, and local regulations and treatment standards for protection of a structure from termites. The establishment of an aerial or above ground colony may require additional treatments to control the termites, as well as landscape modifications, and/or structural repairs to deny termites of a moisture source. Use a 0.05% to 0.1% dilution based on current recommendations. For a typical control situation, use a 0.05% dilution. Use a 0.1% dilution when a severe or persistent infestation exists.

When treating adjacent to an existing structure, the applicator must check the area to be treated, and immediately adjacent areas of the structure, for visible and accessible cracks and holes to prevent any leaks or significant exposures to persons occupying the structure. People present or residing in the structure during application must be advised to remove their pets and themselves from the structure if they see any signs of leakage. After application, the applicator is required to check for leaks. All leaks resulting in the deposition of termiticide in locations other than those prescribed on this label must be cleaned up prior to leaving the application site. Do not allow people or pets to contact contaminated areas or to reoccupy contaminated areas of the structure until the clean up is completed.

Structures that contain wells or cisterns within the foundation of the structure can only be treated using the treated backfill method described in the treatment around wells and cisterns section of this label. Consult state and local specifications for recommended distances of wells from treated area, or if such regulations do not exist, refer to Federal Housing Administration Specifications (H.U.D.) for guidance.

**MIXING:** Refer to **MIXING TABLE** for correct amount of this product to be used.

Follow this procedure for mixing the termiticide dilution:

1. Fill tank to 1/3 full.
2. If using large sprayer, start pump to begin bypass agitation and place end of treating tool in tank to allow circulation through hose.
3. Add appropriate amount of this product. Add remaining amount of water. Let pump run and allow recirculation through the hose for 2 to 3 minutes.

*[Only one of the tables below will be used based upon the size of container offered for sale]  
[For 27.5 fl oz size only]*

MIXING TABLE		
EMULSION CONCENTRATE	GALLONS WATER	AMOUNT OF PRODUCT (fl. oz.)
0.05%	100	27.5
	50	13.8
	25	6.9
	1	0.3
0.1%	50	27.5
	25	13.8
	1	0.6

*[For 2.15 gallon size only]*

MIXING TABLE		
EMULSION CONCENTRATE	GALLONS WATER	AMOUNT OF PRODUCT (fl. oz.)
0.05%	100	27.5
	50	13.8
	25	6.9
	1	0.3
0.1%	100	55.0
	50	27.5
	25	13.8
	1	0.6

**IN-LINE INJECTION:** Use the table below to mix the appropriate amount of this product for the desired

injection volume of finished emulsion.

MIXING TABLE - INJECTOR	
INJECTOR VOLUME	CONCENTRATION
0.3 fl oz/gal	0.05%
0.6 fl oz/gal	0.1%

CONVERSION KEY: 128 fl oz = 1 gal; 16 fl oz = 1 pint; 8 pints = 1 gal; 1 fl oz = 29.5 mL

**APPLICATION VOLUME**

To provide maximum control and protection against termite infestation, apply the specified volume of the finished water emulsion and active ingredient as set forth in the directions for use section of this label. If soil will not accept the labeled application volume, reduce the volume provided there is a corresponding increase in concentration so that the amount of active ingredient applied to the soil remains the same.

**Note:** Large reductions of application volume reduce the ability to obtain a continuous barrier. Variance is allowed when volume and concentration are consistent with label directed rates and a continuous barrier can still be achieved.

**PRE-CONSTRUCTION TREATMENT**

Do not apply at a lower dosage and/or concentration than specified on this label for application prior to installation of the finished grade.

Prior to each application, applicators must notify the general contractor, construction superintendent, or similar responsible party, of the intended termiticide application and intended sites of application and instruct the responsible person to notify construction workers and other individuals to leave the area to be treated during application and until the termiticide is absorbed into the soil.

**CONCRETE SLAB-ON-GROUND OR BASEMENTS:** Apply an overall treatment to the entire surface of soil or other substrate to be covered by the slab including areas to be under carports, porches, basement floor and entrance platforms. Apply at the rate of 1 gallon of solution to accurately and uniformly cover 10 square feet. If fill under slab is gravel or other coarse aggregate, apply at the rate of 1.5 gallons or sufficient volume of solution, to accurately and uniformly cover 10 square feet. In addition, apply 4 gallons of solution (see **APPLICATION VOLUME**) per 10 linear feet to provide a uniform treated zone in soil at critical areas such as along the inside of foundation walls, and around plumbing, bath traps, utility services, and other features that will penetrate the slab.

After completion of grading, make an application by trenching or trenching and rodding around the slab or foundation perimeter. Rodding may be done from the bottom of a shallow trench. When rodding, rod holes must be spaced in a manner that will allow for a continuous chemical treated zone, not to exceed 12 inches, to be deposited along the treated area. Rod holes must not extend below the footing. Apply 4 gallons of solution (see **APPLICATION VOLUME**) per 10 linear feet, per foot of depth to provide a uniform treated zone. When trenching, the trench along the outside foundation must be about 6 inches in width and 6 inches in depth. Use a low pressure spray (not to exceed 25 PSI at the treatment tool when the valve is open) to treat soil which will be placed in the trench after rodding. Mix the spray solution with soil as it is being placed in the trench. When treating voids in hollow masonry units, use 2 gallons of solution per 10 linear feet of wall. Apply solution so it will reach the footing by injecting into the lower areas of the wall, just above the floor or footing.

When treating foundations deeper than 4 feet, apply the termiticide as the backfill is being replaced, or if the construction contractor, fails to notify the applicator to permit this, treat the foundation to a minimum depth of 4 feet after the backfill has been installed. The applicator must trench and rod into the trench or trench along the foundation walls and around pillars and other foundation elements, at the rate prescribed from grade to a minimum depth of 4 feet. When the top of the footing is exposed, the applicator must treat the soil adjacent to the footing to a depth not to exceed the bottom of the footing. Do not treat structures below the footing.

Rodding in trench followed by flooding of trench and treatment of backfill may provide a better opportunity to achieve a continuous chemical treated zone than using soil rodding alone to establish a vertical termiticide treated zone.

**CRAWL SPACES:** Application must be made by trenching or trenching and rodding downward along the

inside and outside of foundation walls, around piers, interior supports in contact with the soil, plumbing, and utility services. Apply 4 gallons of solution (see **APPLICATION VOLUME**) per 10 linear feet, per foot of depth to provide a uniform treated zone. Rodding may be done from the bottom of a shallow trench to top of the footing or a minimum of 4 feet. When rodding, rod holes must be spaced in a manner that will allow for a continuous chemical treated zone to be deposited along the treated area. Rod holes must not extend below the footing. When trenching, the trench must be about 6 inches wide and 6 inches deep. Use a low pressure spray to treat soil which will be placed in the trench, mixing the spray solution with soil as it is being placed in the trench.

**HOLLOW BLOCK FOUNDATIONS OR VOIDS:** Treat hollow block foundations or voids in masonry resting on the footing to provide a continuous chemical treated zone in the voids at the footing. Apply 2 gallons of solution per 10 linear feet to the lower part of the void so that it reaches the top of the footing or soil.

Treatment of voids in block or rubble foundation walls must be closely examined. Applicators must inspect areas of possible runoff as a precaution against application leakage in the treated areas. Some areas may not be treatable or may require mechanical alteration prior to treatment.

All leaks resulting in the deposition of termiticide in locations other than those prescribed on this label must be cleaned up prior to leaving the application site (refer to **PRECAUTIONARY STATEMENTS**). Do not allow people or pets to contact or to reoccupy the contaminated areas of the structure until the clean up is completed.

**POST-CONSTRUCTION TREATMENT**

**CONCRETE SLAB-ON-GROUND:** To apply a treatment under the slab, including attached porches, carports, entrance platforms, garages and similar slab structures, it may be necessary to drill through the slab or exterior foundation. Drill holes must be spaced in a manner that will allow for application of a continuous chemical treated zone. Treat all existing cracks and cold, construction or expansion joints. Also, treat around bath traps, plumbing and utility services which penetrate the slab. Apply 4 gallons of solution (see **APPLICATION VOLUME**) per 10 linear feet per foot of depth to provide a uniform treated zone. **DO NOT MAKE TREATMENT UNTIL LOCATION OF HEAT OR AIR CONDITIONING DUCTS AND VENTS ARE KNOWN AND IDENTIFIED. USE EXTREME CAUTION TO AVOID CONTAMINATION OF DUCTS AND VENTS.** Plug and fill all drilled holes in commonly occupied areas with a suitable sealant. Plugs must be of non-cellulose material or covered by an impervious, non-cellulose material.

Apply by trenching or trenching and rodding around the outside of the foundation wall. Apply 4 gallons of solution (see **APPLICATION VOLUME**) per 10 linear feet per foot of depth to provide a uniform treated zone. When trenching, the trench along the outside foundation must be about 6 inches wide and 6 inches deep. Use a low pressure spray to treat soil as it is being placed in the trench.

Rodding can be done from the bottom of a shallow trench. When rodding, rod holes must be spaced in a manner that will allow for a continuous chemical treated zone, not to exceed 12 inches, to be deposited along the treated area. Rod hole depth must not extend below the footing.

**BATH TRAPS:** Treat exposed soil or soil covered with tar or a similar type sealant beneath and around plumbing and/or drain pipe entry areas with 3 gallons of solution per square foot. Cut and install an access door or inspection vent if not already present. After inspection and removal of any wood or cellulose debris, the soil can be treated by rodding or drenching the soil.

**CRAWL SPACES:** When there is insufficient clearance between floor joists and ground surfaces to allow applicator access, excavate, if possible, and treat according to crawl spaces (refer to **PRE-CONSTRUCTION TREATMENT**). If unable to excavate, use crawl space soil and wood treatment to prevent surface access by termites. Apply 1 gallon of solution (see **APPLICATION VOLUME**) per 10 square feet to provide a uniform chemical treated zone. Use a very coarse spray at a pressure not exceeding 25 PSI at the treatment tool when the valve is open.

Where a crawl space cannot be reached with the application wand, use extension wands or other suitable equipment to apply a coarse spray on the soil, wood and structural members contacting the soil at the above rates. Do not apply to inaccessible crawl space areas using pressures greater than 25 PSI at the treatment tool when the valve is open.

Make treatment also by drilling through the foundation wall or through the floor above and treating the soil perimeter at a rate of 1 gallon of solution per 10 square feet. Drill spacing must be at intervals not to exceed 16 inches. Many states have smaller intervals so check state regulations which may apply.

To prevent subterranean termites from constructing mudtubes between soil and crawl space wood members above, apply an overall soil treatment of this product. Remove all cellulose debris before application. Apply 1 gallon of solution (see **APPLICATION VOLUME**) per 10 square feet to provide a uniform chemical treated zone.

**SHALLOW FOUNDATIONS:** For shallow foundations, one foot or less in depth, dig a narrow trench approximately 6 inches wide and deep along the outside and inside of the foundation walls, being careful not to dig below the bottom of the footings. For foundations with exposed footings, dig a trench alongside the footing taking care not to undermine the footing. Apply 4 gallons of solution (see **APPLICATION VOLUME**) per 10 linear feet to the top of footer to provide a uniform treated zone. The dilution must be applied to the trench and mixed with the soil as it is placed in the trench.

**BASEMENTS - OUTSIDE PERIMETER:** Along the outside of the exterior walls, an application must be made by trenching or rodding within the trench. Rodding depth must be to the top of the footer, or to a minimum of 4 feet or according to state or local regulations. When rodding through a trench, dig a narrow trench about 6 inches wide and 6 inches deep. Apply 4 gallons of solution (see **APPLICATION VOLUME**) per 10 linear feet, per foot of depth to provide a uniform treated zone by rodding through the trench. Use a low pressure spray to treat soil which will be placed into the trench after rodding. Mix spray solution with the soil as it is being placed in the trench.

**BASEMENTS - INSIDE PERIMETER:** If necessary, treat by drilling along the perimeter of the interior walls. Also, make applications around Sewer pipes, floor drains, conduits, expansion joints or any cracks or holes in the basement floor. Apply 4 gallons of solution (see **APPLICATION VOLUME**) per 10 linear feet to provide a uniform treated zone.

Drill holes must be spaced in a manner that will allow for application of a continuous chemical treated zone. Plug and fill all drill holes in commonly occupied areas of the building with a suitable sealant. Plugs must be of non-cellulose material or covered by an impervious, non-cellulose material.

**HOLLOW BLOCK FOUNDATION OR VOIDS:** Treat hollow block foundations or voids in masonry resting on the footing to provide a continuous chemical treated zone in the voids at the footing. Apply 2 gallons of solution per 10 linear feet to the lower part of the void so that it reaches the top of the footing or soil. Drill spacing must be at intervals not to exceed 16 inches. Many states have smaller intervals so check state regulations which may apply.

Treatment of voids in block or rubble foundation walls must be closely examined. Applicators must inspect areas of possible runoff as a precaution against application leakage in the treated areas. Some areas may not be treatable or may require mechanical alteration prior to treatment.

All leaks resulting in the deposition of termiticide in locations other than those prescribed on this label must be cleaned up prior to leaving the application site (refer to **PRECAUTIONARY STATEMENTS**). Do not allow people or pets to contact or to reoccupy the contaminated areas of the structure until the clean up is completed.

**PLENUMS:** For plenum-type structures which use a sealed underfloor space to circulate heated and/or cooled air throughout the structure, apply the dilution at the rate of 4 gallons of solution (see **APPLICATION VOLUME**) per 10 linear feet, per foot of depth of soil to provide a uniform treated zone adjacent to both sides of foundation walls, supporting piers, plumbing and conduits. Treat soil by trenching to a depth of 6 inches or trenching and rodding (where conditions permit) or to the top of the footing. When conditions will not permit trenching or rodding, make a surface application adjacent to interior foundation walls, but the treated strip shall not exceed a width of 18 inches, horizontally, from the foundation walls, piers or pipes. The surface application will be made at a rate of 1.5 gallons of solution per 10 square feet as a very coarse spray under low pressure (not to exceed 25 PSI when measured at the treating tool when valve is on).

When treating plenums, turn off the air circulation system of the structure until application has been completed and all termiticide has been absorbed by the soil.



**TREATMENT AROUND WELLS OR CISTERNS:** Do not contaminate wells or cisterns.

**Structures With Wells/Cisterns Inside Foundations:** Structures that contain wells or cisterns within the foundation of a structure can only be treated using the following techniques:

1. Do not apply within 5 feet of any well or cistern by rodding and/or trenching or by the backfill method. Treat soil between 5 and 10 feet from the well or cistern by the backfill method only. Use the backfill method if treatment of soil adjacent to water pipes is within 3 feet of grade.
  - a. Trench and remove soil to be treated onto heavy plastic sheeting or similar material or into a wheelbarrow.
  - b. Treat the soil at the rate of 4 gallons of solution per 10 linear feet per foot of depth of the trench, or 1 gallon per 1.0 cubic feet of soil. Mix thoroughly into the soil taking care to contain the liquid and prevent runoff or spillage.
  - c. After the treated soil has absorbed the solution, replace the soil into the trench.
2. Treat infested and/or damaged wood in place using an injection technique such as described in the **CONTROL OF WOOD INFESTING PESTS** section of this label.

**Structures With Adjacent Wells/Cisterns and/or Other Water Bodies:** Applicators must inspect all structures with nearby water sources such as wells, cisterns, surface ponds, streams, and other bodies of water and evaluate, at a minimum, the treatment directions listed below prior to making an application.

1. Prior to treatment, if feasible, expose the water pipes) coming from the well to the structure, if the pipes) enter the structure within 3 feet of grade.
2. Prior to treatment applicators are advised to take precautions to limit the risk of applying the termiticide into subsurface drains that could empty into any bodies of water. These precautions include evaluating whether application of the termiticide to the top of the footer may result in contamination of the subsurface drain. Factors such as depth to the drain system and soil type and degree of compaction must be taken into account in determining the depth of treatment.
3. When appropriate (i.e., on the water side of the structure), the treated backfill technique (described above) can also be used to minimize off-site movement of termiticide.

**FOAM APPLICATIONS**

Construction practices, soil subsidence and other factors may create situations in which a continuous chemical treated zone cannot be achieved using conventional treatment alone. In situations where necessary, conventional application methods can be supplemented through use of foam generating equipment, or similar devices, to provide a continuous treated zone.

Make foam application alone or in combination with conventional application methods, provided that the labeled amount of active ingredient per unit area is used.

**Foam Application Use Directions:** Mix appropriate concentration of this product in water and add the manufacturer's specified quantity of foam agent to the product solution (see table for foaming recommendations). Apply a sufficient volume of this product foam alone or in combination with liquid solution to provide a continuous treated zone at the labeled rate for specific application sites.

NOTE: Add the manufacturer's specified quantity of foam agent to the product solution.

**MIXING TABLE - FOAM**

Product (fl oz)	GALLONS OF WATER	FOAM EXPANSION RATIO	FINISHED FOAM (0.05% ai)
6.9	1	25:1	25 gal
	2.5	10:1	
	5	5:1	
13.8	1	50:1	50 gal
	2.5	20:1	
	5	10:1	

Depending on the circumstances, use foam applications alone or in combination with liquid solution applications. Make applications behind veneers, piers, chimney bases, into rubble foundations, into block

voids or structural voids, wall voids, under slabs, stoops, porches, or to the soil in crawlspaces, and other similar voids.

Foam and liquid applications must be consistent with volume and active ingredient instructions in order to ensure proper application has been made. The volume and amount of active ingredient are essential to an effective treatment. At least 75% of the gallons of this product must be applied as a typical liquid treatment. The remaining 25% or fewer gallons are delivered to appropriate locations using a foam application.

NOTE: When foam is used solely to kill subterranean termites in above ground locations (such as feeding galleries in wooden framing, or in voids with framed walls), and whenever the target pest is other than subterranean termites (drywood termites, beetles, ants, etc.) expand dilute solutions of this product by foaming without concentrating the product solution as previously described for soil applications. Add the manufacturers' specified volume of foaming agent to produce foam of the desired expansion ratio. Use application tips and methods suitable to the site and pest.

**CONTROL OF WOOD INFESTING PESTS**

For control of above ground termites and carpenter ants in localized areas, apply a 0.05 to 0.1% solution of sufficient volume of this product foam to voids and galleries in damaged wood, and in spaces between wooden structural members and between the sill plate and foundation where wood is vulnerable. Make applications to inaccessible areas by drilling, and then injecting the suspension or foam with a suitable directional injector into the damaged wood or wall voids. Inject termite carton nests in building voids with a 0.05 to 0.1% suspension or foam. Multiple injection points to varying depths may be necessary. It is desirable to physically remove carton nest material from building voids when such nests are found. Make application, to attics, crawl spaces, unfinished basements, or man-made voids with a coarse fan spray of 0.05 to 0.1% solution or foam to control exposed worker and winged reproductive forms of termites or carpenter ants. This type of application is intended to be a supplemental treatment for control of above ground subterranean termites and carpenter ants.

Remove or prune away any shrubbery, bushes, and tree branches touching the structure. Vegetation touching the structure may offer a route of entry for ants into the structure. This may allow ants to inhabit the structure without coming in contact with the treatment. If nests are found, direct treatment of this product can be made to these nests.

Use a 0.05% to 0.1% solution to control existing infestations of or to prevent infestation by termites or carpenter ants in trees, utility poles, fencing and decking materials, landscape timbers and similar non-structural wood-to-soil contacts. If possible, locate the interior infested cavity and inject a 0.05 to 0.1% solution or sufficient volume of the foam of this product using an appropriate treatment tool with a splashback guard. Also treat these non-structural wood-to-soil contacts by applying a solution to the soil as a spot application or continuous treated zone applied as a drench or by rodding around the base of the post(s) of soil contact(s). Place rod holes approximately 3 inches away from the soil contact point(s) and spaced no more than 12 inches along the perimeter of the soil contact(s). For small poles or posts (< 6 inches in diameter), apply 1 gallon per foot of depth. For larger constructions, apply 4 gallons per 10 linear feet per foot of depth. Retreat as needed to maintain protection.

Inject termite carton nests in trees with a 0.05 to 0.1% solution or sufficient volume of foam using a pointed injection tool. Multiple injection points to varying depths may be necessary. Removal of carton material from trees is desirable but may not be necessary when foam application is used. In some instances, a perimeter application of a 0.05% to 0.1% solution applied to soil around the root flare of the tree may be necessary to prevent reinfestation by termites in the soil. For small trees (< 6 inches in diameter), apply 1 gallon of solution. For larger trees, apply 4 gallons per 10 linear feet (measured as the circumference at the root flare).

For protection of firewood or other wood products stored in contact with soil from carpenter ants and termites, treat soil prior to stacking with a 0.05 to 0.1% solution at 1 gallon per 10 square feet to prevent infestation. Make curative application to the soil around firewood or other wood products stored in contact with soil as described for non-structural wood-to-soil contacts (above).

**Drywood termites and wood-infesting beetles or borers** (such as, but not limited to, powder post beetles, anobiid or deathwatch beetles, false powder post beetles, old house borers, wharf borers, or ambrosia or bark beetles). **Galleries and structure voids** can be treated with sprays, mists, or foams of a 0.05% to 0.1%

solution of this product. Locate galleries by using visual signs (frass or pellets, blistered wood, emergence or clean out holes), the presence of live insects, mechanical sounding techniques, or listening devices (e.g., stethoscopes, acoustic emission detectors). Penetrate the gallery system by drilling holes to receive the injector tip or treatment tool. Distribute drill holes to adequately cover the gallery system. Do not drill where electrical wiring, plumbing lines, etc. are located. Apply solutions of this product as a low pressure (about 20 psi) spray or by misting or, where appropriate, by foaming. It is not necessary to treat to the point where runoff is detected from adjacent holes. Do not apply where electrical shock hazards exist. Seal drill holes after treatment. Also, **wood surfaces** can be sprayed or misted with a 0.05% to 0.1% solution or, where appropriate, use a sufficient volume of foam. For inaccessible surfaces, drill and treat the interior of structural voids. Surfaces treated may include exposed wooden surfaces in crawlspaces, basements, or attics, wooden exterior surfaces such as decks, fencing, or siding, structural voids, channels in damaged wood, in spaces between wooden members of a structure, and junctions between wood and foundations. Apply by brushing or as a coarse, low pressure (about 20 psi) spray to the wood surface; apply sufficient volume to cover the surface to the point of wetness, but avoid applying to the point of runoff. When spraying overhead in living areas, cover surfaces below the treated area with plastic sheeting or similar material, do not allow contact with treated surfaces until spray deposits have dried. Retreat as needed to maintain protection.

**Localized treatment for carpenter bees:** Apply a 0.05% to 0.1% solution as a spray or mist, or sufficient volume of foam, directly into gallery entrance holes. Following treatment, plug entrance holes with small pieces of steel wool or similar material.

**RETREATMENT**

Retreatment for subterranean termites can only be performed if there is clear evidence of reinfestation or disruption of the treated zone due to construction, excavation, or landscaping and/or evidence of the breakdown of the termiticide treated zone in the soil. Re-treat the vulnerable or re-infested areas in accordance with application techniques described in this product's labeling. The timing and type of these retreatments will vary, depending on factors such as termite pressure, soil types, soil conditions and other factors which may reduce the effectiveness of the treated zone. Make retreatment as either a spot or complete treatment.

When a structure is not known to be reinfested and the treated zone is not disturbed, but where the structure was last treated five or more years ago, perform retreatment if, in the judgment of the applicator, it is necessary to ensure adequate protection of the structure. In determining the timing of any retreatment, the applicator must consider efficacy and/or degradation data and/or site-specific conditions and previous experience that indicate a vulnerability of the structure to termite attack.

**Annual retreatment of the structure is prohibited unless there is clear evidence that reinfestation or treated zone disruption has occurred.**

When another registered termite control product/system is used as the primary treatment for prevention or control of subterranean termites and is applied to all label-specified areas, apply this product as a spot application in a secondary treatment to critical areas of the structure including plumbing and utility entry sites, bath traps, expansion joints, foundation cracks. The outside foundation wall, and areas of known or suspected activity at either a pre-construction or post-construction timing. These secondary treatments must be made applied in amounts and concentration in accordance with label directions relevant to the treatment area(s) to receive the secondary treatment.

**PERIMETER PEST CONTROL**

**For control of ants in houses and other structures,** apply a 0.05 to 0.1% solution as a general surface, spot crack and crevice or wall void application. Apply to surfaces on buildings, porches, patios and other structures, around doors and windows, eaves or attic vents. Utility entry points, soffit areas and other exterior openings (including foundation cracks or drilled holes) where the pests enter the structure or where they crawl or hide. Spray into cracks and crevices. Spray, mist or foam into voids where these ants or their nests are present. Apply the volume of spray mist or foam sufficient to cover the area, but do not allow excessive dripping or run-off to occur from vertical or overhead surfaces.

Treat soil, turf or ground cover adjacent to the structure where ants are trailing or may find food or harborage. Apply to flower, shrub or ornamental plant beds adjacent to the structure where ants may find food or forage.

To control ants tunneling in soil apply a 0.05% to 0.1% solution as a drench or soil injection at intervals to establish a continuous treated zone. Treat along the edge of walls, driveways or other hard surfaces where ants are tunneling beneath the surface.

Aerial Nests: If ant nests are located in tree hollows or non-structural wooden construction(e.g.; posts, fences, decks) treat the interior cavity and/or the nest site by injecting a 0.05%-0.01% solution as a spray mist, or sufficient volume of foam.

Apply in sufficient water to cover the foliage and soil area being treated. Maximum application is once per month to maintain control.

Do not allow residents or pets into the immediate area during the application or contact with treated areas until spray has dried. Interior applications for ant control are limited to spot, crack and crevice, or wall void applications only.

Do not use this product against native or imported fire ants, pharaoh or harvester ants.

**NOTE:** In instances of high pest pressures and quick knockdown or elimination at pest entry points is needed, make additional treatments using this product with targeted applications of a pyrethroid to doors and windows, utility entry points, and other places where these pests enter the structure. Read and follow all label directions for use of this companion product.

**RESTRICTIONS**

- After treatment, plug and fill all holes drilled in concrete slab areas of the building with a suitable sealant.
- Do not apply solution until location of heat pipes, ducts, water and sewer lines and electrical conduits are known and identified.
- Caution must be taken to avoid puncturing and injection into the structural elements.
- Do not plant for the purpose of consumption, edible plants into the treated areas of soil.
- Do not contaminate of public and private water supplies.
- Use anti-backflow equipment or an air gap on filling hoses.
- Consult State, Federal, or local authorities for information regarding the approved treatment practices for areas in close proximity to potable water supplies.

**APPLICATION ON TURFGRASS**

Use this product to control insect pests on turfgrass in home lawns, business and office complexes, shopping complexes, multi-family residential complexes, golf courses, airports, cemeteries, parks, playgrounds, and athletic fields.

This product controls soil inhabiting pests such as Northern & Southern masked chafers, *Cyclocephala borealis*, *C. immaculafa*, and/or *C. lurida*; Asiatic garden beetle, *Maladera castanea*; European chafer, *Rhizofroqus majalls*; Green June beetle, *Cotinis nitida*; May or June beetle, *Phyllophaga* spp.; Japanese beetle, *Popillia japonica*; Oriental beetle, *Anomala orientalis*; Billbugs, *Spherophorus* spp.; Annual bluegrass weevil, *Hyperodes* spp.; Black turfgrass ataenius, *Ataenius spretulus* and *Aphodius* spp. European Crane Fly *Tipula paludosa*, and mole crickets, *Scapteriscus* spp. this product can also be used for suppression of cutworms and chinch bugs.

For optimum control, make applications preceding or during the egg laying period of the target pest. The active ingredient in this product has enough residual activity so that applications can be made preceding the egg laying activity. Application timing can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. Most favorable control will be achieved when applications are made prior to egg hatch of the target pests, followed by sufficient irrigation or rainfall to move the active ingredient through the thatch.

Do not make applications when turfgrass areas are waterlogged or the soil is saturated with water. Sufficient distribution of the active ingredient cannot be achieved under these conditions. The treated turf area must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile.

**RESTRICTION:** Do not exceed a total of 1.6 pints (0.4 lb of active ingredient) per acre per year. Do not use on sod farms.

### APPLICATION EQUIPMENT FOR USE ON TURFGRASS

Apply this product in sufficient water to provide adequate distribution in the treated area. The use of accurately calibrated equipment normally used for the application of turfgrass insecticides is required. Use equipment which will produce a uniform, coarse droplet spray, using a low pressure setting to eliminate off target drift. Check calibration periodically to ensure that equipment is working properly.

Do not apply through any irrigation system.

### APPLICATIONS-TURF GRASSES

PEST	RATE	APPLICATION INSTRUCTIONS
Larvae of: Annual bluegrass weevil Asiatic garden beetle Billbug Black turfgrass ataenius Cutworms (suppression) European chafer European crane fly Green June Beetle Japanese beetle Southern Masked chafer Oriental beetle <i>Phyllophaga</i> spp. Southern masked chafer	1.25 to 1.6 pt/A or 0.46 to 0.6 fl. oz. (14 to 17 mL) per 1000 sq. ft.	For best control of grubs, billbugs, annual bluegrass weevil, and European Crane Fly, apply prior to egg hatch of the target pest. Read <b>APPLICATION EQUIPMENT</b> section of this label.
Chinchbugs (suppression) Mole crickets	1.6 pt /A or 0.6 fl. oz. (17 mL) per 1000 sq. ft.	For suppression of chinchbugs, apply before hatching of the first instar nymphs.  To control mole crickets apply before or during the peak egg hatch period.  Follow label instructions for other insecticides if tank-mixing.
Consult your local turf, state Agricultural Experiment Station, or State Extension Service Specialists for more specific information regarding timing of application.		
Irrigation or rainfall must occur within 24 hours after application to move the active ingredient through the thatch. Do not apply more than 1.6 pt (0.4 lb of active ingredient) per acre per year. Do not mow turf or lawn area until after sufficient irrigation or rainfall has occurred so that uniformity of application will not be affected.		

### APPLICATION TO ORNAMENTALS

Use this product on ornamentals in commercial and residential landscapes and interior plantscapes. This product is a systemic product and will be taken up into the plant system from root uptake. To assure optimum effectiveness, the product must be placed where the growing portion of the target plant can absorb the active ingredient. The addition of a nitrogen containing fertilizer, where applicable, into the solution may enhance the uptake of the active ingredient. Application can be made by foliar application or soil applications; including soil injection, drenches, and broadcast sprays. Foliar applications offer locally systemic activity against insect pests.

When making soil applications to plants with woody stems, systemic activity will be delayed until the active ingredient is taken up throughout the plant. In some cases, this translocation delay could take 60 days or longer. For this reason, treat prior to anticipated pest infestation to achieve optimum levels of control.

Do not exceed a total of 1.6 pints (0.4 lb of active ingredient) per acre per year.

**Ant Management Programs**

Use this product to control aphids, scale insects, mealybugs and other sucking pests on ornamentals to limit the honeydew available as a food source for ant populations. Applications of this product can then be supplemented with residual sprays, bait placements or other ant control tactics to further reduce the pest population.

**Not for use in commercial greenhouses, nurseries, sod farms, or on grass grown for seed, or on commercial fruit and nut trees.**

**APPLICATION EQUIPMENT FOR FOLIAR APPLICATIONS**

This product mixes readily with water and may be used in many types of application equipment. Mix product with the required amount of water and apply as desired dependent upon the selected use pattern.

When making foliar applications on hard to wet foliage such as holly, pine, or ivy, add a spreader/ sticker. If concentrate or mist type spray equipment is used, an equivalent amount of product must be used on the area sprayed, as would be used in a dilute application.

This product has been found to be compatible with commonly used fungicides, miticides, liquid fertilizers, and other commonly used insecticides. Check physical compatibility using the correct proportion of products in a small jar test if local experience is unavailable.

Do not apply through any irrigation system.

**APPLICATIONS FOR USE ONLY IN AND ON INDUSTRIAL AND COMMERCIAL BUILDING SITES AND RESIDENTIAL AREAS**

CROP	PEST	RATE	APPLICATION INSTRUCTIONS
Trees Shrubs Evergreens Flowers Foliage plants Groundcovers Interior plantscapes	Adelgids Aphids Japanese beetles Lace bugs Leaf beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassy-winged sharpshooter) Mealybugs Psyllids Sawfly Larvae Thrips (suppression) Whiteflies	1.5 fl. oz. (45 mL) per 100 gal of water	<b>Foliar Applications:</b> Begin applications before the onset of high pest populations and reapply as needed.
	White grub larvae (such as Japanese beetle larvae, Chafers, Phyllophaga spp. Asiatic garden beetle, Oriental beetle)	0.46 to 0.6 fl. oz. (14 to 17 mL) per 1000 sq. ft.	<b>Broadcast Applications:</b> Use enough water to mix the product and thoroughly apply to the treatment area. Do not use less than 2 gallons of water per 1000 sq ft. Irrigate after application to incorporate this product into the upper soil layer.  For additional use directions, refer to the <b>FLOWERS and GROUND COVERS</b> section of this label.

**SOIL INJECTION\* AND SOIL DRENCH APPLICATIONS IN AND ON INDUSTRIAL AND COMMERCIAL BUILDING SITES, RESIDENTIAL AREAS, AND STATE, NATIONAL AND PRIVATE WOODED AND FORESTED AREAS**

**\*No Soil Injection Applications Allowed in Nassau or Suffolk Counties of New York.**

PEST	CROP/RATE	APPLICATION INSTRUCTIONS	REMARKS
Adelgids Aphids Armored scales (suppression) Black vine weevil larvae Eucalyptus longhorned borer Flatheaded borer (including bronze birch and alder borer) Japanese beetles Lace bugs Leaf beetles (including elm and viburnum leaf beetles)	<b>TREES</b>  0.1 to 0.2 fl. oz. (3 to 6 mL) per inch of trunk diameter (D.B.H.)	<b>SOIL INJECTION:</b> <b>Grid System:</b> Space holes in a grid pattern on 2.5 foot centers, extending to the drip line of the tree. <b>Circle System:</b> Apply in holes evenly spaced in circles, (use more than one circle dependent upon the size of the tree) beneath the drip line of the tree extending in from that line. <b>Basal System:</b> Space injection holes evenly around the base of the tree trunk no more than 6 to 12 inches out from the base. <b>Soil Drench:</b> Apply uniformly as a drench around the base of the tree in not less than 10 gallons of water per 1000 square feet. Direct application to the root area. Remove plastic or any other barrier that will stop solution from reaching the root zone.	Use enough water to mix the product and inject an equal amount of solution in each hole. Use low pressure and sufficient solution for distribution of the liquid into the treatment area. Keep the treated area moist for 7 to 10 days.  Do not use less than 4 holes per tree.  <b>For Control of Specified Borers:</b> Trees with existing insect damage and stress may not recover after treatment with this product.  Do not apply to areas that are waterlogged or saturated.
Leafhoppers (including glassy-winged sharpshooter) Leafminers Mealybugs Pine tip moth larvae Psyllids Royal palm bugs Sawfy larvae Soft scales	<b>SHRUBS</b>  0.1 to 0.2 fl. oz. (3 to 6 mL) per foot of shrub height	<b>Soil Injection:</b> Apply at the labeled dosage to each plant. <b>Soil Drench:</b> Apply uniformly as a drench around the base of the tree in not less than 10 gallons of water per 1000 square feet. Direct application to the root area. Remove plastic or any other barrier that will stop solution from reaching the root zone.	Use enough water to mix the product and inject an equal amount of solution in each hole. Use low pressure and sufficient solution for distribution of the liquid into the treatment area. Keep the treated area moist for 7 to 10 days.  Do not use less than 4 holes per shrub.  Do not apply to areas that are waterlogged or saturated.
Thrips (suppression) White grub larvae Whiteflies	<b>FLOWERS AND GROUND-COVERS</b>  0.46 to 0.6 fl. oz. (14 to 17 mL) per 1000 sq ft	Apply as a broadcast treatment before or after planting, or apply after plants are established. Mix into soil. Irrigate thoroughly after application. Do not apply to areas that are waterlogged or saturated.	

**FOLIAR APPLICATIONS FOR USE ONLY IN AND ON RESIDENTIAL AREAS**

CROP	PEST	RATES	APPLICATION INSTRUCTIONS
<p><b>POME FRUITS</b>  Apple  Crabapple  Loquat  Mayhaw  Pear  Pear (oriental)  Quince</p>	<p>Aphids (except Woolly apple aphid)  Leafhoppers (including glassy-winged sharpshooter)  Leafminer  Mealybugs*  San Jose scale*</p>	<p>1.5 fl oz (45 mL) per 100 gal  or  6.0 fl oz/A<sup>1</sup></p>	<p>Apply labeled dosage as foliar spray as needed after petal-fall is complete.</p> <p>For control of rosy apple aphid, apply prior to leafrolling caused by the pest.</p> <p>For first generation leafminer control, make first application as soon as petal-fall is complete. Greatest leafminer control will result from the earliest possible application. For second and succeeding generations of leafminer, optimal control is obtained from applications made early in the adult flight against egg and early instar larvae. A second application may be required 10 days later if severe pressure continues or if generations are overlapping. A single application may result in suppression only. This product will not control late stage larvae.</p> <p>For San Jose Scale, time applications to the crawler stage. Treat each generation.</p> <p>For late season (preharvest) control of leafhopper species, apply this product while most leafhoppers are in the nymphal stage.</p> <p>For control of mealybugs, ensure good spray coverage of the trunk and scaffolding limbs or other resting sites of the mealybug.</p> <p>Do not apply more than 6.0 fluid ounces per acre in a single application. Do not make more than 5 applications per year.</p> <p>Allow 10 or more days between applications. Allow at least 7 days between last application and harvest.</p> <p>* Not for use in California for control on pears.</p>
<p><b>PECAN*</b></p>	<p>Yellow pecan aphid  Black margined aphid  Pecan leaf phylloxera  Pecan spittlebug  Pecan stem phylloxera</p>	<p>1.5 fl. oz. (45 mL) per 100 gal  or  6.0 fl. oz./A<sup>1</sup></p>	<p>Make foliar applications as pests begin to build before populations become extreme. Two applications at a 10 to 14 day interval may be required to achieve control. Scout and retreat if needed.</p> <p>Thorough uniform coverage of foliage is necessary for optimal control. Addition of an organosilicone-based spray adjuvant at a rate not to exceed the adjuvant manufacturer's specified use rate may improve coverage.</p> <p>Do not apply more than a total of 18.0 fluid ounces of this product per acre per year. Do not make more than 3 applications per year.</p> <p>Allow 10 or more days between applications.</p> <p>* Use on pecans not permitted in California unless directed by specific supplemental labeling.</p>

<sup>1</sup> The amount of this product required per acre will depend on tree size and volume of foliage present. The rate per acre is based on a standard of 400 gallons of dilute spray solution per acre for large trees.



### FOLIAR APPLICATIONS FOR USE ONLY IN AND ON INDUSTRIAL AND COMMERCIAL BUILDING SITES AND RESIDENTIAL AREAS

CROP	PEST	RATE	APPLICATION INSTRUCTIONS
GRAPES	Leafhoppers (including glassy-winged sharpshooter) Mealybugs	1.5 fl. oz. (45 mL) per 100 gal or 3.0 fl. oz/A (90 mL/A)	Apply specified dosage as a foliar spray using 200 gallons of water per acre. Do not apply more than a total of 6.0 ounces of this product per acre per year.  Allow at least 14 days between applications. Make applications up to and including day of harvest.

#### RESTRICTIONS

- Do not graze treated areas or use clippings from treated areas for feed or forage.
- Avoid runoff or puddling of irrigation water following application. Keep children and pets off treated area until dry.
- Do not apply this product to areas which are water logged or saturated, which will not allow penetration into the root zone of the plant.
- Do not apply more than 1.6 pt (0.4 lb of active ingredient) per acre per year.

Re-plant treated areas, if necessary, with any crop specified on an imidacloprid label, or with any crop for which a tolerance exists for the active ingredient.

For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12-month plant-back interval must be observed.

### GREENHOUSE & NURSERY APPLICATIONS

#### FOR FOLIAR AND SYSTEMIC INSECT CONTROL

#### FOR USE IN NON-COMMERCIAL GREENHOUSES AND NURSERIES, ORNAMENTALS, FRUIT AND NUT TREES AND VEGETABLE PLANTS

#### APPLICATION TO ORNAMENTALS AND VEGETABLE PLANTS

Use this product to control insect pests on ornamental and vegetable plants in nurseries and greenhouses. Insect protection is achieved because this product is a systemic product and the active ingredient moves upward into the plant system. Apply this product to the growing part of the plant for more absorption of the active ingredient. Add a nitrogen-containing fertilizer to the solution to aid in the uptake of the active ingredient where applicable. Apply this product as a foliar spray or by soil applications such as soil injection, drenches, chemigation and broadcast sprays.

Soil applications to plants with woody stems will require applications of this product before expected pest infestations due to the delay in the uptake of the active ingredient and the time until the product is taken up throughout the plant.

Restriction: Do not exceed a total of 1.6 pints (0.4 lb of active ingredient) per acre per year.

**Bark Media:** The length of protection after treatment with this product may be shortened if the media has 30% or more bark content.

**RESISTANCE:** Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product must conform to resistance management strategies established for the use area. Consult your Cooperative Extension Service for resistance management strategies and recommended pest management practices for your area.

#### Application Equipment for Ornamentals and Vegetable Plants

This product mixes with water and may be applied with different types of application equipment. After mixing with the correct amount of water, follow the application directions for the selected use pattern.

For applications on hard to wet foliage such as holly, pine or ivy, use a spreader/ sticker. For application by concentrate or mist type spray equipment, use the same amount of product on the area sprayed, as would be

used in a dilute application.

This product is compatible with frequently used fungicides, miticides, liquid fertilizers. Test compatibility in a small jar by using the correct proportion of products if compatibility information is not available.

**APPLICATION THROUGH IRRIGATION SYSTEMS**

Apply this product alone or as a tank mixture with other chemicals or pesticides registered for application through irrigation systems. The normal dilution ratio is 1:100 to 1:200, depending on the system. Always meter the product into the irrigation water during the first part of the irrigation cycle. Mix the product separately prior to injection. Agitation may be necessary if the mixture is allowed to stand more than 24 hours.

Remove scale, pesticide residue and other foreign matter from the tank and entire irrigation system.

Apply this product only through micro irrigation (individual spaghetti tubes), drip irrigation, overhead irrigation, ebb and flood, or hand-held or motorized calibrated irrigation equipment.

Do not apply this product through any other type of irrigation system. Crop injury or lack of effectiveness can result from non uniform distribution of treated water.

If you have any questions about calibration, contact your State Extension Service specialist, equipment manufacturers or other experts in this area.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or a person who is under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

**SAFETY DEVICES FOR IRRIGATION SYSTEMS CONNECTED TO PUBLIC WATER SUPPLIES:**

If the source of water for your irrigation system is a public water supply, follow the instructions below:

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

**SAFETY DEVICES FOR IRRIGATION SYSTEMS NOT CONNECTED TO A PUBLIC WATER SUPPLY:**

1. The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from

being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where the pesticide distribution is adversely affected.
6. Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of material that is compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

**APPLICATION TO GRASSY AREAS IN NURSERIES**

Use this product on nursery grass in areas such as under or around field or container grown plants, on roadways or other grassy areas in or around nurseries. This product is not for use on commercial sod farms.

This product controls soil inhabiting pests of grassy areas of nurseries, such as Northern and Southern masked chafers, *Cyclocephala borealis*, *C. immaculata*, and/or *C. lurida*; Asiatic garden beetle, *Maladera castanea*; European chafer, *Rhizotroqus majalis*; Green June beetle, *Cotinis nitida*; May or June beetle, *Phyllophaga* spp.; Japanese beetle, *Popillia japonica*; Oriental beetle, *Anomala orientalis*; Billbugs, *Spherophorus* spp.; Annual bluegrass weevil, *Hyperodes* spp.; Black turfgrass ataenius, *Ataenius spretulus* and *Aphodius* spp. and mole crickets, *Scapteriscus* spp. Also use this product for suppression of cutworms and chinchbugs.

For optimum control, make applications preceding or during the egg laying period of the target pest. The active ingredient in this product has enough residual activity so that applications can be made preceding the egg laying activity. Application timing can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. Most favorable control will be achieved when applications are made prior to egg hatch of the target pests. Follow application with sufficient irrigation or rainfall to move the active ingredient through the thatch.

Do not make applications when grassy areas are waterlogged or the soil is saturated with water. Sufficient distribution of the active ingredient cannot be achieved under these conditions. The treated grassy area must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile. Application cannot exceed a total of 1.6 pints (0.4 lb of active ingredient) per acre per year.

**Application Equipment for Use on Grassy Areas in Nurseries**

Apply this product in enough water to provide sufficient distribution in the treated area. Use accurately calibrated equipment typically used for the application of soil insecticides which will produce a uniform, coarse droplet spray, using a low pressure setting to eliminate off target drift. Check calibration periodically to ensure that equipment is working properly.

**APPLICATION SITES-GRASSY AREAS OF FIELD AND FOREST NURSERIES**

PEST	RATES	APPLICATION INSTRUCTIONS
Larvae of: Annual bluegrass weevil Asiatic garden beetle Billbugs Black turfgrass ataeenius <i>Phyllophaga</i> spp. Cutworms (suppression) European chafer Green June Beetle Japanese beetle Northern masked chafer Oriental beetle Southern masked chafer	19.2 to 25.6 fl. oz. per acre or 0.45 to 0.6 fl. oz. (13 to 17 mL) per 1,000 sq. ft.	For best control of grubs, billbugs and annual bluegrass weevil, make application prior to egg hatch of the target pest.  Make sure to read <b>APPLICATION EQUIPMENT</b> section of this label.  For suppression of chinchbugs, make application prior to the hatching of the first instar nymphs.  For control of mole crickets make application before or during the peak egg hatch period. When adults or large nymphs are present and actively tunneling, include applications of this product with a curative insecticide. Follow label instructions for other insecticides when tank-mixing.
Chinch bugs (suppression) Mole crickets	25.6 oz/A or (17 mL) per 1,000 sq. ft.	Consult your local turf, state Agricultural Experiment Station, or State Extension Service Specialist for more specific information regarding timing of application.  Irrigation or rainfall must occur within 24 hours after application to move the active ingredient through the thatch. Do not apply more than 1.6 pt (0.4 lb of active ingredient) per acre per year. Do not mow grass area until after adequate irrigation or rainfall has occurred so that evenness of application will not be affected.

**ORNAMENTALS-FOLIAR AND SYSTEMIC APPLICATION IN OR ON FIELD-GROWN NURSERY AND CONTAINER STOCK, GREENHOUSE ORNAMENTALS, AND ORNAMENTALS GROWN IN FLAT BENCHES OR BEDS**

PEST	CROP	RATES	APPLICATION INSTRUCTIONS
Adelgids Aphids Japanese beetles (adults) Lacebugs Leaf beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassy-winged sharpshooter) Leafminers Mealybugs Sawfly larvae Thrips (suppression) Whiteflies	Trees (including non-bearing fruit and nut), Shrubs Evergreens Flowers Ground covers Vegetable plants*	1.7 fl. oz. (50 mL) per 100 gal of water	<b>Foliar Applications:</b> Start treatments before high pest pressure is observed and reapply as needed.  For resistance management purposes, do not use a foliar application of this product following a soil application in the same crop.  * Note: For use on vegetable plants intended for resale only including: Broccoli, Chinese Broccoli, Broccoli Raab, Brussels Sprouts, Cabbage, Chinese Cabbage, Cauliflower, Collards, Eggplant, Ground Cherry, Kale, Kohlrabi, Lettuce, Mustard Greens, Pepinos, Peppers, Potatoes, Rape Greens, Sorghum, Sugarbeets, Tomatillo, and Tomato.
White grub larvae (such as Japanese beetle larvae, Chafers, <i>Phyllophaga</i> spp. Asiatic garden beetle, Oriental beetle)		0.45 to 0.6 fl. oz. (13 to 17 mL) per 1,000 sq. ft.	<b>Broadcast Applications:</b> Mix required amount of product in enough water to uniformly and exactly cover the treatment area. Do not use less than 2 gallons of water per 1000 sq. ft. For best control, irrigate carefully to integrate this product into the upper soil level.  Refer to <b>REMARKS</b> section for use directions specific for <b>FLOWERS AND GROUND COVERS</b> concerning additional use directions.

**SOIL INJECTION, SOIL DRENCH AND BROADCAST APPLICATIONS IN RESIDENTIAL NURSERIES AND GREENHOUSES**

PEST	CROP/RATES	APPLICATION INSTRUCTIONS
Adelgids Aphids Armored scales Black vine weevil larvae Eucalyptus longhorned borers Flatheaded borers (including bronze birch and alder borers) Japanese beetles (adults) Lacebugs Leaf beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassy-winged sharpshooter)	<p style="text-align: center;"><b>TREES</b></p> <p>0.1 to 0.2 fl. oz. (3 to 6 mL) per inch of trunk diameter (D.B.H.)</p>	<p><b>Soil Injections:</b>  <b>Grid System:</b> Space holes on 2.5 foot centers, in a grid pattern, extending to the drip line of the tree.  <b>Circle System:</b> Apply in holes evenly spaced in circles, (use more than one circle dependent upon the size of the tree) beneath the drip line of the tree extending in from that line.  <b>Basal System:</b> Space injection holes evenly around the base of the tree trunk no more than 6 to 12 inches out from the base.</p> <p>Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. Keep the treated area moist for 7 to 10 days. Do not use less than 4 holes per tree.</p> <p><b>No Soil Injection Application Allowed in Nassau or Suffolk Counties of New York.</b></p> <p><b>Soil Drench:</b> Uniformly apply the dosage in no less than 10 gallons of water per 1000 square feet as a drench around the base of the tree, directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone.</p> <p><b>For Control of Specified Borers:</b> Application to trees already heavily infested may not prevent the eventual loss of the trees due to existing pest damage and tree stress.</p>
Leafminers Mealybugs Pine Tip moth larvae Psyllids Royal palm bugs Sawfly larvae Soft scales Thrips (suppression) White grub larvae Whiteflies	<p style="text-align: center;"><b>SHRUBS</b></p> <p>0.1 to 0.2 fl. oz. (13 to 17 mL) per foot of shrub height</p>	<p><b>Soil Injection:</b> Apply to individual plants using dosage indicated. Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. Keep the treated area moist for 7 to 10 days. Do not use less than 4 holes per shrub.</p> <p><b>No Soil Injection Application Allowed in Nassau or Suffolk Counties of New York.</b></p> <p><b>Soil Drench:</b> Uniformly apply the dosage in no less than 10 gallons of water per 1,000 square feet as a drench around the base of the tree, directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone.</p>
	<p style="text-align: center;"><b>FLOWERS AND GROUND COVERS</b></p> <p>0.45 to 0.6 fl. oz. (13 to 17 mL) per 1,000 sq. ft.</p>	<p>Apply as a broadcast treatment and incorporate into the soil before planting or apply after plants are established. After application to established plants, irrigated thoroughly.</p>

**EBB & FLOOD APPLICATION**

Prior to treatment, to ensure accurate uptake by the plants, bring at least 10 plants up to a known field capacity and allowed to dry out for one or two days. Once dry, re-wet these plants to determine how much water on average each plant will absorb to bring it back at field capacity. Use the volume absorbed per plant (keeping pot sizes uniform) multiplied by the number of pots being treated. Add to this volume a required minimum to flood your smallest treatment area. This should minimize the return back to the storage tank. Re-use the returned volume with subsequent irrigation or nutrients on the same plants.

PEST	POT SIZE (inches)	Herbaceous species including vegetable plants* (1 or 2 plants per pot)	Woody perennials, Herbaceous species including vegetable plants* (3 or more plants per pot)	APPLICATION INTRUCTIONS
		ML per 100 Plants	ML per 100 Plants	
Adelgids				<p><sup>1</sup> <b>Fungus gnat larvae:</b> Control in the soil by drench or incorporation. This product will not control adult Fungus Gnats.</p> <p><sup>2</sup> <b>Root Mealybug:</b> To obtain control, thoroughly drench the containerized media but do not allow leaching from the bottom of the container. Use the following rate of 1.7 fl oz (50 mL) in 150 gallons of water.</p> <p><sup>3</sup> <b>Citrus Root Weevil:</b> For use on non-bearing citrus nursery stock.</p> <p><sup>4</sup> <b>Thrips:</b> For suppression on foliage only. Thrips in buds and flowers will not be suppressed.</p> <p>Foliar insect control is accomplished by the uptake of this product from a healthy root system. This allows the active ingredient to move up into the plant.</p> <p>* Note: For use on vegetable plants intended for resale only including: Broccoli, Chinese Broccoli, Broccoli Raab, Brussels Sprouts, Cabbage, Chinese Cabbage, Cauliflower, Collards, Eggplant, Ground Cherry, Kale, Kohlrabi, Lettuce, Mustard Greens, Pepinos, Peppers, Potatoes, Rape Greens, Sorghum, Sugarbeets, Tomatillo, and Tomato.</p>
Aphids				
Armored scales				
Fungus Gnats (larvae only) <sup>1</sup>				
Japanese beetles (adults)				
Lacebugs				
Leaf beetles (including elm and viburnum leaf beetles)	2	1.6 mL	2.5 mL	
Leafhoppers (including glassy-winged sharpshooter)	3	2.5 mL	1.7 mL	
Leafminers	4	3.3 mL	5 mL	
Mealybugs	5	4.2 mL	6.3 mL	
Psyllids	6	5 mL	7.7 mL	
Root mealybugs <sup>2</sup>	7	5.9 mL	9.1 mL	
Root Weevil Complex (such as Apopka Weevil, Black Vine Weevil, Citrus Weevil <sup>3</sup> )	8	6.6 mL	10 mL	
Soft scales	9	7.4 mL	11.1 mL	
Thrips (suppression) <sup>4</sup>	10	8.3 mL	12.5 mL	
Whiteflies	11	9 mL	14.3 mL	
White grub larvae (such as Japanese Beetle, Masked Chafers, European Chafer, Oriental Beetle, Asiatic Garden Beetle)	12	10 mL	16.7 mL	

### DRENCH AND IRRIGATION APPLICATIONS

For use only on greenhouse and nursery ornamentals, vegetable plants\*, and interiorscape plants using soil drenches, micro-irrigation, drip irrigation, overhead irrigation, ebb and flood irrigation, or hand-held or motorized calibrated irrigation equipment.

PEST	POT SIZE (inches)	Herbaceous species including vegetable plants* (1 or 2 plants per pot)	Woody perennials, Herbaceous species including vegetable plants* (3 or more plants per pot)	APPLICATION INTRUCTIONS
		No. pots treated with 1.7 fl. oz. (50 mL)	No. pots treated with 1.7 fl. oz. (50 mL)	
Adelgids	2	3,000	2,000	<p>Thoroughly wet most of the potting medium but do not allow runoff or leaching from the bottom of the container.</p> <p>Follow the application with moderate irrigation.</p> <p>During the next 10 days, carefully irrigate to avoid the loss of the active ingredient due to leaching.</p> <p><sup>1</sup> <b>Fungus gnat larvae:</b> Control in the soil by drench or incorporation. This product will not control adult Fungus Gnats.</p> <p><sup>2</sup> <b>Root Mealybug:</b> To obtain control, thoroughly drench the containerized media but do not allow leaching from the bottom of the container. Use the following rate of 1.7 fl oz (50 mL) in 150 gallons of water.</p> <p><sup>3</sup> <b>Citrus Root Weevil:</b> For use on non-bearing citrus nursery stock.</p> <p><sup>4</sup> <b>Thrips:</b> For suppression on foliage only. Thrips in buds and flowers will not be suppressed.</p> <p>Foliar insect control is accomplished by the uptake of this product from a healthy root system. This allows the active ingredient to move up into the plant.</p>
Aphids				
Fungus Gnats (larvae only) <sup>1</sup>				
Japanese beetles (adults)	3	2,000	1,350	
Lacebugs	4	1,500	1,000	
Leaf beetles (including elm and viburnum leaf beetles)				
Leafhoppers (including glassy-winged sharpshooter)	5	1,200	800	
Leafminers	6	1,000	650	
Mealybugs	7	850	550	
Psyllids	8	750	500	
Root mealybugs <sup>2</sup>	9	675	450	
Root Weevil Complex (such as Apopka Weevil, Black Vine Weevil, Citrus Weevil <sup>3</sup> )	10	600	400	
Soft scales	11	550	350	
Thrips (suppression) <sup>4</sup>	12	500	300	
Whiteflies	Ornamental and vegetable plants* grown in flats, benches, or beds		1.7 fl. oz. (50 mL) per 3,000 sq. ft.	<p>Mix the appropriate amount of this product in sufficient water to evenly cover the treatment area.</p> <p>Do not use less than 2 gallons of mixture per 1000 sq. ft.</p> <p>Apply as a broadcast treatment. Before planting, mix into the potting medium or apply after to established plants. Lightly irrigate after application to established plants for best control.</p> <p>Do not allow leaching or runoff for 10 days after application.</p>
White grub larvae (such as Japanese Beetle, Masked Chafers, European Chafer, Oriental Beetle, Asiatic Garden Beetle)				

\* Note: For use on vegetable plants intended for resale only including: Broccoli, Chinese Broccoli, Broccoli Raab, Brussels Sprouts, Cabbage, Chinese Cabbage, Cauliflower, Collards, Eggplant, Ground Cherry, Kale, Kohlrabi, Lettuce, Mustard Greens, Pepinos, Peppers, Potatoes, Rape Greens, Sorghum, Sugarbeets, Tomatillo, and Tomato.

**DRENCH AND IRRIGATION APPLICATIONS (cont'd)**

PEST	Containerized Plants		APPLICATION INTRUCTIONS
	Container Size	No. pots treated with 1.7 fl. oz. (50 mL)	
Adelgids Aphids Fungus Gnats (larvae only) <sup>1</sup> Japanese beetles (adults) Lacebugs Leaf beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassy-winged sharpshooter) Leafminers Mealybugs Psyllids Root mealybugs <sup>2</sup> Root Weevil Complex (such as Apopka Weevil, Black Vine Weevil, Citrus Weevil) <sup>3</sup> Soft scales Thrips (suppression) <sup>4</sup> Whiteflies White grub larvae (such as Japanese Beetle, Masked Chafers, European Chafer, Oriental Beetle, Asiatic Garden Beetle)	1 gallon 2 gallon 3 gallon 5 gallon 7 gallon 10 gallon 15 gallon 20 gallon	340 to 244 280 to 210 220 to 165 160 to 110 100 to 75 60 to 45 40 to 30 20 to 15	<p>Apply in sufficient water to wet the potting medium. For best control, make applications prior to egg hatch of the target pest. Irrigate moderately after application to move the active ingredient into the root zone.</p> <p>To avoid leaching, use 1.7 fl. oz. (50 mL) of this product in the appropriate amount of water to treat the number of pots based on the pot size as stated in the table.</p> <p>Foliar insect control is accomplished by the uptake of this product from a healthy root system. This allows the active ingredient to move up into the plant.</p> <p><sup>1</sup> <b>Fungus gnat larvae:</b> Control in the soil by drench or incorporation. This product will not control adult Fungus Gnats.</p> <p><sup>2</sup> <b>Root Mealybug:</b> To obtain control, thoroughly drench the containerized media but do not allow leaching from the bottom of the container. Use the following rate of 1.7 fl oz (50 mL) in 150 gallons of water.</p> <p><sup>3</sup> <b>Citrus Root Weevil:</b> For use on non-bearing citrus nursery stock.</p> <p><sup>4</sup> <b>Thrips:</b> For suppression on foliage only. Thrips in buds and flowers will not be suppressed.</p>
<b>Field and Forest Nurseries</b>			
White grub larvae (such as Japanese Beetle, Masked Chafers, European Chafer, Oriental Beetle, Asiatic Garden Beetle)	1.7 fl. oz. (50 mL) per 1,000 ft. of row or 3,000 sq. ft.		<p>Before application, mow the vegetation in the treatment area to a height of 3 inches or less. Mow to the lowest height possible.</p> <p>Applications must be made May through July. Treatment must be followed by rainfall or irrigation. Do not use less than 2 gallons of spray volume per 1000 square feet.</p> <p>Apply as a uniform band on either side of the row using a band width six (6) inches wider than the actual root ball diameter to be dug. Do not overlap bands in adjacent rows.</p> <p>For grub control in areas of turf, apply as a broadcast application using 1.35 to 1.7 fl oz (40 to 50 mL) per 3000 sq. ft.</p>

**RESTRICTIONS**

- Do not graze treated areas or use clippings for treated areas for feed or forage. Do not allow runoff or puddling of irrigation water following application.
- Do not apply this product to soils which are water logged or saturated, which will not allow penetration into the root zone of the plants.



- Do not allow leachate run out for the first 10 days after application, in order to retain the product and facilitate full plant uptake of the active ingredient.
- For outdoor ornamentals grown in beds or turf, applications of this product cannot exceed a total of 1.6 pt (0.4 lb of active ingredient) per acre per year.
- Food Crops: Re-plant treated areas, if necessary, with any crop specified on an imidacloprid label, or with any crop for which a tolerance exists for the active ingredient.
- For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12 month plant-back interval must be observed.
- Not for use in commercial greenhouses or nurseries.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE AND SPILL PROCEDURES:** Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spillage or leakages, soak up with an absorbent material such as sand, sawdust, earth, Fuller's earth, etc. Dispose of with chemical waste.

**PESTICIDE DISPOSAL:** Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

**CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying.

(Nonrefillable ≤ 5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(Nonrefillable > 5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Direction for Use of this Product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Envincio LLC or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Envincio LLC and Seller harmless for any claims relating to such factors.

Envincio LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Envincio LLC., and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, ENVINCIO LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall Envincio LLC or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. To the extent consistent with applicable law, **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ENVINCIO LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT,**

**NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ENVINCIO LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Envincio LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Envincio LLC.

Premise and Merit are registered trademarks of Bayer Environmental Science.

[EPA Approval Date]