

87290-13

03-04-2011

1/21



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505C)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

87290-13

Date of Issuance:

MAR 4 2011

Term of Issuance:

Conditional

Name of Pesticide Product:

Willowood Tebucon 3.6 SC

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Willowood, LLC.
c/o Pyxis Regulatory Consulting, Inc.
4110 136th St. NW
Gig Harbor, WA 98332

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

- 1) Submit and/or cite all data required for registration of your product under FIFRA sec 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
- 2) Make the following label change:
 - a) Revise the product registration number to read "EPA Registration Number "87290-13".

(continued on page 2)

Signature of Approving Official:

Mary L. Waller (21)
Fungicide Branch, Registration Division (7505P)

Date:

3/4/2011

- b) On page 2, revise the first sentence of your "Ground Water Advisory" to read "Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use".
- c) On page 2 under your "Directions For Use", revise the second sentence so that it reads "Do not apply this product in a way that will contact workers or other persons, either directly or through drift".
- d) On page 3, revise the first sentence of your "Agricultural Use Requirements" box to read "Use this product in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170".
- e) On page 3, revise your "Pesticide Disposal" statement to read "Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility".
- f) On page 15 under "Soybeans", revise the 2nd – 4th sentences to read "Use the higher rates and shorter spray intervals when disease pressure is severe. The lowest labeled rate of spray surfactant must be tank-mixed with Willowood Tebucon 3.6 SC. Apply Willowood Tebucon 3.6 SC in a minimum for 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons per acre by aircraft spray equipment".

3) Data for storage stability (830.6317) and corrosion characteristics (830.6320) must be submitted within the next 12 months. It is required that the observations be made at 0, 3, 6, 9 and 12 month intervals. Please submit the results in electronic and hard copy format.

Submit one copy of the final printed labeling before the product is released for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. If you have any further questions, please contact Tracy Keigwin of my team at (703) 305-6605 or via e-mail at keigwin.tracy@epa.gov.

Sincerely,


Mary L. Waller
 Product Manager (21)
 Fungicide Branch
 Registration Division (7505P)

Enclosure: Label stamped "Accepted, with comments"

3/21

Willowood Tebucon 3.6 SC

For control of specified diseases on various crops.

ACTIVE INGREDIENT:

Tebuconazole, alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol 38.7%

OTHER INGREDIENTS: 61.3%

TOTAL: 100.0%

Contains 3.6 pounds tebuconazole per gallon

EPA Reg. No. 87290-

EPA Est. No.

STOP - Read the label before use

Keep Out of Reach of Children

CAUTION

FIRST AID

| | |
|--------------------------------|--|
| If swallowed: | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person. |
| If on skin or clothing: | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. |
| If in eyes: | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| If inhaled: | <ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice. |

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 seven days a week, 6:30 am to 4:30 pm Pacific Time or your poison control center at 1-800-222-1222.

NOTE TO PHYSICIAN: No specific antidote. Treat symptomatically. The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.

See inside label booklet for additional Precautionary Statements and Directions for Use.

Manufactured For:

Willowood, LLC
1600 NW Garden Valley Blvd. #130
Roseburg, OR 97471

NET CONTENTS:

**ACCEPTED
with COMMENTS
In EPA Letter Dated**

3/4/2011

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.

87290-13

4
21

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory: Willowood Tebucon 3.6 SC is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water Advisory: Willowood Tebucon 3.6 SC may contaminate water through drift of spray in wind. Willowood Tebucon 3.6 SC has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of Willowood Tebucon 3.6 SC will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply Willowood Tebucon 3.6 SC in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use Willowood Tebucon 3.6 SC only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI). The REI for each crop is listed in the application directions associated with each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store above 28°F or agitate before use.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate water that cannot be used according to label instructions must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

[NONREFILLABLE CONTAINERS]

Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying.

(Nonrefillable container ≤ 5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(Nonrefillable > 5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

[REFILLABLE CONTAINERS]

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

PRODUCT INFORMATION

Read the entire Directions for Use and Conditions of Sale before using this product.

Spray Volume: Willowood Tebucon 3.6 SC may be applied in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied. If necessary, increase the spray volume per acre for complete crop coverage.

Chemigation: Apply Willowood Tebucon 3.6 SC through irrigation equipment only to crops and diseases for which the chemigation use is specified. Apply Willowood Tebucon 3.6 SC only through center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move; or drip (trickle) irrigation systems. Do not apply Willowood Tebucon 3.6 SC through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally dosed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension. Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. Pesticide may be applied continuously for the duration of the water application.

Mixing: Add labeled amount of Willowood Tebucon 3.6 SC into the spray tank while filling with water to the desired level. Operate the agitator while mixing. If other materials are added to the spray tank, the Willowood Tebucon 3.6 SC should be thoroughly dispersed prior to the addition of other materials. Do not tank mix with products containing a prohibition against tank mixing. Follow the most restrictive labeling requirements of any tank mix product.

Compatibility: To determine the compatibility of Willowood Tebucon 3.6 SC with other products, the following procedure should be followed: Pour the recommended proportions of the products into a suitable container of water, mix thoroughly and allow to stand at least five (5) minutes. If the combination remains mixed or can be re-mixed readily, the mixture is considered physically compatible. For further information, contact your local Willowood, LLC representative.

Resistance Management Statement

Willowood Tebucon 3.6 SC is a Group 3 fungicide which exhibits no known cross-resistance to other fungicide groups. However, fungal pathogens are known to develop resistance to products with the same mode of action when used repeatedly. Any fungal population may contain or develop individuals that are resistant to Willowood Tebucon 3.6 SC and other Group 3 fungicides. If Group 3 fungicides are used repeatedly in the same field or in successive years as the primary method of control for targeted diseases, the resistant isolates may eventually dominate the fungal population. Because resistance development cannot be predicted, the use of Willowood Tebucon 3.6 SC should conform to resistance management strategies established for the crop and use area. Such strategies may include rotation and/or tank mixing with products having different modes of action or limiting the total number of applications per season. Contact your local extension specialist, certified crop advisor, and/or manufacturer for fungicide resistance management and/or integrated

disease management recommendations for specific crops and resistant disease populations. Willowood, LLC encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label.

| APPLICATION DIRECTIONS | | |
|---|---|----------------------------------|
| CROP | DISEASE | RATE OF WILLOWOOD TEBUCON 3.6 SC |
| Asparagus | Rust (<i>Puccinia</i> spp.) | 4 to 6 fl. oz. per acre |
| | <p>Notes: Apply Willowood Tebucon 3.6 SC as a foliar spray to the developing ferns after harvest of spears is completed. Apply at the earliest sign of rust pustules or when weather conditions are conducive for rust development. Apply 4 to 6 fl oz of Willowood Tebucon 3.6 SC per acre (0.11 lb ai - 0.17 lb ai per acre) in alternation with another effective fungicide. Under conditions of severe rust pressure, use the higher rate. Repeat applications on a 14-day interval as necessary to maintain control of rust. Do not apply to harvestable spears. Do not apply within 100 days of harvest in California and 180 days in all other states. Do not make more than three foliar applications per season (18 fl oz/acre or 0.51 lb ai/acre).</p> | |
| <p>General Comments: Applications may be made using ground or aerial application equipment. A 50 foot spray drift buffer zone is required for all aerial applications. For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Willowood Tebucon 3.6 SC. Willowood Tebucon 3.6 SC is a sterol demethylation inhibitor (DMI) fungicide (Group 3). Alternating Willowood Tebucon 3.6 SC with other DMI fungicides may lead to resistance.</p> <p>Restricted-entry interval (REI) = 12 hours.</p> | | |

| APPLICATION DIRECTIONS | | |
|--|---|----------------------------------|
| CROP | DISEASE | RATE OF WILLOWOOD TEBUCON 3.6 SC |
| Barley | Rusts (<i>Puccinia</i> spp.) | 4 fl. oz. per acre |
| | Head blight (<i>Fusarium</i> spp.) – Suppression | |
| <p>Notes: Apply Willowood Tebucon 3.6 SC in a minimum of 10 gallons of spray solution per acre by ground or in a minimum of 5 gallons of spray solution per acre by air. A maximum of 4 fl. oz. of Willowood Tebucon 3.6 SC may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw cut after harvest may be fed or used for bedding. Grazing livestock or feeding of green forage is permitted 6 or more days after the last application of Willowood Tebucon 3.6 SC. Barley fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development.</p> <p>Application timing directions: Rusts: Apply Willowood Tebucon 3.6 SC at the earliest sign of rust pustules on foliage. Fusarium head blight: Optimal timing of Willowood Tebucon 3.6 SC for Fusarium head blight suppression is when main stem heads have fully emerged (Feekes 10.5)</p> | | |

on 50% of the plants.

General Comments: For optimum disease control, the lowest specified rate of a spray surfactant should be tank-mixed with Willowood Tebucon 3.6 SC. Willowood Tebucon 3.6 SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6 SC will be resistant to weathering. Willowood Tebucon 3.6 SC is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS

| CROP | DISEASE | RATE OF WILLOWOOD TEBUCON 3.6 SC |
|---|--|----------------------------------|
| Beans (fresh & dry except succulent shelled) | Rust (<i>Uromyces appendiculatus</i>) | 4 to 6 fl. oz. per acre |
| | <p>Notes: Apply Willowood Tebucon 3.6 SC in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 14-day intervals, or as necessary to maintain control. Beans, fresh: Willowood Tebucon 3.6 SC may be applied up to 7 days before harvest. Do not apply more than 24 fl. oz. of Willowood Tebucon 3.6 SC per acre per crop season. Beans, dry: Willowood Tebucon 3.6 SC may be applied up to 14 days before harvest. Do not apply more than 12 fl. oz. of Willowood Tebucon 3.6 SC per acre per crop season.</p> | |

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Willowood Tebucon 3.6 SC. Willowood Tebucon 3.6 SC must have two to four hours of drying time on bean foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6 SC will be resistant to weathering. Willowood Tebucon 3.6 SC is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

| APPLICATION DIRECTIONS | | |
|---|--|----------------------------------|
| CROP | DISEASE | RATE OF WILLOWOOD TEBUCON 3.6 SC |
| Corn (sweet corn, field corn, field corn grown for seed, and popcorn) | Rust (<i>Puccinia</i> spp.) Northern leaf blight (<i>Helminthosporium turcicum</i>) Southern leaf blight (<i>Helminthosporium maydis</i>) Northern leaf spot (<i>Helminthosporium carbonum</i>) Gray leaf spot (<i>Cercospora zeae-maydis</i>) | 4 to 6 fl. oz. per acre |
| | <p>Notes: Apply Willowood Tebucon 3.6 SC in a protective spray schedule or when weather conditions are favorable for disease development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. A maximum of 24 fl. oz. (1.5 pint) of Willowood Tebucon 3.6 SC may be applied per acre per crop season. Sweet corn: Willowood Tebucon 3.6 SC may be applied up to 7 days before the harvest of ears or forage, and 49 days before the harvest of fodder. Field, seed or popcorn: Willowood Tebucon 3.6 SC may be applied up to 21 days before the harvest of forage, and 36 days before the harvest of grain or fodder.</p> | |
| <p>General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Willowood Tebucon 3.6 SC. Willowood Tebucon 3.6 SC must have two to four hours of drying time on corn foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6 SC will be resistant to weathering. Willowood Tebucon 3.6 SC is a demethylation inhibitor (DMI) fungicide (Group 3).</p> | | |
| <p>Restricted-entry interval (REI) for sweet corn = 19 hours.</p> | | |
| <p>Restricted-entry interval (REI) for all corn except sweet corn = 12 hours.</p> | | |

| APPLICATION DIRECTIONS | | |
|--|---|----------------------------------|
| CROP | DISEASE | RATE OF WILLOWOOD TEBUCON 3.6 SC |
| Cotton | Southwestern cotton rust (<i>Puccinia cacabata</i>) | 6 to 8 fl. oz. per acre |
| | <p>Notes: Apply Willowood Tebucon 3.6 SC in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. Willowood Tebucon 3.6 SC may be applied up to 30 days before harvest. Do not apply more than 24 fl. oz. of Willowood Tebucon 3.6 SC per acre per crop season.</p> | |
| <p>General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Willowood Tebucon 3.6 SC. Willowood Tebucon 3.6 SC must have two to four hours of drying time on cotton foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6 SC will be resistant to weathering. Willowood Tebucon 3.6 SC is a demethylation inhibitor (DMI)</p> | | |

fungicide (Group 3).
Restricted-entry interval (REI) = 12 hours.

| APPLICATION DIRECTIONS | | |
|---|--|----------------------------------|
| CROP | DISEASE | RATE OF WILLOWOOD TEBUCON 3.6 SC |
| Cucurbit Vegetables Group Chayote Chinese waxgourd Citron melon Cucumber Gherkin Edible gourd, (includes hyotan, cucuzza, hechima and Chinese okra) <i>Momordica</i> spp. (includes balsam apple, balsam pear, bitter melon and Chinese cucumber) Muskmelon (includes cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon and snake melon) Pumpkin Summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow and zucchini) Winter squash (includes butternut squash, calabaza, hubbard squash, acorn squash and spaghetti squash) Watermelon | Powdery mildew (<i>Sphaerotheca fuliginea</i> / <i>Podosphaera xanthii</i>) (<i>Erysiphe cichoracearum</i>) | 4 to 6 fl. oz. per acre |
| | Gummy stem blight - suppression (<i>Didymella bryoniae</i>) (watermelon, squash, pumpkin, and melons only) | 8 fl. oz. per acre |
| <p>Notes: Apply the specified dosage in a protective spray schedule to foliage and fruit. Repeat applications at 10- to 14-day intervals. Willowood Tebucon 3.6 SC may be applied up to 7 days before harvest. Do not apply more than 24 fl. oz. of Willowood Tebucon 3.6 SC per acre per crop season.</p> | | |

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Willowood Tebucon 3.6 SC. Willowood Tebucon 3.6 SC must have two to four hours of drying time for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6 SC will be resistant to weathering. Willowood Tebucon 3.6 SC is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

| APPLICATION DIRECTIONS | | |
|--|--|--|
| CROP | DISEASE | RATE OF WILLOWOOD TEBUCON 3.6 SC |
| Dry bulb onion Garlic Great-headed (elephant) garlic Welch onion Shallot | White rot (<i>Sclerotium cepivorum</i>) | White rot: 20.5 fl oz per acre applied in a 4 to 6 inch band over/into each furrow. May be applied by chemigation to control white rot. |
| | Rust (<i>Puccinia allii, Puccinia porri</i>) Purple blotch (<i>Alternaria porii</i>) | 4 to 6 fl. oz. per acre |
| | <p>White rot: For the control of white rot, make one application in the furrow at the time of planting. The in-furrow application should be made at the rate of 20.5 fl. oz Willowood Tebucon 3.6 SC per acre. Apply the entire per acre rate in a 4 to 6 inch band over/into each furrow. Additional control may be obtained by including two foliar applications at 4 to 6 fl oz/acre.</p> <p>Rust: For the control of rust make foliar applications at the rate of 4 to 6 fl. oz Willowood Tebucon 3.6 SC per acre per application. Repeat at an interval of 10 to 14 days. Apply Willowood Tebucon 3.6 SC in a protective spray schedule or when weather conditions are favorable for rust development.</p> <p>Notes: Do not apply more than 32.5 fl. oz. Willowood Tebucon 3.6 SC per acre per season if an in-furrow treatment is made. If Willowood Tebucon 3.6 SC is not applied as an in-furrow treatment then do not apply more than 12 fl oz. Willowood Tebucon 3.6 SC per acre per season as a foliar spray. Do not apply within 7 days of harvest (PHI = 7 days).</p> | |
| <p>General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Willowood Tebucon 3.6 SC. Willowood Tebucon 3.6 SC must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6 SC will be resistant to weathering. Willowood Tebucon 3.6 SC is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restricted-entry interval (REI) = 12 hours.</p> | | |

| APPLICATION DIRECTIONS | | |
|--|---|----------------------------------|
| CROP | DISEASE | RATE OF WILLOWOOD TEBUCON 3.6 SC |
| Fruiting Vegetables Group Eggplant Groundcherry Pepino Pepper Tomatillo Tomato | Early blight (<i>Alternaria solani</i>) | 8 fl. oz. per acre |
| | Notes: Apply Willowood Tebucon 3.6 SC as a foliar spray using an interval of 7 days. Do not apply more than 48 fl. oz. of Willowood Tebucon 3.6 SC per acre per season. Do not apply within 7 days of harvest (PHI = 7 days). | |
| <p>General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Willowood Tebucon 3.6 SC. Willowood Tebucon 3.6 SC must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6 SC will be resistant to weathering. Willowood Tebucon 3.6 SC is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restricted-entry interval (REI) = 12 hours.</p> | | |

| APPLICATION DIRECTIONS | | |
|---|---|----------------------------------|
| CROP | DISEASE | RATE OF WILLOWOOD TEBUCON 3.6 SC |
| Grasses Grown For Seed | Rusts (<i>Puccinia</i> spp.) | 4 to 8 fl. oz. per acre |
| | Apply the specified rate of Willowood Tebucon 3.6 SC as soon as weather conditions are favorable for rust development or when first rust pustules are present. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl oz/A and shorter spray intervals. | |
| | Powdery mildew | 4 to 8 fl. oz. per acre |
| | Apply specified rate of Willowood Tebucon 3.6 SC when powdery mildew first appears on the leaves. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl oz/A and shorter spray intervals. | |
| <p>General Comments: Apply the specified rate in a minimum of 20 gallons of water per acre with ground sprayers or in a minimum of 10 gallons of water per acre with aircraft. Thorough coverage is important for optimum disease control.</p> <p>For optimum benefit, the lowest specified rate of a spray surfactant should be tank mixed with Willowood Tebucon 3.6 SC.</p> <p>A maximum of 16 fluid ounces (1 pint) may be applied per acre per crop season. Willowood Tebucon 3.6 SC may be applied up to 4 days before harvest. Chaff, screenings and straw from treated areas may be used for feed purposes; however, do not forage, cut green crop, or use seed for feed purposes. Regrowth may be grazed starting 17 days after last application.</p> <p>Restricted-entry interval (REI) = 12 hours.</p> | | |

| APPLICATION DIRECTIONS | | |
|---|--|----------------------------------|
| CROP | DISEASE | RATE OF WILLOWOOD TEBUCON 3.6 SC |
| Green onion Leek Spring onion Scallion Japanese bunching onion Green shallots Green eschalots | White rot (<i>Sclerotium cepivorum</i>) suppression only | 4 to 6 fl. oz. per acre |
| | Rust (<i>Puccinia allii</i> , <i>Puccinia porri</i>) | |
| | Purple blotch (<i>Alternaria porii</i>) | |

For the control of diseases make foliar applications using an interval of 10 to 14 days. Apply Willowood Tebucon 3.6 SC in a protective spray schedule or when weather conditions are favorable for rust development.

Notes: Do not apply more than 24 fl. oz. Willowood Tebucon 3.6 SC per acre per season. Do not apply within 7 days of harvest (PHI = 7 days).

General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Willowood Tebucon 3.6 SC. Willowood Tebucon 3.6 SC must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6 SC will be resistant to weathering. Willowood Tebucon 3.6 SC is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

| APPLICATION DIRECTIONS | | |
|---|---|----------------------------------|
| CROP | DISEASE | RATE OF WILLOWOOD TEBUCON 3.6 SC |
| Hops | Powdery mildew (<i>Sphaerotheca humuli</i> / <i>Sphaerotheca macularis</i>) | 4 to 8 fl. oz. per acre |
| <p>Notes: Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 10- to 14-day intervals. Willowood Tebucon 3.6 SC may be applied up to 14 days before harvest. Do not apply more than 32 fl. oz. of Willowood Tebucon 3.6 SC per acre per crop season. Increase the spray volume and the application rate as vine growth increases during the season.</p> | | |

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Willowood Tebucon 3.6 SC. Willowood Tebucon 3.6 SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6 SC will be resistant to weathering. Willowood Tebucon 3.6 SC is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

| APPLICATION DIRECTIONS | | |
|--|--|----------------------------------|
| CROP | DISEASE | RATE OF WILLOWOOD TEBUCON 3.6 SC |
| Leafy Brassica Greens Broccoli raab Chinese cabbage (bok choy) Collards Kale Mizuma Mustard greens Mustard spinach Rape greens | Cercospora leaf spot (<i>Cercospora brassicicola</i>) Powdery mildew (<i>Erysiphe cruciferarum</i>) Alternaria leaf spot (<i>Alternaria brassicicola</i>) | 3 to 4 fl. oz. per acre |

| | |
|---------------|--|
| Turnip greens | Notes: Do not apply more than 16 fl. oz. Willowood Tebucon 3.6 SC per acre per season. Do not apply within 7 days of harvest (PHI = 7 days). Do not apply more often than once every 10 days. |
|---------------|--|

General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Willowood Tebucon 3.6 SC. Willowood Tebucon 3.6 SC must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6 SC will be resistant to weathering. Willowood Tebucon 3.6 SC is a demethylation inhibitor (DMI) fungicide (Group 3).

Restriction: Application to turnip greens is limited to East of the Rockies.

Restricted-entry interval (REI) = 12 hours.

| APPLICATION DIRECTIONS | | |
|---|--|----------------------------------|
| CROP | DISEASE | RATE OF WILLOWOOD TEBUCON 3.6 SC |
| Garden beet roots and tops (leaves) | Cercospora leaf spot (<i>Cercospora beticola</i>) | 3 to 7.2 fl. oz. per acre |
| | Notes: Make applications on a 14 day intervals. Do not apply more than 28.8 fl. oz. Willowood Tebucon 3.6 SC per acre per season. Do not apply within 7 days of harvest (PHI = 7 days). | |
| General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Willowood Tebucon 3.6 SC. Willowood Tebucon 3.6 SC must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6 SC will be resistant to weathering. Willowood Tebucon 3.6 SC is a demethylation inhibitor (DMI) fungicide (Group 3). | | |
| Restricted-entry interval (REI) = 12 hours. | | |

| APPLICATION DIRECTIONS | | |
|--|---|----------------------------------|
| CROP | DISEASE | RATE OF WILLOWOOD TEBUCON 3.6 SC |
| Lychee | Anthracoese (<i>Colletotrichum gloesporioides</i>) | 4 to 6 fl. oz. per acre |
| | Notes: Begin first application of Willowood Tebucon 3.6 SC as panicle emerges. Spray up to 6 fl. oz. per acre every 10 days thereafter for a total of 8 sprays. Apply specified dosage in a minimum of 50 gallons of spray solution per acre by ground only. Do not apply more than 48 fl. oz. of Willowood Tebucon 3.6 SC per acre per season. Willowood Tebucon 3.6 SC can be applied up to and including the day of harvest (PHI = 0 days). | |
| General Comments: For optimum disease control, the lowest labeled rate of a non-ionic spray surfactant should be tank-mixed with Willowood Tebucon 3.6 SC. Willowood Tebucon 3.6 SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of | | |

15
21

time, Willowood Tebucon 3.6 SC will be resistant to weathering. Willowood Tebucon 3.6 SC is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 2 days.

| APPLICATION DIRECTIONS | | |
|------------------------|---|----------------------------------|
| CROP | DISEASE | RATE OF WILLOWOOD TEBUCON 3.6 SC |
| Okra | Cercospora leaf spot (<i>Cercospora</i> spp.) | 4 to 6 fl. oz. per acre |
| | <p>Notes: Apply specific dosage of Willowood Tebucon 3.6 SC in a preventative spray program. Use the highest rate when disease conditions are favorable and in areas where high disease pressure is expected. Applications may be repeated at 14-day intervals in order to maintain control of the disease. Apply specified dosage as a foliar spray in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. Applications may be made no closer than 3 days before harvest. Do not apply more than 24 fl. oz. of Willowood Tebucon 3.6 SC per acre per season.</p> | |

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Willowood Tebucon 3.6 SC. Willowood Tebucon 3.6 SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6 SC will be resistant to weathering. Willowood Tebucon 3.6 SC is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

| APPLICATION DIRECTIONS | | |
|------------------------|---|----------------------------------|
| CROP | DISEASE | RATE OF WILLOWOOD TEBUCON 3.6 SC |
| Peanut | <p>SOILBORNE: Sclerotium stem and pod rot (white mold, southern blight, southern stem rot) Rhizoctonia limb rot Rhizoctonia pod rot (Virginia and North Carolina only)</p> <p>FOLIAR: Early leaf spot Late Leaf spot Leaf rust Web blotch (<i>Phoma</i>) Pepper spot (<i>Leptoshaerulina</i>)</p> | 7.2 fl. oz. per acre |
| | <p>FOUR-APPLICATION SPRAY PROGRAM: Apply the specified rate in a preventive spray schedule. See table below for proper timing of applications. Applications of chlorothalonil should be made prior to and following applications of Willowood Tebucon 3.6 SC to discourage development of resistant strains of fungi. For optimum control of foliar diseases such as leaf rust, web blotch, and pepper spot, the lowest label specified rate of a spray surfactant should be tank-mixed with Willowood Tebucon 3.6 SC.</p> | |

LEAF SPOT ADVISORY SCHEDULE: For control of soilborne diseases in an advisory schedule, apply Willowood Tebucon 3.6 SC in the first advisory spray in July and continue Willowood Tebucon 3.6 SC applications at 14-day intervals. Applications after August 15 should be tank mixed with chlorothalonil for resistance management purposes.

GENERAL DIRECTIONS: For optimum control of the specified soilborne diseases, four consecutive applications of Willowood Tebucon 3.6 SC must be made at 14-day intervals.

A maximum of 28.8 fluid ounces of Willowood Tebucon 3.6 SC may be applied per crop season. Willowood Tebucon 3.6 SC may be applied up to 14 days before harvest. Do not feed hay or threshings or allow livestock to graze in treated areas.

Willowood Tebucon 3.6 SC is a sterol demethylation inhibitor (DMI) fungicide. Chlorothalonil may be tank mixed at the rate of 12 ounces of active ingredient with Willowood Tebucon 3.6 SC as a leaf spot resistance management strategy. A spray surfactant is not necessary when Willowood Tebucon 3.6 SC is tank mixed with chlorothalonil. Mixing or alternating Willowood Tebucon 3.6 SC with other DMI fungicides may lead to resistance.

Willowood Tebucon 3.6 SC must be carried by rainfall or irrigation into the root and pod zone for control of root and pod rots caused by *Sclerotium rolfsii* and *Rhizoctonia solani*. Drought conditions will decrease the effectiveness of Willowood Tebucon 3.6 SC against the root and pod rots.

Use Willowood Tebucon 3.6 SC in conjunction with cultural practices that are known to reduce the severity of soilborne diseases, such as proper crop rotation practices.

Restricted-entry interval (REI) = 12 hours.

Timing of Willowood Tebucon 3.6 SC Application for Optimum Control of White Mold and Rhizoctonia Limb and Pod Rot

| Spray Program | Willowood Tebucon 3.6 SC Application No. | Chlorothalonil Application No. |
|----------------|--|--------------------------------|
| 7 Applications | 3,4,5 and 6 | 1,2 and 7 |

APPLICATION DIRECTIONS

| CROP | DISEASE | RATE OF WILLOWOOD TEBUCON 3.6 SC |
|-------|---|----------------------------------|
| Pecan | Brown leaf spot (<i>Sirosporium diffusium</i>) | 4 to 8 fl. oz. per acre |
| | Downy spot (<i>Mycosphaerella caryigena</i>) | |
| | Liver spot (<i>Gnomonia caryae</i>) | |
| | Scab (<i>Cladosporium caryigenum</i>) | |
| | Vein spot (<i>Gnomonia nerviseda</i>) | |
| | Zonate leaf spot (<i>Grovesinia pyramidalis</i>) | |

17
21

Notes: Apply Willowood Tebucon 3.6 SC in a preventive spray schedule beginning at early bud break (young leaves unfolding), and continue applications at 10- to 14-day intervals through the pollination period. Willowood Tebucon 3.6 SC should be applied at 4 fl. oz. per acre in a tank-mix with the recommended rate of Super-Tin® in cover sprays. Follow label directions for the use of SuperTin. Do not add a surfactant to the spray solution when tank-mixing Willowood Tebucon 3.6 SC with SuperTin. Apply Willowood Tebucon 3.6 SC in a spray volume of 15 or more gallons per acre by air or 50 or more gallons per acre by ground. Apply 7 to 8 fl. oz. per acre of Willowood Tebucon 3.6 SC to full-size mature trees, and 4 to 6 fl. oz. per acre of Willowood Tebucon 3.6 SC to smaller trees. Apply the high rate to varieties that are highly susceptible to the indicated diseases, or when severe disease conditions exist. The lowest labeled rate of a surfactant may be added to the spray solution for optimum control of the indicated diseases. Do not apply after shucks begin to split. A maximum of 32 fl. oz. of Willowood Tebucon 3.6 SC may be applied per acre per crop season. Do not cut cover crops in treated areas for feed or allow livestock to graze treated areas.

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Willowood Tebucon 3.6 SC. Willowood Tebucon 3.6 SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6 SC will be resistant to weathering. Willowood Tebucon 3.6 SC is a demethylation inhibitor (DMI) fungicide (Group 3). It may be applied in a tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy.

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS

| CROP | DISEASE(S) | RATE OF WILLOWOOD TEBUCON 3.6 SC |
|---------|---|----------------------------------|
| Soybean | Rust (<i>Phakopsora pachyrhizi</i>) Powdery Mildew (<i>Microsphaera diffusa</i>) | 3 to 4 fl. oz. per acre |

Use Directions: Apply Willowood Tebucon 3.6 SC as a broadcast foliar spray as a preventative spray or at first visible symptoms of disease. Repeat applications on a 10- to 14-day spray interval if environmental conditions are favorable for continued disease development. Use of the higher rates and shorter spray intervals are recommended when disease pressure is severe. The lowest label recommended rate of a spray surfactant must be tank-mixed with Willowood Tebucon 3.6 SC. Willowood Tebucon 3.6 SC should be applied in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons per acre by aircraft spray equipment.

Restrictions: Applications may not be made within 21 days of harvest. Do not apply more than 3 applications per season. Do not apply more than 12 fl. oz/a per use season.

Restricted-entry interval (REI) = 12 hours.

APPLICATION DIRECTIONS

| CROP | DISEASE | RATE OF WILLOWOOD TEBUCON 3.6 SC |
|-----------|------------------------------------|----------------------------------|
| Sunflower | Rust (<i>Puccinia helianthi</i>) | 4 to 6 fl. oz. per acre |

Notes: Apply specific dosage of Willowood Tebucon 3.6 SC at the earliest sign of infection (rust pustules developing) or when weather conditions are favorable for rust development. Apply higher rate to highly susceptible varieties and/or under severe disease conditions. Application may be repeated at 14 days if necessary to maintain control of the disease. Apply specified dosage in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. Do not

apply more than 16 fl. oz. of Willowood Tebucon 3.6 SC per acre per season or within 50 days of harvest.

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Willowood Tebucon 3.6 SC. Contact your state Extension Service or Repair representative for a list of approved surfactants. Willowood Tebucon 3.6 SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6 SC will be resistant to weathering. WILLOWOOD TEBUCONOZOLE 3.6 SC is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

| APPLICATION DIRECTIONS | | |
|--|---|----------------------------------|
| CROP | DISEASE | RATE OF WILLOWOOD TEBUCON 3.6 SC |
| Turnip (Application is limited to East of the Rockies) | Cecospora leaf spot (<i>Cercospora brassicicola</i>) | 4 to 7.2 fl. oz. per acre |
| | <p>Notes: Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 12- to 14-day intervals. Willowood Tebucon 3.6 SC may be applied up to 7 days before harvest. Do not apply more than 28.8 fl. oz. of Willowood Tebucon 3.6 SC per acre per crop season.</p> | |

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Willowood Tebucon 3.6 SC. Willowood Tebucon 3.6 SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6 SC will be resistant to weathering. Willowood Tebucon 3.6 SC is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

| APPLICATION DIRECTIONS | | |
|--|--|----------------------------------|
| CROP | DISEASE | RATE OF WILLOWOOD TEBUCON 3.6 SC |
| Wheat | Rusts leaf, stem, and stripe (<i>Puccinia</i> spp.) | 4 fl. oz. per acre |
| | Head blight or scab (<i>Fusarium</i> spp.) - Suppression | |
| <p>Notes: Wheat fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. A maximum of 4 fl. oz. of Willowood Tebucon 3.6 SC may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw may be fed or used for bedding. Do not allow livestock to graze or feed green forage to livestock prior to 6 days after treatment with Willowood Tebucon 3.6 SC. Apply Willowood Tebucon 3.6 SC F in a minimum of 10 gallons of spray solution per acre by ground, or in a minimum of 5 gallons of spray solution per acre by air.</p> | | |

Application timing directions:
 Rusts: Apply Willowood Tebucon 3.6 SC at the earliest sign of rust pustules on foliage.
 Fusarium head blight: Optimal timing of Willowood Tebucon 3.6 SC for Fusarium head blight suppression is the beginning of flowering on main stem heads (Feekes 10.51).

General Comments: For optimum disease control, the lowest specified rate of a spray surfactant should be tank-mixed with Willowood Tebucon 3.6 SC. Willowood Tebucon 3.6 SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, WILLOWOOD TEBUCONAZOLE 3.6 SC will be resistant to weathering. Willowood Tebucon 3.6 SC is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

SEED TREATMENT - Corn (Sweet Corn, Field Corn, Field Corn Grown For Seed, and Popcorn)
 For control of soilborne and seedborne Fusarium and soilborne and seedborne head smut.

SEED LABELING: To meet U.S. Federal Seed Act requirements, all seed treated with Willowood Tebucon 3.6 SC must be labeled:

TREATED SEED. DO NOT USE FOR FOOD, FEED OR OIL PURPOSES. Treated with Tebuconazole.

USE PRECAUTION: When using formulations that do not contain dye, to comply with 40 CFR 153.155, all seed treated with an economic poison must be colored to distinguish and prevent subsequent inadvertent use as a food for man or feed for animals.

| DISEASE | RATE Fl Oz/CWT | DIRECTIONS FOR USE |
|---|----------------|---|
| Soilborne and Seedborne Fusarium | 0.071 | Apply as a seed treatment using standard slurry or mist-type seed treatment equipment. Uniform application of seed is necessary to ensure seed safety and best disease protection. Seed should be sound and well cured prior to treatment. Product should be diluted with sufficient water to ensure complete seed coverage. Consult a seed treatment specialist regarding slurry rates recommended for the crop to be treated with Willowood Tebucon 3.6 SC. The length of control will vary depending on the rate used. |
| Soilborne and Seedborne Head smut <i>(Sphacelotheca reilana)</i> | 0.27 – 0.54 | |

29
21

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.

Apply only during alternate years in fields adjacent to aquatic areas listed above.

Do not apply by ground or air within 100 feet of aquatic areas listed above.

Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip.

Spray Drift Management: For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter.

Use the largest droplet size consistent with pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure. Apply in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment.

Spray should be released at the lowest possible height consistent with good pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

Make aerial or ground applications when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.

Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

ROTATIONAL CROPS

Treated areas may be replanted with any crop specified on this label as soon as practical after last application. Any crop not specified on this label may be planted into treated areas 120 days after last application.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Willowood, LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Willowood, LLC and Seller harmless for any claims relating to such factors.

Willowood, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Willowood, LLC, and Buyer and User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH

21
21

APPLICABLE LAW, WILLOWOOD, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Willowood, LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF WILLOWOOD, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF WILLOWOOD, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Willowood, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Willowood, LLC.

[EPA approval date]