

87290-13

06/20/2012

1/26



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

Willowood LLC
% Michael Kellogg
Pyxis Regulatory Consulting Inc
4110 136th St NW
Gig Harbor WA 98332

JUN 20 2012

Product Name Willowood Tebucon 3 6SC
EPA Reg No 87290 13
Subject Your notification dated May 3 2012 revised label
OPP Decision Number 465616

Dear Mr Kellogg

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98 10 The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98 10 and finds that the action(s) requested fall within the scope of PRN 98 10

The label submitted with the application has been stamped Notification and will be placed in our records

If you have any questions please contact Robert Westin by phone at (703) 305 5721 or via email at westin.robert@epa.gov

Sincerely

A handwritten signature in cursive script that reads "Mary L. Waller".

Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7504P)



United States
Environmental Protection Agency
 Washington DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide Section I

1 Company/Product Number 87290 13	2 EPA Product Manager M Waller	3 Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4 Company/Product (Name) Willowood LLC / Willowood Tebucon 3 6SC	PM# 21	
5 Name and Address of Applicant (Include ZIP Code) Willowood LLC c/o Pyxis Regulatory Consulting Inc 4110 136th St NW Gig Harbor WA 98332 <input type="checkbox"/> Check if this is a new address	6 Expedited Review In accordance with FIFRA Section 3(c)(3) (b)(i) my product is similar or identical in composition and labeling to EPA Reg No _____ Product Name _____	

Section II

<input type="checkbox"/> Amendment Explain below	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> Me Too Application
<input checked="" type="checkbox"/> Notification Explain below	<input type="checkbox"/> Other Explain below

Explanation Use additional page(s) if necessary (For section I and Section II)
 Notification of minor label revisions per PR Notice 98 10 to facilitate registration in the state of California This notification is consistent with the provisions of PR Notice 98 10 and EPA regulations at 40 CFR 152 46 and no other changes have been made to the labeling or the confidential statement of formula of this product I understand that it is a violation of 18 U S C Sec 1001 to willfully make any false statement to EPA I further understand that if this notification is not consistent with the terms of PR Notice 98 10 and 40 CFR 152 46 this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA

Section III

1 Material This Product Will Be Packaged In				2 Type of Container	
Child Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	If Yes Unit Packaging wgt No per container		<input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If Yes Package wgt	No per container		
3 Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4 Size(s) Retail Container 2.5 110 265 gallons		5 Location of Label Directions <input checked="" type="checkbox"/> O L b l <input type="checkbox"/> O L b l g m p y g p d t	
6 Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph Paper glued Stenciled				<input type="checkbox"/> Other _____	

Section IV

1 Contact Point (Complete items directly below for identification of individual to be contacted if necessary to process this application)		
Name Michael Kellogg	Title Agent	Telephone No (Include Area Code) (253) 853 7369
Certification I certify that the statements I have made on this form and all attachments thereto are true accurate and complete I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law		6 Date Application Received (Stamped)
2 Signature 	3 Title Agent	
4 Typed Name Michael Kellogg	5 Date 5/3/12	

PYXIS REGULATORY CONSULTING, INC

5/31 3/26

4110 136th St NW
Gig Harbor WA 98332

Phone 253 853 7369
Fax 253 853 5516
www.PyxisRC.com

May 3 2012

COURIER DELIVERY

Mary Waller (PM 21)
Document Processing Desk (**NOTIF**)
Office of Pesticide Programs (7504P)
U S Environmental Protection Agency
Room S 4900 One Potomac Yard
2777 S Crystal Drive
Arlington VA 22202

RE Willowood LLC – Willowood Tebucon 3 6SC (EPA Reg No 87290 13)
Notification of Minor Label Revisions per PRN 98 10 to Facilitate Registration in the State of California

Dear Ms Waller


On behalf of Willowood LLC please find the enclosed notification of minor label revisions to the Willowood Tebucon 3 6SC (EPA Reg No 87290 13) labeling to facilitate registration in the State of California The following changes have been made to the Willowood Tebucon 3 6SC labeling

- Added optional referral statement to the label front panel directing the user where to find First Aid Precautionary Statements Directions for Use and Storage and Disposal instructions when booklet labels are used
- Off labeled Fruiting Vegetables Group in California to facilitate California registration

In support of this notification submission we submit the following documents

- 1 Completed Application for Registration (EPA Form 8570 1)
- 2 One (1) copy of the Willowood Tebucon 3 6SC labeling with changes tracked
- 3 One (1) copy of the Willowood Tebucon 3 6SC labeling with changes incorporated
- 4 Certification with Respect to Label Integrity
- 5 One (1) copy of the Willowood Tebucon 3 6SC labeling on CD
- 6 Letter of Authorization

Please feel free to contact me by phone (253) 853 7369 or by email at Mike@PyxisRC.com if you have any questions or need any additional information

Sincerely


Michael Kellogg

Enclosures

cc B Heinze Willowood LLC

4/26

Willowood Tebucon 3.6SC

For control of specified diseases on various crops, golf course turf, field, nursery and container ornamentals and commercial and residential landscapes

ACTIVE INGREDIENT

Tebuconazole alpha [2 (4 chlorophenyl)ethyl] alpha (1 1 dimethylethyl) 1H 1 2 4 triazole 1 ethanol 38 7%

OTHER INGREDIENTS

61 3%

TOTAL

100 0%

Contains 3 6 pounds tebuconazole per gallon

EPA Reg No 87290 13

EPA Est No

STOP - Read the label before use

Keep Out of Reach of Children

CAUTION

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice • Have person sip a glass of water if able to swallow • Do not induce vomiting unless told to do so by a poison control center or doctor • Do not give anything by mouth to an unconscious person
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15 20 minutes • Call a poison control center or doctor for treatment advice
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 20 minutes • Remove contact lenses if present after the first 5 minutes then continue rinsing eye • Call a poison control center or doctor for treatment advice
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air • If person is not breathing call 911 or an ambulance then give artificial respiration preferably by mouth to mouth if possible • Call a poison control center or doctor for treatment advice
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment For emergency information concerning this product call the National Pesticides Information Center (NPIC) at 1 800 858 7378 seven days a week 6 30 am to 4 30 pm Pacific Time or your poison control center at 1 800 222 1222	
NOTE TO PHYSICIAN No specific antidote Treat symptomatically The compound does not cause any definite symptoms that would be diagnostic Contact with the eyes may cause irritation	

See inside label booklet for additional Precautionary Statements and Directions for Use

[Optional referral statement See inside label booklet for First Aid, Precautionary Statements and Directions for Use including Storage and Disposal instructions]

Manufactured For

Willowood LLC
1600 NW Garden Valley Blvd #120
Roseburg OR 97471

NOTIFICATION

JUN 20 2012

NET CONTENTS

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed inhaled or absorbed through skin Avoid contact with skin eyes or clothing Avoid breathing vapor or spray mist

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are listed below If you want more options follow the instructions for category C on an EPA chemical resistance category selection chart

Applicators and other handlers must wear

- Long sleeved shirt and long pants
- Chemical resistant gloves such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE If no such instructions for washables exist use detergent and hot water Keep and wash PPE separately from other laundry

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)] the handler PPE requirements may be reduced or modified as specified in the WPS

USER SAFETY RECOMMENDATIONS

Users should

- Wash hands before eating drinking chewing gum using tobacco or using the toilet
- Remove clothing immediately if pesticide gets inside Then wash thoroughly and put on clean clothing

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals fish and aquatic invertebrates Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark Runoff may be hazardous to aquatic organisms in neighboring areas Do not contaminate water when disposing of equipment washwater or rinsate

Ground Water Advisory Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use Use of this chemical in areas where soils are permeable particularly where the water table is shallow may result in ground water contamination

Surface Water Advisory Willowood Tebucon 3.6SC may contaminate water through drift of spray in wind Willowood Tebucon 3.6SC has a high potential for runoff for several months or more after application Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product A level well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds streams and springs will reduce the potential for contamination of water from rainfall runoff Runoff of Willowood Tebucon 3.6SC will be reduced by avoiding applications when rainfall is forecasted within 48 hours

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling Do not apply this product in a way that will contact workers or other persons either directly or through drift Only protected handlers may be in the area during application For any requirements specific to your State or Tribe consult the agency responsible for pesticide regulation

AGRICULTURAL USE REQUIREMENTS

Use this product in accordance with its labeling and with the Worker Protection Standard 40 CFR part 170 This Standard contains requirements for the protection of agricultural workers on farms forests nurseries and greenhouses and handlers of agricultural pesticides It contains requirements for training decontamination notification and emergency assistance It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) The REI for each crop is listed in the application directions associated with each crop

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants soil or water is

- Coveralls
- Chemical resistant gloves such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

NON AGRICULTURAL USE REQUIREMENTS

The requirements in this box only apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170) The WPS applies when this product is used to produce agricultural plants on farms forests nurseries or greenhouses

Golf Course Turf and Landscape Uses Keep children and pets out of treated areas until sprays have dried

STORAGE AND DISPOSAL

Do not contaminate water food or feed by storage or disposal

PESTICIDE STORAGE Store above 28 F or agitate before use

PESTICIDE DISPOSAL Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility

CONTAINER HANDLING

[NONREFILLABLE CONTAINERS]

Nonrefillable container Do not reuse or refill this container Triple rinse container (or equivalent) promptly after emptying

(Nonrefillable container ≤ 5 gallons) Triple rinse as follows Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip Fill the container 1/4 full with water and recap Shake for 10 seconds Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal Drain for 10 seconds after the flow begins to drip Repeat this procedure two more times Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or if allowed by state and local authorities by burning If burned stay out of smoke

(Nonrefillable > 5 gallons) Triple rinse as follows Empty the remaining contents into application equipment or a mix tank Fill the container 1/4 full with water Replace and tighten closures Tip container on its side and roll it back and forth ensuring at least one complete revolution for 30 seconds Stand the container on its end and tip it back and forth several times Turn the container over onto its other end and tip it back and forth several times Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal Repeat this procedure two more times Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or if allowed by state and local authorities by burning If burned stay out of smoke

[REFILLABLE CONTAINERS]

Refillable container Refill this container with pesticide only Do not reuse this container for any other purpose Cleaning the container before final disposal is the responsibility of the person disposing of the container Cleaning before refilling

is the responsibility of the refiller
 To clean the container before final disposal empty the remaining contents from this container into application equipment or a mix tank Fill the container about 10 percent full with water Agitate vigorously or recirculate water with the pump for 2 minutes Pour or pump rinsate into application equipment or rinsate collection system Repeat this rinsing procedure two more times Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or if allowed by state and local authorities by burning If burned stay out of smoke

PRODUCT INFORMATION

Read the entire Directions for Use and Conditions of Sale before using this product

Spray Volume Willowood Tebucon 3 6SC may be applied in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment Check equipment calibration frequently Complete coverage and uniform application are essential for the most effective results especially when lower spray volumes are applied If necessary increase the spray volume per acre for complete crop coverage

Chemigation Apply Willowood Tebucon 3 6SC through irrigation equipment only to crops and diseases for which the chemigation use is specified Apply Willowood Tebucon 3 6SC only through center pivot lateral move end tow side (wheel) roll traveler big gun solid set or hand move or drip (trickle) irrigation systems Do not apply Willowood Tebucon 3 6SC through any other type of irrigation system Crop injury lack of effectiveness or illegal pesticide residues in the crop can result from non uniform distribution of treated water If you have questions about calibration you should contact State Extension Service specialists equipment manufacturers or other experts Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise

The system must contain a functional check valve vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow The pesticide injection pipeline must contain a functional automatic quick closing check valve to prevent the flow of fluid back toward the injection pump The pesticide injection pipeline must also contain a functional normally closed solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected Systems must use a metering pump such as a positive displacement injection pump (e g diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock Do not apply when wind speed favors drift beyond the area intended for treatment

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water Pesticide may be applied continuously for the duration of the water application

Mixing Add labeled amount of Willowood Tebucon 3 6SC into the spray tank while filling with water to the desired level Operate the agitator while mixing If other materials are added to the spray tank the Willowood Tebucon 3 6SC should be thoroughly dispersed prior to the addition of other materials Do not tank mix with products containing a prohibition against tank mixing Follow the most restrictive labeling requirements of any tank mix product

Compatibility To determine the compatibility of Willowood Tebucon 3 6SC with other products the following procedure should be followed Pour the recommended proportions of the products into a suitable container of water mix thoroughly and allow to stand at least five (5) minutes If the combination remains mixed or can be re mixed readily the mixture is considered physically compatible For further information contact your local Willowood LLC representative

Resistance Management Statement

Willowood Tebucon 3 6SC is a Group 3 fungicide which exhibits no known cross resistance to other fungicide groups

However fungal pathogens are known to develop resistance to products with the same mode of action when used repeatedly. Any fungal population may contain or develop individuals that are resistant to Willowood Tebucon 3 6SC and other Group 3 fungicides. If Group 3 fungicides are used repeatedly in the same field or in successive years as the primary method of control for targeted diseases, the resistant isolates may eventually dominate the fungal population. Because resistance development cannot be predicted, the use of Willowood Tebucon 3 6SC should conform to resistance management strategies established for the crop and use area. Such strategies may include rotation and/or tank mixing with products having different modes of action or limiting the total number of applications per season. Contact your local extension specialist, certified crop advisor, and/or manufacturer for fungicide resistance management and/or integrated disease management recommendations for specific crops and resistant disease populations. Willowood LLC encourages responsible resistance management to ensure effective long term control of the fungal diseases on this label.

DISEASE CONTROL IN CROPS

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3 6SC
Asparagus	Rust (<i>Puccinia</i> spp)	4 to 6 fl oz per acre
	<p>Notes: Apply Willowood Tebucon 3 6SC as a foliar spray to the developing ferns after harvest of spears is completed. Apply at the earliest sign of rust pustules or when weather conditions are conducive for rust development. Apply 4 to 6 fl oz of Willowood Tebucon 3 6SC per acre (0.11 lb ai - 0.17 lb ai per acre) in alternation with another effective fungicide. Under conditions of severe rust pressure, use the higher rate. Repeat applications on a 14 day interval as necessary to maintain control of rust. Do not apply to harvestable spears. Do not apply within 100 days of harvest in California and 180 days in all other states. Do not make more than three foliar applications per season (18 fl oz/acre or 0.51 lb ai/acre).</p>	
<p>General Comments: Applications may be made using ground or aerial application equipment. A 50 foot spray drift buffer zone is required for all aerial applications. For optimum disease control, the lowest labeled rate of a spray surfactant should be tank mixed with Willowood Tebucon 3 6SC. Willowood Tebucon 3 6SC is a sterol demethylation inhibitor (DMI) fungicide (Group 3). Alternating Willowood Tebucon 3 6SC with other DMI fungicides may lead to resistance.</p>		
<p>Restricted entry interval (REI) = 12 hours</p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3 6SC
Barley	Rusts (<i>Puccinia</i> spp)	4 fl oz per acre
	Head blight (<i>Fusarium</i> spp) – Suppression	
<p>Notes: Apply Willowood Tebucon 3 6SC in a minimum of 10 gallons of spray solution per acre by ground or in a minimum of 5 gallons of spray solution per acre by air. A maximum of 4 fl oz of Willowood Tebucon 3 6SC may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw cut after harvest may be fed or</p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3 6SC
	<p>used for bedding Grazing livestock or feeding of green forage is permitted 6 or more days after the last application of Willowood Tebucon 3 6SC Barley fields should be observed closely for early disease symptoms particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development</p> <p>Application timing directions Rusts Apply Willowood Tebucon 3 6SC at the earliest sign of rust pustules on foliage Fusarium head blight Optimal timing of Willowood Tebucon 3 6SC for Fusarium head blight suppression is when main stem heads have fully emerged (Feekes 10 5) on 50% of the plants</p>	
<p>General Comments For optimum disease control the lowest specified rate of a spray surfactant should be tank mixed with Willowood Tebucon 3 6SC Willowood Tebucon 3 6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs After this period of time Willowood Tebucon 3 6SC will be resistant to weathering Willowood Tebucon 3 6SC is a demethylation inhibitor (DMI) fungicide (Group 3)</p> <p>Restricted entry interval (REI) = 12 hours</p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3 6SC
Beans (fresh & dry except succulent shelled)	Rust (<i>Uromyces appendiculatus</i>)	4 to 6 fl oz per acre
	<p>Notes Apply Willowood Tebucon 3 6SC in a protective spray schedule or when weather conditions are favorable for rust development Repeat applications at 14 day intervals or as necessary to maintain control Beans fresh Willowood Tebucon 3 6SC may be applied up to 7 days before harvest Do not apply more than 24 fl oz of Willowood Tebucon 3 6SC per acre per crop season Beans dry Willowood Tebucon 3 6SC may be applied up to 14 days before harvest Do not apply more than 12 fl oz of Willowood Tebucon 3 6SC per acre per crop season</p>	
<p>General Comments For optimum disease control the lowest labeled rate of a spray surfactant should be tank mixed with Willowood Tebucon 3 6SC Willowood Tebucon 3 6SC must have two to four hours of drying time on bean foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs After this period of time Willowood Tebucon 3 6SC will be resistant to weathering Willowood Tebucon 3 6SC is a demethylation inhibitor (DMI) fungicide (Group 3)</p> <p>Restricted entry interval (REI) = 12 hours</p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3 6SC
Corn (sweet corn field corn field corn grown for seed and popcorn)	Rust (<i>Puccinia</i> spp) Northern leaf blight (<i>Helminthosporium turcicum</i>) Southern leaf blight (<i>Helminthosporium maydis</i>) Northern leaf spot (<i>Helminthosporium carbonum</i>) Gray leaf spot (<i>Cercospora zeae maydis</i>)	4 to 6 fl oz per acre
	<p>Notes Apply Willowood Tebucon 3 6SC in a protective spray schedule or when weather conditions are favorable for disease development Repeat applications at 7 to 14 day intervals or as necessary to maintain control A maximum of 24 fl oz (1 5 pint) of Willowood Tebucon 3 6SC may be applied per acre per crop season Sweet corn Willowood Tebucon 3 6SC may be applied up to 7 days before the harvest of ears or forage and 49 days before the harvest of fodder Field seed or popcorn Willowood Tebucon 3 6SC may be applied up to 21 days before the harvest of forage and 36 days before the harvest of grain or fodder</p>	
<p>General Comments For optimum disease control the lowest labeled rate of a spray surfactant should be tank mixed with Willowood Tebucon 3 6SC Willowood Tebucon 3 6SC must have two to four hours of drying time on corn foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs After this period of time Willowood Tebucon 3 6SC will be resistant to weathering Willowood Tebucon 3 6SC is a demethylation inhibitor (DMI) fungicide (Group 3)</p> <p>Restricted entry interval (REI) for sweet corn = 19 hours</p> <p>Restricted entry interval (REI) for all corn except sweet corn = 12 hours</p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3 6SC
Cotton	Southwestern cotton rust (<i>Puccinia cacabata</i>)	6 to 8 fl oz per acre
	<p>Notes Apply Willowood Tebucon 3 6SC in a protective spray schedule or when weather conditions are favorable for rust development Repeat applications at 7 to 14 day intervals or as necessary to maintain control Willowood Tebucon 3 6SC may be applied up to 30 days before harvest Do not apply more than 24 fl oz of Willowood Tebucon 3 6SC per acre per crop season</p>	
<p>General Comments For optimum disease control the lowest labeled rate of a spray surfactant should be tank mixed with Willowood Tebucon 3 6SC Willowood Tebucon 3 6SC must have two to four hours of drying time on cotton foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs After this period of time Willowood Tebucon 3 6SC will be resistant to weathering Willowood Tebucon 3 6SC is a demethylation inhibitor (DMI) fungicide</p>		

(Group 3)

Restricted entry interval (REI) = 12 hours

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3 6SC
Cucurbit Vegetables Group Chayote Chinese waxgourd Citron melon Cucumber Gherkin Edible gourd (includes hyotan cucuzza hechima and Chinese okra) <i>Momordica</i> spp (includes balsam apple balsam pear bitter melon and Chinese cucumber) Muskmelon (includes cantaloupe casaba crenshaw melon golden pershaw melon honeydew melon honey balls mango melon Persian melon pineapple melon Santa Claus melon and snake melon) Pumpkin Summer squash (includes crookneck squash scallop squash straightneck squash vegetable marrow and zucchini) Winter squash (includes butternut squash calabaza hubbard squash acorn squash and spaghetti squash) Watermelon	Powdery mildew (<i>Sphaerotheca fuliginea</i> / <i>Podosphaera xanthii</i>) (<i>Erysiphe cichoracearum</i>)	4 to 6 fl oz per acre
	Gummy stem blight suppression (<i>Didymella bryoniae</i>) (watermelon squash pumpkin and melons only)	8 fl oz per acre
<p>Notes Apply the specified dosage in a protective spray schedule to foliage and fruit Repeat applications at 10 to 14 day intervals Willowood Tebucon 3 6SC may be applied up to 7 days before harvest Do not apply more than 24 fl oz of Willowood Tebucon 3 6SC per acre per crop season</p>		

General Comments For optimum disease control the lowest labeled rate of a spray surfactant should be tank mixed with Willowood Tebucon 3 6SC Willowood Tebucon 3 6SC must have two to four hours of drying time for the active ingredient to move systemically into plant tissue before rain or irrigation occurs After this period of time Willowood Tebucon 3 6SC will be resistant to weathering Willowood Tebucon 3 6SC is a demethylation inhibitor (DMI) fungicide (Group 3)

Restricted entry interval (REI) = 12 hours

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3 6SC
Dry bulb onion Garlic Great headed (elephant) garlic Welch onion Shallot	White rot (<i>Sclerotium cepivorum</i>)	White rot 20.5 fl oz per acre applied in a 4 to 6 inch band over/into each furrow May be applied by chemigation to control white rot
	Rust (<i>Puccinia allii</i> <i>Puccinia porri</i>)	4 to 6 fl oz per acre

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3 6SC
	Purple blotch (<i>Alternaria porri</i>)	
	<p>White rot For the control of white rot make one application in the furrow at the time of planting The in furrow application should be made at the rate of 20.5 fl oz Willowood Tebucon 3 6SC per acre Apply the entire per acre rate in a 4 to 6 inch band over/into each furrow Additional control may be obtained by including two foliar applications at 4 to 6 fl oz/acre</p> <p>Rust For the control of rust make foliar applications at the rate of 4 to 6 fl oz Willowood Tebucon 3 6SC per acre per application Repeat at an interval of 10 to 14 days Apply Willowood Tebucon 3 6SC in a protective spray schedule or when weather conditions are favorable for rust development</p> <p>Notes Do not apply more than 32.5 fl oz Willowood Tebucon 3 6SC per acre per season if an in furrow treatment is made If Willowood Tebucon 3 6SC is not applied as an in furrow treatment then do not apply more than 12 fl oz Willowood Tebucon 3 6SC per acre per season as a foliar spray Do not apply within 7 days of harvest (PHI = 7 days)</p>	
<p>General Comments For optimum results use as a preventative treatment Begin applications as soon as crop and/or environmental conditions become favorable for disease development The lowest recommended rate of a spray surfactant may be tank mixed with Willowood Tebucon 3 6SC Willowood Tebucon 3 6SC must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs After this period of time Willowood Tebucon 3 6SC will be resistant to weathering Willowood Tebucon 3 6SC is a demethylation inhibitor (DMI) fungicide (Group 3)</p> <p>Restricted entry interval (REI) = 12 hours</p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3 6SC
Fruiting Vegetables Group_ Eggplant Groundcherry Pepino Pepper Tomatillo Tomato <u>Not registered for this use in California</u>	Early blight (<i>Alternaria solani</i>)	8 fl oz per acre
	<p>Notes Apply Willowood Tebucon 3 6SC as a foliar spray using an interval of 7 days Do not apply more than 48 fl oz of Willowood Tebucon 3 6SC per acre per season Do not apply within 7 days of harvest (PHI = 7 days)</p>	
<p>General Comments For optimum results use as a preventative treatment Begin applications as soon as crop and/or environmental conditions become favorable for disease development The lowest recommended rate of a spray surfactant may be tank mixed with Willowood Tebucon 3 6SC Willowood Tebucon 3 6SC must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs After this period of time Willowood Tebucon 3 6SC will be resistant to weathering Willowood Tebucon 3 6SC is a demethylation inhibitor (DMI) fungicide (Group 3)</p> <p>Restricted entry interval (REI) = 12 hours</p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3 6SC
Grasses Grown For Seed	Rusts (<i>Puccinia spp</i>)	4 to 8 fl oz per acre
	Apply the specified rate of Willowood Tebucon 3 6SC as soon as weather conditions are favorable for rust development or when first rust pustules are present Repeat applications at 14 to 16 day intervals Under heavy disease pressure use 6 to 8 fl oz/A and shorter spray intervals	
	Powdery mildew	4 to 8 fl oz per acre
	Apply specified rate of Willowood Tebucon 3 6SC when powdery mildew first appears on the leaves Repeat applications at 14 to 16 day intervals Under heavy disease pressure use 6 to 8 fl oz/A and shorter spray intervals	
<p>General Comments Apply the specified rate in a minimum of 20 gallons of water per acre with ground sprayers or in a minimum of 10 gallons of water per acre with aircraft Thorough coverage is important for optimum disease control</p> <p>For optimum benefit the lowest specified rate of a spray surfactant should be tank mixed with Willowood Tebucon 3 6SC</p> <p>A maximum of 16 fluid ounces (1 pint) may be applied per acre per crop season Willowood Tebucon 3 6SC may be applied up to 4 days before harvest Chaff screenings and straw from treated areas may be used for feed purposes however do not forage cut green crop or use seed for feed purposes Regrowth may be grazed starting 17 days after last application</p> <p>Restricted entry interval (REI) = 12 hours</p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3 6SC
Green onion Leek Spring onion Scallion Japanese bunching onion Green shallots Green eschalots	White rot (<i>Sclerotium cepivorum</i>) suppression only	4 to 6 fl oz per acre
	Rust (<i>Puccinia allii Puccinia porri</i>)	
	Purple blotch (<i>Alternaria porri</i>)	
<p>For the control of diseases make foliar applications using an interval of 10 to 14 days Apply Willowood Tebucon 3 6SC in a protective spray schedule or when weather conditions are favorable for rust development</p> <p>Notes Do not apply more than 24 fl oz Willowood Tebucon 3 6SC per acre per season Do not apply within 7 days of harvest (PHI = 7 days)</p>		
<p>General Comments For optimum results use as a preventative treatment Begin applications as soon as crop and/or environmental conditions become favorable for disease development The lowest recommended rate of a spray surfactant may be tank mixed with Willowood Tebucon 3 6SC Willowood Tebucon 3 6SC must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs After this period of time Willowood Tebucon 3 6SC will be resistant to weathering Willowood Tebucon 3 6SC is a demethylation inhibitor (DMI) fungicide (Group 3)</p> <p>Restricted entry interval (REI) = 12 hours</p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3 6SC
Hops	Powdery mildew (<i>Sphaerotheca humuli</i> / <i>Sphaerotheca macularis</i>)	4 to 8 fl oz per acre
	<p>Notes Apply the specified dosage in a protective spray schedule to foliage Repeat applications at 10 to 14 day intervals Willowood Tebucon 3 6SC may be applied up to 14 days before harvest Do not apply more than 32 fl oz of Willowood Tebucon 3 6SC per acre per crop season Increase the spray volume and the application rate as vine growth increases during the season</p>	
<p>General Comments For optimum disease control the lowest labeled rate of a spray surfactant should be tank mixed with Willowood Tebucon 3 6SC Willowood Tebucon 3 6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs After this period of time Willowood Tebucon 3 6SC will be resistant to weathering Willowood Tebucon 3 6SC is a demethylation inhibitor (DMI) fungicide (Group 3)</p>		
<p>Restricted entry interval (REI) = 12 hours</p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3 6SC
Leafy Brassica Greens Broccoli raab Chinese cabbage (bok choy) Collards Kale Mizuna Mustard greens Mustard spinach Rape greens Turnip greens	Cercospora leaf spot (<i>Cercospora brassicicola</i>)	3 to 4 fl oz per acre
	Powdery mildew (<i>Erysiphe cruciferarum</i>)	
	Alternaria leaf spot (<i>Alternaria brassicicola</i>)	
<p>Notes Do not apply more than 16 fl oz Willowood Tebucon 3 6SC per acre per season Do not apply within 7 days of harvest (PHI = 7 days) Do not apply more often than once every 10 days</p>		
<p>General Comments For optimum results use as a preventative treatment Begin applications as soon as crop and/or environmental conditions become favorable for disease development The lowest recommended rate of a spray surfactant may be tank mixed with Willowood Tebucon 3 6SC Willowood Tebucon 3 6SC must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs After this period of time Willowood Tebucon 3 6SC will be resistant to weathering Willowood Tebucon 3 6SC is a demethylation inhibitor (DMI) fungicide (Group 3)</p>		
<p>Restriction Application to turnip greens is limited to East of the Rockies</p>		
<p>Restricted entry interval (REI) = 12 hours</p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3 6SC
Garden beet roots and tops (leaves)	Cercospora leaf spot (<i>Cercospora beticola</i>)	3 to 7.2 fl oz per acre
	Notes Make applications on a 14 day interval Do not apply more than 28.8 fl oz Willowood Tebucon 3 6SC per acre per season Do not apply within 7 days of harvest (PHI = 7 days)	
General Comments For optimum results use as a preventative treatment Begin applications as soon as crop and/or environmental conditions become favorable for disease development The lowest recommended rate of a spray surfactant may be tank mixed with Willowood Tebucon 3 6SC Willowood Tebucon 3 6SC must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs After this period of time Willowood Tebucon 3 6SC will be resistant to weathering Willowood Tebucon 3 6SC is a demethylation inhibitor (DMI) fungicide (Group 3)		
Restricted entry interval (REI) = 12 hours		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3 6SC
Lychee	Anthracoise (<i>Colletotrichum gloeosporioides</i>)	4 to 6 fl oz per acre
	Notes Begin first application of Willowood Tebucon 3 6SC as panicle emerges Spray up to 6 fl oz per acre every 10 days thereafter for a total of 8 sprays Apply specified dosage in a minimum of 50 gallons of spray solution per acre by ground only Do not apply more than 48 fl oz of Willowood Tebucon 3 6SC per acre per season Willowood Tebucon 3 6SC can be applied up to and including the day of harvest (PHI = 0 days)	
General Comments For optimum disease control the lowest labeled rate of a non ionic spray surfactant should be tank mixed with Willowood Tebucon 3 6SC Willowood Tebucon 3 6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs After this period of time Willowood Tebucon 3 6SC will be resistant to weathering Willowood Tebucon 3 6SC is a demethylation inhibitor (DMI) fungicide (Group 3)		
Restricted entry interval (REI) = 2 days		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3 6SC
Okra	Cercospora leaf spot (<i>Cercospora spp</i>)	4 to 6 fl oz per acre
	Notes Apply specific dosage of Willowood Tebucon 3 6SC in a preventative spray program Use the highest rate when disease conditions are favorable and in areas where high disease pressure is expected Applications may be repeated at 14 day	



intervals in order to maintain control of the disease Apply specified dosage as a foliar spray in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air Applications may be made no closer than 3 days before harvest Do not apply more than 24 fl oz of Willowood Tebucon 3 6SC per acre per season

General Comments For optimum disease control the lowest labeled rate of a spray surfactant should be tank mixed with Willowood Tebucon 3 6SC Willowood Tebucon 3 6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs After this period of time Willowood Tebucon 3 6SC will be resistant to weathering Willowood Tebucon 3 6SC is a demethylation inhibitor (DMI) fungicide (Group 3)

Restricted entry interval (REI) = 12 hours

APPLICATION DIRECTIONS

CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3 6SC
Peanut	<p>SOILBORNE Sclerotium stem and pod rot (white mold southern blight southern stem rot) Rhizoctonia limb rot Rhizoctonia pod rot (Virginia and North Carolina only)</p> <p>FOLIAR Early leaf spot Late Leaf spot Leaf rust Web blotch (<i>Phoma</i>) Pepper spot (<i>Leptoshaerulina</i>)</p>	7 2 fl oz per acre
<p>FOUR APPLICATION SPRAY PROGRAM Apply the specified rate in a preventive spray schedule See table below for proper timing of applications Applications of chlorothalonil should be made prior to and following applications of Willowood Tebucon 3 6SC to discourage development of resistant strains of fungi For optimum control of foliar diseases such as leaf rust web blotch and pepper spot the lowest label specified rate of a spray surfactant should be tank mixed with Willowood Tebucon 3 6SC</p> <p>LEAF SPOT ADVISORY SCHEDULE For control of soilborne diseases in an advisory schedule apply Willowood Tebucon 3 6SC in the first advisory spray in July and continue Willowood Tebucon 3 6SC applications at 14 day intervals Applications after August 15 should be tank mixed with chlorothalonil for resistance management purposes</p>		

GENERAL DIRECTIONS For optimum control of the specified soilborne diseases four consecutive applications of Willowood Tebucon 3 6SC must be made at 14 day intervals

A maximum of 28 8 fluid ounces of Willowood Tebucon 3 6SC may be applied per crop season Willowood Tebucon 3 6SC may be applied up to 14 days before harvest Do not feed hay or threshings or allow livestock to graze in treated areas

Willowood Tebucon 3 6SC is a sterol demethylation inhibitor (DMI) fungicide Chlorothalonil may be tank mixed at the rate of 12 ounces of active ingredient with Willowood Tebucon 3 6SC as a leaf spot resistance management strategy A spray surfactant is not necessary when Willowood Tebucon 3 6SC is tank mixed with chlorothalonil Mixing or alternating

Willowood Tebucon 3 6SC with other DMI fungicides may lead to resistance

Willowood Tebucon 3 6SC must be carried by rainfall or irrigation into the root and pod zone for control of root and pod rots caused by *Sclerotium rolfsii* and *Rhizoctonia solani*. Drought conditions will decrease the effectiveness of Willowood Tebucon 3 6SC against the root and pod rots

Use Willowood Tebucon 3 6SC in conjunction with cultural practices that are known to reduce the severity of soilborne diseases such as proper crop rotation practices

Restricted entry interval (REI) = 12 hours

Timing of Willowood Tebucon 3 6SC Application for Optimum Control of White Mold and Rhizoctonia Limb and Pod Rot

Spray Program	Willowood Tebucon 3 6SC Application No	Chlorothalonil Application No
7 Applications	3 4 5 and 6	1 2 and 7

APPLICATION DIRECTIONS

CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3 6SC
Pecan	Brown leaf spot (<i>Sirosporium diffusum</i>)	4 to 8 fl oz per acre
	Downy spot (<i>Mycosphaerella caryigena</i>)	
	Liver spot (<i>Gnomonia caryae</i>)	
	Scab (<i>Cladosporium caryigenum</i>)	
	Vein spot (<i>Gnomonia nerviseda</i>)	
	Zonate leaf spot (<i>Grovesinia pyramidalis</i>)	

Notes Apply Willowood Tebucon 3 6SC in a preventive spray schedule beginning at early bud break (young leaves unfolding) and continue applications at 10 to 14 day intervals through the pollination period. Willowood Tebucon 3 6SC should be applied at 4 fl oz per acre in a tank mix with the recommended rate of Super-Tin® in cover sprays. Follow label directions for the use of SuperTin. Do not add a surfactant to the spray solution when tank mixing Willowood Tebucon 3 6SC with SuperTin. Apply Willowood Tebucon 3 6SC in a spray volume of 15 or more gallons per acre by air or 50 or more gallons per acre by ground. Apply 7 to 8 fl oz per acre of Willowood Tebucon 3 6SC to full size mature trees and 4 to 6 fl oz per acre of Willowood Tebucon 3 6SC to smaller trees. Apply the high rate to varieties that are highly susceptible to the indicated diseases or when severe disease conditions exist. The lowest labeled rate of a surfactant may be added to the spray solution for optimum control of the indicated diseases. Do not apply after shucks begin to split. A maximum of 32 fl oz of Willowood Tebucon 3 6SC may be applied per acre per crop season.

Do not cut cover crops in treated areas for feed or allow livestock to graze treated areas

General Comments For optimum disease control the lowest labeled rate of a spray surfactant should be tank mixed with Willowood Tebucon 3 6SC Willowood Tebucon 3 6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs After this period of time Willowood Tebucon 3 6SC will be resistant to weathering Willowood Tebucon 3 6SC is a demethylation inhibitor (DMI) fungicide (Group 3) It may be applied in a tank mix or alternated (every other spray application) with a non DMI fungicide as a resistance management strategy

Restricted entry interval (REI) = 12 hours

APPLICATION DIRECTIONS

CROP	DISEASE(S)	RATE OF WILLOWOOD TEBUCON 3 6SC
Soybean	Rust (<i>Phakopsora pachyrhizi</i>) Powdery Mildew (<i>Microspheera diffusa</i>)	3 to 4 fl oz per acre

Use Directions Apply Willowood Tebucon 3 6SC as a broadcast foliar spray as a preventative spray or at first visible symptoms of disease Repeat applications on a 10 to 14 day spray interval if environmental conditions are favorable for continued disease development Use the higher rates and shorter spray intervals when disease pressure is severe The lowest labeled rate of spray surfactant must be tank mixed with Willowood Tebucon 3 6SC Apply Willowood Tebucon 3 6SC in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons per acre by aircraft spray equipment

Restrictions Applications may not be made within 21 days of harvest Do not apply more than 3 applications per season Do not apply more than 12 fl oz/a per use season

Restricted entry interval (REI) = 12 hours

APPLICATION DIRECTIONS

CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3 6SC
Sunflower	Rust (<i>Puccinia helianthi</i>)	4 to 6 fl oz per acre

Notes Apply specific dosage of Willowood Tebucon 3 6SC at the earliest sign of infection (rust pustules developing) or when weather conditions are favorable for rust development Apply higher rate to highly susceptible varieties and/or under severe disease conditions Application may be repeated at 14 days if necessary to maintain control of the disease Apply specified dosage in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air Do not apply more than 16 fl oz of Willowood Tebucon 3 6SC per acre per season or within 50 days of harvest

General Comments For optimum disease control the lowest labeled rate of a spray surfactant should be tank mixed with Willowood Tebucon 3 6SC Contact your state Extension Service or Willowood LLC representative for a list of approved surfactants Willowood Tebucon 3 6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs After this period of time Willowood Tebucon 3 6SC will be resistant to weathering WILLOWOOD TEBUCONAZOLE 3 6SC is a demethylation inhibitor (DMI) fungicide (Group 3)

Restricted entry interval (REI) = 12 hours

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3 6SC
Turnip (Application is limited to East of the Rockies)	Cercospora leaf spot (<i>Cercospora brassicicola</i>)	4 to 7.2 fl oz per acre
	<p>Notes Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 12 to 14 day intervals. Willowood Tebucon 3 6SC may be applied up to 7 days before harvest. Do not apply more than 28.8 fl oz of Willowood Tebucon 3 6SC per acre per crop season.</p>	
<p>General Comments For optimum disease control, the lowest labeled rate of a spray surfactant should be tank mixed with Willowood Tebucon 3 6SC. Willowood Tebucon 3 6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3 6SC will be resistant to weathering. Willowood Tebucon 3 6SC is a demethylation inhibitor (DMI) fungicide (Group 3).</p>		
<p>Restricted entry interval (REI) = 12 hours</p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3 6SC
Wheat	Rusts leaf, stem, and stripe (<i>Puccinia</i> spp.)	4 fl oz per acre
	Head blight or scab (<i>Fusarium</i> spp.) Suppression	
<p>Notes Wheat fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. A maximum of 4 fl oz of Willowood Tebucon 3 6SC may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw may be fed or used for bedding. Do not allow livestock to graze or feed green forage to livestock prior to 6 days after treatment with Willowood Tebucon 3 6SC. Apply Willowood Tebucon 3 6SC in a minimum of 10 gallons of spray solution per acre by ground, or in a minimum of 5 gallons of spray solution per acre by air.</p>		
<p>Application timing directions Rusts: Apply Willowood Tebucon 3 6SC at the earliest sign of rust pustules on foliage. Fusarium head blight: Optimal timing of Willowood Tebucon 3 6SC for Fusarium head blight suppression is the beginning of flowering on main stem heads (Feekes 10.51).</p>		
<p>General Comments For optimum disease control, the lowest specified rate of a spray surfactant should be tank mixed with Willowood Tebucon 3 6SC. Willowood Tebucon 3 6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3 6SC will be resistant to weathering. Willowood Tebucon 3 6SC is a demethylation inhibitor (DMI) fungicide (Group 3).</p>		
<p>Restricted entry interval (REI) = 12 hours</p>		

SEED TREATMENT Corn (Sweet Corn Field Corn Field Corn Grown For Seed and Popcorn)
 For control of soilborne and seedborne Fusarium and soilborne and seedborne head smut

SEED LABELING To meet U S Federal Seed Act requirements all seed treated with Willowood Tebucon 3 6SC must be labeled

TREATED SEED DO NOT USE FOR FOOD FEED OR OIL PURPOSES Treated with Tebuconazole

USE PRECAUTION When using formulations that do not contain dye to comply with 40 CFR 153 155 all seed treated with an economic poison must be colored to distinguish and prevent subsequent inadvertent use as a food for man or feed for animals

DISEASE	RATE Fl Oz/CWT	DIRECTIONS FOR USE
Soilborne and Seedborne Fusarium	0.071	Apply as a seed treatment using standard slurry or mist type seed treatment equipment Uniform application of seed is necessary to ensure seed safety and best disease protection Seed should be sound and well cured prior to treatment Product should be diluted with sufficient water to ensure complete seed coverage Consult a seed treatment specialist regarding slurry rates recommended for the crop to be treated with Willowood Tebucon 3 6SC The length of control will vary depending on the rate used
Soilborne and Seedborne Head smut <i>(Sphacelotheca reilana)</i>	0.27 – 0.54	

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES RESERVOIRS RIVERS PERMANENT STREAMS MARSHES OR NATURAL PONDS AND ESTUARIES

Apply only during alternate years in fields adjacent to aquatic areas listed above

Do not apply by ground or air within 100 feet of aquatic areas listed above

Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip

Spray Drift Management For aerial applications the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices The minimum practical boom length should be used and must not exceed 75% of the wing span or rotor diameter

Use the largest droplet size consistent with pest control Formation of very small droplets may be minimized by appropriate nozzle selection by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure Apply in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment

Spray should be released at the lowest possible height consistent with good pest control and flight safety Applications more than 10 feet above the crop canopy should be avoided

Make aerial or ground applications when wind velocity favors on target product deposition (approximately 3 to 10 mph) Do not apply when wind velocity exceeds 15 mph Avoid applications when wind gusts approach 15 mph

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas Avoid spraying during conditions of low humidity and/or high temperature

Do not make aerial or ground applications during temperature inversions Inversions are characterized by stable air and increasing temperatures with height above the ground Mist or fog may indicate the presence of an inversion in humid areas The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface

ROTATIONAL CROPS

Treated areas may be replanted with any crop specified on this label as soon as practical after last application Any crop not specified on this label may be planted into treated areas 120 days after last application

DISEASE CONTROL IN GOLF COURSE TURF, FIELD, NURSERY AND CONTAINER ORNAMENTALS AND COMMERCIAL AND RESIDENTIAL LANDSCAPES

Chemigation Do not apply this product through any type of irrigation system

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES RESERVOIRS RIVERS PERMANENT STREAMS MARSHES OR NATURAL PONDS AND ESTUARIES

- Do not apply within 100 feet of the aquatic areas listed above
- Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetation filter strip
- See Spray Drift Management section for further information

Spray Drift Management

Make ground application when wind velocity favors on target product deposition (approximately 3 to 10 mph) Do not apply when wind velocity exceeds 15 mph Avoid applications when wind gusts approach 15 mph

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperatures.

Do not make ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Spray Volume For best results Willowood Tebucon 3 6SC may be applied in 66-132 gallons of water per acre for turf using ground based equipment. For ornamentals 50-300 gallons of finished spray per acre are recommended depending upon the equipment, plant species and plant growth stage at time of application. For the most effective results, equipment calibration should be checked regularly. When using lower spray volumes, be sure to maintain uniform application and full crop coverage so as to ensure effective control. Increase spray volume to ensure proper application, if required.

Compatibility Test for Mix Components

Before mixing components, always perform a compatibility jar test. For 66 gallons per acre spray volume, use 5 cups of water in a clear, clean mixing jar. For other spray volumes, adjust accordingly. Only use water from the intended source at the source temperature. Add components in the sequence indicated below in Mixing Order using 3 teaspoons for each pound of dry product or 1 1/2 teaspoon for each pint of liquid product of recommended label rate per acre. Always cap the jar and invert 10 cycles between component additions. When the components have all been added to the jar and fully mixed, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent and use the compatibility agent as directed on its label.

Mixing Continuous agitation is required during mixing. When mixing this product and water, use the specified application rates as listed for each crop on this label. Before combining any other substances with the mixture, ensure that Willowood Tebucon 3 6SC is completely dispersed in the mixture.

Recommended Mixing Procedure

- 1 Water: Add three quarters of the required volume to a thoroughly clean sprayer tank.
- 2 Agitation: Start agitation and maintain constant agitation throughout mixing and application.
- 3 Inductor: If an inductor is used, rinse it thoroughly after each component has been added.
- 4 Products in PVA Bags: Place any product contained in water soluble PVA bags into the mixing tank. Wait until all water soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
- 5 Water Dispersible Products: Including dry flowables (DF), wettable powders (WP), suspension concentrates (SC) or suspo emulsions (SE).
- 6 Water soluble products
- 7 Emulsifiable concentrates (such as oil concentrates when applicable)
- 8 Water soluble additives (such as AMS or UAN when applicable)
- 9 Remaining quantity of water

DISEASE CONTROL IN GOLF COURSE TURF

Turf Use Restrictions and Precautions

For use on golf course turf only.
 Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (i.e. elementary, middle and high school), campgrounds, churches, and theme parks.
 Not for homeowner use.
 Not for use on turf being grown for sale or commercial use as sod.
 Do not use clippings for animal feed.
 Do not exceed 3.6 fl. oz. of Willowood Tebucon 3 6SC per 1,000 sq. ft. per year.
 Do not apply more than 6 applications per year.

Product Information

For use on all golf turf applications of cool season and warm season grasses (such as Bentgrasses, Bluegrasses, Fescues, Ryegrasses, St. Augustine grasses, and Zoysia) or their mixtures. Willowood Tebucon 3 6SC is not phytotoxic to any of the above mentioned grasses when used in accordance with the label.

Note Bermudagrass can be sensitive to Willowood Tebucon 3 6SC under certain conditions Do not apply consecutive applications during or just after dormancy break Avoid applications when temperatures are expected to exceed 85 degrees F

Willowood Tebucon 3 6SC can be used for the prevention and control of the diseases mentioned in the table below Begin applications when conditions favor disease development and repeat applications as long as these conditions persist Preventative treatments can be applied using 28 day intervals as indicated When treating golf greens always treat aprons and approaches Spray uniformly over the area to be treated with properly calibrated equipment

Apply the specified amount of Willowood Tebucon 3 6SC in sufficient water for thorough coverage A volume of 66 132 gallons per acre (1 5 3 0 gallons per 1 000 sq ft) is recommended Apply using properly calibrated low volume hand held mechanical or motorized ground broadcast equipment Application to small areas may be made with low pressure handwand or backpack equipment Maintain constant agitation during application

Depending on the disease Willowood Tebucon 3 6SC should be watered into the crown and active root zone for best results Make all applications after mowing and allow foliage to dry thoroughly before irrigation For best results use spray mixture the same day it is prepared

Golf Course Turf Disease Control

DISEASE	RATE of Willowood Tebucon 3 6SC (Fl oz/1000 Sq Ft)	NOTES
Dollar Spot (<i>Sclerotinia homoeocarpa</i>) Copper Spot (<i>Gloeocercospora sorghi</i>) Powdery Mildew (<i>Erysiphe graminis</i>) Corticium Red Tread (<i>Laetisaria fuciformis</i>) Rusts (<i>Puccinia</i> spp)	0 6	For prevention begin applications when conditions are favorable for disease development Do not make two consecutive applications of Willowood Tebucon 3 6SC Alternate with another fungicide with a different mode of action A second application may be made after 28 days
Brown Patch/Rhizoctonia Blight Large Patch (<i>Rhizoctonia solani</i>) Brown Ring Patch (<i>R circinata</i>)	0 6	For prevention begin applications when conditions are favorable for disease development Do not make two consecutive applications of Willowood Tebucon 3 6SC Alternate with a different mode of action A second application may be made after 28 days
Anthracnose Basal and Foliar (<i>Colletotrichum cereal</i>) Red Thread (<i>Laetisaria fuciformis</i>) Pink Patch (<i>Limonomyces rosipellis</i>)	0 6	Immediately after fungicide is applied irrigate the area with sufficient water to move the active ingredient down into the crown and root zone of the turf The amount of water is dependent on the depth of root zone For prevention begin applications two or four weeks prior to the historical appearance of disease symptoms Initiate cultural control practices at the same time the fungicide is applied Refer to your local County Extension Service for this information Appl, subsequent application at 28 day intervals

DISEASE	RATE of Willowood Tebucon 3 6SC (Fl oz/1000 Sq Ft)	NOTES
Take All Patch (<i>Gaeumannomyces graminis</i>)	0.6	For prevention apply in the fall when soil temperature reaches 55-65° F and again in the spring under similar soil temperature conditions. Applications in both fall and spring may be necessary. Immediately after fungicide is applied irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
Gray Leaf Spot (<i>Pyricularia grisea</i>)	0.6	Apply when conditions are favorable for disease development at 28 day intervals.
Stipe Smut (<i>Ustilago striiformis</i>)	0.6	Make a single application to historical disease areas in spring as grass growth begins.
Spring Dead Spot (<i>Leptosphaeria korrea</i> , <i>L. narmari</i> , <i>Ophiosphaerella herpotricha</i> , <i>Gaeumannomyces graminis</i>) Necrotic Ring Spot (<i>Leptosphaeria korrea</i>)	0.6	For prevention apply in fall when soil temperatures reach 65° F and again in spring under similar soil temp conditions or after dormancy break. Immediately after fungicide is applied irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
Fusarium Patch (<i>Fusarium roseum</i>)	0.6	Apply first application in mid June or 28 days prior to time this blight normally becomes evident. Make applications at no less than 28 day intervals.
Summer Patch (<i>Magnaporthe poae</i>)	0.6	Apply beginning in the spring. Do not make two consecutive applications of Willowood Tebucon 3 6SC. Alternate with another fungicide with a different mode of action. Second and third applications may be made at 28 day intervals. See local university recommendations for suggested timing. Immediately after fungicide is applied irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
Zoysia Patch Large Patch of zoysia (<i>Rhizoctonia solani</i>)	0.6	Make first application in early fall (mid September to mid October) prior to development of disease symptoms. A second application in early spring may be necessary in areas where disease pressure is known to be heavy.
Gray Snow Mold/Typhula Blight (<i>Typhula incarnate</i>) Pink Snow Mold/Microdochium Patch	0.6	Apply in the fall before anticipated turf dormancy and before first snow cover. If turf breaks dormancy during winter months a second application may be made. Do not apply over a snow cover or when turf is

DISEASE	RATE of Willowood Tebucon 3 6SC (Fl oz/1000 Sq Ft)	NOTES
(<i>Microdochium nivale</i>)		dormant It is recommended that Willowood Tebucon 3 6SC be tank mixed with other registered snow mold products for best season long results
NOTE Apply the specified amount of Willowood Tebucon 3 6SC in 1 5 to 3 0 gallons of water per 1000 sq ft Make all applications after mowing and allow foliage to dry thoroughly before irrigation Do not use clippings for animal feed Do not exceed 3 6 fl oz of Willowood Tebucon 3 6SC per 1000 sq ft per year Do not exceed 6 applications per year		

DISEASE CONTROL IN FIELD NURSERY AND CONTAINER ORNAMENTALS AND COMMERCIAL and RESIDENTIAL LANDSCAPES

Ornamental Use Restrictions and Precautions

For use on ornamental plants only not for use on woodlands or forest management

Not for homeowner use

Do not apply more than 10 fl oz per acre in a single application

Do not apply more than 0 94 gallons (120 fl oz) of Willowood Tebucon 3 6SC (equal to 3 38 lbs of tebuconazole) per acre per year

Do not make more than 4 applications per year at highest rate

Do not apply to bearing fruit trees or vegetables

Willowood Tebucon 3 6SC can be used in a preventative and curative disease control program for the listed plant types and disease in the table below Optimum disease management is obtained when Willowood Tebucon 3 6SC is used in conjunction with sound disease management practices

Apply material with properly calibrated hand held mechanical or motorized spray equipment Begin applications when disease first appears and repeat at 14 21 day intervals during the growing season Use the shortest interval when conditions are unusually favorable for the development of disease For hand held mechanical or motorized applications mix as directed below and apply as a full coverage spray to drip for the prevention and control of the diseases listed below Choose a finished spray volume appropriate for the size of the plants and amount of foliage which will provide thorough coverage throughout the canopy Allow sprays to dry before overhead irrigation is applied

Apply Willowood Tebucon 3 6SC at rates of 4 10 fl oz per acre in 100 gallons of water Spray volume may range from 50 up to 300 gallons of finished spray per acre depending upon equipment plant species and plant growth stage at time of application

Note The Directions for Use of this product reflect the cumulative inputs from both historical field use and product testing programs However it is impossible to test this product on all species and cultivars A preliminary trial is suggested on a small scale before a full treatment is applied to any plant type not shown on this label but found in a similar use site with a listed disease problem Wait 5 7 days after treatment to evaluate results This product is not recommended for use on African Violets Begonias Boston Fern and Geraniums

Ornamentals Disease Control

PLANTS	DISEASE	APPLICATION	
		To Prevent Diseases	To Treat Existing Disease
Roses	Black Spot Powdery Mildew Rust	Apply every 14 21 days during the growing season starting when leaves first appear	Apply every 14 days for a total of 3 applications beginning at the first sign of disease
Flowers	Leaf Spot Powdery Mildew Rust Southern Blight	Apply at least 3 times per year 14 21 days apart beginning with Spring bud break Rotation or Tank mixing with barrier protectant fungicides is	
Crabapples (Ornamental)	Anthracoise Leaf Spot		

PLANTS	DISEASE	APPLICATION	
		To Prevent Diseases	To Treat Existing Disease
Dogwoods and Other Landscape (Ornamental) Trees	Powdery Mildew Rust Scab	recommended for resistance management	
Azaleas Camellias Rhododendrons and Other Landscape (Ornamental) Shrubs	Anthraxnose Black Spot Leaf Spot Petal Blight Powdery Mildew Rust	Petal Blight – Apply 2 3 times per week into the flowers as they open and develop color	
Ground Covers and Vines	Southern Blight		
HOW MUCH TO USE FOR SMALL PLANTINGS ADD 1 TEASPOON TO 2 5 GALLONS OF WATER			

Pump Style Sprayers

- 1 Add the appropriate amounts of concentrate and water to the sprayer tank
- 2 Close the sprayer shake well and pressurize
- 3 Adjust nozzle to a coarse spray pattern and apply
- 4 Occasionally re pressurize the sprayer if needed to maintain a good spray pattern

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable return the product at once unopened and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product which are beyond the control of Willowood LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Willowood LLC and Seller harmless for any claims relating to such factors.

Willowood LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Willowood LLC, and Buyer and User assume the risk of any such use. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, WILLOWOOD LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

To the extent consistent with applicable law, neither Willowood LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF WILLOWOOD LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR AT THE ELECTION OF WILLOWOOD LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Willowood LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Willowood LLC.