



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

May 17, 2024

Victoria Smith
Agent for Willowood, LLC
c/o Generic Crop Science, LLC
1887 Whitney Mesa Drive #9740
Henderson, NV 89014-2069

Subject: Label Amendment – Add me-too use on Tree Nut Crop Group 14 and state-specific restrictions.
Product Name: Willowood Tebucon 3.6 SC
EPA Registration Number: 87290-13
Application Date: 03/24/2023
Case Number: 479808

Dear Victoria Smith:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find

or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Senedu Alemu at alemu.senedu@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to be 'CGP', with a long horizontal line extending from the top of the signature.

for

Cynthia Giles-Parker
Chief, Fungicide Branch
Registration Division (7505T)
Office of Pesticide Programs

Enclosure

[MASTER LABEL]

TEBUCONAZOLE GROUP 3 FUNGICIDE

Willowood Tebucon 3.6SC

[ABN: Willowood Tebucon 4SC]
 [ABN: Farmers First Tebuconazole 4SC]
 [ABN: Farmers First Tebucon 4SC]
 [ABN: GCS Tebuconazole 4SC]
 [ABN: GCS Tebucon 4SC]

For control of specified diseases on various crops, golf course turf, and field, and nursery and container ornamentals in commercial and residential landscapes.

ACTIVE INGREDIENT:

Tebuconazole, alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol38.7%

OTHER INGREDIENTS: 61.3%**TOTAL:** 100.0%

Contains 3.6 pounds tebuconazole per gallon.

EPA Reg. No. 87290-13

EPA Est. No.

STOP - Read the label before use
Keep Out of Reach of Children
CAUTION

FIRST AID

If swallowed:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. DO NOT induce vomiting unless told to do so by a poison control center or doctor. DO NOT give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at **1-800-858-7378** Mon.-Fri., 8:00 am to 12:00 pm Pacific Time or your poison control center at **1-800-222-1222**.

NOTE TO PHYSICIAN: No specific antidote. Treat symptomatically. The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.

[See inside label booklet for additional Precautionary Statements and Directions for Use.]

[Optional referral statement: See inside label booklet for First Aid, Precautionary Statements and Directions for Use including Storage and Disposal instructions.]

Manufactured For:
 Willowood, LLC
 1887 Whitney Mesa Drive #9740
 Henderson, NV 89014

NET CONTENTS:

[Contains tebuconazole, the active ingredient used in Folicur® 3.6F.]

[Willowood Tebucon 3.6SC is not manufactured, or distributed by Bayer CropScience, seller of Folicur® 3.6F.]

ACCEPTED**05/17/2024**

Under the Federal Insecticide, Fungicide
 and Rodenticide Act as amended, for the
 pesticide registered under
 EPA Reg. No.

87290-13

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with skin, eyes, or clothing. Avoid breathing vapor or spray mist. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves (barrier laminate, or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton)
- Shoes plus socks

User Safety Requirements:

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish, and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory: Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water Advisory: Willowood Tebucon 3.6SC may contaminate water through drift of spray in wind. **Willowood Tebucon 3.6SC** has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of **Willowood Tebucon 3.6SC** will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

[Aerial application is prohibited in New York State.]

AGRICULTURAL USE REQUIREMENTS

Use this product in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI). The REI for each crop is listed in the application directions associated with each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves (barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton)
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box only apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Golf Course Turf and Landscape Uses: Keep children and pets out of treated areas until sprays have dried.

PRODUCT INFORMATION

Read the entire Directions for Use and Conditions of Sale before using this product.

Spray Volume: Willowood Tebucon 3.6SC may be applied in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied. If necessary, increase the spray volume per acre for complete crop coverage.

Chemigation: Apply Willowood Tebucon 3.6SC through irrigation equipment only to crops and diseases for which the chemigation use is specified. Apply Willowood Tebucon 3.6SC only through center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move; or drip (trickle) irrigation systems. **DO NOT** apply Willowood Tebucon 3.6SC through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts. **DO NOT** connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments if the need arises.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, for example, a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension. Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. Pesticide may be applied continuously for the duration of the water application.

Mixing: Add labeled amount of Willowood Tebucon 3.6SC into the spray tank while filling with water to the desired level. Operate the agitator while mixing. If other materials are added to the spray tank, disperse Willowood Tebucon 3.6SC thoroughly prior to the addition of other materials. **DO NOT** tank mix with products containing a prohibition against tank mixing. Follow the most restrictive labeling requirements of any tank mix product.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Compatibility: To determine the compatibility of Willowood Tebucon 3.6SC with other products, the following procedure must be followed: Pour the directed proportions of the products into a suitable container of water, mix thoroughly, and allow to stand at least five (5) minutes. If the combination remains mixed or can be re-mixed readily, the mixture is considered physically compatible. For information, contact your local Willowood, LLC representative.

Resistance Management Statement

Willowood Tebucon 3.6SC is a Group 3 fungicide which exhibits no known cross-resistance to other fungicide groups. However, fungal pathogens are known to develop resistance to products with the same mode of action when used repeatedly. Any fungal population may contain or develop individuals that are resistant to Willowood Tebucon 3.6SC and other Group 3 fungicides. If Group 3 fungicides are used repeatedly in the same field or in successive years as the primary method of control for targeted diseases, the resistant isolates may eventually dominate the fungal population. Because resistance development cannot be predicted, the use of Willowood Tebucon 3.6SC must conform to resistance management strategies established for the crop and use area. Such strategies may include rotation and/or tank mixing with products having different modes of action or limiting the total number of applications per year. Contact your local extension specialist, certified crop advisor, and/or manufacturer for fungicide resistance management and/or integrated disease management guidelines for specific crops and resistant disease populations. Willowood, LLC encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label.

ROTATIONAL CROPS

Treated areas may be replanted with any crop specified on this label as soon as practical after last application. Any crop not specified on this label may be planted into treated areas 120 days after last application.

DISEASE CONTROL IN CROPS

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Almond[*]	Brown rot blossom blight (<i>Monilinia laxa</i> , <i>M. fructicola</i>)	8 fl. oz. (0.225 lb. a.i.) per acre
<p>Application Instructions: Begin application at pink bud. If the bloom period is extended, and/or severe disease conditions exist, make a second application at full bloom. If conditions remain favorable for disease, make another application at petal fall. Willowood Tebucon 3.6SC in a minimum spray volume of 15 gallons per acre by air or 50 gallons per acre by ground. Reduce the application interval for varieties that are highly susceptible to the indicated diseases or when severe disease conditions exist. The use of ground application after petal fall is preferred because of difficulty in penetrating the canopy and obtaining thorough coverage of the foliage and fruit by air.</p> <p>For optimum disease control, tank mix the lowest labeled rate of a spray surfactant with Willowood Tebucon 3.6SC. Willowood Tebucon 3.6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6SC will be resistant to weathering. Willowood Tebucon 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3). It may be applied in a tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy.</p> <p>Precautions:</p> <ul style="list-style-type: none"> Tank mixing with other DMI fungicides is not advised. <p>Restrictions:</p> <ul style="list-style-type: none"> A maximum of 32 fl. oz. (0.9 lb. a.i.) of Willowood Tebucon 3.6SC may be applied per acre per year. A maximum of 8 fl. oz. (0.225 lb. a.i.) of Willowood Tebucon 3.6SC may be applied per application. DO NOT make more than 4 applications per year. DO NOT cut cover crops in treated areas for feed or allow livestock to graze treated areas. DO NOT apply within 35 days of harvest (PHI = 35 days). Restricted-entry interval (REI) = 12 hours. Minimum retreatment interval is 7 days. <p>[*Not for use in California.]</p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Asparagus	Rust (<i>Puccinia</i> spp.)	4 to 6 fl. oz. (0.113 – 0.14 lb. a.i.) per acre
<p>Application Instructions: Apply Willowood Tebucon 3.6SC as a foliar spray to the developing ferns after harvest of spears is completed. Apply at the earliest sign of rust pustules or when weather conditions are conducive for rust development. Apply 4 to 6 fl. oz. of Willowood Tebucon 3.6SC per acre (0.11 lb. a.i. – 0.17 lb. a.i. per acre) in alternation with another effective fungicide. Under conditions of severe rust pressure, use the higher specified rate. Repeat applications on a 14-day interval as necessary to maintain control of rust.</p> <p>Applications may be made using ground or aerial application equipment. For optimum disease control, tank mix the lowest labeled rate of a spray surfactant with Willowood Tebucon 3.6SC. Willowood Tebucon 3.6SC is a sterol demethylation inhibitor (DMI) fungicide (Group 3). Alternating Willowood Tebucon 3.6SC with other DMI fungicides may lead to resistance.</p> <p>Restrictions:</p> <ul style="list-style-type: none"> A 50-foot spray drift buffer zone is required for all aerial applications. A maximum of 6 fl. oz. (0.14 lb. a.i.) of Willowood Tebucon 3.6SC may be applied per application. DO NOT apply to harvestable spears. DO NOT apply within 100 days of harvest in California and 180 days in all other states. DO NOT make more than three foliar applications (18 fl. oz./acre or 0.51 lb. a.i./acre) per year. DO NOT apply more than 18 fl. oz./acre (0.51 lb. a.i./acre) per year. Restricted-entry interval (REI) = 12 hours. Minimum retreatment interval is 14 days. 		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Barley	Rusts (<i>Puccinia</i> spp.)	4 fl. oz. (0.113 lb. a.i.) per acre
	Head blight (<i>Fusarium</i> spp.) – Suppression	
	Application Instructions: Apply Willowood Tebucon 3.6SC in a minimum of 10 gallons of	

	<p>spray solution per acre by ground or in a minimum of 5 gallons of spray solution per acre by air. Observe barley fields closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development.</p> <p>Application timing directions: Rusts: Apply Willowood Tebucon 3.6SC at the earliest sign of rust pustules on foliage. Fusarium head blight: Optimal timing of Willowood Tebucon 3.6SC for Fusarium head blight suppression is when main stem heads have fully emerged (Feekes 10.5) on 50% of the plants.</p>
<p>For optimum disease control, tank mix the lowest specified rate of a spray surfactant with Willowood Tebucon 3.6SC. Willowood Tebucon 3.6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6SC will be resistant to weathering. Willowood Tebucon 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restrictions:</p> <ul style="list-style-type: none"> • A maximum of 4 fl. oz. (0.113 lb. a.i.) of Willowood Tebucon 3.6SC may be applied per acre per year. • A maximum of 4 fl. oz. (0.113 lb. a.i.) of Willowood Tebucon 3.6SC may be applied per application. • DO NOT apply within 30 days of harvest (PHI = 30 days). • DO NOT make more than one application per year at the highest rate. • Straw cut after harvest may be fed or used for bedding. • Grazing livestock or feeding of green forage is permitted 6 or more days after the last application of Willowood Tebucon 3.6SC. • Restricted-entry interval (REI) = 12 hours. 	

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Beans (fresh & dry except succulent shelled)	Rust (<i>Uromyces appendiculatus</i>)	4 to 6 fl. oz. (0.113 – 0.14 lb. a.i.) per acre
	Application Instructions: Apply Willowood Tebucon 3.6SC in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 14-day intervals, or as necessary to maintain control.	
For optimum disease control, tank mix the lowest labeled rate of a spray surfactant with Willowood Tebucon 3.6SC . Willowood Tebucon 3.6SC must have two to four hours of drying time on bean foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6SC will be resistant to weathering. Willowood Tebucon 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).		
Restrictions:		
<ul style="list-style-type: none">• Minimum retreatment interval is 14 days.		
<u>Beans, fresh:</u>		
<ul style="list-style-type: none">• Willowood Tebucon 3.6SC may be applied up to 7 days before harvest.• DO NOT apply more than 24 fl. oz. (0.675 lb. a.i.) of Willowood Tebucon 3.6SC per acre per year.• DO NOT apply more than 6 fl. oz. (0.14 lb. a.i.) of Willowood Tebucon 3.6SC per application.• DO NOT make more than 4 applications per year at the highest application rate.		
<u>Beans, dry:</u>		
<ul style="list-style-type: none">• Willowood Tebucon 3.6SC may be applied up to 14 days before harvest.• DO NOT apply more than 12 fl. oz.(0.34 lb. a.i.) of Willowood Tebucon 3.6SC per acre per year.• DO NOT make more than 2 applications per year at the highest application rate.• Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Corn[*] (sweet corn, field corn, field corn grown for seed, and popcorn)	Rust (<i>Puccinia</i> spp.) Northern leaf blight (<i>Helminthosporium turcicum</i>) Southern leaf blight (<i>Helminthosporium maydis</i>) Northern leaf spot (<i>Helminthosporium carbonum</i>) Gray leaf spot (<i>Cercospora zeae-maydis</i>)	4 to 6 fl. oz. (0.113 – 0.14 lb. a.i.) per acre
	Application Instructions: Apply Willowood Tebucon 3.6SC in a protective spray schedule or when weather conditions are favorable for disease development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control.	
For optimum disease control, tank mix the lowest labeled rate of a spray surfactant with Willowood Tebucon 3.6SC . Willowood Tebucon 3.6SC must have two to four hours of drying time on corn foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6SC will be resistant to weathering. Willowood Tebucon 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).		
Restrictions: <ul style="list-style-type: none">• A maximum of 24 fl. oz. (0.675 lb. a.i.) of Willowood Tebucon 3.6SC may be applied per acre per year.• A maximum of 6 fl. oz. (0.14 lb. a.i.) of Willowood Tebucon 3.6SC may be applied per application.• DO NOT make more than 4 applications per year.• Sweet corn: Willowood Tebucon 3.6SC may be applied up to 7 days before the harvest of ears or forage, and		

49 days before the harvest of fodder.

- Field, seed, or popcorn: **Willowood Tebucon 3.6SC** may be applied up to 21 days before the harvest of forage, and 36 days before the harvest of grain or fodder.
- **Restricted-entry interval (REI) for sweet corn** = 19 days.
- **Restricted-entry interval (REI) for all corn except sweet corn** = 12 hours.
- Minimum retreatment interval is 7 days.

[*Not for use in New York State.]

SEED TREATMENT - Corn (Sweet Corn, Field Corn, Field Corn Grown For Seed, and Popcorn)[*]

For control of soilborne and seedborne Fusarium and soilborne and seedborne head smut.

[*Not for use on corn in New York State.]

USE RESTRICTION: When using formulations that **DO NOT** contain dye, to comply with 40 CFR 153.155, all seed treated with an economic poison must be colored to distinguish and prevent subsequent inadvertent use as a food for man or feed for animals.

The Federal Seed Act requires that containers containing treated seeds shall be labeled with the following statements: This seed has been treated with tebuconazole fungicide. **DO NOT** use treated seed for feed, food, or oil purposes.

The U.S. Environmental Protection Agency requires the following statements on containers containing seed treated with tebuconazole:

- Store treated seed away from food and feedstuffs.
- **DO NOT** allow children, pets, or livestock to have access to treated seeds.
- Wear long pants, long-sleeved shirt and protective gloves when handling treated seed.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting.
- Dispose of all excess treated seed by burying seed away from bodies of water.
- **DO NOT** contaminate bodies of water when disposing of planting equipment wash water.
- Dispose of seed packaging or containers in accordance with local requirements.
- Excess treated seed may be used for ethanol production if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.

DISEASE	RATE FL. Oz./CWT	DIRECTIONS FOR USE
Soilborne and Seedborne Fusarium	0.071 (0.002 lb. a.i.)	Apply as a seed treatment using standard slurry or mist-type seed treatment equipment. Uniform application of seed is necessary to ensure seed safety and best disease protection. Seed must be sound and well cured prior to treatment. Product must be diluted with sufficient water to ensure complete seed coverage. Consult a seed treatment specialist regarding slurry rates specified for the crop to be treated with Willowood Tebucon 3.6SC . The length of control will vary depending on the rate used.
Soilborne and Seedborne Head smut (<i>Sphacelotheca reiliana</i>)	0.27 to 0.54 (0.0076 – 0.015 lb. a.i.)	

APPLICATION DIRECTIONS

CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Cotton	Southwestern cotton rust (<i>Puccinia cacabata</i>)	6 to 8 fl. oz. (0.14 – 0.225 lb. a.i.) per acre
	Application Instructions: Apply Willowood Tebucon 3.6SC in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control.	

For optimum disease control, tank mix the lowest labeled rate of a spray surfactant with **Willowood Tebucon 3.6SC**. **Willowood Tebucon 3.6SC** must have two to four hours of drying time on cotton foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, **Willowood Tebucon 3.6SC** will be resistant to weathering. **Willowood Tebucon 3.6SC** is a demethylation inhibitor (DMI) fungicide (Group 3).

Restrictions:

- **Willowood Tebucon 3.6SC** may be applied up to 30 days before harvest.
- **DO NOT** apply more than 24 fl. oz. (0.675 lb. a.i.) of **Willowood Tebucon 3.6SC** per acre per year.
- **DO NOT** apply more than 8 fl. oz. (0.225 lb. a.i.) of **Willowood Tebucon 3.6SC** per application.
- **DO NOT** make more than 3 applications per year at the maximum listed rate.
- **Restricted-entry interval (REI)** = 12 hours.
- Minimum retreatment interval is 7 days.

APPLICATION DIRECTIONS

CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Cucurbit Vegetables Group Chayote Chinese waxgourd Citron melon Cucumber Gherkin Edible gourd, (hyotan, cucuzza,	Powdery mildew (<i>Sphaerotheca fuliginea</i> / <i>Podosphaera xanthii</i>) (<i>Erysiphe cichoracearum</i>)	4 to 6 fl. oz. (0.113 – 0.14 lb. a.i.) per acre
	Gummy stem blight – Suppression (<i>Didymella bryoniae</i>) (watermelon, squash, pumpkin, and melons only)	8 fl. oz. (0.225 lb. a.i.) per acre

hechima, and Chinese okra) <i>Momordica</i> spp. (balsam apple, balsam pear, bitter melon, and Chinese cucumber) Muskmelon (cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon) Pumpkin Summer squash (crookneck squash, scallop squash, straightneck squash, vegetable marrow, and zucchini) Winter squash (butternut squash, calabaza, hubbard squash, acorn squash, and spaghetti squash) Watermelon	Application Instructions: Apply the specified dosage in a protective spray schedule to foliage and fruit. Repeat applications at 10- to 14-day intervals.
For optimum disease control, tank mix the lowest labeled rate of a spray surfactant with Willowood Tebucon 3.6SC . Willowood Tebucon 3.6SC must have two to four hours of drying time for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6SC will be resistant to weathering. Willowood Tebucon 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).	
Restrictions: <ul style="list-style-type: none"> Willowood Tebucon 3.6SC may be applied up to 7 days before harvest. DO NOT apply more than 24 fl. oz. (0.675 lb. a.i.) of Willowood Tebucon 3.6SC per acre per year. DO NOT apply more than 8.0 fl. oz. (0.225 lb. a.i.) of Willowood Tebucon 3.6SC per acre in a single application. Restricted-entry interval (REI) = 12 hours. Minimum retreatment interval is 10 days. DO NOT make more than 3 applications per year at the maximum listed rate. 	

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Dry bulb onion Garlic Great-headed (elephant) garlic Welch onion Shallot	White rot (<i>Sclerotium cepivorum</i>)	White rot: 20.5 fl. oz. (0.58 lb. a.i.) per acre applied in a 4-to-6-inch band over/into each furrow. May be applied by chemigation to control white rot.
	Rust (<i>Puccinia allii</i> , <i>Puccinia porri</i>)	4 to 6 fl. oz. (0.113 – 0.14 lb. a.i.) per acre
	Purple blotch (<i>Alternaria porri</i>)	
	White rot: For the control of white rot, make one application in the furrow at the time of planting. Apply the in-furrow application at the rate of 20.5 fl. oz. Willowood Tebucon 3.6SC per acre. Apply the entire per acre rate in a 4 to 6 inch band over/into each furrow. Additional control may be obtained by including two foliar applications at 4 to 6 fl. oz./acre (0.113 – 0.14 lb. a.i.).	
	Rust: For the control of rust make foliar applications at the rate of 4 to 6 fl. oz. Willowood Tebucon 3.6SC per acre per application. Repeat at an interval of 10 to 14 days. Apply Willowood Tebucon 3.6SC in a protective spray schedule or when weather conditions are favorable for rust development.	

For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest specified rate of a spray surfactant may be tank-mixed with **Willowood Tebucon 3.6SC**. **Willowood Tebucon 3.6SC** must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, **Willowood Tebucon 3.6SC** will be resistant to weathering. **Willowood Tebucon 3.6SC** is a demethylation inhibitor (DMI) fungicide (Group 3).

Restrictions:

- DO NOT** apply more than 32.5 fl. oz. (0.914 lb. a.i.) **Willowood Tebucon 3.6SC** per acre per year if an in-furrow treatment is made.
- If **Willowood Tebucon 3.6SC** is not applied as an in-furrow treatment, then **DO NOT** apply more than 12 fl. oz. (0.3375 lb. a.i.) **Willowood Tebucon 3.6SC** per acre per year as a foliar spray.
- DO NOT** apply within 7 days of harvest (PHI = 7 days).
- DO NOT** apply more than 20.5 fl. oz. (0.58 lb. a.i.) in a single application.
- Restricted-entry interval (REI)** = 12 hours.
- Minimum retreatment interval is 10 days.

- **DO NOT** make more than 5 applications per year at the maximum listed rate.

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Filbert (Hazelnut)[*]	Eastern filbert blight (<i>Anisogramma anomala</i>)	8 fl. oz. (0.225 lb. a.i.) per acre
<p>Application Instructions: Apply Willowood Tebucon 3.6SC in a preventive spray schedule when conditions are favorable for disease (including budbreak when the first green tissue is visible). Continue applications of Willowood Tebucon 3.6SC at 7- to 14-day intervals if weather conditions are conducive to disease development. Use shorter spray intervals during budbreak and rapid shoot elongation. The expanding shoot tip must be protected. Reduce the application interval for varieties that are highly susceptible to the indicated disease or when severe disease conditions exist. Apply Willowood Tebucon 3.6SC in sufficient spray volume for thorough coverage. Tank mixing the lowest labeled rate of a spray surfactant with Willowood Tebucon 3.6SC may improve spray coverage and penetration of the active ingredient into plant tissue. The use of ground application is preferred because of the difficulty in penetrating the canopy and obtaining thorough coverage of the foliage and stems by air. Willowood Tebucon 3.6SC is a sterol demethylation inhibitor (DMI) fungicide. It may be applied in tank mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy.</p> <p>For optimum disease control, tank mix the lowest labeled rate of a spray surfactant with Willowood Tebucon 3.6SC. Willowood Tebucon 3.6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6SC will be resistant to weathering. Willowood Tebucon 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3). It may be applied in a tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy.</p> <p>Restrictions:</p> <ul style="list-style-type: none"> • A maximum of 32 fl. oz. (0.9 lb. a.i.) of Willowood Tebucon 3.6SC may be applied per acre per year. • DO NOT cut cover crops in treated areas for feed or allow livestock to graze treated areas. • DO NOT apply within 35 days of harvest (PHI = 35 days). • DO NOT apply more than 8 fl. oz. (0.225 lb. a.i.) in a single application. • DO NOT make more than 4 applications per year. • Tank mixing with other DMI fungicides is not advised. • Restricted-entry interval (REI) = 12 hours. • Minimum retreatment interval is 7 days. <p>[*Not for use in California.]</p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Fruiting Vegetables Group (except Okra)[*] African eggplant Bush tomato Bell pepper Cocona Currant tomato Garden huckleberry Goji berry Groundcherry Martynia Naranjilla Pea eggplant Pepino Non-bell pepper Roselle Scarlet eggplant Sunberry Tomatillo Tomato Tree tomato Cultivars, varieties and/or hybrids of these	Early blight (<i>Alternaria solani</i>) Application Instructions: Apply Willowood Tebucon 3.6SC as a foliar spray using an interval of 7 days.	8 fl. oz. (0.225 lb. a.i.) per acre
<p>For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest specified rate of a spray surfactant may be tank-mixed with Willowood Tebucon 3.6SC. Willowood Tebucon 3.6SC must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6SC will be resistant to weathering. Willowood Tebucon 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).</p>		

Restrictions:

- **DO NOT** apply more than 48 fl. oz. (1.35 lb. a.i.) of **Willowood Tebucon 3.6SC** per acre per year.
- **DO NOT** apply within 7 days of harvest (PHI = 7 days).
- **Restricted-entry interval (REI)** = 12 hours
- **DO NOT** apply more than 8 fl. oz. (0.225 lb. a.i.) in a single application.
- Minimum retreatment interval is 7 days.
- **DO NOT** make more than 6 applications per year at the maximum listed rate.

[*Not for use in California.]

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Grasses Grown For Seed	Rusts (<i>Puccinia</i> spp.)	4 to 8 fl. oz. (0.113 – 0.225 lb. a.i.) per acre
	Apply the specified rate of Willowood Tebucon 3.6SC as soon as weather conditions are favorable for rust development or when first rust pustules are present. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl. oz./A (0.14 – 0.225 lb. a.i.) and shorter specified spray intervals.	
	Powdery mildew	4 to 8 fl. oz. (0.113 – 0.225 lb. a.i.) per acre
	Apply specified rate of Willowood Tebucon 3.6SC when powdery mildew first appears on the leaves. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl. oz./A (0.14 – 0.225 lb. a.i.) and shorter specified spray intervals.	

Apply the specified rate in a minimum of 20 gallons of water per acre with ground sprayers or in a minimum of 10 gallons of water per acre with aircraft. Thorough coverage is important for optimum disease control.

For optimum benefit, tank mix the lowest specified rate of a spray surfactant with **Willowood Tebucon 3.6SC**.

Restrictions:

- A maximum of 16 fluid ounces (0.45 lb. a.i.) may be applied per acre per year.
- **Willowood Tebucon 3.6SC** may be applied up to 4 days before harvest.
- Chaff, screenings, and straw from treated areas may be used for feed purposes; however, **DO NOT** forage, cut green crop, or use seed for feed purposes.
- Regrowth may be grazed starting 17 days after last application.
- **DO NOT** apply more than 8 fl. oz. (0.225 lb. a.i.) in a single application.
- **DO NOT** make more than 2 applications per year.
- **Restricted-entry interval (REI)** = 12 hours.
- Minimum retreatment interval is 14 days.

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Green onion Leek Spring onion Scallion Japanese bunching onion Green shallots Green eschalots	White rot (<i>Sclerotium cepivorum</i>) – Suppression only Rust (<i>Puccinia allii</i> , <i>Puccinia porri</i>) Purple blotch (<i>Alternaria porri</i>)	4 to 6 fl. oz. (0.113 – 0.14 lb. a.i.) per acre
For the control of diseases make foliar applications using an interval of 10 to 14 days. Apply Willowood Tebucon 3.6SC in a protective spray schedule or when weather conditions are favorable for rust development.		

For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest specified rate of a spray surfactant may be tank-mixed with **Willowood Tebucon 3.6SC**. **Willowood Tebucon 3.6SC** must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, **Willowood Tebucon 3.6SC** will be resistant to weathering. **Willowood Tebucon 3.6SC** is a demethylation inhibitor (DMI) fungicide (Group 3).

Restrictions:

- **DO NOT** apply more than 24 fl. oz. (0.675 lb. a.i.) **Willowood Tebucon 3.6SC** per acre per year.
- **DO NOT** apply within 7 days of harvest (PHI = 7 days).
- **DO NOT** apply more than 6 fl. oz. (0.14 lb. a.i.) in a single application.
- **Restricted-entry interval (REI)** = 12 hours.
- Minimum retreatment interval is 10 days.
- **DO NOT** make more than 4 applications per year at the maximum listed rate.

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Hops	Powdery mildew (<i>Sphaerotheca humuli</i> / <i>Sphaerotheca macularis</i>)	4 to 8 fl. oz. (0.113 – 0.225 lb. a.i.) per acre
Application Instructions: Apply the specified dosage in a protective spray schedule to foliage. Repeat		

applications at 10- to 14-day intervals. Increase the spray volume and the application rate as vine growth increases during the season.

For optimum disease control, tank mix the lowest labeled rate of a spray surfactant with **Willowood Tebucon 3.6SC**. **Willowood Tebucon 3.6SC** must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, **Willowood Tebucon 3.6SC** will be resistant to weathering. **Willowood Tebucon 3.6SC** is a demethylation inhibitor (DMI) fungicide (Group 3).

Restrictions:

- **Willowood Tebucon 3.6SC** may be applied up to 14 days before harvest.
- **DO NOT** apply more than 32 fl. oz. (0.9 lb. a.i.) of **Willowood Tebucon 3.6SC** per acre per year.
- **DO NOT** apply more than 8 fl. oz. (0.225 lb. a.i.) in a single application.
- **Restricted-entry interval (REI)** = 12 hours.
- Minimum retreatment interval is 10 days.
- **DO NOT** make more than 4 applications per year.

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Leafy Brassica Greens Broccoli raab Chinese cabbage (bok choy) Collards Kale Mizuna Mustard greens Mustard spinach Rape greens Turnip greens[*]	Cercospora leaf spot (<i>Cercospora brassicicola</i>) Powdery mildew (<i>Erysiphe cruciferarum</i>) Alternaria leaf spot (<i>Alternaria brassicicola</i>)	3 to 4 fl. oz. (0.084 – 0.113 lb. a.i.) per acre

For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest specified rate of a spray surfactant may be tank-mixed with **Willowood Tebucon 3.6SC**. **Willowood Tebucon 3.6SC** must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, **Willowood Tebucon 3.6SC** will be resistant to weathering. **Willowood Tebucon 3.6SC** is a demethylation inhibitor (DMI) fungicide (Group 3).

Restrictions:

- Application to turnip greens is limited to East of the Rockies.
- **DO NOT** apply more than 16 fl. oz. (0.45 lb. a.i.) **Willowood Tebucon 3.6SC** per acre per year.
- **DO NOT** apply more than 4.0 fl. oz. (0.113 lb. a.i.) of **Willowood Tebucon 3.6SC** per acre in a single application.
- **DO NOT** apply within 7 days of harvest (PHI = 7 days).
- **DO NOT** apply more often than once every 10 days.
- **Restricted-entry interval (REI)** = 12 hours.
- **DO NOT** make more than 4 applications per year.

[*Application to turnip greens is limited to East of the Rockies.]

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Garden beet roots and tops (leaves)	Cercospora leaf spot (<i>Cercospora beticola</i>)	3 to 7.2 fl. oz. (0.084 – 0.203 lb. a.i.) per acre

For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest specified rate of a spray surfactant may be tank-mixed with **Willowood Tebucon 3.6SC**. **Willowood Tebucon 3.6SC** must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, **Willowood Tebucon 3.6SC** will be resistant to weathering. **Willowood Tebucon 3.6SC** is a demethylation inhibitor (DMI) fungicide (Group 3).

Restrictions:

- Make applications on a 14-day interval.
- **DO NOT** apply more than 28.8 fl. oz. (0.81 lb. a.i.) **Willowood Tebucon 3.6SC** per acre per year.
- **DO NOT** apply more than 7.2 fl. oz. (0.203 lb. a.i.) of **Willowood Tebucon 3.6SC** per acre in a single application.
- **DO NOT** apply within 7 days of harvest (PHI = 7 days).
- **Restricted-entry interval (REI)** = 12 hours.
- **DO NOT** make more than 4 applications per year.

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Lychee	Anthrachnose (<i>Colletotrichum gloeosporioides</i>)	4 to 6 fl. oz. (0.113 – 0.14 lb. a.i.) per acre

Application Instructions: Begin first application of **Willowood Tebucon 3.6SC** as panicle emerges. Spray up to 6 fl. oz. (0.14 lb. a.i.) per acre every 10 days thereafter for a total of 8 sprays. Apply specified dosage in a minimum of 50 gallons of spray solution per acre by ground only.

For optimum disease control, tank mix the lowest labeled rate of a non-ionic spray surfactant with **Willowood Tebucon 3.6SC**. **Willowood Tebucon 3.6SC** must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, **Willowood Tebucon 3.6SC** will be resistant to weathering. **Willowood Tebucon 3.6SC** is a demethylation inhibitor (DMI) fungicide (Group 3).

Restrictions:

- **DO NOT** apply more than 48 fl. oz. (1.35 lb. a.i.) of **Willowood Tebucon 3.6SC** per acre per year.
- **DO NOT** apply more than 6.0 fl. oz. (0.14 lb. a.i.) of **Willowood Tebucon 3.6SC** per acre in a single application.
- **Willowood Tebucon 3.6SC** can be applied up to and including the day of harvest (PHI = 0 days).
- **Restricted-entry interval (REI)** = 2 days.
- Minimum retreatment interval is 10 days.
- **DO NOT** make more than 8 applications per year at the maximum listed rate.

APPLICATION DIRECTIONS

CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Okra	Cercospora leaf spot (<i>Cercospora</i> spp.)	4 to 6 fl. oz. (0.113 – 0.14 lb. a.i.) per acre
	Apply specific dosage of Willowood Tebucon 3.6SC in a preventative spray program. Use the highest rate when disease conditions are favorable and in areas where high disease pressure is expected. Applications may be repeated at 14-day intervals in order to maintain control of the disease. Apply specified dosage as a foliar spray in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air.	

For optimum disease control, tank mix the lowest labeled rate of a spray surfactant with **Willowood Tebucon 3.6SC**. **Willowood Tebucon 3.6SC** must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, **Willowood Tebucon 3.6SC** will be resistant to weathering. **Willowood Tebucon 3.6SC** is a demethylation inhibitor (DMI) fungicide (Group 3).

Restrictions:

- Applications may be made no closer than 3 days before harvest.
- **DO NOT** apply more than 24 fl. oz. (0.675 lb. a.i.) of **Willowood Tebucon 3.6SC** per acre per year.
- **DO NOT** apply more than 6.0 fl. oz. (0.14 lb. a.i.) of **Willowood Tebucon 3.6SC** per acre in a single application.
- **Restricted-entry interval (REI)** = 12 hours.
- minimum retreatment interval is 14 days.
- **DO NOT** make more than 4 applications per year at the maximum listed rate.

APPLICATION DIRECTIONS

CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Peanut	SOILBORNE: Sclerotium stem and pod rot (white mold, southern blight, southern stem rot) Rhizoctonia limb rot Rhizoctonia pod rot (Virginia and North Carolina only)	7.2 fl. oz. (0.203 lb. a.i.) per acre
	FOLIAR: Early leaf spot Late Leaf spot Leaf rust Web blotch (<i>Phoma</i>) Pepper spot (<i>Leptosphaerulina</i>)	
	FOUR-APPLICATION SPRAY PROGRAM: Apply the specified rate in a preventive spray schedule. See table below for proper timing of applications. Make applications of chlorothalonil prior to and following applications of Willowood Tebucon 3.6SC to discourage development of resistant strains of fungi. For optimum control of foliar diseases including leaf rust, web blotch, and pepper spot, tank mix the lowest label specified rate of a spray surfactant with Willowood Tebucon 3.6SC . LEAF SPOT ADVISORY SCHEDULE: For control of soilborne diseases in an advisory schedule, apply Willowood Tebucon 3.6SC in the first advisory spray in July and continue Willowood Tebucon 3.6SC applications at 14-day intervals. Tank mix applications after August 15 with chlorothalonil for resistance management purposes.	

DIRECTIONS: For optimum control of the specified soilborne diseases, four consecutive applications of **Willowood Tebucon 3.6SC** must be made at 14-day intervals.

Willowood Tebucon 3.6SC is a sterol demethylation inhibitor (DMI) fungicide. Chlorothalonil may be tank mixed at the rate specified on the tank mix label with **Willowood Tebucon 3.6SC** as a leaf spot resistance management strategy. A spray surfactant is not necessary when **Willowood Tebucon 3.6SC** is tank mixed with chlorothalonil. Mixing or alternating **Willowood Tebucon 3.6SC** with other DMI fungicides may lead to resistance.

Willowood Tebucon 3.6SC must be carried by rainfall or irrigation into the root and pod zone for control of root and pod rots caused by *Sclerotium rolfsii* and *Rhizoctonia solani*. Drought conditions will decrease the effectiveness of **Willowood Tebucon 3.6SC** against the root and pod rots.

Use **Willowood Tebucon 3.6SC** in conjunction with cultural practices that are known to reduce the severity of soilborne diseases, including proper crop rotation practices.

Restrictions:

- A maximum of 28.8 fl. oz. (0.81 lb. a.i.) of **Willowood Tebucon 3.6SC** may be applied per year.
- **DO NOT** apply more than 7.2 fl. oz. (0.203 lb. a.i.) of **Willowood Tebucon 3.6SC** per acre in a single application.
- **Willowood Tebucon 3.6SC** may be applied up to 14 days before harvest.
- **DO NOT** feed hay or threshings or allow livestock to graze in treated areas.
- **Restricted-entry interval (REI)** = 12 hours.
- **DO NOT** make more than 4 applications per year.
- Minimum retreatment interval is 14 days.

Timing of Willowood Tebucon 3.6SC Application for Optimum Control of White Mold and Rhizoctonia Limb and Pod Rot

Spray Program	Willowood Tebucon 3.6SC Application No.	Chlorothalonil Application No.
7 Applications	3,4,5 and 6	1,2 and 7

APPLICATION DIRECTIONS

CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Pecan	Brown leaf spot (<i>Sirosporium diffusum</i>) Downy spot (<i>Mycosphaerella caryigena</i>) Liver spot (<i>Gnomonia caryae</i>) Scab (<i>Cladosporium caryigenum</i>) Vein spot (<i>Gnomonia nerviseda</i>) Zonate leaf spot (<i>Grovesinia pyramidalis</i>)	4 to 8 fl. oz. (0.113 – 0.225 lb. a.i.) per acre
Application Instructions: Apply Willowood Tebucon 3.6SC in a preventive spray schedule beginning at early bud break (young leaves unfolding), and continue applications at 10- to 14-day intervals through the pollination period. Apply Willowood Tebucon 3.6SC at 4 fl. oz. (0.113 lb. a.i.) per acre in a tank-mix with the rate of registered products containing Fentin hydroxide in cover sprays. Follow label directions for the use of registered products containing Fentin hydroxide. DO NOT add a surfactant to the spray solution when tank-mixing Willowood Tebucon 3.6SC with SuperTin (EPA Reg. No. 70506-12). Apply Willowood Tebucon 3.6SC in a spray volume of 15 or more gallons per acre by air or 50 or more gallons per acre by ground. Apply 7 to 8 fl. oz. (0.197 - 0.225 lb. a.i.) per acre of Willowood Tebucon 3.6SC to full-size mature trees, and 4 to 6 fl. oz. (0.113 – 0.14 lb. a.i.) per acre of Willowood Tebucon 3.6SC to smaller trees. Apply the high rate to varieties that are highly susceptible to the indicated diseases, or when severe disease conditions exist. The lowest labeled rate of a surfactant may be added to the spray solution for optimum control of the indicated diseases.		

For optimum disease control, tank mix the lowest labeled rate of a spray surfactant with **Willowood Tebucon 3.6SC**. **Willowood Tebucon 3.6SC** must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, **Willowood Tebucon 3.6SC** will be resistant to weathering. **Willowood Tebucon 3.6SC** is a demethylation inhibitor (DMI) fungicide (Group 3). It may be applied in a tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy.

Restrictions:

- **DO NOT** apply after shucks begin to split.
- A maximum of 32 fl. oz. (0.9 lb. a.i.) of **Willowood Tebucon 3.6SC** may be applied per acre per year.
- **DO NOT** apply more than 8.0 fl. oz. (0.225 lb. a.i.) of **Willowood Tebucon 3.6SC** per acre in a single application.
- **DO NOT** cut cover crops in treated areas for feed or allow livestock to graze treated areas.
- **Restricted-entry interval (REI)** = 12 hours.
- **DO NOT** make more than 4 applications per year at the maximum listed rate.
- Minimum retreatment interval is 10 days.

APPLICATION DIRECTIONS

CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Pistachio[*]	Botryosphaeria panicle and shoot blight (<i>Botryosphaeria dothidea</i>)	8 fl. oz. (0.225 lb. a.i.) per acre

	<p>Application Instructions: Begin application at 25 to 50% bloom and repeat again 10 to 14 days later to protect young flower clusters and fruit. Make two additional applications of Willowood Tebucon 3.6SC 14 days apart beginning 49 days before harvest.[*] Apply Willowood Tebucon 3.6SC in a minimum spray volume of 15 gallons per acre by air or 50 gallons per acre by ground. Reduce the application interval for varieties that are highly susceptible to the indicated diseases or when severe disease conditions exist. The use of ground application after flowering is preferred because of difficulty in penetrating the canopy and obtaining thorough coverage of the foliage and fruit by air. Control of insect vectors and a thorough pruning program to remove plant tissue infected by <i>Botryosphaeria</i> are critical for optimum control of this disease.</p>
<p>For optimum disease control, tank mix the lowest labeled rate of a spray surfactant with Willowood Tebucon 3.6SC. Willowood Tebucon 3.6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6SC will be resistant to weathering. Willowood Tebucon 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3). It may be applied in a tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy.</p>	
<p>Precautions:</p>	
<ul style="list-style-type: none"> • Tank mixing with other DMI fungicides is not advised. 	
<p>Restrictions:</p>	
<ul style="list-style-type: none"> • A maximum of 32 fl. oz. (0.9 lb. a.i.) of Willowood Tebucon 3.6SC may be applied per acre per year. • DO NOT apply more than 8.0 fl. oz. (0.225 lb. a.i.) of Willowood Tebucon 3.6SC per acre in a single application. • DO NOT cut cover crops in treated areas for feed or allow livestock to graze treated areas. • DO NOT apply within 35 days of harvest (PHI = 35 days). • Restricted-entry interval (REI) = 12 hours. • DO NOT make more than 4 applications per year at the maximum listed application rate. • Minimum retreatment interval is 10 days. 	
<p>[*Not for use in California.]</p>	

APPLICATION DIRECTIONS		
CROP	DISEASE(S)	RATE OF WILLOWOOD TEBUCON 3.6SC
Soybean[*]	Rust (<i>Phakopsora pachyrhizi</i>) Powdery Mildew (<i>Microsphaera diffusa</i>)	3 to 4 fl. oz. (0.084 – 0.113 lb. a.i.) per acre
<p>Use Directions: Apply Willowood Tebucon 3.6SC as a broadcast foliar spray as a preventative spray or at first visible symptoms of disease. Repeat applications on a 10- to 14-day spray interval if environmental conditions are favorable for continued disease development. Use the higher specified rates and shorter spray intervals when disease pressure is severe. The lowest labeled rate of spray surfactant must be tank-mixed with Willowood Tebucon 3.6SC. Apply Willowood Tebucon 3.6SC in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons per acre by aircraft spray equipment.</p>		
<p>Restrictions:</p>		
<ul style="list-style-type: none"> • Applications may not be made within 21 days of harvest. • DO NOT apply more than 3 applications per year. • DO NOT apply more than 12 fl. oz. (0.3375 lb. a.i.) per year. • DO NOT apply more than 4.0 fl. oz. (0.113 lb. a.i.) of Willowood Tebucon 3.6SC per acre in a single application. • Restricted-entry interval (REI) = 12 hours. • Minimum retreatment interval is 10 days. • DO NOT make more than 3 applications at the maximum listed application rate. 		
<p>[*Not for use in New York State.]</p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Sunflower	Rust (<i>Puccinia 13elanthin</i>) Application Instructions: Apply specific dosage of Willowood Tebucon 3.6SC at the earliest sign of infection (rust pustules developing) or when weather conditions are favorable for rust development. Apply higher rate to highly susceptible varieties and/or under severe disease conditions. Application may be repeated at 14 days if necessary to maintain control of the disease. Apply specified dosage in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air.	4 to 6 fl. oz. (0.113 – 0.14 lb. a.i.) per acre
<p>For optimum disease control, tank mix the lowest labeled rate of a spray surfactant with Willowood Tebucon 3.6SC. Contact your State Extension Service or Willowood, LLC representative for a list of approved surfactants. Willowood Tebucon 3.6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6SC will be resistant to weathering. Willowood Tebucon 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).</p>		
<p>Restrictions:</p>		
<ul style="list-style-type: none"> • DO NOT apply more than 16 fl. oz. (0.45 lb. a.i.) of Willowood Tebucon 3.6SC per acre per year or within 50 days of harvest. 		

- **DO NOT** apply more than 6.0 fl. oz. (0.14 lb. a.i.) of **Willowood Tebucon 3.6SC** per acre in a single application.
- **Restricted-entry interval (REI)** = 12 hours.
- Minimum retreatment interval is 14 days.
- **DO NOT** make more than 2 applications per year at the maximum listed rate.

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Turnip (Application is limited to East of the Rockies)	Cercospora leaf spot (<i>Cercospora brassicicola</i>)	4 to 7.2 fl. oz. (0.113 – 0.203 lb. a.i.) per acre
	Application Instructions: Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 12- to 14-day intervals.	
For optimum disease control, tank mix the lowest labeled rate of a spray surfactant with Willowood Tebucon 3.6SC . Willowood Tebucon 3.6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6SC will be resistant to weathering. Willowood Tebucon 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).		
Restrictions: <ul style="list-style-type: none">• Willowood Tebucon 3.6SC may be applied up to 7 days before harvest.• DO NOT apply more than 28.8 fl. oz. of Willowood Tebucon 3.6SC per acre per year.• DO NOT apply more than 7.2 fl. oz. (0.203 lb. a.i.) of Willowood Tebucon 3.6SC per acre in a single application.• Restricted-entry interval (REI) = 12 hours.• Pre-Harvest interval is 7 days.		
DO NOT make more than 3 applications per year at the maximum listed rate. Minimum retreatment interval is 12 days.		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Tree Nut Group 14[*] (See separate use direction sections for Pecan, Almond, Pistachio and Filbert) Beechnut (<i>Fagus</i> spp.) Brazil nut (<i>Bertholletia excelsa</i>) Butternut (<i>Juglans cinerea</i>) Cashew (<i>Anacardium occidentale</i>) Chestnut (<i>Castanea</i> spp.) Chinquapin (<i>Castanea pumila</i>) Hickorynut (<i>Carya</i> spp.) Macadamia nut (bush nut) (<i>Macadamia</i> spp.) Walnut, black and English (Persian) (<i>Juglans</i> spp.) Cultivars, varieties, and/or hybrids of these	Brown rot blossom blight (<i>Monilinia taxa</i> , <i>M. fructicola</i>) Botryosphaeria panicle and shoot blight (<i>Botryosphaeria dothidea</i>) Eastern filbert blight (<i>Anisogramma anomala</i>)	8 fl. oz. (0.225 lb. a.i.) per acre
For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. Apply on a 7- to 14-day spray schedule. The lowest specified rate of a spray surfactant may be tank-mixed with Willowood Tebucon 3.6SC . Willowood Tebucon 3.6SC must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6SC will be resistant to weathering. Willowood Tebucon 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).		
Precautions:		
<ul style="list-style-type: none"> • Tank mixing with other DMI fungicides is not advised. 		
Restrictions:		
<ul style="list-style-type: none"> • DO NOT cut cover crops in treated areas for feed, or allow livestock to graze treated areas. • DO NOT apply more than 32 fl. oz. (0.9 lb. a.i.) Willowood Tebucon 3.6SC per acre per year. • DO NOT apply within 35 days of harvest (PHI = 35 days). • Restricted-entry interval (REI) = 12 hours. • DO NOT make more than 4 applications per year. • DO NOT apply more than 8 fl. oz. (0.225 lb. a.i.) in a single application. • Minimum retreatment interval is 7 days. 		
[*Not for use in California.]		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Wheat	Rusts leaf, stem, and stripe (<i>Puccinia</i> spp.) Head blight or scab (<i>Fusarium</i> spp.) – Suppression	4 fl. oz. (0.113 lb. a.i.) per acre

Application Instructions: Observe wheat fields closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development.

Application timing directions:

Rusts: Apply **Willowood Tebucon 3.6SC** at the earliest sign of rust pustules on foliage. Fusarium head blight: Optimal timing of **Willowood Tebucon 3.6SC** for Fusarium head blight suppression is the beginning of flowering on main stem heads (Feekes 10.51).

For optimum disease control, tank mix the lowest specified rate of a spray surfactant with **Willowood Tebucon 3.6SC**. **Willowood Tebucon 3.6SC** must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, **Willowood Tebucon 3.6SC** will be resistant to weathering. **Willowood Tebucon 3.6SC** is a demethylation inhibitor (DMI) fungicide (Group 3).

Restrictions:

- A maximum of 4 fl. oz. (0.113 lb. a.i.) of **Willowood Tebucon 3.6SC** may be applied per acre per year.
- **DO NOT** apply within 30 days of harvest.
- Straw may be fed or used for bedding.
- **DO NOT** allow livestock to graze or feed green forage to livestock prior to 6 days after treatment with **Willowood Tebucon 3.6SC**.
- Apply **Willowood Tebucon 3.6SC** in a minimum of 10 gallons of spray solution per acre by ground, or in a minimum of 5 gallons of spray solution per acre by air.
- **Restricted-entry interval (REI)** = 12 hours.
- **DO NOT** make more than one application per year.

OBSERVE THE FOLLOWING RESTRICTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS INCLUDING LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.

Apply only during alternate years in fields adjacent to aquatic areas listed above. **DO NOT** apply by ground or air within 100 feet of aquatic areas listed above.

DO NOT cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip.

Spray Drift Management: For aerial applications, mount the spray boom on the aircraft so as to minimize drift caused by wing tip vortices. Use the minimum practical boom length, and must not exceed 75% of the wing span or rotor diameter.

Use the largest droplet size consistent with pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure. Apply in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment.

Release spray at the lowest possible height consistent with good pest control and flight safety. Avoid applications more than 10 feet above the crop canopy.

Make aerial or ground applications when wind velocity favors on-target product deposition (approximately 3 to 10 mph). **DO NOT** apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.

DO NOT make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

DISEASE CONTROL IN GOLF COURSE TURF, AND FIELD, AND NURSERY AND CONTAINER ORNAMENTALS IN COMMERCIAL AND RESIDENTIAL LANDSCAPES

Chemigation: **DO NOT** apply this product through any type of irrigation system.

OBSERVE THE FOLLOWING RESTRICTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS INCLUDING LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES

- **DO NOT** apply within 100 feet of the aquatic areas listed above.
- **DO NOT** cultivate within 10 feet of an aquatic area to allow growth of a vegetation filter strip.
- See **Spray Drift Management** section for information.

Spray Drift Management

Make ground application when wind velocity favors on-target product deposition (approximately 3 to 10 mph). **DO NOT** apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperatures.

DO NOT make ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Spray Volume: For best results **Willowood Tebucon 3.6SC** may be applied in 66-132 gallons of water per acre for turf using ground-based equipment. For ornamentals, use 50-300 gallons of finished spray per acre depending upon the equipment, plant species and plant growth stage at time of application. For the most effective results, check equipment calibration regularly. When using lower spray volumes, be sure to maintain uniform application and full crop coverage so as to ensure effective control. Increase spray volume to ensure proper application, if required.

Compatibility Test for Mix Components:

Before mixing components, always perform a compatibility jar test. For 66 gallons per acre spray volume, use 5 cups of water in a clear, clean mixing jar. For other spray volumes adjust accordingly. Only use water from the intended source at the source temperature. Add components in the sequence indicated below in Mixing Order using 3 teaspoons for each pound of dry product or 1 ½ teaspoon for each pint of liquid product of directed label rate per acre. Always cap the jar and invert 10 cycles between component additions. When the components have all been added to the jar and fully mixed, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution must not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent and use the compatibility agent as directed on its label.

Mixing: Continuous agitation is required during mixing. When mixing this product and water, use the specified application rates as listed for each crop on this label. Before combining any other substances with the mixture, ensure that **Willowood Tebucon 3.6SC** is completely dispersed in the mixture.

Mixing Procedure:

1. Water. Add three-quarters of the required volume to a thoroughly clean sprayer tank.
2. Agitation. Start agitation and maintain constant agitation throughout mixing and application.
3. Inductor. If an inductor is used, rinse it thoroughly after each component has been added.
4. Products in PVA Bags. Place any product contained in water soluble PVA bags into the mixing tank. Wait until all water soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5. Water Dispersible Products. Including dry flowables (DF), wettable powders (WP), suspension concentrates (SC), or suspo-emulsions (SE).
6. Water-soluble products.
7. Emulsifiable concentrates (for example, oil concentrates when applicable).
8. Water soluble additives (for example, AMS or UAN when applicable)
9. Remaining quantity of water.

DISEASE CONTROL IN GOLF COURSE TURF

Turf Use Restrictions

- For use on golf course turf only.
- **DO NOT** use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (i.e., elementary, middle, and high school), campgrounds, churches, and theme parks.
- Not for residential use.
- Not for use on turf being grown for sale or commercial use as sod. **DO NOT** use clippings for animal feed.
- **DO NOT** exceed 3.6 fl. oz. (0.101 lb. a.i.) of **Willowood Tebucon 3.6SC** per 1,000 sq. ft. per year.
- **DO NOT** apply more than 6 applications per year.
- Minimum retreatment interval is 21 days.
- **DO NOT** apply more than 1.1 fl. oz. (0.03 lb. a.i.) in a single application.
- In the State of New York, **DO NOT** apply more than 3 applications per year.
- **DO NOT** exceed 1.8 fl. oz. (0.05 lb. a.i.) of this product per 1,000 sq. ft. (2.2 lbs. a.i./A) per year.

Product Information

For use on all golf turf applications of cool season and warm season grasses (including Bentgrasses, Bluegrasses, Fescues, Ryegrasses, St. Augustine grasses, and Zoysia) or their mixtures. **Willowood Tebucon 3.6SC** is not phytotoxic to any of the above-mentioned grasses when used in accordance with the label.

Bermudagrass can be sensitive to **Willowood Tebucon 3.6SC** under certain conditions. **DO NOT** apply consecutive applications during or just after dormancy break. Avoid applications when temperatures are expected to exceed 85°F.

Willowood Tebucon 3.6SC can be used for the prevention and control of the diseases mentioned in the table below. Begin applications when conditions favor disease development and repeat applications as long as these conditions persist.

Preventative treatments can be applied using 28-day intervals as indicated. When treating golf greens, always treat aprons and approaches. Spray uniformly over the area to be treated with properly calibrated equipment.

Apply the specified amount of **Willowood Tebucon 3.6SC** in sufficient water for thorough coverage. A volume of 66-132 gallons per acre (1.5-3.0 gallons per 1,000 sq. ft.) is advised. Apply using properly calibrated low volume, hand-held, mechanical, or motorized ground broadcast equipment. Application to small areas may be made with low-pressure hand wand or backpack equipment. Maintain constant agitation during application.

Depending on the disease, **Willowood Tebucon 3.6SC** must be watered into the crown and active root zone for best results. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. For best results use spray mixture the same day it is prepared.

Golf Course Turf Disease Control

DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC (Fl. Oz./1,000 Sq. Ft.)	APPLICATION INSTRUCTIONS
Dollar Spot (<i>Sclerotinia homoeocarpa</i>) Copper Spot (<i>Gloeocercospora sorghi</i>) Powdery Mildew (<i>Erysiphe graminis</i>) Corticium Red Tread (<i>Laetisaria fuciformis</i>) Rusts (<i>Puccinia</i> spp.) Red Thread (<i>Laetisaria fuciformis</i>) Pink Patch (<i>Limonomyces roseipellis</i>)	0.6-1.1 (0.016 - 0.03 lb. a.i.)	For prevention, begin applications when conditions are favorable for disease development using the low to medium rate. For curative treatments, use the medium to high rate. DO NOT make two consecutive applications of Willowood Tebucon 3.6SC or other DMI containing fungicides. Alternate with another fungicide with a different mode of action. A second application may be made after 21 days.
Brown Ring Patch (<i>Waitea circinata</i>)	0.6-1.1 (0.016 - 0.03 lb. a.i.)	For prevention, begin applications when conditions are favorable for disease development. Make two applications using the low to medium rate. Repeat application at 21-day interval.
Brown Patch/Rhizoctonia Blight, Large Patch (<i>Rhizoctonia solani</i>) Anthracnose - Basal and Foliar (<i>Colletotrichum cereal</i>)	0.6-1.1 (0.016 - 0.03 lb. a.i.)	For prevention, begin applications when conditions are favorable for disease development using the low to medium rate. For curative treatments, use the medium to high rate. DO NOT make two consecutive applications of this product or other DMI containing fungicides. Alternate with another fungicide with a different mode of action. A second application may be made after 21 days.
Bermuda Grass decline (<i>Gaeumannomyces graminis</i> var. <i>graminis</i>)	0.6-1.1 (0.016 - 0.03 lb. a.i.)	Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and root zone of the turf. The amount of water is dependent on the depth of root zone. For prevention, begin applications two or four weeks prior to the historical appearance of disease symptoms. Make two applications at the low to medium rate, 21 days apart or one application at the high rate. Alternate with another fungicide with a different mode of action as needed. Initiate cultural control practices at the same time the fungicide is applied. Refer to your local County Extension Service for this information.
Take-All Patch (<i>Gaeumannomyces graminis</i>)	0.6-1.1 (0.016 - 0.03 lb. a.i.)	For prevention, make two applications at the low rate, 21 days apart or one application at the high rate in the fall when root zone soil temperatures reach 75°F and again in the spring at 55-60°F root zone soil temperatures. Applications in both fall and spring may be necessary. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
Gray Leaf Spot (<i>Pyricularia grisea</i>)	0.6-1.1 (0.016 - 0.03 lb. a.i.)	For prevention, begin applications when conditions are favorable for disease development (July-Aug) using the low to medium rate. For curative treatments, use the medium to high rate. DO NOT make two consecutive

		applications of this product or other DMI containing fungicides. Alternate with another fungicide with a different mode of action. A second application may be made after 21 days.
Stipe Smut (<i>Ustilago striiformis</i>)	0.6-1.1 (0.016 - 0.03 lb. a.i.)	Make a single application to historical disease areas in spring as grass growth begins. For curative treatments, use the medium to high rate.
Spring Dead Spot (<i>Leptosphaeria korrae</i> , <i>L. narmari</i> , <i>Ophiosphaerella herpotricha</i> , <i>Gaeumannomyces graminis</i>) Necrotic Ring Spot (<i>Leptosphaeria korrae</i>)	0.6-1.1 (0.016 - 0.03 lb. a.i.)	For prevention, make two applications at the low to medium rate, 21 days apart or one application at the high rate in the fall when root zone soil temperatures reach 75°F. A spring application may be necessary in areas where disease pressure is known to be heavy. Apply using low to medium rate when root zone soil temperature reaches 55-60°F. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
Fusarium Patch (<i>Fusarium roseum</i>)	0.6-1.1 (0.016 - 0.03 lb. a.i.)	Apply first application in mid-June or 2-4 weeks prior to time this blight normally becomes evident. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone. Alternate with another fungicide with a different mode of action. A second application may be made after 21 days.
Summer Patch (<i>Magnaporthe poae</i>)	0.6-1.1 (0.016 - 0.03 lb. a.i.)	For prevention, apply beginning in the spring when root zone soil temperatures reach 55- 60°F. DO NOT make two consecutive applications of this product or other DMI containing fungicides. Alternate with another fungicide with a different mode of action. Make additional applications at 21-day intervals. See local university guidelines for suggested timing. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
Zoysia Patch, Large Patch of zoysia (<i>Rhizoctonia solani</i>)	0.6-1.1 (0.016 - 0.03 lb. a.i.)	Alternate with another fungicide with a different mode of action. Second and third applications may be made at 21-day intervals. See local university guidelines for additional information. An additional application in early spring may be necessary in areas where disease pressure is known to be heavy.
Gray Snow Mold/Typhula Blight (<i>Typhula incarnate</i>) Pink Snow Mold/Microdochium Patch (<i>Microdochium nivalis</i>)	0.6-1.1 (0.016 - 0.03 lb. a.i.)	Apply in the fall, before anticipated turf dormancy and before first snow cover. If turf breaks dormancy during winter months a second application may be made. DO NOT apply over a snow cover, or when turf is dormant. It is advised that Willowood Tebucon 3.6SC be tank-mixed with other registered snow mold products for best season long results.
Fairy Ring <i>Chlorophyllum</i> (<i>Lepiota</i>), <i>Lycoperdon</i> , <i>Marasmius</i>	0.6-1.1 (0.016 - 0.03 lb. a.i.)	For prevention in Cool season turf, make two applications at the low-medium rate in the spring when root zone soil temperatures reach 55-60°F. Make a second application using a 21-day interval. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone. DO NOT use a wetting agent unless hydrophobic soil conditions exist. For curative treatment, use the medium to high rate. For Warm season turf breaking dormancy, DO NOT make two consecutive applications of this product or other DMI containing fungicides. Alternate with another fungicide with a different mode of action. For hydrophobic areas, use an appropriate wetting agent to effectively penetrate the hydrophobic zone commonly created with this disease.

Apply the specified amount of **Willowood Tebucon 3.6SC** in 1.0 to 3.0 gallons of water per 1000 sq. ft. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. **DO NOT** use clippings for animal feed. **DO NOT** exceed 3.6 fl. oz. of **Willowood Tebucon 3.6SC** per 1,000 sq. ft. per year. **DO NOT** exceed 6 applications per year. In the State of New York, **DO NOT** apply more than 3 applications per year. **DO NOT** exceed 1.8 fl. oz. of this product per

1,000 sq. ft. (2.2 lbs. a.i./A) per year.

DISEASE CONTROL IN FIELD, AND NURSERY AND CONTAINER ORNAMENTALS IN COMMERCIAL AND RESIDENTIAL LANDSCAPES**Ornamental Use Restrictions**

- For use on ornamental plants only; not for use on woodlands or forest management. Intended for use by professional applicators.
- Minimum retreatment interval is 21 days.
- DO NOT** apply more than 10 fl. oz. (0.28 lb. a.i.) per acre in a single application.
- DO NOT** apply more than 40 fl. oz. of **Willowood Tebucon 3.6SC** (equal to 1.13 lbs. of tebuconazole) per acre per year.
- DO NOT** make more than 4 applications per year at highest rate. **DO NOT** apply to bearing fruit trees or vegetables.

Willowood Tebucon 3.6SC can be used in a preventative and curative disease control program for the listed plant types and disease in the table below. Optimum disease management is obtained when **Willowood Tebucon 3.6SC** is used in conjunction with sound disease management practices.

Apply material with properly calibrated hand-held, mechanical, or motorized spray equipment. Begin applications when disease first appears and repeat at 14-21 day intervals during the growing season. Use the shortest interval when conditions are unusually favorable for the development of disease. For hand-held, mechanical, or motorized applications, mix as directed below and apply as a full coverage spray to drip for the prevention and control of the diseases listed below. Choose a finished spray volume appropriate for the size of the plants and amount of foliage, which will provide thorough coverage throughout the canopy. Allow sprays to dry before overhead irrigation is applied.

Apply **Willowood Tebucon 3.6SC** at rates of 4-10 fl. oz. (0.113 – 0.28 lb. a.i.) per acre in 100 gallons of water. Spray volume may range from 50 up to 300 gallons of finished spray per acre depending upon equipment, plant species and plant growth stage at time of application.

The “**DIRECTIONS FOR USE**” of this product reflect the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied to any plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results. This product is not advised for use on African Violets, Begonias, Boston Fern, and Geraniums.

Ornamentals Disease Control

PLANTS	DISEASE	APPLICATION	
		To Prevent Diseases	To Treat Existing Disease
Roses	Black Spot Powdery Mildew Rust	Apply every 14-21 days during the growing season, starting when leaves first appear.	Apply every 14 days for a total of 3 applications beginning at the first sign of disease.
Flowers	Leaf Spot Powdery Mildew Rust Southern Blight	Apply at least 3 times per year, 14-21 days apart, beginning with Spring bud break. Rotation or Tank mixing with barrier protectant fungicides is advised for resistance management.	
Crabapples (Ornamental), Dogwoods, and Other Landscape (Ornamental) Trees	Anthrachnose Leaf Spot Powdery Mildew Rust Scab		
Azaleas, Camellias, Rhododendrons, and Other Landscape (Ornamental) Shrubs	Anthrachnose Black Spot Leaf Spot Petal Blight	Petal Blight – Apply 2-3 times per week into the flowers as they open and develop color.	
Ground Covers and Vines	Powdery Mildew Rust Southern Blight		
HOW MUCH TO USE FOR SMALL PLANTINGS: ADD 1 TEASPOON TO 2.5 GALLONS OF WATER.			
Restricted-entry interval (REI) = 12 hours			

Pump Style Sprayers

- Add the appropriate amounts of concentrate and water to the sprayer tank.
- Close the sprayer, shake well, and pressurize.
- Adjust nozzle to a coarse spray pattern and apply.
- Occasionally re-pressurize the sprayer if needed to maintain a good spray pattern.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store above 28°F or agitate before use.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

[NONREFILLABLE CONTAINERS]

Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying.

(Nonrefillable container ≤ 5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(Nonrefillable > 5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

[REFILLABLE CONTAINERS]

Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Willowood, LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Willowood, LLC and Seller harmless for any claims relating to such factors.

Willowood, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Willowood, LLC, and Buyer and User assume the risk of any such use. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, WILLOWOOD, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**










To the extent consistent with applicable law, neither Willowood, LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF WILLOWOOD, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF WILLOWOOD, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Willowood, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Willowood, LLC.

[Farmers First is a trademark of Farmers Business Network, Inc.]

[<BRAND>™ or ® is a trademark of <TRADEMARK HOLDER>.]

[All trademarks are the property of their respective owners.]

<p>[www.FBN.com][FBN.com] [Available at www.FBN.com] [844-200-FARM (3276)] [Farmers First]</p>		<p>{any color}</p> 
		
		
 <p>[FUNGICIDE] [Fungicide]</p>		

{Note to Reviewer: The mention of the product name throughout this Master Label may be updated/replaced with the term "This product" on the Market Label.}

[EPA approval date]

