



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

April 6, 2020

Anna Armstrong
Agent for Willowood, LLC
c/o Wagner Regulatory Associates, Inc.
PO Box 640
Hockessin, DE 19707

Subject: Label Amendment – Add water-soluble packaging formulation
Product Name: Willowood Propanil 80DF
EPA Registration Number: 87290-17
Application Date: November 14, 2018
Decision Number: 547438

Dear Ms. Armstrong:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2
EPA Reg. No. 87290-17
Decision No. 547438

with FIFRA section 6. If you have any questions, please contact Lydia Crawford by phone at 703-347-0622, or via email at Crawford.Lydia@epa.gov.

Sincerely,



Emily Schmid, Product Manager 25
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

PROPANIL	GROUP	7	HERBICIDE
----------	-------	----------	-----------

[Master Label]

Willowood Propanil 80DF; ABN: Willowood Propanil 80CHS

[Note to reviewer: language to be used when packaged in water soluble packaging:

Water Soluble Packaging

Do not sell the soluble bags individually.]

[Note to reviewer: Language to be used when packaged in the LOCK'nLOAD system: **LOCK'nLOAD.**]

For Postemergence Control of Broadleaf and Grass Weeds in Rice Fields

Active Ingredient:

Propanil: 3', 4'-Dichloropropionanilide81.00%

Other Ingredients:19.00%


TOTAL:100.00%

This product contains 0.8 lb. of active ingredient per pound of formulated product.

EPA Reg. No. 87290-17

EPA Est. No. _____

Net Contents: _____ [lbs.]



**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID
<p>IF ON SKIN OR CLOTHING:</p> <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
<p>IF IN EYES:</p> <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
<p>IF SWALLOWED:</p> <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
<p>IF INHALED:</p> <ul style="list-style-type: none"> • Move person to fresh air. • If person in not breathing, call 911 or an ambulance, then give artificial respiration, preferable mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment.</p>

Manufactured For:

Willowood LLC

385 Interlocken Crescent, Suite 240, Broomfield, CO 80021

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Caution. Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing.

Personal Protective Equipment (PPE)

Mixers, loaders, and other handlers exposed to the concentrate must wear:

- Coveralls over long-sleeve shirt and long pants
- Chemical-resistant gloves made of barrier laminate, nitrile rubber > 14 mils, butyl rubber > 14 mils, or Viton > 14 mils
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear, if overhead exposure
- Chemical resistant apron

Applicators and other handlers exposed to the dilute must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of barrier laminate, nitrile rubber > 14 mils, butyl rubber > 14 mils, or Viton > 14 mils

See **Engineering Controls** for additional requirements and options.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing or other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Engineering Controls:

When handlers use closed systems in a manner that meet the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(6)].

Human flagging prohibited. Flagging to support aerial application is limited to use of the Global Positioning System (GPS) or mechanical flaggers.

[Water soluble packets, when used correctly, qualify as a closed mixing/loading system under the Worker Protection Standard [40 CFR 170.240(d)(4)]. Handlers handling this product while it is enclosed in intact water soluble packets may elect to wear reduced PPE of long-sleeved shirt, long pants, shoes and socks instead of listed PPE.

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment break-down.]

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water except as specified on this label. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical prior to flooding may result in shallow groundwater contamination due to cracks in the subsoil of the rice paddy.

This product may contaminate water through runoff following rainfall events and by seepage through levees. This product has a high potential for runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Levees should be constructed with adequate time prior to chemical application so that they are compacted to reduce seepage and to hold a 3 to 6 inch flood.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on the label. If terms are unacceptable, return at once unopened.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Chemical-resistant gloves made of barrier laminate, nitrile rubber > 14 mils, butyl rubber > 14 mils, or Viton > 14 mils
- Chemical-resistant footwear plus socks
- Protective eyewear

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

[Keep in original packaging during storage. Store above freezing in a dry, well-ventilated area. Prolonged storage at temperatures below 0°F may cause the soluble bag to become brittle.]

Pesticide Storage: Do not use, pour, spill or store near heat or open flame. If the entire contents are not used, protect remaining material from moisture and heat. [Resealing by rolling top down and storage under roof are recommended.] Keep containers closed when not in use.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA region office for guidance.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then, offer for recycling, if available or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. [For products packaged in Water Soluble Bags (WSB), offer container and inner liner for recycling or dispose of them in the trash as long as the WSB have not broken in the container or liner. If WSB is broken, empty the remaining contents in the container and liner into the application equipment or a mix tank and then dispose of container and liner at an approved waste disposal facility.]

Container Handling Refillable Container: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Drum with Inner Liner: Nonrefillable container. Do not reuse or refill this container. Inner Liner: Completely empty removable liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Offer for recycling, if available or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

DRUMS: Offer for recycling if available or dispose of drum in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. If drum is contaminated and cannot be reused, dispose of in the same manner required for its liner.

SUPERSACK: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or by other procedures approved by state and local authorities.

Steps to be taken in Case Material is Released or Spilled: Sweep or shovel into containers for disposal or reworking. Keep dusting to a minimum. Flush contaminated area with a large amount of water to a chemical or sanitary sewer containing a settling pit.

Product Information

Willowood Propanil 80CHS for postemergence weed control in rice is formulated as an 80% active dry flowable formulation. Willowood Propanil 80CHS is not a hormone-type herbicide, but kills susceptible weeds by direct contact action. For this reason, thorough coverage of emerged weeds is essential for best results. Willowood Propanil 80CHS has no preemergence or residual herbicidal activity. Only weeds that have emerged and are exposed at time of application will be controlled. Apply Willowood Propanil 80CHS only to fields that have been drained of floodwater. Willowood Propanil 80CHS is most effective if applied when susceptible grasses and broadleaf weeds are small and growing actively under favorable soil

moisture and weather conditions. Early weed control removes weed competition from the rice crop, saves moisture, and generally contributes to increased yields.

[Note to reviewer: Language to be used when packaged in LOCK'nLOAD system.]

[Willowood Propanil 80CHS is packaged in LOCK'nLOAD® containers and must only be dispensed into the mix tank from the LOCK'nLOAD® container as directed below.

LOCK'nLOAD® is an EPA approved closed handling system consisting of an HDPE bottle with a male threaded valve attached to the bottle and a female valve assembly that is to be attached to the lid of the mix tank.

- Install female valve assembly on mix tank lid using template provided.
- Remove LOCK'nLOAD® container from pallet.
- Remove protective lid from male valve on the LOCK'nLOAD® container.
- Raise the protective cap on the female valve assembly.
- Invert LOCK'nLOAD® container placing male valve on top of female valve.
- Rotate mix tank container clockwise to empty contents.
- Once empty, rotate container counter clockwise to remove empty LOCK'nLOAD® container from the valve assembly.
- Close the protective cap lid on female valve assembly.
- Replace cap on LOCK'nLOAD® container and place back into pallet.
- Repeat process as needed.]

[LOCK'nLOAD® is a registered trademark of Amvac Chemical Corporation.]

Read Mixing and Equipment label instructions before application. When tank mixing, always read all individual manufacturers' labels. In interpreting all labels for the tank mixture, the most restrictive situations must apply.

[Note to reviewer: Language to be used when packaged in water soluble bags.]

[Willowood Propanil 80CHS is packaged in 10 "Water Soluble Bags". Each water-soluble bag contains 2.5 lbs product (2.0 lbs active ingredient).

Do not touch water-soluble bags with wet gloves, as the bags will dissolve.

Do not open water-soluble bags.]

Chemigation: Do not apply this product through any type of irrigation system

WEED RESISTANCE MANAGEMENT

Willowood Propanil 80CHS contains the active ingredient propanil. Propanil is classified as a Group 7 herbicide (amide chemical family) and is an inhibitor of photosystem II site A.

Herbicide resistance is defined as the inherited ability of a plant to survive and reproduce following exposure to a dose of herbicide normally lethal to the wild type. In a plant, resistance may be naturally occurring or induced by such techniques as genetic engineering or selection of variants produced by tissue culture or mutagenesis. Any weed population may contain or develop plants that are naturally resistant to Willowood Propanil 80CHS and other Group 7 herbicides. Weed species with acquired resistance to Group 7 herbicides may eventually dominate the weed population if Group 7 herbicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by Willowood Propanil 80CHS or other Group 7 herbicides.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of Willowood Propanil 80CHS or other Group 7 herbicides within a

growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.

- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Willowood LLC. In addition to the guidance above, registrants are encouraged to incorporate the appropriate elements of Best Management Practices from HRAC and WSSA on the label.

Contact your local sales representative, extension agent, or certified crop advisors to find out if suspected resistant weeds to this MOA have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions. Tank mix products so that there are multiple effective mechanisms of action for each target weed.

SPRAY DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator and the grower. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from ground and aerial applications to agricultural field crops.

- Apply only when the wind speed is less than or equal to 10 mph at the application site.
- Apply as a medium or coarser spray (ASABE standard 572.1).
- For ground applications, apply using a nozzle height of no more than 4 feet above the ground or crop canopy.
- For aerial applications, do not apply by air if drift can occur to sensitive non-target crops

or plants that are within 100 feet of the application site. Do not release spray at a height greater than 14 feet above the ground or crop canopy. The boom length must not exceed 70% of the wingspan or 85% of the rotor blade diameter. Do not make any type of application into temperature inversions. When applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

Where states have more stringent regulations, they must be followed.

The applicator must be familiar with and take into account the information covered in the following Aerial Drift Reduction Advisory.

AERIAL DRIFT REDUCTION ADVISORY

Information on Droplet Size: The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

Controlling Droplet Size:

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of Nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

Application Height: Do not make applications at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment: When applications are made with a cross wind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Increase swath adjustment increase with increasing drift potential (higher wind, smaller drops, etc.).

Wind: Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Do not apply at wind speeds below 2 mph due to variable wind direction and high inversion potential.

Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity: When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions: Do not apply during a local, low level temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of the smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas: The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

**For Use on Rice Grown in California and Rice Grown in Southern United States:
Arkansas, Florida, Louisiana, Mississippi, Missouri, South Carolina, Tennessee, and
Texas**

Restrictions

- **Preharvest Interval:** Do not apply this product within 60 days of rice harvest.
- **Chemigation:** Do not apply this product through any type of irrigation system.
- **Do not** apply more than a maximum of 7.5 lbs. of Willowood Propanil 80CHS (6 lbs. active ingredient) per acre in a single application or exceed 10 lbs. of Willowood Propanil 80CHS (8 lbs. of active ingredient) per acre total dosage per season.
- **In California:** Use Willowood Propanil 80CHS only where rice fields are completely drained or a minimal amount of water remains. If high water level is desired, re-flood field after 12 hours and before 7 days after treatment. This will discourage new weed infestations.
- **Do not** apply this product to any crop other than rice. Willowood Propanil 80CHS will cause injury to most crops except cereal grains and perennial grasses.
- **Do not** apply this product (directly or indirectly) to wild rice (*Zizania spp.*).
- Avoid drift or accidental application from turning aircraft on beans, cucurbits, cotton, soybeans, corn, safflower, seedling legumes, vegetables, orchards, vineyards, gardens, shrubs, and ornamentals. Once applied, Willowood Propanil 80CHS does not release fumes hazardous to nearby crops.
- Applications to fields where catfish farming is practiced and drain water from treated fields into areas where catfish farming is practiced is prohibited.
- **Do not** graze treated fields or feed treated forage within 60 days of the last application.
- **Do not** rotate treated land to other crops or transplant to crops other than rice for 60 days following treatment of this product.
- **Do not** apply this product within 14 days before or after carbamate or organophosphate insecticide applications. Otherwise, serious injuries to rice may occur.
- Water drained from treated rice fields must not be used to irrigate other crops or released within 1/2 miles upstream of a potable water intake in flowing water (e.g., river, stream, etc.) or within 1/2 miles of a potable water intake in a standing body of water, such as a lake, pond, or reservoir.
- **Do not** apply when weather conditions favor drift from area to be treated.

Emergency Release Provisions – California Only

Do not discharge water from treated rice paddies in California following treatment, unless

excessive rainfall completely submerges the rice crop and forces premature release, for:

- 7 days in dry seeded rice in California.
- 7 days for water-seeded rice in California.

Emergency Release Provisions – For Rice Grown in Southern United States: Arkansas, Florida, Louisiana, Mississippi, Missouri, South Carolina, Tennessee, and Texas

Water holding (discharge) intervals for flood water from treated rice paddies following treatment in the United States:

- For delayed flood (water-seeded) rice grown south of Interstate Highway 10 from the Texas/Louisiana border to Houston and east of State Highway 35 from Houston to Port Lavaca- Flood water must be held for 10 days after application, unless excessive rainfall completely submerges the rice crop and forces premature release. For Texas rice grown in areas north or west of these boundaries, the water holding interval will be 7 days.
- For delayed flood (water-seeded) rice in southern Louisiana south of Highway 14 – Flood water must be held for 15 days after propanil application, unless excessive rainfall completely submerges the rice crop and forces premature release. For delayed flood (water-seeded) rice in Louisiana, north of the Highway 14 boundary, the water holding interval is 7 days.
- For rice in CA and all other parts of the United States not mentioned above – Flood water must be held for 7 days after application unless excessive rainfall completely submerges the rice crop and forces premature release.

Weeds Controlled - California Only

Annual sedges	<i>Cyperus spp.</i>
Barnyardgrass ^a	<i>Echinochloa crus-galli</i>
Crabgrass	<i>Digitaria spp.</i>
Early watergrass ^{a,b}	<i>Echinochloa oryzoides</i>
Junglerice ^a	<i>E. colonum</i>
Late watergrass ^{a,b}	<i>Echinochloa phyllopogon</i>
Ricefield bullrush	<i>Scirpus mucronatus</i>
Rice flatsedge	<i>Cyperus iria</i>
Smallflower umbrella plant	<i>Cyperus difformis</i>

^aIn isolated instances, biotypes of barnyardgrass/watergrass may develop that cannot be effectively controlled by propanil alone. Where these biotypes are known or suspected to be present, and are found in a mixed weed population in which Willowood Propanil 80CHS is effective, tank mix Willowood Propanil 80CHS at labeled rate with other rice herbicides that are recommended for control of barnyardgrass/watergrass (up to the 3-leaf stage).

^bApplications to early and late watergrass made past the 4-leaf stage will result in partial control.

Weeds Controlled– For Rice Grown in Southern United States: Arkansas, Florida, Louisiana, Mississippi, Missouri, South Carolina, Tennessee, and Texas

Willowood Propanil 80CHS provides selective postemergence control of the following weeds in rice:

Annual sedges	<i>Cyperus spp.</i>
Barnyardgrass ^a	<i>Echinochloa crus-galli</i>
Beakrush (spearhead)	<i>Rhynchospora corniculata</i>
Broadleaf signalgrass	<i>Brachiaria platyphylla</i>

Crabgrass	<i>Digitaria</i> spp.
Curly dock	<i>Rumex crispus</i>
Foxtail	<i>Setaria</i> spp.
Goosegrass	<i>Eleusine indica</i>
Gulf cockspur	<i>Echinochloa crus-pavonis</i>
Hemp sesbania (coffeebean)	<i>Sesbania herbacea</i>
Hoorahgrass	<i>Fimbristylis miliaceae</i>
Junglerice ^a	<i>E. colonum</i>
Mexicanweed	<i>Caperonia castaneifolia</i>
Paragrass	<i>Panicum purpurascens</i>
Redroot pigweed	<i>Amaranthus retroflexus</i>
Redweed	<i>Melochia corchorifolia</i>
Rice flatsedge	<i>Cyperus iria</i>
Smallflower umbrella plant	<i>Cyperus difformis</i>
Spikerush (wiregrass)	<i>Eleocharis</i> spp.
Texas panicum	<i>Panicum texanum</i>
Watergrass ^a	<i>Echinochloa</i> spp.
Woolly croton	<i>Croton</i> spp.

^a In isolated instances, biotypes of barnyardgrass/watergrass may develop that cannot be effectively controlled by propanil alone. Where these biotypes are known or suspected to be present, and are found in a mixed weed population in which Willowood Propanil 80CHS is effective, tank mix Willowood Propanil 80CHS at labeled rate with other rice herbicides that are recommended for control of barnyardgrass/watergrass (up to the 3-leaf stage).

Timing and Dosage

Early Timing and Rates

Apply Willowood Propanil 80CHS when a satisfactory stand of rice has been established that will tolerate flooding. The amount of Willowood Propanil 80CHS to apply depends upon the growth stage and condition of the target weeds. Willowood Propanil 80CHS is most effective if applied when susceptible grasses and broadleaf weeds are small and actively growing under favorable soil moisture and weather conditions. Use a higher rate in the rate range for heavy weed infestations, weeds in advanced stages of growth, or when growing conditions are less than optimum. Emergency treatments made to weeds in advanced growth stages, such as when grass weeds are tillering, must occur at least 60 days before harvest.

For best results, apply Willowood Propanil 80CHS at the rate of 3.75 to 5 lbs. (3 to 4 lbs. active ingredient) per acre when the grasses are actively growing in the 1- to early 4-leaf stage. This rate will also control many seedling broadleaf and aquatic weeds. Generally, this will be 15 to 25 days after planting.

Mid-Timing and Rates

Willowood Propanil 80CHS can be applied at the rate of 5 to 7.5 lbs. (4 to 6 lbs. active ingredient) per acre to actively growing grasses in the 4- to 6-leaf and early tillering stage, or when they are in the 2- to 4-leaf stage but stressed under dry soil conditions. Generally, this will be 20 to 30 days after planting.

Use of Surfactants: The addition of a crop oil concentrate at 1 to 2 pints per acre, or other 80% active nonionic surfactant at a rate of 1 to 2 pints per 100 gallons of spray mixture, is recommended.

Rescue Timing and Rates

Apply Willowood Propanil 80CHS at the rate of 6.25 to 7.5 lbs. (5 to 6 lbs. active ingredient) in 15 gallons of spray per acre for emergency control of older tillering grass. Generally, this will be 30 to 40 days after planting. If the field is already flooded, the water should be lowered or

drained before spraying to expose more of the grass and weeds. Emergency treatment should be considered as a salvage operation only and cannot be relied upon for total control of grass and weeds.

[Note to reviewer: Language to be used when packaged in water soluble bags.]

[Water Soluble Packaging]

Product Table			
Use Rate Lbs. per Acre	Number Water Soluble Bags Required		
	10 Acre	20 Acre	30 Acre
3.75	15	30	45
5	20	40	60
6.25	25	50	75
7.5	30	60	90

1

Mixing Directions

Willowood Propanil 80CHS will disperse more quickly if water temperature is 50°F or warmer. Use only clean water for spraying. With the pump and agitator running, slowly add the specified amount of Willowood Propanil 80CHS into a partially filled mix tank. The jet or tank agitators must be positioned to create a rippling or rolling action on the liquid surface and to provide complete agitation at the bottom of the tank, preventing dead spots where the material can accumulate. A centrifugal pump is suggested to provide additional propeller shear action for dispersing and mixing this product. To avoid foaming, keep filling and bypass lines below the liquid surface. Willowood Propanil 80CHS must be completely dispersed and mixed prior to application.

[Language to be used when packaged in water soluble bags.]

[Mixing for Water Soluble Bags]

MIXING PROCEDURE:

1. Ensure that the sprayer is totally clean.
2. Fill the spray tank three quarters full with water. Engage gentle agitation.
3. Ensure the agitation system is working properly and that it creates a rippling or rolling action on the water surface.
4. Add the appropriate number of water-soluble bags of Willowood Propanil 80CHS directly into the spray tank. Do not touch water-soluble bags with wet gloves. Allow eight (8) minutes for complete mixing. The water-soluble bag may become brittle with age and exposure to cold temperatures. Longer mixing time may be required if the bag is brittle or if the water is cold.
5. Ensure Willowood Propanil 80CHS is completely in suspension before adding other tank mix partners.
6. Add tank mix partners in the following order: Water soluble, Emulsifiable concentrates, Water soluble additives (such as AMS or UAN when applicable)
7. Continue agitation while completing the filling of the sprayer.
8. Continuous agitation is required to keep Willowood Propanil 80CHS in suspension. Do not allow the spray mixture to stand without agitation.
9. Use the spray suspension as soon as it is prepared.
10. Do not mix, load or clean spray equipment where there is a potential to contaminate wells or aquatic systems.

NOTE: Growers using a sprayer with by-pass agitation should allow the water-soluble bags to completely dissolve before engaging the by-pass. Otherwise, undissolved bags could be sucked into the by-pass and plug the main screen.]

If a tank mixture is to be applied, always conduct a compatibility test prior to use by mixing proportional amounts of all spray ingredients in a test vessel (jar). The order of addition to water should be dry flowables or wettable powders first, flowables second, liquid formulations third, and crop oil concentrate last. Allow for each material to go into solution prior to the addition of the next material. Shake the mixture vigorously and allow it to stand for fifteen minutes. Rapid precipitation of the ingredients and failure to re-suspend when shaken indicates that the mixture is incompatible and should not be applied.

As each material is added to the spray mixture, always allow for complete mixing before adding the next ingredient. Add crop oil concentrate last and continue agitation while filling the mixing tank to the desired spray volume.

Do not add Willowood Propanil 80CHS directly to the spray tank of aircraft. Once properly dispersed in mix tank, pump spray mixture to aircraft spray tank (include rinsate from mix tank). To ensure uniformity of sprays, maintain good agitation throughout application.

Application Equipment

Aircraft

Fixed wing aircraft or helicopters should have well-designed spray systems that produce a uniform pattern of medium or coarser spray droplets. Apply Willowood Propanil 80CHS in no less than 10 gallons of total spray per acre with boom-nozzle sprayers. Increase volume to 12 to 15 gallons per acre for larger or denser stands of grass or during periods of low humidity.

The optimum effective spray swath width depends on operating conditions and type of aircraft being used. For uniform spray coverage with fixed-wing aircraft or helicopter, spray swath width should not exceed the width of wing span or rotor plus 10%. Measure the swaths accurately for flagging.

Ground Sprayers

Use standard low-pressure herbicide boom sprayers equipped with flat fan nozzles. Use nozzle sizes that deliver medium or coarser droplets in 15 to 20 gallons total spray per acre at 40 to 50 psi and at ground speeds not in excess of 3 to 4 mph. Adjust boom height so nozzle spray patterns meet uniformity. Avoid raising boom too high.

Flush all equipment with clear water after each day's use. Clean all equipment, including nurse tanks, used for Willowood Propanil 80CHS, with detergent wash followed by a water rinse, **before and after** spraying other pesticides or other crops.

Sprayer Cleanup

Before using equipment exposed to this product to treat another crop, clean the sprayer and any other equipment (loading hoses, batch tanks, etc.) using the following procedure:

1. Steam-clean tank using a non-chlorine-based detergent, taking care to remove all physical residues.
2. Thoroughly rinse sprayer, tanks, boom, and hoses with clean water (free of sediment and agricultural chemicals).
3. Fill the tank one-half full with clean water and add Nutrasol at 32 oz. per 100 gallons water. Fill the tank to capacity with clean water. Flush the nozzles, boom, and hoses, and agitate (and recirculate, if possible) the sprayer for 15 minutes. Drain the equipment, taking care to flush the boom and hoses thoroughly.
4. Rinse tanks, hoses and nozzles with clean water to remove Nutrasol.
5. Fill the tank one-half full with clean water and add 1 gallon 21% ammonia or 7 gallons 3% ammonia per 100 gallon water. Fill the tank to capacity with clean water. Flush the nozzles, boom, and hoses and agitate (and recirculate, if possible) the sprayer for 15

- minutes. Drain the equipment, taking care to flush the boom and hoses thoroughly.
6. Remove nozzles, screens, and strainers, and clean them separately.
 7. Rinse tanks, booms, and hoses with clean water.
 8. Repeat steps 5 and 7 an additional 3 times.
 9. Rinse tanks, booms, and hoses to remove all traces of ammonia.
 10. Water rinses may be applied to rice fields. Dispose of bleach rinses at an approved waste disposal facility.

NOTE: When applying multiple loads of this product several days in a row, the following procedure must be performed at the end of each day; partially fill the tank with fresh water, flush the boom and hoses, and allow to set overnight.

ATTENTION:

- Do not use chlorine bleach with ammonia. All traces of liquid fertilizer containing ammonia, ammonium nitrate or ammonium sulphate must be rinsed from the mixing and application equipment using water before adding chlorine bleach solution. Failure to do so will release a gas with a musty chlorine odor that can cause eye, nose, and throat and lung irritation.
- Do not clean equipment in an enclosed area.

Perform cleanup procedures on batch tanks and any other mixing equipment separately from aircraft hoppers. Take care to clean loading hoses and any other equipment or surfaces exposed to this product.

Crop Tolerance and Growing Conditions

All leading commercial varieties of rice are exceptionally tolerant to Willowood Propanil 80CHS. A temporary yellowing or tip burn of rice may be noted after treatment, but new growth is normal. Severe leaf burn and partial killing of rice may occur if the product is applied when rice is under stress and in a weakened growth condition due to disease or insect infestations, excessive soil salts, overwatering, or prolonged drought and extremely hot weather. Growers are cautioned not to spray under such conditions and/or when maximum daily temperatures have been or are expected to exceed 100°F.

Effect of Climatic Conditions and Cultural Practices on Weed Control

Field and Seedbed Preparation

Fields should be accurately leveled and contoured and have well-prepared seedbeds free of clods. Such conditions encourage uniform and rapid emergence of rice, grass and broadleaf weeds, allowing more accurate timing and coverage of Willowood Propanil 80CHS sprays for optimum weed control.

Water Management

Before application of Willowood Propanil 80CHS, drained or dry planted fields should be flushed as often as necessary to prevent drying and crusting. Flushing encourages uniform emergence and vigorous growth of grass, broadleaf weeds and rice, which is essential for optimum weed control. Flushing of fields should occur when weeds and rice are actively growing at time of treatment. Make sure the field is drained prior to treatment so that grasses and broadleaf weeds are fully exposed. Weeds that are partially submerged in standing water at time of application will not be satisfactorily controlled.

Treated fields should be flooded before a second infestation of grass develops. To prevent additional grass weed seed from germinating, rice fields should be flooded within 24 hours after spraying, or as soon as possible after 24 hours.

Temperature

The temperature a few days before and after applying Willowood Propanil 80CHS has an important effect on the weed-killing activity. The activity increases as daily maximum temperatures increase above 75°F and decreases as the daily maximum temperatures decline below 75°F. Do not apply Willowood Propanil 80CHS when maximum temperatures have been or are expected to stay below 65°F or exceed 100°F. Less than optimum temperature at time of application is not critical so long as the temperature exceeds 75°F during the day.

Relative Humidity and Rain

Grasses and weeds are more responsive to Willowood Propanil 80CHS during periods of high humidity when the foliage is moist or covered by dew. When the humidity is very low, spray tends to evaporate before reaching weed foliage. For best results under low relative humidity conditions, increase spray volume to 12 to 15 gallons per acre. **Do not** spray if rain is expected within 8 hours to avoid loss of deposited spray and herbicide adsorption by the weeds.

Wind

Do not apply when the wind speed exceeds 10 mph to avoid drift hazard to sensitive crops and the possibility of uneven (streaked) applications.

Compatibility with Other Chemicals

Tank mix applications of Willowood Propanil 80CHS with other herbicides, insecticides, spray adjuvants, or liquid fertilizers may reduce crop tolerance and/or weed control or impair mixing properties. Use of these products in tank mix applications with Willowood Propanil 80CHS is done at the user's risk.

Adverse Reaction to Insecticides

Rice plants may be severely injured or killed if Willowood Propanil 80CHS is applied in tank mix combinations or sequentially before or after certain insecticides. Do not combine Willowood Propanil 80CHS with carbamate insecticides such as carbaryl, etc., or organophosphorus insecticides (such as malathion and methyl parathion, etc.). Do not apply any of the carbamate or organophosphorus insecticides to rice fields within 14 days before or after Willowood Propanil 80CHS.

Do not apply Willowood Propanil 80CHS to rice fields planted with rice seed treated with bird repellents containing methiocarb. Consult local Extension specialist for current recommendations of approved insecticides on rice.

LIMIT OF WARRANTY AND LIABILITY**IMPORTANT: READ BEFORE USE**

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully.

However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Willowood LLC. To the extent consistent with applicable law,

all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, WILLOWOOD LLC MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. To the extent consistent with applicable law, no agent of Willowood LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, WILLOWOOD LLC DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT WILLOWOOD LLC'S ELECTION, THE REPLACEMENT OF PRODUCT.

[Language to be used when packaged in water soluble bags.]
[Water Soluble Packaging (WSP) Unit Label]

[Willowood Propanil 80CHS
81% Propanil: 3', 4'-dichloropropionanilide

CAUTION

Keep Out of Reach of Children
EPA Reg. No. – 87290-17
Refer to full label for complete
Directions for Use]