

87290-20

08/03/2012

1/22



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D C 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

Ms Lori Kohler, Agent  
Willowood, LLC  
c/o Wagner Regulatory Associates, Inc  
P O Box 640  
7217 Lancaster Pike, Suite A  
Hockessin, DE 19707

AUG - 3 2012

Subject Label Notification Per PR Notice 98-10,  
Action Adding Miami-Dade County Use Prohibition  
Product Name Willowood Fomesafen 1 88 SL  
EPA Registration Number 87290-20  
Submission Date May 22, 2011  
Decision Number 468164

Dear Ms Kohler

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration (PR) Notice 98-10 dated May 22, 2012 for the product referenced above. The Registration Division has conducted a review of this request for its applicability under PR Notice 98-10 and finds that the action requested falls within the scope of PR Notice 98-10. The label submitted with the application has been stamped "NOTIFICATION" and will be placed in our records.

If you have any questions regarding this letter, please contact Jasmine Branch at (703) 347-0351 or [branch.jasmine@epa.gov](mailto:branch.jasmine@epa.gov)

Sincerely,

*MICHAEL WALTER*

Kathryn V Montague *KVM*  
Product Manager (23)  
Herbicide Branch  
Registration Division (7505P)

Enclosure



United States  
**Environmental Protection Agency**  
 Washington DC 20460

**Registration**  
 Amendment  
 **Other**

OPP Identifier Number  
**NOTIFICATION**  
**AUG - 3 2012**

**Application for Pesticide - Section I**

1 Company/Product Number 87290 20	2 EPA Product Manager Kathryn Montague	3 Proposed Classification
4 Company/Product (Name) Willowood LLC/Willowood Fomesafen 1 88 SL	PM# 23	<input checked="" type="checkbox"/> None      Restricted
5 Name and Address of Applicant (Include Zip Code) Willowood LLC c/o Wagner Regulatory Associates Inc P O Box 640 Hockessin DE 19707 <input type="checkbox"/> Check if this is a new address	6 <b>Expedited Review</b> In accordance with FIFRA Section 3(c)(3) (b)(l) my product is similar or identical in composition and labeling to <b>NOTIFICATION</b> EPA Reg No Product Name <b>AUG - 3 2012</b>	

**Section - II**

<input type="checkbox"/> Amendment Explain below	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> Me Too Application
<input checked="" type="checkbox"/> Notification Explain below	<input type="checkbox"/> Other Explain below

**Explanation** Use additional page(s) if necessary (For Section I and Section II)  
 This notification is consistent with the provisions of PR Notice 98 10 and EPA regulations at 40 CFR 152 46 and no other changes have been made to the labeling or the confidential statement of formula of this product I understand that it is a violation of 18 U S C Sec 1001 to willfully make any false statement to EPA I further understand that if this notification is not consistent with the terms of PR Notice 98 10 and 40 CFR 152 46 this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA

**Section - III**

**1 Material This Product Will Be Packaged In**

Child Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No If Yes      No per Unit Packaging wgt      container	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No If Yes      No per Package wgt      container	2 Type of Container <input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) HDPE lined bags
3 Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container	4 Size(s) Retail Container 2 5 gallons	5 Location of Label Directions <input type="checkbox"/> On Label <input checked="" type="checkbox"/> On Labeling accompanying product	
6 Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other _____ <input checked="" type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			

**Section - IV**

**1 Contact Point (Complete items directly below for identification of individual to be contacted if necessary to process this application)**

Name Lori Kohler	Title Agent for Willowood LLC	Telephone No (Include Area Code) (302) 635 7281
<b>Certification</b> I certify that the statements I have made on this form and all attachments thereto are true accurate and complete I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law		6 Date Application Received  (Stamped)
2 Signature 	3 Title Agent for Willowood LLC	
4 Typed Name Lori Kohler	5 Date May 22 2012	



**WRA**

Wagner Regulatory Associates, Inc.  
P.O. Box 640  
7217 Lancaster Pike Suite A  
Hockessin, Delaware 19707

May 22, 2012

Document Processing Desk (NOTIF)  
ATTN: Kathryn Montague PM Team 23  
Registration Division  
U.S. Environmental Protection Agency  
Office of Pesticide Programs (7504P)  
Room 5-4900 One Potomac Yard  
2777 South Crystal Drive  
Arlington, Virginia 22202-4501

Dear Ms. Montague

**Subject: Notification of Addition of State Requested Language per PR Notice 98-10  
Willowood Fomcsafen 1 88 SL (EPA Reg No 87290-20)**

Wagner Regulatory Associates, Inc. as agent for Willowood LLC, submits the enclosed notification for the addition of state requested language for the above referenced product. The State of Florida requested that use in Miami-Dade County be restricted. No other changes were made to this label.

In support of this notification, the following documents are attached:

- Letter from Willowood LLC appointing Wagner Regulatory Associates, Inc. as its agent
- Application for Pesticide Notification (8570-1)
- Label with highlighted change and CD
- Certification with Respect to Label Integrity

If you have any questions about this submission, please contact the undersigned at [lori@wagnerreg.com](mailto:lori@wagnerreg.com) or (302) 635-7281.

Respectfully submitted,

Lori Kohler  
Agent for Willowood LLC

cc: [illegible]

4/22



# WILLOWOOD FOMESAFEN 1.88 SL

For Control of Weeds in Soybeans

Active Ingredient

Sodium salt of Fomesafen 5-[2-chloro-4-(trifluoromethyl)phenoxy]-N-(methylsulfonyl)-2-nitrobenzamide 22.1%\*

Other Ingredients 77.9%

---

Total 100.0%

\*Equivalent to 21.0% fomesafen or 1.88 lbs fomesafen active ingredient per gal

KEEP OUT OF REACH OF CHILDREN  
**WARNING/AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle  
(If you do not understand the label, find someone to explain it to you in detail)

See additional precautionary statements and directions for use in the attached booklet

EPA Reg No 87290-20

EPA Est No \_\_\_\_\_

Net Contents

2.5 gallons

\_\_\_\_\_ gallons (bulk)

Manufactured for  
Willowood LLC  
1600 NW Garden Valley Blvd Suite 120  
Roseburg, OR 97471

**NOTIFICATION**

**AUG - 3 2012**

cc  
cc  
cc  
cc  
cc  
cc  
cc  
cc  
cc  
cc  
cc  
cc  
cc

5/22

<b>FIRST AID</b>	
If on skin or clothing	<ul style="list-style-type: none"> <li>• Take off contaminated clothing</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes</li> <li>• Call a Poison Control Center or doctor for treatment advice</li> </ul>
If in eyes	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing</li> <li>• Call a Poison Control Center or doctor for treatment advice</li> </ul>
If swallowed	<ul style="list-style-type: none"> <li>• Call a Poison Control Center or doctor immediately for treatment advice</li> <li>• Have person sip a glass of water if able to swallow</li> <li>• Do not induce vomiting unless told to by a Poison Control Center or doctor</li> <li>• Do not give anything by mouth to an unconscious person</li> </ul>
If inhaled	<ul style="list-style-type: none"> <li>• Move person to fresh air</li> <li>• If person is not breathing call 911 or an ambulance then give artificial respiration preferably mouth-to-mouth, if possible</li> <li>• Call a Poison Control Center or doctor for further treatment advice</li> </ul>
<b>NOTE TO PHYSICIAN</b>	
Probable mucosal damage may contraindicate the use of gastric lavage	
Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment	
<b>HOT LINE NUMBER</b>	
For Emergency Information concerning this product call the National Pesticides Information Center (NPIC) at 1-800-858-7378 seven days a week 6 30 am to 4 30 pm Pacific Time, or your poison control center at 1-800-222-1222	

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**WARNING/AVISO**

This product contains fomesafen which has been determined to cause tumors in laboratory animals (mice) Risks can be reduced by closely following use directions and precautions and by wearing the protective clothing specified elsewhere on this label

**CAUSES SKIN IRRITATION CAUSES MODERATE EYE IRRITATION HARMFUL IF SWALLOWED OR ABSORBED THROUGH SKIN** Do not get on skin or on clothing Avoid breathing vapor or spray mist Avoid contact with eyes Prolonged or repeated skin contact may cause allergic reactions in some individuals **WASH THOROUGHLY WITH SOAP AND WATER AFTER HANDLING**

**Personal Protective Equipment (PPE)**

Some materials that are chemical-resistant to this product are listed below If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart

Applicators and other handlers must wear

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves such as barrier laminate, nitrile rubber, neoprene rubber, or Viton®
- Chemical-resistant footwear plus socks
- Chemical-resistant apron when cleaning equipment, mixing or loading

In addition for aerial applications mixers and loaders handling more than 150 gallons of Willowood Fomesafen 1 88 SL in any single work day must wear

Dust/mist filtering NIOSH-approved respirator with any N, R, P, or HE filter

### User Safety Recommendations

#### Users should

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

- Remove PPE immediately after handling this product. Wash the outside of gloves before removing.

As soon as possible, wash thoroughly and change into clean clothing.

#### Environmental Hazards

Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters. Do not apply when weather conditions favor drift from target area.

This chemical is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

### CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Willowood, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold WILLOWOOD, LLC and Seller harmless for any claims relating to such factors.

WILLOWOOD, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law, (1) this warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or WILLOWOOD, LLC and (2) Buyer and User assume the risk of any such use. TO THE EXTENT PERMITTED BY APPLICABLE LAW, WILLOWOOD, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL. To the extent permitted by applicable law, in no event shall WILLOWOOD, LLC be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF WILLOWOOD, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF WILLOWOOD, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

WILLOWOOD, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of WILLOWOOD, LLC

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves such as barrier laminate, nitrile rubber, neoprene rubber, or Viton
- Chemical-resistant footwear plus socks

### **INFORMATION**

Read all label directions before using.

Willowood Fomesafen 1 88 SL is a selective herbicide which may be applied preplant, preemergence or postemergence for control or suppression of broadleaf weeds, grasses and sedges in soybeans.

Willowood Fomesafen 1 88 SL is generally most effective and consistent when used postemergence, working through contact action. Therefore, emerged weeds must have thorough spray coverage for effective control. Some bronzing, crinkling or spotting of soybean leaves may occur following a postemergent application, but soybeans soon outgrow these effects and develop normally.

Optimum weed control is achieved by postemergent applications of Willowood Fomesafen 1 88 SL to young actively growing broadleaf weeds that are not under stress from moisture, temperature, low soil fertility, mechanical or chemical injury.

Certain germinating broadleaf weeds, grasses and sedges may be controlled or suppressed by soil residual activity from either preplant, preemergent or postemergent applications if rainfall occurs shortly after application. The extent and consistency of soil activity is dependent upon soil characteristics, ground cover, amount of rainfall following application and the rate of Willowood Fomesafen 1 88 SL used.

**Information on Weed Resistance**

Naturally occurring biotypes of certain broadleaf species with resistance to this herbicide and related products (same mode of action) are known to exist. Selection of resistant biotypes, through repeated use of these herbicides, may result in control failures.

If poor performance cannot be attributed to adverse weather conditions or improper application methods, a resistant biotype may be present. In such a case, additional treatments with this herbicide or similar mode of action products are not recommended. Consult your local company representative or agricultural advisor for assistance.

**APPLICATION DIRECTIONS**

**Application Timing**

Best broad spectrum postemergence control of susceptible broadleaf weeds is obtained when Willowood Fomesafen 1 88 SL is applied early to actively growing weeds. This usually occurs 14 to 28 days after planting. Refer to the weed control tables for specific recommendations on weed growth stages and rates.

**Spray Additives**

Only spray additives cleared for use on growing crops under 40 CFR 180.1001 may be used in the spray mixture. For best broad spectrum postemergence control of susceptible broadleaf weeds in Regions 2,3,4 and 5 (see Regional Use Maps), Willowood Fomesafen 1 88 SL should be used with 1.0-2.5% v/v liquid nitrogen (28% or similar) or a minimum of 8.5 lbs ammonium sulfate per 100 gals, of spray volume.

**For Postemergence Applications Always Add One of the Following except in tank mix with products prohibiting spray additives - (See Tank Mix Directions for Use)**

**Crop Oil Concentrate (COC) or Methylated Seed Oil (MSO)** Use a nonphytotoxic COC or MOS containing 15-20% approved emulsifier at 0.5-1% v/v (2-4 qts/100 gals) of finished spray volume. COC or MSO can improve weed control but may slightly reduce crop tolerance.

**Nonionic Surfactant (NIS)** Use NIS containing at least 80% active ingredient at 0.25-0.5% v/v (2-4 qts/100 gals) of finished spray volume (Region 1 and East of Interstates 79 and 77 for Regions 2 and 3).

**Other Adjuvants** Adjuvants other than COC or NIS may be used providing the product meets the following criteria:

- 1. Contains only EPA exempt ingredients
- 2. Is nonphytotoxic to the target crop
- 3. Is compatible in mixture (May be established through a jar test)
- 4. Is supported locally for use with Willowood Fomesafen 1 88 SL on the target crop through proven field trials and through university and extension recommendations.

**Note** no adjuvants are needed for preplant or preemergence applications unless Willowood Fomesafen 1 88 SL is being used in a burndown.

**Recommended Mixing Order**

- 1. Fill spray tank with half the required amount of water and begin agitation \*
- 2. Add fertilizer (UAN, AMS)



- 3 Add dry pesticide formulations
  - 4 Add Willowood Fomesafen 1 88 SL
  - 5 Add liquid pesticide formulation
  - 6 Add adjuvant (MSO, COC or NIS)
  - 7 Add remainder of water and then maintain constant agitation
- \*Compatibility agent, 1 gal/500 gals, of water or 0.2% v/v, may be added as needed

**Ground Application**

Use sufficient spray volume and pressure to ensure complete coverage of the target. A minimum spray volume of 15 gals/A and 30-60 psi at the nozzle tip is recommended. On large weeds and/or dense foliage, use 60 psi and a minimum of 20 gals /A to ensure coverage of weed foliage.

The use of flat fan nozzles will result in the most effective postemergence application of Willowood Fomesafen 1 88 SL. The sprayer must be calibrated to provide the proper volume and rate per acre. In addition, the boom and nozzle height must be adjusted to provide complete coverage of target weeds.

**DO NOT USE FLOOD TYPE OR OTHER SPRAY NOZZLES, WHICH DELIVER COARSE, LARGE DROPLET SPRAYS**

**Band Applications**

Thorough weed coverage is important for postemergent control. Best coverage is obtained with a minimum of two nozzles, one directed to each side of the planted row. Application with a single nozzle directed over the top of the row is not recommended for postemergence applications but is suitable for preemergence applications. Cultivation of untreated areas may be needed following band applications. When making postemergence band applications and cultivating in the same operation, position nozzles in advance of the cultivation device. This will reduce dust in the spray area. Dust can intercept spray, reducing weed coverage, resulting in less than adequate weed control.

Calculate the amount of herbicide and water volume needed for postemergence band treatment by the following formulas:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast rate per acre} = \text{Band herbicide rate per acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast volume per acre} = \text{Band herbicide volume per acre}$$

**Aerial Application**

Use sufficient spray volume and pressure to ensure complete coverage of the target. A minimum of 5 gals/A of spray mixture should be applied with a maximum of 40 PSI pressure. When broadleaf weed foliage is dense, use a minimum of 10 gals/A to ensure coverage of weed foliage.

**DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM**

**Cultivation** Cultivation prior to application is not recommended. Cultivation may put weeds under stress, reducing weed control. Timely cultivation 1-3 weeks after applying Willowood Fomesafen 1 88 SL may assist weed control.

**Rainfastness** Willowood Fomesafen 1 88 SL requires a 1 hour rain-free period for best results when applied postemergence

### PRECAUTIONS

- A maximum of 1 6 pts of Willowood Fomesafen 1 88 SL (or a maximum of 0 375 lbs a /A of fomesafen from any product containing fomesafen) may be applied per acre per year in Region 1 (see Regional Use Map)
- A maximum of 1 6 pts of Willowood Fomesafen 1 88 SL (or a maximum of 0 375 lbs a /A of fomesafen from any product containing fomesafen) may be applied per acre in alternate years in Region 2 (see Regional Use Map)
- A maximum of 1 3 pts of Willowood Fomesafen 1 88 SL (or a maximum of 0 313 lbs a /A of fomesafen from any product containing fomesafen) may be applied per acre in alternate years in Region 3 (see Regional Use Map)
- A maximum of 1 pt of Willowood Fomesafen 1 88 SL (or a maximum of 0 25 lbs a /A of fomesafen from any product containing fomesafen) may be applied per acre in alternate years in Region 4 (see Regional Use Map)
- A maximum of 0 75 pt of Willowood Fomesafen 1 88 SL (or a maximum of 0 1875 lbs a /A of fomesafen from any product containing fomesafen) may be applied per acre in alternate years in Region 5 (see Regional Use Map)
- Thoroughly clean the spray system with water and a commercial tank cleaner before and after each use
- Tank mixes of Willowood Fomesafen 1 88 SL with other pesticides, fertilizers or any other additives except as specified on this label or other approved Willowood supplemental labels may result in tank mix incompatibility, unsatisfactory performance and/or unsatisfactory crop injury
- Apply postemergence to actively growing weeds Avoid applying Willowood Fomesafen 1 88 SL to weeds or soybeans which are under stress from moisture, temperature low soil fertility, mechanical or chemical injury, as reduced weed control and/ or increased crop injury may result
- Avoid overlapping spray swaths, as injury may occur to rotational crops
- To provide adequate coverage, it is recommended that ground speed not exceed 10 MPH during application
- Do not graze treated areas or harvest for forage or hay
- Do not apply within 45 days of soybean harvest

### ROTATIONAL CROP RESTRICTIONS

The following rotational crops may be planted after applying Willowood Fomesafen 1 88 SL at recommended rates in soybeans

Crop To Be Planted	Minimum Rotation Interval (Months After Last Willowood Fomesafen 1 88 SL Application)
Dry beans, snap beans, soybeans and cotton	0
Small grains such as wheat, barley, rye	4
Corn*, peanuts, peas, rice, seed corn	10
To avoid crop injury do not plant alfalfa, sunflowers, sugar beets sorghum** or any other crop within	18

- \* Use 12 month minimum rotation interval for popcorn in the states of Ohio, Kentucky, Illinois, Indiana, Iowa and Region 4 when applied at a rate of 1.0 pt/A or more
- \* Use 18 month minimum rotation interval for sweet corn in the states of Connecticut, Maine, Massachusetts, New Hampshire, New York, Rhode Island, Vermont and Region 5
- \*\* Sorghum may be planted back after 10 months in Region 1

Do not graze rotated small grain crops or harvest forage or straw for livestock


**Replanting**

If replanting is necessary in fields previously treated with Willowood Fomesafen 1 88 SL the field may be replanted to cotton, dry beans, snap beans or soybeans. Do not apply a second application of Willowood Fomesafen 1 88 SL or other fomesafen-containing product as crop injury or illegal residues may occur in harvested crops. If tank-mix combinations were used, refer to product labels for any additional replanting instructions.

**WILLOWOOD FOMESAFEN 1.88 SL - USE RATES AND WEEDS CONTROLLED**

REFER TO MAP FOR DEFINITION OF SPECIFIED GEOGRAPHIC REGIONS


WILLOWOOD FOMESAFEN 1 88 SL REGIONAL USE MAP



**REGION 1**

**(Maximum Rate 1 6 pts /A per year)**

**REGION 1** Includes the following states or portion of states where Willowood Fomesafen 1 88 SL may be applied Alabama, Arkansas, Florida (except Miami-Dade county), Georgia, Louisiana, Mississippi, Missouri (Counties of Bollinger, Butler, Cape Girardeau, Dunklin, Madison, Mississippi, New Madrid, Pemiscot, Perry Ripley, Scott, Stoddard and Wayne), North Carolina, Oklahoma (East of U S Highway 75 and East of Indian Nation Parkway), South Carolina, Tennessee and Texas (all areas East of U S Highway 77 to State Road 239, including all of Calhoun County)



**REGION 2**

**(Maximum Rate 1 6 pts /A, alternate years)**


**REGION 2** Includes the following states or portion of states where Willowood Fomesafen 1 88 SL may be applied Delaware, Kentucky, Maryland, Virginia and West Virginia South of Interstate 70 in the following states Illinois, Indiana and Ohio and in Pennsylvania (all areas South of Interstate 80 to the intersection of U S Highway 15 and East of U S Highway 15 and U S Highway 522)



**REGION 3**

**(Maximum Rate 1 3 pts /A, alternate years)**


**REGION 3** Includes the following states or portion of states where Willowood Fomesafen 1 88 SL may be applied Connecticut, Iowa, Maine, Massachusetts, Missouri (all counties except for those listed in Region 1), New Hampshire, New Jersey, New York, Pennsylvania (all areas except those listed in Region 2), Rhode Island, Vermont, Wisconsin (South of U S Highway 18 between Prairie Du Chien and Madison, and South of Interstate 94 between Madison and Milwaukee) and North of Interstate 70 in the following states Illinois, Indiana and Ohio



**REGION 4**

**(Maximum Rate 1 pt/A, alternate years)**

**REGION 4** Includes the following states or portion of states where Willowood Fomesafen 1 88 SL may be applied Kansas (all counties East of or intersected by U S Highway 281), Michigan (Southern Peninsula), Minnesota (all areas South of Interstate 94) Nebraska (all counties East of or intersected by U S Highway 281), and Wisconsin (all areas except those in Region 3, South of Interstate 94 from Minnesota state line to Eau Claire and South of U S Highway 29 from Eau Claire to Green Bay plus Barron, Chippewa, Clark, Door, Dunn, Eau Claire, Kewaunee, Marathon, Menominee, Oconto, Polk, Shawano, and St Croix counties) The following counties are excluded Adams, Marquette, Portage, Waupaca, Waushara and Wood) North Dakota (all areas East of Interstate 29 from Fargo South to the South Dakota state line), South Dakota (all areas East of Interstate 29 from the North Dakota state line to Watertown, all areas East of Highway 81 from Watertown to Madison and all areas East and South of State Road 34 and U S Highway 281 to the Nebraska state line)



14/22

**REGION 5****(Maximum Rate 0 75 pts /A, alternate years)**

**REGION 5** Includes the following states or portion of states where Willowood Fomesafen 1 88 SL may be applied North Dakota (all areas East of U S Highway 281 except those areas in Region 4), South Dakota (all areas East of U S Highway 281 except those areas in Region 4) and Minnesota (all areas South of U S Highway 2 except those areas in Region 4)

**APPLICATION RATES FOR WEED GROWTH STAGES**

Weed	Willowood Fomesafen 1 88 SL Rate (pts /A) Maximum Growth Stage Controlled At			
	3/4 pts /A # of True Leaves	1 pt/A # of True Leaves	1 25 pts /A # of True Leaves	1 5 pts /A # of True Leaves
Anoda, Spurred	—	2*	2	4
Balloonvine	—	—	2	4
Carpetweed	—	8" Diameter Size	Unlimited Size	Unlimited Size
Citron (Wild Watermelon)	—	2	4	4
Cocklebur, Common	2	4	6	8
Copperleaf, Hophornbeam	—	4	4	6
Copperleaf, Virginia	—	4	4	6
Crotalaria, Showy	—	6	6	8
Croton, Tropic	—	4	4	6
Cucumber, Volunteer	—	4	6	8
Eclipta	—	2	4	4
Groundcherry, Cutleaf	—	4	6	8
Hemp	—	4	6	6
Horsenettle	—	2*	4*	4*
Jimsonweed	4	6	8	8
Ladysthumb	2*	2	4	6
Lambsquarters, Common	2*	2*	2*	2*

15/22

Weed	Willowood Fomesafen 1 88 SL Rate (pts /A) Maximum Growth Stage Controlled At			
	3/4 pts /A # of True Leaves	1 pt/A # of True Leaves	1 25 pts /A # of True Leaves	1 5 pts /A # of True Leaves
Mexicanweed	—	2*	2	4
Morningglory				
Cypressvine	2	4	6	6
Entireleaf var	3*	3	4	5
Ivyleaf	3*	3	4	5
Purple Moonflower	3*	3	5	6
Red (Scarlet)	3*	3	6	6
Smallflower	3*	3	4	6
Pitted (Smallwhite)	4*	4	6	6
Tall (Common)	2*	2	3	5
Palmleaf (Willowleaf)	3*	3	6	6
Mustard, Wild	4	6	8	8
Nightshade, Black	2	4	6	6
Nutsedge, Yellow	—	—	*	*
Pigweed, spp				
Amaranth, Palmer	2	4	6	6
Amaranth, Spiny	2	2	4	6
Redroot	2	4	6	8
Smooth	2	4	6	6
Waterhemp, Common	2*	2	4	6
Waterhemp, Tall	2*	2	4	6
Poinsettia, Wild	-	2	4	6
Purslane, Common	-	Multi-Leaf 6" Diameter	Multi-Leaf 8" Diameter	Multi-Leaf 8" Diameter
Pusley, Florida	-	2	2	4
Ragweed, Common	4*	4	6	8
Ragweed, Giant	4*	4	6	8
Redweed	-	-	2*	3*
Sesbania, Hemp		8	12	12
Sicklepod		-	Cotyledon*	Cotyledon*
Sida, Prickly	-	2*	2	4
Smartweed, Pennsylvania	4*	4	6	6
Smellmelon	-	2	2	4
Spurge, Prostrate	-	-	1" Diameter*	2" Diameter*
Spurge, Spotted	-	-	2*	2*
Starbur, Bristly	-	4	4	6

16/22

Weed	Willowood Fomesafen 1 88 SL Rate (pts /A) Maximum Growth Stage Controlled At			
	3/4 pts /A # of True Leaves	1 pt/A # of True Leaves	1 25 pts /A # of True Leaves	1 5 pts /A # of True Leaves
Sunflower, Common	-	-	2	4
Velvetleaf	-	2	4	4
Venice Mallow	4	6	6	8
Witchweed	-	Multi-leaf Up to 7"	Multi-leaf Up to 10"	Multi-leaf Up to 10"
Yellow Rocket	4	4	6	8

\*Suppression Only

**SPECIAL USE DIRECTIONS FOR ADDITIONAL WEED PROBLEMS**

**Suppression of Annual Grasses**

The grasses listed below may be suppressed by postemergence applications and controlled or suppressed by preemergence applications of Willowood Fomesafen 1 88 SL at 1-1 5 pts/A Consult Use Rate Table for maximum rate in each region For full-season broad-spectrum annual grass control, Fusilade® DX or Fusion® herbicide should be used alone or in tank mix with Willowood Fomesafen 1 88 SL Consult tank mix section

- Barnyardgrass
- Broadleaf Signalgrass
- Crabgrass
  - Foxtail
  - Giant
  - Green
  - Yellow
- Goosegrass
- Johnsongrass Seedling
- Panicum Fall
- Panicum, Texas

**Suppression of Perennial Weeds**

Use of Willowood Fomesafen 1 88 SL at postemergence rates of 1-1 5 pts/A will aid in suppressing the above-ground portions of the weeds listed below until crop canopy can assist in suppression Perennial weeds continue to regrow from underground rootstocks even if above-ground foliage is temporarily controlled or retarded Even though Willowood Fomesafen 1 88 SL and crop competition can suppress perennial weeds for a growing season, the rootstocks will continue to live and reestablishment will occur in subsequent years

- Milkweed, Climbing
- Milkweed Honeyvine
- Bindweed, Field
- Bindweed, Hedge
- Trumpet creeper



**TANK MIX AND SEQUENTIAL APPLICATIONS FOR SOYBEANS**

Willowood Fomesafen 1 88 SL can be used sequentially or in tank mix with one or more of the following products Assure II®, Basagran®, Butyrac®, Classic®, FirstRate®, Fusilade DX, Fusion, Ignite®, Glyphosate (such as Touchdown®, Roundup®, Glyphomax™), Gramoxone® Inteon, Harmony®, Poast®, Poast Plus®, Pursuit®, Raptor®, Resource®, Scepter®, Select®, and Synchrony® STS®

Under certain conditions the mixture of Willowood Fomesafen 1 88 SL with one or more of the above mentioned broadleaf herbicides may cause a reduction in activity of any postemergence grass herbicide in the mixture

For sequential applications allow 2-3 days after the application of the grass herbicide before applying Willowood Fomesafen 1 88 SL or Willowood Fomesafen 1 88 SL mixtures Where Willowood Fomesafen 1 88 SL or the Willowood Fomesafen 1 88 SL mixture is applied first, apply the grass herbicide when grass weeds begin to develop new leaves (generally around 7 days)

- Tank mix applications can result in increased crop injury as compared to either product used alone
- Do not exceed 1 fl oz of Butyrac per acre in mixture with Willowood Fomesafen 1 88 SL
- Do not exceed 0.25 oz /A of Synchrony STS herbicide in the tank with labeled rates of Willowood Fomesafen 1 88 SL on non-STs varieties This tank mix can be applied postemergence to any soybean variety for additional broadleaf weed control Refer to the Synchrony STS label for more information and crop rotation restrictions
- Always read and follow the recommendations restrictions and limitations for all products whether used alone, sequentially or in a tank mix The most restrictive labeling of any product used applies

**GLYPHOSATE TOLERANT SOYBEAN TANK MIXES**

Willowood Fomesafen 1 88 SL at 6-12 oz /A, can be tank mixed with glyphosate products (such as Touchdown or Roundup) that are labeled for glyphosate tolerant soybeans for improved postemergence control of many weeds such as morning-glory spp , hemp sesbania, waterhemp, and black nightshade which are known to have tolerance to glyphosate but are susceptible to Willowood Fomesafen 1 88 SL

**FOLLOW THE RECOMMENDATIONS ON THE GLYPHOSATE PRODUCT LABEL FOR THE USE OF SPRAY ADDITIVES IN THIS TANK MIX**

Do not allow this tank mix to move off target as contact by even minute quantities can cause severe damage or death to any non-target vegetation

**Note** Postemergence application of this tank mix on soybean varieties which do not contain the glyphosate tolerant gene will result in severe crop injury or death of the soybean crop Always read and follow the recommendations, restrictions and limitations for all products used. The most restrictive labeling of any product applies

**AERIAL SPRAY DRIFT MANAGEMENT ADVISORY****SPRAY DRIFT MANAGEMENT**

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR. The interaction of many equipment-and-weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

- 1 The distance of the outer most nozzles on the boom must not exceed  $\frac{3}{4}$  the length of the wingspan or rotor.
- 2 Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the AERIAL DRIFT REDUCTION ADVISORY.

**AERIAL DRIFT REDUCTION ADVISORY**

[This section is advisory in nature and does not supersede the mandatory label requirements.]

**INFORMATION ON DROPLET SIZE**

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (See Wind, Temperature and Humidity, and Temperature Inversions).

**CONTROLLING DROPLET SIZE**

- **Volume** Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of Nozzles** Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lower drift.

**BOOM LENGTH**

For some use patterns, reducing the effective boom length to less than  $\frac{3}{4}$  of the wingspan or rotor length may further reduce drift without reducing swath width.

**APPLICATION HEIGHT**

Applications should not be made at a height greater than 10 ft above the top of the target plants unless a greater height is required for aircraft safety Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind

**SWATH ADJUSTMENT**

When applications are made with a crosswind, the swath will be displaced downwind Therefore, on the up and downwind edges of the field, the applicator should compensate for this displacement by adjusting the path of the aircraft upwind Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc )

**WIND**

Drift potential is lowest between winds speeds of 2-10 mph However, many factors, including droplet size and equipment type determine drift potential at any given speed Application should be avoided below 2 mph due to variable wind direction and high inversion potential NOTE Local terrain can influence wind patterns Every applicator should be familiar with local wind patterns and how they affect spray drift

**TEMPERATURE AND HUMIDITY**

When making applications in low relative humidity set up equipment to produce larger droplets to compensate for evaporation Droplet evaporation is most severe when conditions are both hot and dry

**TEMPERATURE INVERSIONS**

Applications should not occur during a temperature inversion because drift potential is high Temperature inversions restrict vertical air mixing which causes small suspended droplets to remain in a concentrated cloud This cloud can move in unpredictable directions due to the light variable winds common during inversions Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind They begin to form as the sun sets and often continue into the morning Their presence can be indicated by ground fog, however, if fog is not present inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion while smoke that moves upward and rapidly dissipates indicates good vertical air mixing

**SENSITIVE AREAS**

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g residential areas, bodies of water, known habitat for threatened or endangered species, nontarget crops) is minimal (e.g when wind is blowing away from the sensitive areas)

**APPENDIX**

Scientific names are listed for those weeds referred to in the Willowood Fomesafen 1 88 SL label

<b>COMMON NAME</b>	<b>SCIENTIFIC NAME</b>
Amaranth, Palmer	<i>Amaranthus palmeri</i>
Amaranth, Spiny	<i>Amaranthus spinosus</i>
Anoda, Spurred	<i>Anoda cristata</i>
Balloonvine	<i>Cardiospermum halicacabum</i>
Barnyardgrass	<i>Echinochloa crus-galli</i>
Bindweed, Field	<i>Convolvulus arvensis</i>
Bindweed, Hedge	<i>Calystegia sepium</i>
Broadleaf Signalgrass	<i>Brachiaria platyphylla</i>
Carpetweed	<i>Mollugo verticillata</i>
Citron (Wild Watermelon)	<i>Citrullus vulgaris</i>
Cocklebur, Common	<i>Xanthium strumarium</i>
Copperleaf, Hophornbeam	<i>Acalypha qstryifolia</i>
Copperleaf, Virginia	<i>Acalypha virginica</i>
Crabgrass	<i>Digitaria spp</i>
Crotalaria, Showy	<i>Crotalaria spectabilis</i>
Croton, Tropic	<i>Croton glandulosus</i>
Cucumber, Volunteer	<i>Cucumissativas</i>
Eclipta	<i>Eclipta prostrata</i>
Foxtail Giant	<i>Setaria faberi</i>
Foxtail, Green	<i>Setaria viridis</i>
Foxtail, Yellow	<i>Setaria pumila</i>
Goosegrass	<i>Eleusine indica</i>
Groundcherry, Cutleaf	<i>Physalis angulata</i>
Hemp	<i>Cannabis sativa</i>
Horsenettle	<i>Solanum carolinense</i>
Jimsonweed	<i>Datura stramonium</i>
Johnsongrass, Seedling	<i>Sorghum halepense</i>
Ladysthumb	<i>Polygonum persicaria</i>
Lambsquarters, Common	<i>Chenopodium album</i>
Mexicanweed	<i>Caperonia castaniifolia</i>
Milkweed, Climbing	<i>Sarcostemma cyanchoides</i>
Milkweed, Honeyvine	<i>Ampelamus albidus</i>
Morningglory, Cypressvine	<i>Ipomoea quamodit</i>
Entireleaf	<i>Ipomoea hederacea var integruscula</i>
Ivyleaf	<i>Ipomoea hederacea var hederacea</i>
Purple Moonflower	<i>Ipomoea turbinata</i>
Red (Scarlet)	<i>Ipomoea coccinea</i>
Smallflower	<i>Jacquemontia tamnifolia</i>
Pitted (Small White)	<i>Ipomoea lacunosa</i>
Tall (Common)	<i>Ipomoea purpurea</i>
Palmleaf (Willowleaf)	<i>Ipomoea wrightii</i>
Mustard, Wild	<i>Brassica kaber</i>
Nightshade, Black	<i>Solanum nigrum</i>

COMMON NAME	SCIENTIFIC NAME
Nutsedge, Yellow	<i>Cyperus esculentus</i>
Panicum, Fall	<i>Panicum dichotomiflorum</i>
Panicum, Texas	<i>Panicum texanum</i>
Pigweed, Redroot	<i>Amaranthus retroflexus</i>
Pigweed, Smooth	<i>Amaranthus hybridus</i>
Poinsettia, Wild	<i>Euphorbia heterophylla</i>
Purslane, Common	<i>Portulaca oleracea</i>
Pusley, Florida	<i>Mchardia scabra</i>
Ragweed, Common	<i>Ambrosia artemisiifolia</i>
Ragweed, Giant	<i>Ambrosia trifida</i>
Redweed	<i>Melochia corchorifolia</i>
Sesbania, Hemp	<i>Sesbania exaltata</i>
Sicklepod	<i>Cassia obtusifolia</i>
Sida, Prickly	<i>Sida spinosa</i>
Smartweed Pennsylvania	<i>Polygonum pennsylvanicum</i>
Smellmelon	<i>Cucumis melo</i>
Spurge, Prostrate	<i>Euphorbia humistrata</i>
Spurge, Spotted	<i>Euphorbia maculata</i>
Starbur, Bristly	<i>Acanthospermum hispidum</i>
Sunflower, Common	<i>Helianthus annuus</i>
Trumpet creeper	<i>Campsis radicans</i>
Velvetleaf	<i>Abutilon theophrasti</i>
Venice Mallow	<i>Hibiscus trionum</i>
Waterhemp, Common	<i>Amaranthus rudis</i>
Waterhemp, Tall	<i>Amaranthus tuberculatos</i>
Witchweed	<i>Striga asatica</i>
Yellow Rocket	<i>Barbarea vulgaris</i>

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal

**Prohibitions**

Open dumping is prohibited Do not reuse empty container

**Pesticide Storage**

Store above 32°F in original containers only If product solidifies, return to room temperature and agitate to reconstitute Keep container closed when not in use Do not store near food or feed In case of spill or leak on floor or paved surfaces, soak up with sand, earth or synthetic absorbent Remove to chemical waste area

**Pesticide Disposal**

Pesticide wastes are toxic Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance

**Container Handling [Less Than 5 Gallons]**

Non-refillable container Do not reuse or refill this container Offer for recycling if available Triple rinse container (or equivalent) promptly after emptying Triple rinse as follows Empty the remaining contents into application equipment or mix tank Drain for 10 seconds after the flow begins to drip Fill the container 1/4 full with water and recap Shake for 10 seconds Pour rinsate into application equipment or mix tank or store rinsate for later use and disposal Drain for 10 seconds after the flow begins to drip Repeat this procedure two more times Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration or, if allowed by state and local authorities, by burning If burned, stay out of smoke

**Container Handling [For Bulk and Mini-Bulk Containers]**

Refillable container Refill this container with pesticide only Do not use this container for any other purpose Cleaning the container before final disposal is the responsibility of the person disposing of the container Cleaning before refilling is the responsibility of the person refilling To clean container before final disposal empty the remaining contents from this container into application equipment or mix tank Fill the container about 10 percent full with water Agitate vigorously or recirculate water with the pump for 2 minutes Pour or pump rinsate into application equipment or rinsate collection system Repeat this rinsing procedure two more times Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities **CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER**

Fusilade®, Fusion®, Gramoxone®, Touchdown®, are trademarks of the Syngenta Group Company  
Basagran®, Poast®, Poast Plus®, Pursuit®, Raptor®, and Scepter® are trademarks of BASF Ag Products

Assure II®, Classic®, Harmony®, Synchrony® STS® and Viton® are trademarks of E I DuPont de Nemours & Co , Inc

Select® and Resource® are trademarks of Valent Chemical Co

Butyrac® is a trademark of Albaugh Inc

Roundup® is a trademark of Monsanto Company

FirstRate® and Glyphomax™ are trademarks of Dow Agro Sciences

Ignite® is a trademark of Bayer CropScience