



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
Washington, D.C. 20460

OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

NOV 10 2011

Leanne Pruett
Willowood, LLC
c/o Pyxis Regulatory Consulting, Inc.
4110 136th Street, NW
Gig Harbor, WA 98332

Subject: Supplemental Label (Application to Head Lettuce, Endive, Escarole,
Radicchio Greens Including Single Application and Split Application
and Chemigation)
Willowood Pronamide 3.3SC
EPA Reg. No. 87290-22
Application Dated November 1, 2011

Dear Ms. Pruett:

The supplemental labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable.

The supplemental labeling must be incorporated into the main product labeling within eighteen (18) months of the date of this letter.

A stamped copy of your label is enclosed for your records. You must submit one (1) copy of the final printed label before you release the product for shipment. Products shipped after eighteen (18) months from the date of this letter must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA §6(e). Your release for shipment of the product constitutes acceptance of these conditions.

If you have any questions, please contact Mindy Ondish at (703)605-0723 or at ondish.mindy@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Kable Bo Davis", written over a horizontal line.

Kable Bo Davis
Product Manager 25
Herbicide Branch
Registration Division (7505P)

RESTRICTED USE PESTICIDE

Because Pronamide has produced tumors in laboratory animals, this product is for retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR WILLOWOOD PRONAMIDE 3.3SC BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Willowood Pronamide 3.3 SC and this supplement.

Willowood Pronamide 3.3SC

EPA Reg. No. 87290-22

Expiration Date: This supplemental label expires on December 31, 2014 and must not be used or distributed after this date.

APPLICATION TO HEAD LETTUCE / ENDIVE / ESCAROLE / RADICCHIO GREENS INCLUDING SINGLE APPLICATION AND SPLIT APPLICATION AND CHEMIGATION

KEEP OUT OF REACH OF CHILDREN

CAUTION

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Willowood Pronamide 3.3SC is a selective herbicide for the control of certain annual grasses and broadleaf weeds in direct seeded or transplanted head lettuce, endive, escarole and radicchio greens.

Weeds Controlled

Willowood Pronamide 3.3SC is effective at 2.5 to 5.0 pints of product (1 to 2 lbs. active ingredient) per treated acre for the preemergence control of the following weeds:

ACCEPTED
NOV 10 2011
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 87290-22

Grasses

- barley, foxtail
- barley, volunteer
- barnyardgrass
- bluegrass, annual
- brome, downy (cheatgrass)
- canarygrass
- crabgrass
- foxtail, yellow
- goosegrass
- lovegrass
- oats, volunteer
- panicum, fall
- ryegrass, Italian
- rye, volunteer
- wheat, volunteer

Broadleaf Weeds

- carpetweed
- chickweed, common
- goosefoot, nettleleaf
- henbit
- knotweed
- lambsquarters, common
- mornignglory, annual
- mustard, wild
- nettle, burning
- nightshade, black
- nightshade, hairy
- purslane, common
- rocket, London
- shepherdspurse
- smartweed, pale
- tomato, volunteer

Dosage

Willowood Pronamide 3.3SC may be applied at the rate of 2.5 to 5.0 pints of product (1 to 2 lbs. active ingredient) per acre broadcast application. The dosage rate required is dependent on soil texture and method of irrigation. At rates specified on this label, Willowood Pronamide 3.3SC may not be applied for weed control on highly organic (peat and muck) soils.

For head lettuce, endive, escarole and radicchio greens, follow the dosage instructions listed in chart below:

Pints of Willowood Pronamide 3.3SC Per Broadcast Acre¹			
Weeds	Dependable Rainfall Or Overhead Irrigation	Less Dependable Rainfall or Furrow Irrigation	Soil Texture Group²
susceptible annual grasses	2.5-3.5 (surface application)	3.5-5.0 (soil incorporation)	Coarse and medium Textured soils
broadleaf weeds	3.5-5.0 (surface application)	5.0 (soil incorporation)	Fine textured soils

¹ Reduce dosage rate accordingly for banded applications.

² Soil Texture Group

Coarse: sand, loamy sand, sandy loam

Medium: loam, silt loam, silt, sandy clay loam

Fine: silty clay loam, clay loam, sandy clay, silty clay, clay

Crop Tolerance

Most varieties of head lettuce are highly tolerant of the specified rates of Willowood Pronamide 3.3SC. Do not use more than 3.75 pints of Willowood Pronamide 3.3SC on val temp, grande verde and prima verde varieties of crisp head lettuce, or on endive, escarole and radicchio greens.

Timing and Application

Willowood Pronamide 3.3SC can be applied either pre-plant, post-plant or postemergence to head lettuce, endive, escarole or radicchio greens in banded, bed-topped or broadcast applications. Most applications will be made preemergence to the crop just before or after planting and preemergence to the weeds. Applications can be made before or after thinning of head lettuce but must be made prior to weed emergence.

Split Application:

Willowood Pronamide 3.3SC application can be split so that part of the maximum application rate of the product can be initially applied to head lettuce, endive, escarole or radicchio greens, and the balance of the maximum application rate can be applied up to 10 days later. Total amount of Pronamide 3.3SC applied must not exceed the maximum rates indicated above, up to 5 pints of product (2 lb. active ingredient) per acre per crop season. An example of this split application would be an application of 1.25 pints product (1/2 lb active ingredient) per acre at planting and 3.75 pints product (1 1/2 lb active ingredient) per acre 7 to 10 days post plant.

Do not apply Willowood Pronamide 3.3SC to head lettuce within 55 days of harvest and do not make more than two applications to each crop of head lettuce, endive, escarole or radicchio greens.

Mix the specified amount of Willowood Pronamide 3.3SC in clean water and apply uniformly with a ground sprayer in 20 to 50 gallons of water per treated acre. Reduce dosage and volume accordingly for banded treatments. Use a standard low pressure sprayer equipped with flat fan nozzles that provide uniform spray distribution.

Application Moisture Requirements

Willowood Pronamide 3.3SC acts mainly through root absorption, therefore it is necessary to move Willowood Pronamide 3.3SC into the root zone of germinating weeds to provide effective control. This can be accomplished by overhead sprinkler irrigation, by rainfall or by shallow mechanical incorporation.

Sprinkler Irrigation

Willowood Pronamide 3.3SC can be applied to the soil surface without mechanical incorporation after planting or transplanting if overhead irrigation is used. An initial irrigation of 1 to 2 inches must promptly follow the application of Willowood Pronamide 3.3SC, especially in hot weather.

Applications Dependent on Natural Rainfall

In areas of dependable natural rainfall, Willowood Pronamide 3.3SC can be applied as a surface treatment preemergence to the weeds. Applications to direct seeded or transplanted head lettuce, endive, escarole or radicchio greens are most successful when followed by 1/2 to 1 inch of rainfall within two to three days after application.

Furrow Irrigation – Mechanical Incorporation

Where rainfall is not dependable or supplementary overhead irrigation is not used, shallow pre-plant incorporation is required. PTO-driven incorporators or rolling cultivators that thoroughly mix Willowood Pronamide 3.3SC into the top 2 inches of soil are suggested.

Incorporation must be simultaneous or immediately after application of Willowood Pronamide 3.3SC, especially in hot weather. Irrigation must be started as soon as possible.

Where furrow irrigation is used, spray application and mechanical incorporation must be made after beds have been formed. Willowood Pronamide 3.3SC will not be as effective if disced in prior to bed shaping.

Hoing, thinning or shallow cultivation of soil treated with Willowood Pronamide 3.3SC will not destroy its herbicidal activity.

Temperature

Willowood Pronamide 3.3SC is not highly volatile, but it may degrade rather quickly if left exposed on the soil surface in warm weather. If applied when air temperatures exceed 85°F it must be shallow incorporated or watered into the soil as soon as possible, preferably within 1 or 2 days.

Rotation Crops

Follow the directions given in the Product Information section of this label under Rotation Crop Planting Information.

Head Lettuce/Endive/Escarole/Radicchio Greens – Specific Use Restrictions

- Do not apply Willowood Pronamide 3.3SC to head lettuce, endive, escarole, radicchio varieties that will be harvested less than 55 days after treatment.
- Do not apply more than two applications of Willowood Pronamide 3.3SC to each crop of head lettuce, endive, escarole or radicchio greens.
- Do not apply Willowood Pronamide 3.3SC to leaf lettuce.
- Do not apply more than 5.0 pints of Willowood Pronamide 3.3 SC (2 lb active ingredient) per acre per crop season

**For Grass and Weed Control in Head Lettuce – Aerial Application
(For Use in Arizona and California)**

Willowood Pronamide 3.3SC may be applied by aircraft for preemergence control of susceptible grasses and broadleaf weeds in head lettuce. Willowood Pronamide 3.3SC must be applied at the dosage rate of 2.5 to 5.0 pints of product (1.0 to 2.0 pounds active ingredient) per treated acre depending on soil type (refer to comments under Dosage chart in **HEAD LETTUCE/ENDIVE/ESCAROLE/RADICCHIO GREENS** section). Mix the specified amount of this product in 10 to 20 gallons of water per acre for aerial application. Avoid drift to all other crops and non-target areas.

**Chemigation Application on Head Lettuce, Endive, Escarole or Radicchio
(For Use in Arizona and California)**

Apply Willowood Pronamide 3.3SC by chemigation for weed control in direct seeded or transplanted head lettuce, endive, escarole or radicchio. Application must be made prior to weed emergence. Application may be made preemergence to head lettuce, endive, escarole, or radicchio greens or postemergence to head lettuce. **Do not apply postemergence to endive, escarole, or radicchio greens.**

Application Rate: Apply Willowood Pronamide 3.3SC at the rate of 1.25 to 2.5 pints per acre (0.5 to 1 lb. active ingredient per acre) depending upon soil type, weed species and level of infestation.

Weeds	Pints Willowood Pronamide 3.3SC Per Acre Broadcast Application ¹	Soil Texture Group ¹
Susceptible annual grasses and broadleaf weeds	1.25-2.5 (Surface application)	Coarse and medium textured soils
	1.25-2.5 (Surface application)	Fine textured soils

¹Soil Texture Group

Coarse: sand, loamy sand, sandy loam

Medium: loam, silt loam, silt, sandy clay loam

Fine: silty clay loam, clay loam, sandy clay, silty clay, clay

Split Application:

Willowood Pronamide 3.3SC chemigation application can be split, so that part of the maximum application rate of the product can be initially applied to head lettuce, endive, escarole or radicchio greens, and the balance of the maximum application rate can be applied up to 10 days later. Total amount of Pronamide 3.3SC applied must not exceed 2.5 pints product (1 lb. active ingredient) per acre per crop season. The Pre-harvest interval (PHI) of 55 days must be observed. An example of this split application would be application of 0.625 pint product (1/4 lb. active ingredient) per acre at planting and 1.875 pints product (3/4 lb active ingredient) per acre 7 to 10 days post plant.

Application Moisture Requirements: Willowood Pronamide 3.3SC acts mainly through root absorption; therefore, it is necessary to move Willowood Pronamide 3.3SC into the root zone of germinating weeds to provide effective control. This can be accomplished by applying a minimum of 0.75 inch of overhead sprinkler irrigation when applied by chemigation to fields that have been pre-irrigated.

Time of Treatment: Applying Willowood Pronamide 3.3SC following initial irrigation of the crop will limit movement of the herbicide below the root zone of germinating weeds and improve weed control. The optimal application timing following pre-irrigation will vary depending on season, weed species present and environmental conditions. The following information is provided as a general guideline:

Timing	Date	Application Timing (Days After Starting Sprinklers)
Early	Sept. 1 to Oct. 15	1-3
Mid	Oct. 15 to Dec. 15	3-6
Late	Dec. 15 to Jan.	5-6

Chemigation Equipment: Willowood Pronamide 3.3SC may be applied through center pivot, lateral move, solid set or hand move systems capable of uniform delivery of the herbicide. Solid set or hand move systems should be capable of delivering a uniform pressure of 60 to 70 psi at all nozzles. Pipes and nozzles must be positioned to provide uniform coverage of the treatment area. Placement of nozzles in a diamond shaped (♦) pattern will provide more uniform coverage. Do not apply when wind velocity is sufficient to distort uniformity of coverage or cause drift to susceptible non-target plants.

The injection-metering pump must be calibrated as specified by the manufacturer and checked periodically during application to ensure proper operation. Pesticide injection hoses, which connect chemigation-metering equipment to the sprinkler irrigation system, must be of braided reinforced construction with an internal tube made of nylon, cross-linked polyethylene, or high-density polyethylene.

Mixing: Mixing tanks should be large enough to contain the entire amount of herbicide mixture for the area to be treated. Use a minimum of 3 gallons of water per 1.25 pints of Willowood Pronamide 3.3SC. Agitation of the herbicide mixture is required at all times during mixing and application (injection).

Application: For hand move or solid set systems set up to deliver about 1/10 inch of water per hour, Willowood Pronamide 3.3SC should be injected over a period of 1 to 2 hours. Once the herbicide has been injected, continue irrigation for at least the time required to flush the system and deliver additional irrigation sufficient to incorporate the herbicide into the upper inch of soil.

Chemigation Use Restrictions for Head Lettuce, Endive, Escarole or Radicchio Greens

- Do not apply Willowood Pronamide 3.3SC to direct seeded varieties of head lettuce, endive, escarole and radicchio greens that will be harvested less than 55 days after treatment or transplanted head lettuce that will be harvested less than 35 days after application.
- Do not apply more than two applications of Willowood Pronamide 3.3SC per season to head lettuce, endive, escarole, or radicchio greens.

- Do not apply Willowood Pronamide 3.3SC postemergence to endive, escarole, or radicchio greens.
- Do not apply more than 2.5 pints of Willowood Pronamide 3.3SC (1 lb. active ingredient) per acre.

Chemigation Instructions

Do not apply this product through any irrigation system unless the instructions for chemigation are followed. Apply this product only through continuously moving center pivot, lateral move end tow, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.

Do not connect an irrigation system (including greenhouse system) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems on the pesticide container label are in place.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Sprinkler Chemigation

1. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system.
2. The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back-flow.
3. The pesticide injection pipeline must contain a functional automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump stops.
6. The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point that pesticide distribution is adversely affected.
7. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
8. Do not apply when wind speed favors drift beyond the area intended for treatment.

Read the "CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY" in the label booklet for Willowood Pronamide 3.3SC before using.

Willowood, LLC
 1600 NW Garden Valley Blvd. #130
 Roseburg, OR 97471

[EPA Approval Date]