

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

July 9, 2013

OFFICE OF CHEMICAL SAFETY AND POLITION PREVENTION

Emory Oleochemicals, LLC 4900 Este Avenue Cincinnati, Ohio 45232

Subject: Notification for product Brand Name change, consistent with the provisions of

PR Notice 98-10.

Deletion of CAMCO Chemicals, Inc. and APCO packaging as producers, and addition of Schirm USA, Inc. and Adjuvants Unlimited, LLC as producers of Emerion7020 Concentrate on the CSF.

Product Brand Name: Emery Oleochemicals, LLC. New Brand Name: Emerion 7020 Concentrate.

EPA Registration Number: 87663-3. Your Submission dated, June 11, 2013.

Dear Mr. Sam Ghantous:

The Biopesticides and Pollution Prevention Division is in receipt of your application for Notification under Pesticide Registration Notice (PRN) 98-10 dated above. A screen of this request has been conducted for its applicability under PRN 98-10, and it has been determined that the actions requested fall within the scope of this Notice. Our records have been duly noted, and the label submitted with this application has been stamped "Notification Accepted" and will be placed accordingly in our records.

Three (3) copies of final printed labeling must be submitted to the Agency before your product as modified, may be sold or distributed [PR Notice 82-2 and 40 CFR 156.10(a) (6)].

If you have any questions regarding this action, you may contact Mr. Sylvester George at (703) 603-0688 or via e-mail at george.sylvester@epa.gov.

Sincerely,

Linda A. Hollis

Linda A. Hollis, Chief Biochemical Pesticide Branch Biopesticides and Pollution Prevention Division (7511P)

Please read instructions on revers	e before completing form		Forn	Approved, OME	No. 2070-00	60, Approval expires 5-31-98
	Unite	ed States	·	Registrat	ion	OPP Identifier Number
.O.EDA				X Amendm	ent	
\$EPA	Environmental i	Protection A	gency _		J.1.C	
	Washingt	on, DC 20460		Other		
			esticide - Section	1		
1. Company/Product Number			2. EPA Product Mai		3. Propo	sed Classification
Emery Oleochemica			Gina Burnett			
4. Company/Product (Name)			PM#		X No	ne Restricted
	als LLC / Emerion 7020 Co	oncentrate	BPPD-23 6. Expedited Revie	u. In cocardo	ooo with EIE	TPA Continu 2/0//2/
Name and Address of Appl Emery Oleochemica			(b)(i), my product is			
4900 Este Ave	10 120		to:	on mar or room.	out in comp	conton and laboring
Cincinnati, Ohio 452	:32		EPA Reg No.			
	-l		Draduat Nama			
Check if this is a new ad	dress		Product Name			
		Soot	ion II			
V Amondanant Evaluis be		Sect	ion – II	labala in reana		
X Amendment – Explain be	HOW		Final printed	labels in respoi	Notif	ication Accepte
		•	Agency letter			reacion Accepte
Resubmission in respons	se to Agency letter dated		"Me Too" App	olication	Data:	1/10/2013
					vate:	The land
X Notification – Explain bel	.ow		Other – Expla	ain below	Review	7/10/2013 er: Sylvester George
Explanation: Use additional p	page(s) if necessary. (For	Section I and Se	ection II.)			
1 Notification apporain	ng to Pesticide Registration	n Notice (DD) 09	10 of product Broad Na	ma ahanaa N	ow product	name is Emerica 7020
 Notification according Concentrate. 	g to resticide Registration	ii Notice (PK) 96-	TO OF Product Brand Na	me change. N	ew product	name is cinenon 7020
	Chemicals, Inc. and APC	O Packaging as p	producers, and addition	of Schirm USA	, Inc. and A	djuvants Unlimited, LLC
as producers of Eme	erion 7020 Concentrate or	the CSF.				-
		Cook				
Material This Product Will be	o Dockogod in:	Sec	tion III			
Child-Resistant Packaging	Unit Packaging	Tw	ater Soluble Packaging		2 Type	of Container
Yes	Yes		Yes		,	etal
X No	X No	 	K No		X Pla	astic
				<u></u>	_ Gi	ass
*Certification must	If "Yes"	• •	"Yes"	No. per	·	per
be submitted	Unit Packaging wgt.	container Pa	ackage wgt	container	Ot	her (Specify)
Location of Net Contents In	formation	4. Size(s) Reta	il Container	15.10	cation of la	bel directions
X Label	Container		z, 1, 2.5,5,55,270 & 600		On Label	DCT directions
		, ,	. , . , ,			ccompanying product
6. Manner in Which Label is A	ffixed to Product	Lithograpi	h	Other		
		X Paper glu	ed			
		Stenciled				
1.0	'A		tion IV	***************************************		C C C C
Contact Person (Complete Name		entification of ind itle	ividual to be contacted,	if necessary, to	T	is application:) le No. (Include Area
Name Sam Ghantous			ional Product Safety an	d Regulations	Code)	c (Include Area
Sum Shamous			ional i rodaot odiroty an		ccccc	513-762-2750
					((
		tification		<u> </u>		Date Application
I certify that the statements I had been supported as that any knowing					((((Received `` (Stainped)
I acknowledge that any knowin both under applicable law	gry raise or misreading stat	ement may be put	usuavie by tine or impri	Somment of	ر د ز د د د	(Staniped)
2. Signature		3. Title				((((((((((((((((((((
	4		Regional Product Safety	and Regulation	ns	
Sam Gha	wlow		•			(((
4. Typed Name		5. Date	 			
Sam Gh	antous		June 10, 2013			



June 11, 2013

Ms. Gina Burnett
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
One Potomac Yard
2777 S. Crystal Drive
Arlington, VA 22202

Subject:

Emery Oleochemicals LLC

EPA Reg. # 87663-3 Registration Amendment

Dear Ms. Burnett:

Emery Oleochemicals LLC is hereby submitting this request to amend the existing registration of *Emery Agro 7020 Concentrate*, *EPA Reg # 87663-3*.

The requested changes are:

- 1. Notification according to Pesticide Registration Notice (PR) 98-10 of product Brand Name change. The new product Brand Name will be Emerion® 7020 Concentrate.
- 2. Deletion of CAMCO Chemicals, Inc. and APCO Packaging as producers, and addition of Schirm USA, Inc. and Adjuvants Unlimited, LLC (three facilities) as producers of Emerion 7020 Concentrate on the CSF.

In support of these amendments, Emery Oleochemicals is submitting the following document:

- Application for Registration Form 8570-1.
- Revised Confidential Statement of Formula (CSF) form 8570-4.
- Revised label reflecting the new product Brand Name.

In the meantime, if you have any questions or require any additional information, please feel free to contact me @ (513) 762-2750 or by email, <u>Sam.Ghantous@Emeryoleo.com</u>. As always, thank you for your attention to our registration activities.

Sincerely,

Sam Ghantous

Sam Chant

Sr. Manager, Regional Product Safety & Regulations

Emery Oleochemicals LLC

Enclosures (3)

4906 Este Avenue Cincinnati, OH 45232 www.emervoleo.com Emery Oleochemicals NA Headquarters



Emerion 7020 Concentrate

Non-selective Herbicide

Sa FOR ORGANIC PRODUCTION

NOT FOR FOOD USE

MASTER LABEL

SUBLABEL A: FOR AGRICULTURAL USE

FOR CONTACT SPRAY CONTROL OR BURNDOWN OF WEEDS AND GRASSES FOR ORNAMENTALS, TURF, TREES IN AND AROUND FARMS, FORESTS, NURSERIES AND GREENHOUSES

SUBLABEL B: FOR COMMERCIAL USE

FOR CONTACT SPRAY CONTROL OR BURNDOWN OF WEEDS AND GRASSES FOR USE IN STRUCTURAL PEST CONTROL, CONTROL OF VEGETATION IN PASTURE AND RANGELAND FOR USE IN PARKS, GOLF COURSES AND PUBLIC AND PRIVATE LAWNS AND GROUNDS FOR AESTHETIC PURPOSES FOR PLANTS IN HABITATIONS, RESIDENTIAL FRUIT AND VEGETABLE GARDENS; RESIDENTIAL GREENHOUSES

Active ingredient:	د د د د
Ammonium Nonanoate	40.0 wt% .
Other ingredients	. <u>60.0 wt.%</u>
Total	. 100.00 wt.%

cecece

SUBLABEL A: FOR AGRICULTURAL USE

FOR CONTACT SPRAY CONTROL OR BURNDOWN OF WEEDS AND GRASSES FOR ORNAMENTALS, TURF, TREES IN AND AROUND FARMS, FORESTS, NURSERIES AND GREENHOUSES

Emerion 7020 Concentrate

Non-selective Herbicide

FOR ORGANIC PRODUCTION

NOT FOR FOOD USE

Active ingredient:	
Ammonium Nonanoate	.40.0 wt.%
Other ingredients	60.0 wt.%

EMERION 7020 CONCENTRATE contains 3.3 lbs. of ammonium nonanoate per US gallon

KEEP OUT OF REACH OF CHILDREN

WARNING "AVISO"

"Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)"

	FIRST AID			
If in eyes	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continuring eye. Call a poison control center or doctor for treatment advice. 			
If inhaled	 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. 			
If on skin or clothing	• Take off contaminated clothing • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.			
	EMERGENCY NUMBER			
FOR 24-HOUR EM	ERGENCY MEDICAL ASSISTANCE, CALL THE NATIONAL POISON CONTROL CENTER 1-800-222-1222.			
Have the product contain	ner or label with you when calling a poison control center or doctor.			

This product is protected by U. S. Patent No. 6,323,156

EPA Reg. No. 87663-3

EPA Establishment No. 72038-DE-001, 39578-TX-001, 064784-OK-001, 89402-OK-001, 89402-OK-001, 89402-OK-001

Lot No. xxxxxxxx

Net Contents: 3,4,16 & 32 fl oz, 1, 2.5,5, 55, 270 & 6000 gals

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Warning. Causes substantial, but temporary eye irritation. Do not get in eyes or on clothing. Wear goggles. Harmful if inhaled. Avoid breathing spray mist. Causes moderate skin irritation. Avoid contact with skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Coveralls worn over short-sleeve shirt and short pants
- Socks and chemical resistant footwear
- Chemical resistant gloves
- Protective eyewear
- When mixing and loading wear a chemical resistant apron

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergents and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses; this pesticide is toxic to fish and aquatic invertebrates. Use care when applying in areas adjacent to any body of water. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. This product is toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product if bees are visiting the treatment area.

For containers greater than 5 gallon

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State/Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water is:

- Coveralls worn over short-sleeve shirt and short pants
- Socks and chemical resistant footwear
- Chemical resistant gloves
- Protective eyewear

METHODS OF USE AND GENERAL APPLICATION INSTRUCTIONS

General Instructions:

EMERION 7020 CONCENTRATE is a contact non-selective herbicide for spray application only to undesirable plant growth. Do not allow spray to contact any green plant parts of desirable plants. **EMERION 7020 CONCENTRATE** provides control and burndown suppression of annual and perennial broadleaf and grass weeds. Spore producing plants such as mosses and liverworts are also controlled. The amount of burndown and the duration of weed suppression may be reduced when weed growth conditions are unfavorable or when plants are mature.

EMERION 7020 CONCENTRATE is non-volatile and water soluble for foliar application in tractor powered field-type sprayers or manual pump sprayers. Complete and uniform coverage of weeds by the spray solution is required for the best weed control. Plant foliage will change from a green color to brown/black necrotic tissue within one to two hours after spray application of EMERION 7020 CONCENTRATE diluted with water. EMERION 7020 CONCENTRATE effect on plant tissue may be more rapid in warm weather than in cold conditions. However, weed control is normally unaffected by temperature.

EMERION 7020 CONCENTRATE herbicide is a soap product which penetrates the cell walls of plants to disrupt the cellular organization of physiological functions which are compartmentalized by membranes within the cell walls. Plant growth ceases when cellular contents are mixed which causes brown necrotic plant tissue.

EMERION 7020 CONCENTRATE does not migrate through the soil and is not translocated in plants. To ensure satisfactory control, plant leaves must be thoroughly and uniformly covered with the spray solution. EMERION 7020 CONCENTRATE does not provide any residual weed control in soil to affect germinating weed seeds.

Mixing and Application Instructions:

For use, EMERION 7020 CONCENTRATE is diluted with water to the specified concentration for effective control of the undesirable vegetation. Apply using standard methods of liquid herbicide application. Dilution must be in accordance with label instructions. Do not apply this product through any type of irrigation system.

A 12% v/v dilution is recommended for most weed control situations and based on the results, applicators may increase or decrease spray concentrations as discussed in the following text and tables to obtain better control or to reduce herbicide use.

The degree of dilution for application is based on the concentration of active ingredient needed for the size of vegetation to be suppressed or the rate of herbicidal effect desired. The larger the vegetation, the higher the concentration (less dilution of EMERION 7020 CONCENTRATE) required for rapid action. See required concentration for variously sized weeds and grasses in Dosage and Application Rates Section. Hard to control or suppress weeds (perennial weeds and some grasses) may require several applications for complete control.

Spray equipment options include hand-held, boom sprayers, pressure sprayers and hose-end sprayers. Spray nozzles that produce a uniform spray will give maximum coverage of the leaves and improved weed control. **EMERION 7020 CONCENTRATE** is completely soluble in water and when uniformly mixed, no additional mixing or agitation is required.

Application Precautions:

- Do not apply to weeds when wet from dew, rain or irrigation.
- Do not irrigate within 2 hours after application
- Do not apply if rainfall is expected within 2 hours.

During application, some foaming may occur; however, weed control is unaffected if the foam is deposited on the plant surface and is not blown away as drift. Foaming can be reduced by using the minimum spray pressure required for a uniform application to the target weeds. Use low spray pressure to reduce foaming and avoid contact with desirable plants. Most spray nozzles are designed to operate at 10 to 15 psi and provide uniform spray coverage of weeds.

Mixing:

Fill sprayer tank with half the required amount of water, add the full amount of EMERION 7020 CONCENTRATE to be used (see Dilution Factors below), then fill the sprayer tank with the remainder of the water required for the desired final concentration. Since EMERION 7020 CONCENTRATE is completely soluble in water and when uniformly dispersed in water, continuous mixing or agitation is not required.

Broadcast Application with Field Sprayer Boom Equipment

The amount of weed vegetation will determine the spray volume required for complete coverage of undesired plants (weeds). Weed vegetation conditions that affect spray coverage are number of weeds present, leaf shape, weed size and weed species. For weeds of over 1 inch height do not use less than 30 gal/acre. Large weeds of 12 to 18 inch height may require 80 to 125 gal/acre or more for control.

Hand-Held Equipment

Thorough saturation of the foliage and stems is required for control, but stop sprays when run-off from weed leaves occurs. Use low spray pressure to reduce foaming and avoid contact with desirable plants. Most spray nozzles are designed to operate at 10 to 15 psi and provide uniform spray coverage of weeds.

Directed Spray Equipment

Use a shielded sprayer to prevent spray contact on desirable plants. Avoid spray contact of green plant stems or green bark of young trees and shrubs.

NOTE: In areas of hard water, the final mixture may appear milky. This condition does not change the effectiveness of the treatment. Some visible foam may appear on the leaves as the plants ere sprayed. Overspray or drift onto desirable plants is usually not a serious problem because leaves and stems require a thoroughly wet spray coverage for significant injury. Repeat applications as often as necessary to obtain desired control.

Dosage and Application Rates:

For general weed and grass control, rates are based on the size of the plants and/or the desired speed of kill. The larger the plants, the higher the dosage rates needed to ensure maximum herbicidal activity. Also, the higher the concentration, the quicker the plants wilt and turn brown. Apply EMERION 7020 CONCENTRATE spray solutions only when weed surfaces are dry.

The rate table presents the suggested percent volume/volume solutions of EMERION 7020 CONCENTRATE to use for application as follows:

- Use a 6 to 8 % V/V **EMERION 7020 CONCENTRATE** spray solution for control of annual weeds of 1 inch height or less and for control of liverworts and mosses.
- Use an 8 to 10% V/V **EMERION 7020 CONCENTRATE** spray solution for control of annual weeds over 1 inch height and up to 4 inch height.
- Use a 10 to 13% V/V **EMERION 7020 CONCENTRATE** spray solution for weeds over 4 inch height and for hard to control weeds.
- A 15% V/V **EMERION 7020 CONCENTRATE** spray solution is the maximum labeled rate. Use this rate for hard to control perennial weeds or extremely dense weed growth.

Repeat spray applications as necessary to obtain the desired control or suppression of weeds from newly germinated weed seeds and regrowth from roots or stems.

RATE TABLE

Final Spray Volume	Amount o	Amount of EMERION 7020 CONCENTRATE for Percent V/V (Volume/Volume) Solution				
(gallons)	6%	8%	10%	13%	15%	
1	8 fl oz	10 fl oz	13 fl oz	1 pt	1.2 pt	
2	1 pt	1.3 pt	1.3 pt	2 pt	2.4 pt	
5	2.5 pt	3.3 pt	4 pt	5 pt	6 pt	
10	5 pt	6.5 pt	1 gal	1.3 gal	1.5 gal	
20	1.3 gal	1.6 gal	2 gal	2.6 gal	3.0 galcies	

Height of Plants	Spray		
to be Controlled	Solution (% V/V)	(ι (((((((((((((((((((
1 inch or less	6 to 8%	, ,	ر (رزد
1 to 3 inches	8 to 10%	6 L C L L L L L L L L L L L L L L L L L	(((((((((((((((((((
Above 3 inches	10 to 13%		((

NOTE: Do not use spray concentrations higher than 15% in a pressure sprayer since unacceptable foaming and bubble formation may occur at the nozzle heads.

Use Methods

Determine the weed control situation and select the use method required as follows:

- **A. Vegetative Burndown:** Broadcast spray for weed control for no-till planting or seedbed preparation to control weeds prior to seeding or transplanting. Spot sprays may be used in ornamentals and turf.
- **B. Directed and shielded sprays:** Spray nozzle type or configuration for directed spray or a shield placed around the nozzle to prevent spray contact on the foliage or green stems or bark. Directed/shielded spray applications to area between plastic mulch strips and staked crops for weed control.

PESTS

I. Weeds controlled or suppressed by **EMERION 7020 CONCENTRATE**:

COMMON NAMES

TAXONOMIC NAMES

Broadleaf Weeds:			
Algae	Gloeocapsa magma		
Bittercress, hairy	Cardamine hirsuta		
Chickweed, common	Stellaria media		
Chickweed, mouse-ear	Cerastium vulgatum		
Cocklebur, common	Xanthium strumarium		
Corn spurry	Spergula arvensis		
Cudweed, purple	Gnaphalium purpureum		
Groundsel	Senecio spp.		
Lambsquarters, common	Chenopodium album		
Liverwort	Machantia spp.		
Marestail or Horseweed rosettes	Conyza canadensis		
Morningglory, annual	Ipomoea spp.		
Moss	Bryophyta		
Mustards	Brassica spp.		
Oxalis or Woodsorrel	Oxalis stricta		
Pansy, wild	Viola tricolor		
Plantain	Plantago spp.		
Pigweed, smooth and redroot	Amaranthus spp.		
Mallow, roundleaved	Malva spp.		
Moneywort	Lysimachia nummularia		
Shepherdspurse	Capsella bursa-pastoris		
Sorrel, sheep	Rumex acetosella		
Spurge, spotted	Euphorbia maculata		
Field pennycress	Thiaspi arvense		
Velvetleaf	Abutilon theophrasti	,	
Grass and Other Weeds:	•		
Bentgrass, colonial	Agrostis tenuis		
Bluegrass, annual	Poa annua	•	
Crabgrass, large	Digitaria sanguinalis		
Fescue, creeping red	Festuca rubra		
Fescue, hard	Festuca ovina		(((
Nimblewill	Muhlenbergia scheberi	ης η ((((((((((((((((((
Onion, wild	Allium canadense	((((((((((((((((((((((
Ryegrass, perennial	Lolium perenne	ι (((((((((((((((((((•
Star-of-Bethlehem	Ornithogalum nutans	(((6 6	(c t c c c c c c c c c c c c c c c c c
II. Weeds Moderately Difficult to Control ¹		(((((((((((((((((((((
Bermudagrass (Wireweed)	Cynodon dactylon		(((((((((((((((((((
Bindweed, field	Convolvulus arvensis		, , , , , , , , , , , , , , , , , , ,
Dandelion	Taraxacum officinale		
Nutsedge, yellow Ragweed, common	Cyperus esculentus Ambrosia artemisiifolia		
	amorasia ariomisiitaila		

STORAGE AND HANDLING

DO NOT contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store container in cool place until used. Store at temperatures above 32° F.

Pesticide Disposal: Wastes resulting from use of this product must be disposed of on-site or at an approved waste disposal facility.

Container Handling:

Non-refillable Containers: Do not reuse or refill this container. Offer for recycling if available.

Clean container promptly after emptying.

Non-refillable container equal to or less than 50 lbs or 5 gals: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Non-refillable Container greater than 50 lbs or 5-55 gals: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Refillable Container (50 lbs or larger): Refill this container with this pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

For non-medical emergencies or spills, see **EMERION 7020 CONCENTRATE** MSDS or call CHEMTREC at 800-424-9300.

Terms and Conditions of Use:

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty and Disclaimer Notice:

To the extent consistent with applicable law, Emery Oleochemicals LLC makes no warranty or guarantee of any kind, expressed or implied concerning the effects of use of this product, other than those specified on this label. Buyers or users accept all responsibility for results due to misuse or improper handling of this product.

Inherent Risks of Use:

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to the label instructions (including adverse conditions noted on the label, such as unfavorable temperatures, wind, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Emery Oleochemicals LLC or the seller. To the extent consistent with applicable law, all such risks shall be assumed by the buyer.

Limitation of Remedies:

To the extent consistent with applicable law, the exclusive remedy for losses or damages resulting from the use of this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Emery Oleochemicals LLC's election, one of the following:

- 1. Refund of purchase price paid by buyer or user for product bought, or
- 2. Replacement of amount of product used.

To the extent allowable by state law, Emery Oleochemicals LLC shall not be liable for losses or damages resulting from handling or use of this product unless Emery Oleochemicals LLC is promptly notified of such loss or damage in writing. In no case shall Emery Oleochemicals LLC be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Emery Oleochemicals LLC or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

Produced for: Emery Oleochemicals LLC 4900 Este Avenue Cincinnati, OH 45232 513-762-2500 www.Emeryoleo.com

SUBLABEL B: FOR COMMERCIAL USE

FOR CONTACT SPRAY CONTROL OR BURNDOWN OF WEEDS AND GRASSES FOR USE IN STRUCTURAL PEST CONTROL, CONTROL OF VEGETATION IN PASTURE AND RANGELAND FOR USE IN PARKS, GOLF COURSES AND PUBLIC AND PRIVATE LAWNS AND GROUNDS FOR AESTHETIC PURPOSES FOR PLANTS IN HABITATIONS, RESIDENTIAL FRUIT AND VEGETABLE GARDENS; RESIDENTIAL GREENHOUSES

Emerion 7020 Concentrate

Non-selective Herbicide

TO FOR ORGANIC PRODUCTION

NOT FOR FOOD USE

Δ	ctive	in	ored	ient:	
_	CUIVE		21 CU	TCHT.	

EMERION 7020 CONCENTRATE contains 3.3 lbs. of ammonium nonanoate per US gallon

KEEP OUT OF REACH OF CHILDREN

WARNING "AVISO"

"Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)"

	FIRST AID				
If in eyes	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. 				
If inhaled	 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. 				
If on skin or clothing	 Take off contaminated clothing Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. 				
	EMERGENCY NUMBER				
FOR 24-HOUR EM	ERGENCY MEDICAL ASSISTANCE, CALL THE NATIONAL POISON				

CONTROL CENTER 1-800-222-1222.

Have the product container or label with you when calling a poison control center or doctor.

This product is protected by U. S. Patent No. 6,323,156

EPA Reg. No. 87663-3

EPA Establishment No. 72038-DE-001, 39578-TX-001, 064784-OK-001, 89402-OK-001, 89402-OK-001, 89402-OK-001

Lot No. xxxxxxxx

Net Contents: 3,4,16 & 32 fl oz, 1, 2.5 & 5 gal

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Warning. Causes substantial, but temporary eye irritation. Do not get in eyes or on clothing. Wear goggles. Harmful if inhaled. Avoid breathing spray mist. Causes moderate skin irritation. Avoid contact with skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergents and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses; this pesticide is toxic to fish and aquatic invertebrates. Use care when applying in areas adjacent to any body of water. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate. Drift and runoff may be hazardous to aquatic torganisms in water adjacent to treated areas. This product is toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product if bees are visiting the treatment area.

For containers greater than 5 gallon

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, occans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State/Tribal agency responsible for pesticide regulation.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried.

METHODS OF USE AND GENERAL APPLICATION INSTRUCTIONS

General Instructions:

EMERION 7020 CONCENTRATE is a contact non-selective herbicide for spray application only to undesirable plant growth. Do not allow spray to contact any green plant parts of desirable plants. **EMERION 7020 CONCENTRATE** provides control and burndown suppression of annual and perennial broadleaf and grass weeds. Spore producing plants such as mosses and liverworts are also controlled. The amount of burndown and the duration of weed suppression may be reduced when weed growth conditions are unfavorable or when plants are mature.

EMERION 7020 CONCENTRATE is non-volatile and water soluble for foliar application in tractor powered field-type sprayers or manual pump sprayers. Complete and uniform coverage of weeds by the spray solution is required for the best weed control. Plant foliage will change from a green color to brown/black necrotic tissue within one to two hours after spray application of EMERION 7020 CONCENTRATE diluted with water. EMERION 7020 CONCENTRATE effect on plant tissue may be more rapid in warm weather than in cold conditions. However, weed control is normally unaffected by temperature.

EMERION 7020 CONCENTRATE herbicide is a soap product which penetrates the cell walls of plants to disrupt the cellular organization of physiological functions which are compartmentalized by membranes within the cell walls. Plant growth ceases when cellular contents are mixed which causes brown necrotic plant tissue.

EMERION 7020 CONCENTRATE does not migrate through the soil and is not trenslocated in plants. To ensure satisfactory control, plant leaves must be thoroughly and uniformly covered with the spray solution. EMERION 7020 CONCENTRATE does not provide any residual weed control in soil to affect germinating weed seeds.

Mixing and Application Instructions:

For use, EMERION 7020 CONCENTRATE is diluted with water to the specified concentration for effective control of the undesirable vegetation. Apply using standard methods of liquid herbicide application. Dilution must be in accordance with label instructions. Do not apply this product through any type of irrigation system.

A 12% v/v dilution is recommended for most weed control situations and based on the results, applicators may increase or decrease spray concentrations as discussed in the following text and tables to obtain better control or to reduce herbicide use.

The degree of dilution for application is based on the concentration of active ingredient needed for the size of vegetation to be suppressed or the rate of herbicidal effect desired. The larger the vegetation, the higher the concentration (less dilution of EMERION 7020 CONCENTRATE) required for rapid action. See required concentration for variously sized weeds and grasses in Dosage and Application Rates Section. Hard to control or suppress weeds (perennial weeds and some grasses) may require several applications for complete control.

Spray equipment options include hand-held, boom sprayers, pressure sprayers and hose-end sprayers. Spray nozzles that produce a uniform spray will give maximum coverage of the leaves and improved weed control. **EMERION 7020 CONCENTRATE** is completely soluble in water and when uniformly mixed, no additional mixing or agitation is required.

Application Precautions:

- Do not apply to weeds when wet from dew, rain or irrigation.
- Do not irrigate within 2 hours after application
- Do not apply if rainfall is expected within 2 hours.

During application, some foaming may occur; however, weed control is unaffected if the foam is deposited on the plant surface and is not blown away as drift. Foaming can be reduced by using the minimum spray pressure required for a uniform application to the target weeds. Use low spray pressure to reduce foaming and avoid contact with desirable plants. Most spray nozzles are designed to operate at 10 to 15 psi and provide uniform spray coverage of weeds.

Mixing:

Fill sprayer tank with half the required amount of water, add the full amount of EMERION 7020 CONCENTRATE to be used (see Dilution Factors below), then fill the sprayer tank with the remainder of the water required for the desired final concentration. Since EMERION 7620 CONCENTRATE is completely soluble in water and when uniformly dispersed in water, continuous mixing or agitation is not required.

Broadcast Application with Field Sprayer Boom Equipment

The amount of weed vegetation will determine the spray volume required for complete coverage of undesired plants (weeds). Weed vegetation conditions that affect spray coverage are number of weeds present, leaf shape, weed size and weed species. For weeds of over 1 inch height do not use less than 30 gal/acre. Large weeds of 12 to 18 inch height may require 80 to 125 gal/acre or more for control.

Hand-Held Equipment

Thorough saturation of the foliage and stems is required for control, but stop sprays when run-off from weed leaves occurs. Use low spray pressure to reduce foaming and avoid contact with desirable plants. Most spray nozzles are designed to operate at 10 to 15 psi and provide uniform spray coverage of weeds.

Directed Spray Equipment

Use a shielded sprayer to prevent spray contact on desirable plants. Avoid spray contact of green plant stems or green bark of young trees and shrubs.

NOTE: In areas of hard water, the final mixture may appear milky. This condition does not change the effectiveness of the treatment. Some visible foam may appear on the leaves as the plants are sprayed. Overspray or drift onto desirable plants is usually not a serious problem because leaves and stems require a thoroughly wet spray coverage for significant injury. Repeat applications as often as necessary to obtain desired control.

Dosage and Application Rates:

For general weed and grass control, rates are based on the size of the plants and/or the desired speed of kill. The larger the plants, the higher the dosage rates needed to ensure maximum herbicidal activity. Also, the higher the concentration, the quicker the plants wilt and turn brown. Apply **EMERION 7020 CONCENTRATE** spray solutions only when weed surfaces are dry.

The rate table presents the suggested percent volume/volume solutions of EMERION 7020 CONCENTRATE to use for application as follows:

- Use a 6 to 8 % V/V **EMERION 7020 CONCENTRATE** spray solution for control of annual weeds of 1 inch height or less and for control of liverworts and mosses.
- Use an 8 to 10% V/V **EMERION 7020 CONCENTRATE** spray solution for control of annual weeds over 1 inch height and up to 4 inch height.
- Use a 10 to 13% V/V EMERION 7020 CONCENTRATE spray solution for weeds over 4 inch height and for hard to control weeds.
- A 15% V/V EMERION 7020 CONCENTRATE spray solution is the maximum labeled rate. Use this rate for hard to control perennial weeds or extremely dense weed growth.

Repeat spray applications as necessary to obtain the desired control or suppression of weeds from newly germinated weed seeds and regrowth from roots or stems.

RATE TABLE

Final Spray	Amount of EMERION 7020 CONCENTRATE for Percent V/V					
Volume		(Volume/Volume) Solution				
_ (gallons)	6%	8%	10%	13%	15%	
1	8 fl oz	10 fl oz	13 fl oz	1 pt	1.2 pt	
2	1 pt	1.3 pt	1.3 pt	2 pt	2.4 pt	
5	2.5 pt	3.3 pt	4 pt	5 pt	6 pt	
10	5 pt	6.5 pt	1 gal	1.3 gal	1.5 gal	
20	1.3 gal	1.6 gal	2 gal	2.6 gal	3.0 gal	

Height of Plants to be Controlled	Spray Solution (% V/V)
1 inch or less	6 to 8%
1 to 3 inches	8 to 10%
Above 3 inches	10 to 13%

NOTE: Do not use spray concentrations higher than 15% in a pressure sprayer since unacceptable foaming and bubble formation may occur at the nozzle heads.

Use Methods

Determine the weed control situation and select the use method required as follows:

- **A. Vegetative Burndown:** Broadcast spray for weed control for no-till planting or seedbed preparation to control weeds prior to seeding or transplanting. Spot sprays may be used in ornamentals and turf.
- **B. Directed and shielded sprays:** Spray nozzle type or configuration for directed spray or a shield placed around the nozzle to prevent spray contact on the foliage or green stems or bark. Directed/shielded spray applications to area between plastic mulch strips and staked crops for weed control.

PESTS

I. Weeds controlled or suppressed by **EMERION 7020 CONCENTRATE**:

COMMON NAMES

TAXONOMIC NAMES

Broadleaf Weeds:			
Algae	Gloeocapsa magma		
Bittercress, hairy	Cardamine hirsuta		
Chickweed, common	Stellaria media	•	
Chickweed, mouse-ear	Cerastium vulgatum		
Cocklebur, common	Xanthium strumarium		
Corn spurry	Spergula arvensis		
Cudweed, purple	Gnaphalium purpureum		
Groundsel	Senecio spp.		
Lambsquarters, common	Chenopodium album		
Liverwort	Machantia spp.		
Marestail or Horseweed rosettes	Conyza canadensis		ř
Morningglory, annual	Ipomoea spp.		
Moss	Bryophyta		
Mustards	Brassica spp.		
Oxalis or Woodsorrel	Oxalis stricta		
Pansy, wild	Viola tricolor		
Plantain	Plantago spp.		
Pigweed, smooth and redroot	Amaranthus spp.		
Mallow, roundleaved	Malva spp.		
Moneywort	Lysimachia nummularia		
Shepherdspurse	Capsella bursa-pastoris		
Sorrel, sheep	Rumex acetosella		
Spurge, spotted	Euphorbia maculata	•	
Field pennycress	Thiaspi arvense		
Velvetleaf	Abutilon theophrasti		
Grass and Other Weeds:		•	
Bentgrass, colonial	Agrostis tenuis		
Bluegrass, annual	Poa annua		
Crabgrass, large	Digitaria sanguinalis		
Fescue, creeping red	Festuca rubra		
Fescue, hard	Festuca ovina		
Nimblewill	Muhlenbergia scheberi	,,,,,,	((6
Onion, wild	Allium canadense	• • • • • • • • • • • • • • • • • • •	•
Ryegrass, perennial	Lolium perenne		(
Star-of-Bethlehem	Ornithogalum nutans	ξιξείς (, ,
Stat-of-Definement	отинодиит пишт	ږ د د د د	ι (ε (ε (ε
II. Weeds Moderately Difficult to Control ¹	•	(i.
Bermudagrass (Wireweed)	Cynodon dactylon		((((((((((((((((((((
Bindweed, field	Convolvulus arvensis		
Dandelion	Taraxacum officinale		((((
Nutsedge, yellow	Cyperus esculentus		
Ragweed, common	Ambrosia artemisiifolia		
Use 13 to 15% V/V EMERION 7020 CONCENTRATE for control or			

STORAGE AND HANDLING

DO NOT contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store container in cool place until used. Store at temperatures above 32° F.

Pesticide Disposal: Wastes resulting from use of this product must be disposed of on-site or at an approved waste disposal facility.

Container Handling:

Non-refillable Containers: Do not reuse or refill this container. Offer for recycling if available.

Clean container promptly after emptying.

Non-refillable container equal to or less than 50 lbs or 5 gals: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Non-refillable Container greater than 50 lbs or 5-55 gals: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Refillable Container (50 lbs or larger): Refill this container with this pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

For non-medical emergencies or spills, see **EMERION 7020 CONCENTRATE** MSDS or call CHEMTREC at 1-800-424-9300.

Terms and Conditions of Use:

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty and Disclaimer Notice:

To the extent consistent with applicable law, Emery Oleochemicals LLC makes no warranty or guarantee of any kind, expressed or implied concerning the effects of use of this product, other than those specified on this label. Buyers or users accept all responsibility for results due to misuse or improper handling of this product.

Inherent Risks of Use:

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to the label instructions (including adverse conditions noted on the label, such as unfavorable temperatures, wind, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Emery Oleochemicals LLC or the seller. To the extent consistent with applicable law, all such risks shall be assumed by the buyer.

Limitation of Remedies:

To the extent consistent with applicable law, the exclusive remedy for losses or damages resulting from the use of this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Emery Oleochemicals LLC's election, one of the following:

- 1. Refund of purchase price paid by buyer or user for product bought, or
- 2. Replacement of amount of product used.

To the extent allowable by state law, Emery Oleochemicals LLC shall not be liable for losses or damages resulting from handling or use of this product unless Emery Oleochemicals LLC is promptly notified of such loss or damage in writing. In no case shall Emery Oleochemicals LLC be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Emery Oleochemicals LLC or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

Produced for: Emery Oleochemicals LLC 4900 Este Avenue Cincinnati, OH 45232 513-762-2500 www.Emeryoleo.com