



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

February 27, 2023

Sam Ghantous
Global Head, Product Safety and Regulations
Emery Oleochemicals LLC
4900 Este Avenue, Bldg. 6A-3
Cincinnati, Ohio 45232

Subject: Non-PRIA (Pesticide Registration Improvement Act) Labeling Amendment – Adding
USDA Bio-preferred logo, OMRI Logo and footer label version and date change.
Product Name: FL-AN640OG
EPA Registration Number: 87663-14
EPA Receipt Date: 09/02/2021
Action Case Number: 00319754

Dear Mr. Ghantous:

The amended labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable.

This approval does not affect any terms or conditions that were previously imposed on this registration. You continue to be subject to existing terms or conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release this product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR § 152.3.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review the U.S. Environmental Protection Agency (EPA). If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should EPA find or if it is brought to our attention that a website contains statements or claims substantially


Page 2 of 2
EPA Reg. No. 87663-14
Action Case No. 00319754

differing from statements or claims made in connection with obtaining a FIFRA section 3 registration, the website will be referred to EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6.


If you have any questions, please contact Ms. Menyon Adams by phone at (202) 566-1604 or via email at @epa.gov.

Sincerely,



James Parker, Lead
Biochemical Pesticides Branch
Biopesticides and Pollution
Prevention Division (7511M)
Office of Pesticide Programs

Enclosure



FL-AN640OG

(Alternate Brand Name “EMERION® W 40 SL OG”)

Non-selective Herbicide

 **FOR ORGANIC GARDENING**

NOT FOR FOOD USE

MASTER LABEL

SUBLABEL A: FOR COMMERCIAL USE

FOR CONTACT SPRAY CONTROL OR BURNDOWN OF WEEDS AND GRASSES
FOR ORNAMENTALS, TURF, LANDSCAPES, INTERIORSCAPES, RESIDENTIAL
GREENHOUSES AND RESIDENTIAL SITES

SUBLABEL B: FOR RESIDENTIAL USE

FOR CONTACT SPRAY CONTROL OR BURNDOWN OF WEEDS AND GRASSES
FOR HOMES AND GARDENS

Active ingredient:

Ammonium Nonanoate	40.0 wt. %
Other ingredients	<u>60.0 wt. %</u>
Total	100.00 wt. %

Optional Label Claims:

- Fast Acting
- Acts on Contact
- Acts on Contact to Kill Green Leafy Tissues
- Effective Weed Control in Cooler Climates
- Quicker Action in Warm / Hot Weather Conditions
- Rain Fast in 2 Hours
- For Control or Burndown of Weeds and Grasses.
- Grass and Weed Knockdown
- Controls Annual and Perennial weeds
- Ideal For Spot Weed Control Around Ornamental Trees, Bushes, Flowers and Landscaping Beds
- Contains Fatty Acids Derived from Fats and Oils
- Quickly Kills Green Plant Tissue on Contact
- Degrades Rapidly
- Will Not Leach Through the Soil to Harm Desirable Plants
- Reseed immediately after use

Optional Graphics:

- USDA Bio-Preferred Program label [logo]:



This mark is not an indication of safety. Read and follow all label instructions

- OMRI Listed [logo]:



This mark is not an indication of safety. Read and follow all label instructions

SUBLABEL A: FOR COMMERCIAL USE

FOR CONTACT SPRAY CONTROL OR BURNDOWN OF WEEDS AND GRASSES
FOR ORNAMENTALS, TURF, LANDSCAPES, INTERIORSCAPES, RESIDENTIAL
GREENHOUSES AND RESIDENTIAL SITES

FL-AN640OG

(Alternate Brand Name "EMERION® W 40 SL OG")

Non-selective Herbicide



FOR ORGANIC GARDENING

NOT FOR FOOD USE

Active ingredient:

Ammonium Nonanoate 40.0 wt. %
Other ingredients 60.0 wt. %
Total 100.00 wt. %

FL-AN640OG contains 3.3 lbs. of ammonium nonanoate per US gallon

KEEP OUT OF REACH OF CHILDREN

WARNING "AVISO"

"Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)"

FIRST AID	
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
EMERGENCY NUMBER	
FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL THE NATIONAL POISON CONTROL CENTER 1-800-222-1222.	
Have the product container or label with you when calling a poison control center or doctor.	

This product is protected by U. S. Patent No. 6,323,156

EPA Reg. No. 87663-14

EPA Establishment No. 87663-OH-001,72038-DE-001, 6574-KY-001, 064784-OK-001

Lot No. xxxxxxxxx

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Warning. Causes substantial, but temporary eye irritation. Do not get in eyes or on clothing. Wear goggles. Harmful if inhaled. Avoid breathing spray mist. Causes moderate skin irritation. Avoid contact with skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

For terrestrial uses: This pesticide is toxic to fish and aquatic invertebrates. Use care when applying in areas adjacent to any body of water. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

This product is toxic to bees exposed to direct treatment or residues on blooming gardens or weeds. Do not apply this product if bees are visiting the treatment area.

For containers greater than 5 gallon

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State/Tribal agency responsible for pesticide regulation.

METHODS OF USE AND GENERAL APPLICATION INSTRUCTIONS

General Instructions:

FL-AN640OG is a contact non-selective herbicide for spray application only to undesirable plant growth. Do not allow spray to contact any green plant parts of desirable plants. **FL-**

AN640OG provides control and burndown suppression of annual and perennial broadleaf and grass weeds. Spore producing plants such as mosses and liverworts are also controlled. The amount of burndown and the duration of weed suppression may be reduced when weed growth conditions are unfavorable or when plants are mature.

FL-AN640OG is non-volatile and water soluble for foliar application in manual pump sprayers. Complete and uniform coverage of weeds by the spray solution is required for the best weed control. Plant foliage will change from a green color to brown/black necrotic tissue within one to two hours after spray application of **FL-AN640OG** diluted with water. **FL-AN640OG** effect on plant tissue may be more rapid in warm weather than in cold conditions. However, weed control is normally unaffected by temperature.

FL-AN640OG herbicide is a soap product which penetrates the cell walls of plants to disrupt the cellular organization of physiological functions which are compartmentalized by membranes within the cell walls. Plant growth ceases when cellular contents are mixed which causes brown necrotic plant tissue.

FL-AN640OG does not migrate through the soil and is not translocated in plants. **To ensure satisfactory control, plant leaves must be thoroughly and uniformly covered with the spray solution.** **FL-AN640OG** does not provide any residual weed control in soil to affect germinating weed seeds.

Mixing and Application Instructions:

For use, **FL-AN640OG** is diluted with water to the specified concentration for effective control of the undesirable vegetation. Apply using standard methods of liquid herbicide application. Dilution must be in accordance with label instructions. Do not apply this product through any type of irrigation system.

A 12% v/v dilution is recommended for most weed control situations and based on the results, applicators may increase or decrease spray concentrations as discussed in the following text and tables to obtain better control or to reduce herbicide use.

The degree of dilution for application is based on the concentration of active ingredient needed for the size of vegetation to be suppressed or the rate of herbicidal effect desired. The larger the vegetation, the higher the concentration (less dilution of FL-AN640OG) required for rapid action. See required concentration for variously sized weeds and grasses in Dosage and Application Rates Section. Hard to control or suppress weeds (perennial weeds and some grasses) may require several applications for complete control.

Spray equipment options include hand-held and hose-end sprayers. Spray nozzles that produce a uniform spray will give maximum coverage of the leaves and improved weed control.

FL-AN640OG is completely soluble in water and when uniformly mixed, no additional mixing or agitation is required.

Application Precautions:

- Do not apply to weeds when wet from dew, rain or water.

- Do not water within 2 hours after application
- Do not apply if rainfall is expected within 2 hours.

During application, some foaming may occur; however, weed control is unaffected if the foam is deposited on the plant surface and is not blown away as drift. Foaming can be reduced by using the minimum spray pressure required for a uniform application to the target weeds. Use low spray pressure to reduce foaming and avoid contact with desirable plants. Most spray nozzles are designed to operate at 10 to 15 psi and provide uniform spray coverage of weeds.

Mixing:

Fill sprayer tank with half the required amount of water, add the full amount of **FL-AN640OG** to be used (see Dilution Factors below), then fill the sprayer tank with the remainder of the water required for the desired final concentration. Since **FL-AN640OG** is completely soluble in water and when uniformly dispersed in water, continuous mixing or agitation is not required.

Hand-Held Equipment

Thorough saturation of the foliage and stems is required for control, but stop sprays when run-off from weed leaves occurs. Use low spray pressure to reduce foaming and avoid contact with desirable plants. Most spray nozzles are designed to operate at 10 to 15 psi and provide uniform spray coverage of weeds.

Directed Spray Equipment

Use a shielded sprayer to prevent spray contact on desirable plants. Avoid spray contact of green plant stems or green bark of young trees and shrubs.

NOTE: In areas of hard water, the final mixture may appear milky. This condition does not change the effectiveness of the treatment. Some visible foam may appear on the leaves as the plants are sprayed. Overspray or drift onto desirable plants is usually not a serious problem because leaves and stems require a thoroughly wet spray coverage for significant injury. Repeat applications as often as necessary to obtain desired control.

Dosage and Application Rates:

For general weed and grass control, rates are based on the size of the plants and/or the desired speed of kill. The larger the plants, the higher the dosage rates needed to ensure maximum herbicidal activity. Also, the higher the concentration, the quicker the plants wilt and turn brown. Apply **FL-AN640OG** spray solutions only when weed surfaces are dry.

The rate table presents the suggested percent volume/volume solutions of **FL-AN640OG** to use for application as follows:

- Use a 6 to 8 % V/V **FL-AN640OG** spray solution for control of annual weeds of 1 inch height or less and for control of liverworts and mosses.
- Use an 8 to 10% V/V **FL-AN640OG** spray solution for control of annual weeds over 1 inch height and up to 4 inch height.

- Use a 10 to 13% V/V **FL-AN6400G** spray solution for weeds over 4 inch height and for hard to control weeds.
- A 15% V/V **FL-AN6400G** spray solution is the maximum labeled rate. Use this rate for hard to control perennial weeds or extremely dense weed growth.

Repeat spray applications as necessary to obtain the desired control or suppression of weeds from newly germinated weed seeds and regrowth from roots or stems.

RATE TABLE

Final Spray Volume (gallons)	Amount of FL-AN6400G for Percent V/V (Volume/Volume) Solution				
	6%	8%	10%	13%	15%
1	8 fl oz	10 fl oz	13 fl oz	1 pt	1.2 pt
2	1 pt	1.3 pt	1.6 pt	2 pt	2.4 pt
5	2.5 pt	3.3 pt	4 pt	5 pt	6 pt
10	5 pt	6.5 pt	1 gal	1.3 gal	1.5 gal
20	1.3 gal	1.6 gal	2 gal	2.6 gal	3.0 gal

<u>Height of Plants to be Controlled</u>	<u>Spray Solution (% V/V)</u>
1 inch or less	6 to 8%
1 to 4 inches	8 to 10%
Above 4 inches	10 to 13%

NOTE: Do not use spray concentrations higher than 15% in a pressure sprayer since unacceptable foaming and bubble formation may occur at the nozzle heads.

Use Methods

Determine the weed control situation and select the use method required as follows:

A. Vegetative Burndown: Broadcast spray for weed control for no-till planting or seedbed preparation to control weeds prior to seeding or transplanting. Spot sprays may be used in ornamentals and turf.

B. Directed and shielded sprays: Spray nozzle type or configuration for directed spray or a shield placed around the nozzle to prevent spray contact on the foliage or green stems or bark. Directed/shielded spray applications to area between plastic mulch strips and staked gardens for weed control.

PESTS

I. Weeds controlled or suppressed by FL-AN640OG:

COMMON NAMES	TAXONOMIC NAMES
Broadleaf Weeds:	
Algae	<i>Gloeocapsa magma</i>
Bittercress, hairy	<i>Cardamine hirsuta</i>
Chickweed, common	<i>Stellaria media</i>
Chickweed, mouse-ear	<i>Cerastium vulgatum</i>
Cocklebur, common	<i>Xanthium strumarium</i>
Corn spurry	<i>Spergula arvensis</i>
Cudweed, purple	<i>Gnaphalium purpureum</i>
Groundsel	<i>Senecio spp.</i>
Lambsquarters, common	<i>Chenopodium album</i>
Liverwort	<i>Machantia spp.</i>
Marestail or Horseweed rosettes	<i>Conyza canadensis</i>
Morningglory, annual	<i>Ipomoea spp.</i>
Moss	<i>Bryophyta</i>
Mustards	<i>Brassica spp.</i>
Oxalis or Woodsorrel	<i>Oxalis stricta</i>
Pansy, wild	<i>Viola tricolor</i>
Plantain	<i>Plantago spp.</i>
Pigweed, smooth and redroot	<i>Amaranthus spp.</i>
Mallow, roundleaved	<i>Malva spp.</i>
Moneywort	<i>Lysimachia nummularia</i>
Shepherdspurse	<i>Capsella bursa-pastoris</i>
Sorrel, sheep	<i>Rumex acetosella</i>
Spurge, spotted	<i>Euphorbia maculata</i>
Field pennycress	<i>Thiaspi arvense</i>
Velvetleaf	<i>Abutilon theophrasti</i>
Grass and Other Weeds:	
Bentgrass, colonial	<i>Agrostis tenuis</i>
Bluegrass, annual	<i>Poa annua</i>
Crabgrass, large	<i>Digitaria sanguinalis</i>
Fescue, creeping red	<i>Festuca rubra</i>
Fescue, hard	<i>Festuca ovina</i>
Nimblewill	<i>Muhlenbergia scheberi</i>
Onion, wild	<i>Allium canadense</i>
Ryegrass, perennial	<i>Lolium perenne</i>
Star-of-Bethlehem	<i>Ornithogalum nutans</i>
II. Weeds Moderately Difficult to Control¹	
Bermudagrass (Wireweed)	<i>Cynodon dactylon</i>
Bindweed, field	<i>Convolvulus arvensis</i>
Dandelion	<i>Taraxacum officinale</i>
Nutsedge, yellow	<i>Cyperus esculentus</i>
Ragweed, common	<i>Ambrosia artemisiifolia</i>

¹Use 13 to 15% V/V FL-AN640OG for control or suppression

STORAGE AND HANDLING

DO NOT contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store container in cool place until used. Store at temperatures above 32° F.

Pesticide Disposal: Wastes resulting from use of this product must be disposed of on-site or at an approved waste disposal facility.

Container Handling:

Non-refillable Containers: Do not reuse or refill this container. Offer for recycling if available.

Clean container promptly after emptying (liquid formulations).

Non-refillable container equal to or less than 5 gals: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Non-refillable container between 5- 55 gals: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Refillable container 5 gals to bulk: Refill this container with this pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. . Repeat this rinsing procedure two more times. Dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

For non-medical emergencies or spills, see **FL-AN640OG** MSDS or call CHEMTREC at 800-424-9300.

Warranty and Disclaimer Notice:

To the extent consistent with applicable law, Emery Oleochemicals LLC, makes no warranty or guarantee of any kind, expressed or implied concerning the effects of use of this product, other than those specified on this label. Buyers or users accept all responsibility for results due to misuse or improper handling of this product.

Note to PM: these statements are for exclusive use by The Ortho Group

- **NOTICE:** To the extent consistent with applicable law, the buyer assumes all risks of use, storage or handling of this product not in accordance with label directions.

- **Ortho Guarantee**

If for any reason you, the consumer, are not satisfied with this product, mail us your original proof of purchase to obtain a full refund of your purchase price.

- **Ortho & Your Environment**

Your yard is a place for family and pets to enjoy. That's why Ortho products are designed with care to provide effective solutions to weed problems. Follow instructions for appropriate usage, storage and disposal.

Manufactured by:

Emery Oleochemicals LLC

4900 Este Avenue

Cincinnati, OH 45232

513-762-2500

www.Emeryoleo.com

SUBLABEL B: FOR RESIDENTIAL USE

FOR CONTACT SPRAY CONTROL OR BURNDOWN OF WEEDS AND GRASSES
FOR HOMES AND GARDENS

FL-AN640OG

(Alternate Brand Name "EMERION® W 40 SL OG")
Non-selective Herbicide

 **FOR ORGANIC GARDENING**

NOT FOR FOOD USE

Active ingredient:

Ammonium Nonanoate 40.0 wt.%
Other ingredients 60.0 wt.%
Total 100.00 wt.%

FL-AN640OG contains 3.3 lbs. of ammonium nonanoate per US gallon

KEEP OUT OF REACH OF CHILDREN

WARNING "AVISO"

"Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)"

FIRST AID	
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
EMERGENCY NUMBER	
FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL THE NATIONAL POISON CONTROL CENTER 1-800-222-1222.	
Have the product container or label with you when calling a poison control center or doctor.	

This product is protected by U. S. Patent No. 6,323,156
EPA Reg. No. 87663-14
EPA Establishment No. 87663-OH-001, 72038-DE-001, 6574-KY-001, 064784-OK-001
Lot No. xxxxxxxxx
Net Contents: 3,4,16 & 32 fl oz, 1, 2.5 & 5 gals

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Warning. Causes substantial, but temporary eye irritation. Do not get in eyes or on clothing. Wear goggles. Harmful if inhaled. Avoid breathing spray mist. Causes moderate skin irritation. Avoid contact with skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

For terrestrial uses: This pesticide is toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, or gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. This product is toxic to bees exposed to direct treatment or residues on blooming garden or weeds. Do not apply this product if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State/Tribal agency responsible for pesticide regulation.

METHODS OF USE AND GENERAL APPLICATION INSTRUCTIONS

General Instructions:

FL-AN640OG is a contact non-selective herbicide for spray application only to undesirable plant growth. Do not allow spray to contact any green plant parts of desirable plants. **FL-AN640OG** provides control and burndown suppression of annual and perennial broadleaf and grass weeds. Spore producing plants such as mosses and liverworts are also controlled. The amount of burndown and the duration of weed suppression may be reduced when weed growth conditions are unfavorable or when plants are mature.

FL-AN640OG is non-volatile and water soluble for foliar application in manual pump sprayers. Complete and uniform coverage of weeds by the spray solution is required for the best weed control. Plant foliage will change from a green color to brown/black necrotic tissue within one to two hours after spray application of **FL-AN640OG** diluted with water. **FL-AN640OG** effect on plant tissue may be more rapid in warm weather than in cold conditions. However, weed control is normally unaffected by temperature.

FL-AN640OG herbicide is a soap product which penetrates the cell walls of plants to disrupt the cellular organization of physiological functions which are compartmentalized by membranes within the cell walls. Plant growth ceases when cellular contents are mixed which causes brown necrotic plant tissue.

FL-AN640OG does not migrate through the soil and is not translocated in plants. **To ensure satisfactory control, plant leaves must be thoroughly and uniformly covered with the spray solution.** **FL-AN640OG** does not provide any residual weed control in soil to affect germinating weed seeds.

Mixing and Application Instructions:

For use, **FL-AN640OG** is diluted with water to the specified concentration for effective control of the undesirable vegetation. Apply using standard methods of liquid herbicide application. Dilution must be in accordance with label instructions. Do not apply this product through any type of irrigation system.

A 12% v/v dilution is recommended for most weed control situations and based on the results, applicators may increase or decrease spray concentrations as discussed in the following text and tables to obtain better control or to reduce herbicide use.

The degree of dilution for application is based on the concentration of active ingredient needed for the size of vegetation to be suppressed or the rate of herbicidal effect desired. The larger the vegetation, the higher the concentration (less dilution of FL-AN640OG) required for rapid action. See required concentration for variously sized weeds and grasses in Dosage and Application Rates Section. Hard to control or suppress weeds (perennial weeds and some grasses) may require several applications for complete control.

Spray equipment options include hand-held and hose-end sprayers. Spray nozzles that produce a uniform spray will give maximum coverage of the leaves and improved weed control. **FL-AN640OG** is completely soluble in water and when uniformly mixed, no additional mixing or agitation is required.

Application Precautions:

- Do not apply to weeds when wet from dew, rain or water.
- Do not water within 2 hours after application
- Do not apply if rainfall is expected within 2 hours.

During application, some foaming may occur; however, weed control is unaffected if the foam is deposited on the plant surface and is not blown away as drift. Foaming can be reduced by using the minimum spray pressure required for a uniform application to the target weeds. Use low spray pressure to reduce foaming and avoid contact with desirable plants. Most spray nozzles are designed to operate at 10 to 15 psi and provide uniform spray coverage of weeds.

Mixing:

Fill sprayer tank with half the required amount of water, add the full amount of **FL-AN640OG** to be used (see Dilution Factors below), then fill the sprayer tank with the remainder of the water required for the desired final concentration. Since **FL-AN640OG** is completely soluble in water and when uniformly dispersed in water, continuous mixing or agitation is not required.

Hand-Held Equipment

Thorough saturation of the foliage and stems is required for control, but stop sprays when run-off from weed leaves occurs. Use low spray pressure to reduce foaming and avoid contact with desirable plants. Most spray nozzles are designed to operate at 10 to 15 psi and provide uniform spray coverage of weeds.

Directed Spray Equipment

Use a shielded sprayer to prevent spray contact on desirable plants. Avoid spray contact of green plant stems or green bark of young trees and shrubs.

NOTE: In areas of hard water, the final mixture may appear milky. This condition does not change the effectiveness of the treatment. Some visible foam may appear on the leaves as the plants are sprayed. Overspray or drift onto desirable plants is usually not a serious problem because leaves and stems require a thoroughly wet spray coverage for significant injury. Repeat applications as often as necessary to obtain desired control.

Dosage and Application Rates:

For general weed and grass control, rates are based on the size of the plants and/or the desired speed of kill. The larger the plants, the higher the dosage rates needed to ensure maximum herbicidal activity. Also, the higher the concentration, the quicker the plants wilt and turn brown. Apply **FL-AN640OG** spray solutions only when weed surfaces are dry.

The rate table presents the suggested percent volume/volume solutions of **FL-AN640OG** to use for application as follows:

- Use a 6 to 8 % V/V **FL-AN640OG** spray solution for control of annual weeds of 1 inch height or less and for control of liverworts and mosses.
- Use an 8 to 10% V/V **FL-AN640OG** spray solution for control of annual weeds over 1 inch height and up to 4 inch height.
- Use a 10 to 13% V/V **FL-AN640OG** spray solution for weeds over 4 inch height and for hard to control weeds.
- A 15% V/V **FL-AN640OG** spray solution is the maximum labeled rate. Use this rate for hard to control perennial weeds or extremely dense weed growth.

Repeat spray applications as necessary to obtain the desired control or suppression of weeds from newly germinated weed seeds and regrowth from roots or stems.

RATE TABLE

Final Spray Volume (gallons)	Amount of FL-AN640OG for Percent V/V (Volume/Volume) Solution				
	6%	8%	10%	13%	15%
1	8 fl oz	10 fl oz	13 fl oz	1 pt	1.2 pt
2	1 pt	1.3 pt	1.6 pt	2 pt	2.4 pt
5	2.5 pt	3.3 pt	4 pt	5 pt	6 pt
10	5 pt	6.5 pt	1 gal	1.3 gal	1.5 gal
20	1.3 gal	1.6 gal	2 gal	2.6 gal	3.0 gal

Height of Plants
to be Controlled
1 inch or less

Spray
Solution (% V/V)
6 to 8%

1 to 4 inches

8 to 10%

Above 4 inches

10 to 13%

NOTE: Do not use spray concentrations higher than 15% in a pressure sprayer since unacceptable foaming and bubble formation may occur at the nozzle heads.

Use Methods

Determine the weed control situation and select the use method required as follows:

A. Vegetative Burndown: Broadcast spray for weed control for no-till planting or seedbed preparation to control weeds prior to seeding or transplanting. Spot sprays may be used in ornamentals and turf.

B. Directed and shielded sprays: Spray nozzle type or configuration for directed spray or a shield placed around the nozzle to prevent spray contact on the foliage or green stems or bark. Directed/shielded spray applications to area between plastic mulch strips and staked gardens for weed control.

PESTS

I. Weeds controlled or suppressed by FL-AN640OG:

COMMON NAMES	TAXONOMIC NAMES
Broadleaf Weeds:	
Algae	<i>Gloeocapsa magma</i>
Bittercress, hairy	<i>Cardamine hirsuta</i>
Chickweed, common	<i>Stellaria media</i>
Chickweed, mouse-ear	<i>Cerastium vulgatum</i>
Cocklebur, common	<i>Xanthium strumarium</i>
Corn spurry	<i>Spergula arvensis</i>
Cudweed, purple	<i>Gnaphalium purpureum</i>
Groundsel	<i>Senecio spp.</i>
Lambsquarters, common	<i>Chenopodium album</i>
Liverwort	<i>Machantia spp.</i>
Marestail or Horseweed rosettes	<i>Conyza canadensis</i>
Morningglory, annual	<i>Ipomoea spp.</i>
Moss	<i>Bryophyta</i>
Mustards	<i>Brassica spp.</i>
Oxalis or Woodsorrel	<i>Oxalis stricta</i>
Pansy, wild	<i>Viola tricolor</i>
Plantain	<i>Plantago spp.</i>
Pigweed, smooth and redroot	<i>Amaranthus spp.</i>
Mallow, roundleaved	<i>Malva spp.</i>
Moneywort	<i>Lysimachia nummularia</i>
Shepherdspurse	<i>Capsella bursa-pastoris</i>
Sorrel, sheep	<i>Rumex acetosella</i>
Spurge, spotted	<i>Euphorbia maculata</i>
Field pennycress	<i>Thiaspi arvense</i>
Velvetleaf	<i>Abutilon theophrasti</i>
Grass and Other Weeds:	
Bentgrass, colonial	<i>Agrostis tenuis</i>
Bluegrass, annual	<i>Poa annua</i>
Crabgrass, large	<i>Digitaria sanguinalis</i>
Fescue, creeping red	<i>Festuca rubra</i>
Fescue, hard	<i>Festuca ovina</i>
Nimblewill	<i>Muhlenbergia scheberi</i>
Onion, wild	<i>Allium canadense</i>
Ryegrass, perennial	<i>Lolium perenne</i>
Star-of-Bethlehem	<i>Ornithogalum nutans</i>
II. Weeds Moderately Difficult to Control¹	
Bermudagrass (Wireweed)	<i>Cynodon dactylon</i>
Bindweed, field	<i>Convolvulus arvensis</i>
Dandelion	<i>Taraxacum officinale</i>
Nutsedge, yellow	<i>Cyperus esculentus</i>
Ragweed, common	<i>Ambrosia artemisiifolia</i>

¹Use 13 to 15% V/V FL-AN640OG for control or suppression

STORAGE AND HANDLING

DO NOT contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store container in cool place until used. Store at temperatures above 32° F.

Pesticide Disposal: Wastes resulting from use of this product must be disposed of on-site or at an approved waste disposal facility.

Container Disposal:

If empty: Non-refillable - Do not reuse or refill this container. Offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

For non-medical emergencies or spills, see **FL-AN640OG** MSDS or call CHEMTREC at 800-424-9300.

Warranty and Disclaimer Notice:

To the extent consistent with applicable law, Emery Oleochemicals LLC makes no warranty or guarantee of any kind, expressed or implied concerning the effects of use of this product, other than those specified on this label. Buyers or users accept all responsibility for results due to misuse or improper handling of this product.

Note to PM: these statements are for exclusive use by The Ortho Group

- **NOTICE:** To the extent consistent with applicable law, the buyer assumes all risks of use, storage or handling of this product not in accordance with label directions.
- **Ortho Guarantee**
If for any reason you, the consumer, are not satisfied with this product, mail us your original proof of purchase to obtain a full refund of your purchase price.
- **Ortho & Your Environment**

Your yard is a place for family and pets to enjoy. That's why Ortho products are designed with care to provide effective solutions to weed problems. Follow instructions for appropriate usage, storage and disposal.

Manufactured by:

Emery Oleochemicals LLC
4900 Este Avenue
Cincinnati, OH 45232
513-762-2500
www.Emeryoleo.com