



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

June 18, 2019

Cynthia Ann Smith
Agent for Crop Protection Technology, Pte Ltd.
Conn& Smith, Inc.
6713 CATSKILL ROAD
LORTON, VA 22079

Subject: Non-PRIA (Pesticide Registration Improvement Act) Labeling Amendment –
Amendment to Revise some California Prohibitions and to Make Small Corrections
Product Name: Trabon EC
EPA Registration Number: 87749-1
Application Date: 1/28/19
OPP Decision Number: 547553

Dear Ms. Smith:

The amended labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable.

The alternate brand name - Rango – has been added to the registration, and our records have been updated accordingly. This approval does not affect any terms or conditions that were previously imposed on this registration. You continue to be subject to existing terms or conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release this product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. “To distribute or sell” is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR § 152.3.

Should you wish to add/retain a reference to your company’s website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the U.S. Environmental Protection Agency (EPA). If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration

Page 2 of 2
EPA Reg. No. 87749-1
OPP Decision No. 547553

process. Therefore, should the EPA find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA-approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6.


If you have any questions, please contact Chris Pfeifer of my team by phone at (703) 308-0031 or via email at pfeifer.chris@epa.gov.

Sincerely,



Andrew Bryceland, Team Leader
Biochemical Pesticides Branch
Biopesticides and Pollution
Prevention Division (7511P)
Office of Pesticide Programs

Enclosure



TRABON™ EC Master Label

Contents

Page

RESIDENTIAL USE	2
Container front panel.	2
Container back panel booklet, visible without opening the booklet.	3
Container back panel booklet, NOT visible without opening the booklet	4
COMMERCIAL USE	10
Container front panel.	10
Container back panel booklet, visible without opening booklet	11
Container back panel booklet, NOT visible without opening booklet	13

{RESIDENTIAL USE: Container front panel}

TRABON™ EC

[or Alternate brand name:

DYNABLOCK CPT™

CLEAR CROP™

CROP CLEAN™

AGROCLARA™

HARVESTGUARD™]

FOR RESIDENTIAL USE

[Optional Statements:]

FOR RESIDENTIAL HOME AND GARDEN USE

FOR USE AROUND HOMES, DAY CARE CENTERS, SCHOOLS,
PARKS, OFFICE BUILDINGS, AND CHURCHES

TRABON EC [or Alternate brand name] is a Broad Spectrum Contact [Insecticide] [and] [Fungicide] for use against pests in/on [fruits,] [vegetables,] [flowers,] [and] [ornamental] [shrubs] [and] [trees].

Controls a variety of soft-bodied pests including [aphids,] [whiteflies,] [leafhoppers] [and] [mites].
Provides [curative] [and] [preventative control] of powdery mildew.

Active Ingredient:

Soybean oil 45.0%

Other Ingredients 55.0%

Total 100.0%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Booklet for User Recommendations and Directions for Use.

NET CONTENTS:

3.38 fl. oz. (100 mL)

8.45 fl. oz. (250 mL)

16.9 fl. oz. (500 mL)

PATENT PENDING

EPA Reg. No. 87749-1

EPA Est. No. 87749-MYS-1

MANUFACTURED BY:

Crop Protection Technology, Pte Ltd
71 Bukit Batok Crescent
#07-03 Prestige Centre
Singapore 658071

Label Version No. _____

TRABON EC (EPA Reg. No. 87749-1)

January 28, 2019 proposed amendments to the 09/18/2012 Accepted Label with 11/15/2016 Notification

Revises "Not for use in California" notations to reflect uses CDPR has agreed to accept.

{RESIDENTIAL USE: Container back panel booklet, visible without opening the booklet}

TRABON™ EC
[or Alternate brand name]

USER SAFETY RECOMMENDATIONS

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. This product may be toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product if bees are visiting the treatment area.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Do not reuse empty container. Store in tightly closed container in a cool dry place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container.

PATENT PENDING

EPA Reg. No. 87749-1

EPA Est. No. 87749-MYS-1

MANUFACTURED BY:

Crop Protection Technology, Pte Ltd

71 Bukit Batok Crescent

#07-03 Prestige Centre

Singapore 658071

™ Trademark of Crop Protection Technology, Pte Ltd.

Label Version No. _____

{RESIDENTIAL USE: Container back panel booklet, NOT visible without opening the booklet}

GENERAL INFORMATION

TRABON EC is an emulsified suspension of soybean oil that when mixed with water according to the directions and applied directly to pests at the specified rates will control the listed pests.

TRABON EC is a contact pesticide. For maximum pest control, direct contact of sprays with pests is essential. Best applied in early morning or late afternoon when pests tend to be less active. The frequency of application should be adjusted to maintain sufficient pest control. Application frequency may vary depending on crop type, geographical region and local weather conditions. Consult with your state Agricultural Extension Specialist for specific recommendations regarding the crop type and pest control recommendations for your region.

Use the low application rates on the label when pest densities are relatively low. Use the progressively higher application rates on the label for progressively higher pest populations.

TRABON EC has not been tested on all plant varieties under all growing conditions. Test a small number of plants for damage before applying TRABON EC to the entire crop.

INTEGRATED PEST MANAGEMENT AND RESISTANCE MANAGEMENT

Use an effective IPM program. Avoid repeated or sequential use of products in the same mode of pesticidal action. Monitor crop for pest pressure and for loss of efficacy. Contact your extension specialist, certified crop consultant, or manufacturer for the latest resistance management information. Contact the producer to report loss of efficacy.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

TRABON EC may be applied using a watering can or a hand-held pressure sprayer.

Use sufficient water to ensure thorough and uniform coverage to infested plant material. Add enough water to allow mixing or agitation. Apply solution quickly. If application is delayed, stir contents of watering can or shake or roll spray tank to mix contents before applying solution. Triple rinse and flush spray equipment thoroughly following each use.

Keep unprotected persons out of treated areas until sprays have dried.

APPLICATION RATE AND FREQUENCY

AVOCADO, BANANA, DATE, FIG, MANGO, PAPAYA, POMEGRANATE, PINEAPPLE [†]			
INSECT PESTS: Whiteflies, Spider mites, Russet mites, Broad mites, Leafhoppers, Mealybug, Flea beetles, Thrips larvae, Scale insects, Melon aphids, Potato aphid	FUNGAL PESTS: Powdery mildew	FOLIAR APPLICATION RATE: Mix 1.0 teaspoon to 1.0 Tablespoon TRABON EC with 1 gallon of water. Apply 2.5 to 10.0 gallons solution per 1000 sq. ft. Ensure thorough penetration and coverage of target pest or disease for maximum effectiveness.	APPLICATION TIMING: Apply when conditions favor pest development or at first signs of damage or infestation. Repeat every 7 to 14 days as necessary. TRABON EC may be applied up to and including the day of harvest.
† Not for use in California.			

BERRY AND SMALL FRUIT GROUP:			
Blackberry (<i>Rubus</i> spp.) (including Andean blackberry, arctic blackberry, bingleberry, black satin berry, boysenberry, brombeere, California blackberry, Chesterberry, Cherokee blackberry, Cheyenne blackberry, common blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, evergreen blackberry, Himalayaberry, hullberry, lavacaberry, loganberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, mora, mures deronce, nectarberry, Northern dewberry, olallieberry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, Southern dewberry, tayberry, youngberry, zarzamora, and cultivars, varieties and/or hybrids of these); Blueberry, highbush (<i>Vaccinium</i> spp.); Blueberry, lowbush (<i>Vaccinium angustifolium</i> Aiton); Grape (<i>Vitis</i> spp.); Kiwifruit, fuzzy (<i>Actinidia deliciosa</i> A. Chev.) (C.F. Liang and A.R. Fergusons, <i>Actinida chinensis</i> Planch.); Kiwifruit, hardy (<i>Actinidia arguta</i> (Siebold and Zucc.) Planch. ex Miq); Raspberry, black and red (<i>Rubus</i> spp.); Strawberry (<i>Fragaria x ananassa</i> Duchesne); and Cultivars, varieties, and/or hybrids of these.			
INSECT PESTS[†]: Whiteflies, Spider mites, Russet mites, Broad mites, Leafhoppers, Mealybug, Thrips larvae, Scale insects, Melon aphids, Potato aphid	FUNGAL PESTS: Powdery mildew	FOLIAR APPLICATION RATE: Mix 1.0 teaspoon to 1.0 Tablespoon TRABON EC with 1 gallon of water. Apply 1.0 to 5.0 gallons solution per 1000 sq. ft. Ensure thorough penetration and coverage of target pest or disease for maximum effectiveness.	APPLICATION TIMING: Apply when conditions favor pest development or at first signs of damage or infestation. Repeat every 7 to 14 days as necessary. TRABON EC may be applied up to and including the day of harvest.
† Not for use in California.			

BRASSICA (COLE) LEAFY VEGETABLES:

Broccoli (*Brassica oleracea* var. *botrytis*); Broccoli, Chinese (gai lon) (*Brassica alboglabra*); Cabbage (*Brassica oleracea*); Cabbage, Chinese (bok choy) (*Brassica chinensis*); Cabbage, Chinese mustard (gai choy) (*Brassica campestris*); Cauliflower (*Brassica oleracea* var. *botrytis*); Collards (*Brassica oleracea* var. *acephala*); Kale (*Brassica oleracea* var. *acephala*); and Mustard greens (*Brassica juncea*).

BULB VEGETABLES (Allium app.) †:

Chive, fresh leaves (*Allium schoenoprasum* L.); Chive, Chinese, fresh leaves (*Allium tuberosum* Rottler ex Spreng); Garlic, bulb (*Allium sativum* L. var. *sativum*) (*A. sativum* Common Garlic Group); Leek *Allium porrum* L. (syn: *A. ampeloprasum* L. var. *porrum* (L.) J. Gay) (*A. ampeloprasum* Leek Group); Lily, bulb (*Lilium* spp. (*Lilium Leichtlinii* var. *maximowiczii*, *Lilium lancifolium*)); Onion, bulb (*Allium cepa* L. var. *cepa*) (*A. cepa* Common Onion Group); Shallot, fresh leaves (*Allium cepa* var. *aggregatum* G. Don.); and Cultivars, varieties, and/or hybrids of these.

CUCURBIT VEGETABLES:

Chinese waxgourd (Chinese preserving melon) (*Benincasa hispida*); Citron melon (*Citrullus lanatus* var. *citroides*); Cucumber (*Cucumis sativus*); Gourd, edible (*Lagenaria* spp.) (includes hyotan, cucuzza); (*Luffa acutangula*, *L. cylindrica*) (includes hechima, Chinese okra); *Momordica* spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber); Muskmelon (hybrids and/or cultivars of *Cucumis melo*) (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon); Pumpkin (*Cucurbita* spp.); and Squash, summer (*Cucurbita pepo* var. *meloepo*) (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini).

FRUITING VEGETABLES (Except Cucurbits):

Bush tomato, *Solanum centrale* J.M. Black; Currant tomato, *Lycopersicon pimpinellifolium* L.; Eggplant, *Solanum melongena* L.; Pepper, bell, *Capsicum annuum* L. var. *annuum*, *Capsicum* spp.; Pepper, nonbell, *Capsicum chinese* Jacq., *C. annuum* L. var. *annuum*, *C. frutescens* L., *C. baccatum* L., *C. pubescens* Ruiz & Pav., *Capsicum* spp.; Roselle, *Hibiscus sabdariffa* L.; Tomatillo, *Physalis philadelphica* Lam; Tomato, *Solanum lycopersicum* L., *Solanum lycopersicum* L. var. *lycopersicum*; and Cultivars, varieties and/or hybrids of these.

HERBS AND SPICES †:

Basil (*Ocimum basilicum*); Caraway (*Carum carvi*); Catnip (*Nepeta cataria*); Chive, Chinese (*Allium tuberosum*); Coriander (cilantro or Chinese parsley) (leaf) (*Coriandrum sativum*); Coriander (cilantro) (seed) (*Coriandrum sativum*); Curry (leaf) (*Murraya koenigii*); Dill (dillweed) (*Anethum graveolens*); Dill (seed) (*Anethum graveolens*); Lemongrass (*Cymbopogon citratus*); Mustard (seed) (*Brassica juncea*, *B. hirta*, *B. nigra*); Parsley (dried) (*Petroselinum crispum*); Rosemary (*Rosemarinus officinalis*); Sage (*Salvia officinalis*); Tarragon (*Artemisia dracunculus*); and Thyme (*Thymus* spp.).

LEAFY VEGETABLES (Except Brassica Vegetables) †:

Arugula (Roquette) (*Eruca sativa*); Celery (*Apium graveolens* var. *dulce*); Celery, Chinese (*Apium graveolens* var. *secalinum*); Endive (escarole) (*Cichorium endivia*); Fennel, Florence (finocchio) (*Foeniculum vulgare* Azoricum Group); Lettuce, head and leaf (*Lactuca sativa*); Purslane, garden (*Portulaca oleracea*); Radicchio (red chicory) (*Cichorium intybus*); Rhubarb (*Rheum rhabarbarum*); Spinach (*Spinacia oleracea*); and Swiss chard (*Beta vulgaris* var. *cicla*).

LEGUME VEGETABLES (Succulent or Dried) †:

Bean (*Phaseolus* spp.) (includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean); Bean (*Vigna* spp.) (includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean); Broad bean (fava bean) (*Vicia faba*); Pea (*Pisum* spp.) (includes dwarf pea, edible-pod pea, English pea, field pea, garden pea, green pea, snow pea, sugar snap pea); and Soybean (*Glycine max*).

TRABON EC (EPA Reg. No. 87749-1)

January 28, 2019 proposed amendments to the 09/18/2012 Accepted Label with 11/15/2016 Notification

Revises "Not for use in California" notations to reflect uses CDPR has agreed to accept.

ROOT AND TUBER VEGETABLES †:

Beet, garden (*Beta vulgaris*); Carrot (*Daucus carota*); Chervil, turnip-rooted (*Chaerophyllum bulbosum*); Parsnip (*Pastinaca sativa*); Radish (*Raphanus sativus*); Rutabaga (*Brassica campestris* var. *napobrassica*); Sweet potato (*Ipomoea batatas*); and Yam, true (*Dioscorea* spp.).

INSECT PESTS:	FUNGAL PESTS †:	FOLIAR APPLICATION RATE:	APPLICATION TIMING:
Whiteflies, Spider mites, Russet mites, Broad mites, Leafhoppers, Mealybug, Flea beetles, Thrips larvae, Scale insects, Melon aphids, Potato aphid	Powdery mildew	Mix 1.0 teaspoon to 1.0 Tablespoon TRABON EC with 1 gallon of water. Apply 1 to 2.5 gallons solution per 1000 sq. ft. Ensure thorough penetration and coverage of target pest or disease for maximum effectiveness.	Apply when conditions favor pest development or at first signs of damage or infestation. Repeat every 7 to 21 days as necessary. TRABON EC may be applied up to and including the day of harvest.

† Not for use in California.

CITRUS FRUITS †:

Citrus spp. *Eremocitrus* spp., *Fortunella* spp., *Microcitrus* spp., and *Poncirus* spp; Grapefruit, *Citrus natsudaoidai* Hayata; Kumquat, *Fortunella* spp; Lemon, *Citrus limon* (L.) Burm. f; Lime, *Citrus aurantiifolia* (Christm.) Swingle; Mediterranean mandarin, *Citrus deliciosa* Ten; Mount White lime, *Microcitrus garrowayae* (F.M. Bailey) *Citrus aurantium* L; Orange, sweet, *Citrus sinensis* (L.); Osbeck Pummelo, *Citrus maxima* (Burm.); *Microcitrus inodora* (F.M. Bailey) Swingle; Satsuma mandarin; Tanaka Tangelo, *Citrus x tangelo* J.W. Ingram & H.E. Moore; Tangerine (Mandarin); and Cultivars, varieties and/or hybrids of these.

POME FRUITS:

Apple, *Malus domestica* Borkh.; Loquat, *Eriobotrya japonica* (Thunb.) Lindl.; Pear, *Pyrus communis* L.; Pear, Asian, *Pyrus pyrifolia* (Burm. f.) Nakai var. *culta* (Makino) Nakai; *Pseudocyonia sinensis* (Thouin) C.K. Schneid.; and Cultivars, varieties and/or hybrids of these.

STONE FRUITS †:

Apricot (*Prunus armeniaca*); Cherry, sweet (*Prunus avium*); Nectarine (*Prunus persica*); Peach (*Prunus persica*); Plum (*Prunus domestica*, *Prunus* spp.); Plum, Japanese (*Prunus salicina*); Plumcot (*Prunus armeniaca* P. *domestica*); and Prune (fresh) (*Prunus domestica*, *Prunus* spp.).

INSECT PESTS †:	FUNGAL PESTS:	FOLIAR APPLICATION RATE:	APPLICATION TIMING:
Whiteflies, Spider mites, Leafhoppers, Mealybug, Thrips larvae, Scale insects, Melon aphids, Potato aphid	Powdery mildew	Mix 1.0 teaspoon to 1.0 Tablespoon TRABON EC with 1 gallon of water. Apply 2.5 to 10.0 gallons solution per 1000 sq. ft. Ensure thorough penetration and coverage of target pest or disease for maximum effectiveness.	Apply when conditions favor pest development or at first signs of damage or infestation. Repeat every 7 to 21 days as necessary. TRABON EC may be applied up to and including the day of harvest.

† Not for use in California.

TRABON EC (EPA Reg. No. 87749-1)

January 28, 2019 proposed amendments to the 09/18/2012 Accepted Label with 11/15/2016 Notification

Revises "Not for use in California" notations to reflect uses CDPR has agreed to accept.

CORN (<i>Zea mays</i>)[†]			
INSECT PESTS: Whiteflies, Spider mites, Russet mites, Broad mites, Leafhoppers, Mealybug, Flea beetles, Thrips larvae, Melon aphids, Potato aphid	FUNGAL PESTS: Powdery mildew	FOLIAR APPLICATION RATE: Mix 1.0 teaspoon to 1.0 Tablespoon TRABON EC with 1 gallon of water. Apply 0.23 to 1.1 gallons solution per 1000 sq. ft. Ensure thorough penetration and coverage of target pest or disease for maximum effectiveness.	APPLICATION TIMING: Apply when conditions favor pest development or at first signs of damage or infestation. Repeat every 7 to 14 days as necessary. TRABON EC may be applied up to and including the day of harvest.
† Not for use in California.			

FLOWERS AND ORNAMENTAL SHRUBS:			
Azalea (<i>Rhododendron spp.</i>); Begonia (<i>Begonia spp.</i>); Camellias (<i>Camellia spp.</i>); Carnation (<i>Dianthus caryophyllus</i>); Chrysanthemum (<i>Chrysanthemum indicum</i> , <i>C. morifolium</i>); Cotoneaster (<i>Cotoneaster spp.</i>); Cyclamen (<i>Cyclamen persicum</i>); Fern (<i>Osmunda spp.</i>); Gerbera (<i>Gerbera hybrida</i>); Grape vines (<i>Litis spp.</i>) (non-bearing); Hibiscus (<i>Hibiscus spp.</i>); Hydrangea (<i>Hydrangea spp.</i>); Juniper (<i>Juniperus spp.</i>); Kalanchoe (<i>Kalanchoe blossfeldiana</i>); Lavender (<i>Lavandula spp.</i>); Orchids (<i>Arachnis spp.</i> , <i>Cattleya spp.</i> , <i>Cymbidium spp.</i> , <i>Dendrobium spp.</i> , <i>Phalaenopsis spp.</i>); Lilac (<i>Syringa spp.</i>); Pansy (<i>Viola spp.</i>); Poinsettia (<i>Euphorbia pulcherrima</i>); Roses (<i>Rosa spp.</i>); and Snapdragon (<i>Antirrhinum majus</i>).			
INSECT PESTS: Whiteflies, Spider mites, Russet mites, Broad mites, Leafhoppers, Mealybug, Flea beetles, Thrips larvae, Scale insects, Melon aphids, Potato aphid	FUNGAL PESTS[†]: Powdery mildew	FOLIAR APPLICATION RATE: Mix 1.0 teaspoon to 1.0 Tablespoon TRABON EC with 1 gallon of water. Apply 5 to 30 gallons solution per 1000 sq. ft. Ensure thorough penetration and coverage of target pest or disease for maximum effectiveness.	APPLICATION TIMING: Apply when conditions favor pest development or at first signs of damage or infestation. Repeat every 7 to 14 days as necessary.
† Not for use in California.			

ORNAMENTAL TREES[†]:			
Ash (<i>Fraxinus spp.</i>); Buckeye (<i>Aesculus spp.</i>); Cedar (<i>Cedrus spp.</i>); Crabapple (<i>Malus spp.</i>); Crape myrtle (<i>Lagerstroemia spp.</i>); Cypress (<i>Cupressaceae spp.</i>); Dogwood (<i>Cornus spp.</i>); Elm (<i>Ulmus spp.</i>); Fir (<i>Abies spp.</i>); Magnolia (<i>Grandiflora spp.</i>); Maple (<i>Aces spp.</i>); Redbud (<i>Cercis spp.</i>); Pine (<i>Pinus spp.</i>); Poplar (<i>Polpulus spp.</i>); Spruce (<i>Picae spp.</i>); and Sycamore (<i>Platanus spp.</i>).			
INSECT PESTS: Whiteflies, Spider mites, Russet mites, Broad mites, Leafhoppers, Mealybug, Thrips larvae, Scale insects	FUNGAL PESTS: Powdery mildew	FOLIAR APPLICATION RATE: Mix 1.0 teaspoon to 1.0 Tablespoon TRABON EC with 1 gallon of water. Apply 10 to 30 gallons solution per 1000 sq. ft. Ensure thorough penetration and coverage of target pest or disease for maximum effectiveness.	APPLICATION TIMING: Apply when conditions favor pest development or at first signs of damage or infestation. Repeat every 7 to 14 days as necessary.
† Not for use in California.			

CONDITIONS OF SALE AND WARRANTY**IMPORTANT: READ BEFORE USE**

Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The Directions for Use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of the Manufacturer and Seller. All such risks shall be assumed by the User or Buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, the Manufacturer and Seller make no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Manufacturer and Seller is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Manufacturer and Seller disclaim any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Manufacturer's or Seller's election, the replacement of product.

TRABON™ is a trademark of Crop Protection Technology, Pte Ltd.

DYNABLOCK CPT™ is a trademark of Crop Protection Technology, Pte Ltd.

CLEAR CROP™ is a trademark of Crop Protection Technology, Pte Ltd.

CROP CLEAN™ is a trademark of Crop Protection Technology, Pte Ltd.

AGROCLARA™ is a trademark of Crop Protection Technology, Pte Ltd.

HARVESTGUARD™ is a trademark of Crop Protection Technology, Pte Ltd.

TRABON EC (EPA Reg. No. 87749-1)

January 28, 2019 proposed amendments to the 09/18/2012 Accepted Label with 11/15/2016 Notification

Revises "Not for use in California" notations to reflect uses CDPR has agreed to accept.

{COMMERCIAL USE: Container front panel}

TRABON™ EC
FOR COMMERCIAL USE

[OPTIONAL STATEMENTS]

FOR [COMMERCIAL FARM], [FORESTRY], [NURSERY], [GREENHOUSE], AND [COMMERCIAL APPLICATOR]
USE

FOR COMMERCIAL USE IN VITICULTURE

TRABON EC is a Broad Spectrum Contact [Insecticide] [and] [Fungicide]
for use against pests in [row crops,] [tree fruits,] [berries,] [grapes,] [tropical fruits,]
[horticultural,] [and] [greenhouse crops].

Controls a variety of soft-bodied pests including [aphids,] [whiteflies,] [leafhoppers] [and] [mites].
Provides [curative] [and] [preventative control] of powdery mildew.

Active Ingredient:

Soybean oil 45.0%

Other Ingredients 55.0%

Total 100.0%

KEEP OUT OF REACH OF CHILDREN
CAUTION

See Back Panel for Additional Precautionary Statements. See Booklet for Directions for Use.

NET CONTENTS:

- 1.06 Quart (1.00 Liter)
- 1.32 Gallon (5.00 Liter)
- 2.64 Gallon (10.0 Liter)
- 5.28 Gallon (20.0 Liter)
- 52.8 Gallon (200.0 Liter)
- 264.2 Gallon (1000 Liter)

PATENT PENDING

EPA Registration No. 87749-1
EPA Establishment No. 87749-MYS-1

MANUFACTURED BY:

Crop Protection Technology, Pte Ltd
71 Bukit Batok Crescent
#07-03 Prestige Centre
Singapore 658071

Label Version No. _____

{COMMERCIAL USE: Container back panel booklet, visible without opening booklet}

TRABON™ EC

PERSONAL PROTECTION EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirt, long pants, shoes and socks. Follow manufacturer's instructions for cleaning and maintaining PPE. If no instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE after handling this product. Wash thoroughly and change into clean clothing.

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

[For containers < 5 gallons]

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. This product may be toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product if bees are visiting the treatment area.

[For containers ≥ 5 gallons]

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. This product may be toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product if bees are visiting the treatment area. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Do not reuse empty container. Store in tightly closed container in a cool dry place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

[For 1, 5, 10, 20, and 200 liter containers]

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available.

[For 1000 liter container]

CONTAINER HANDLING: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

SPILLS, LEAKS OR OTHER EMERGENCY

For minor spills or leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire or other emergency, call CHEMTREC at 1-800-262-8200 day or night.

{*COMMERCIAL USE: Container back panel booklet, NOT visible without opening booklet*}

GENERAL INFORMATION

TRABON EC is an emulsified suspension of soybean oil that when mixed with water according to the directions and applied directly to pests at the specified rates will control the listed pests.

TRABON EC is a contact pesticide. For maximum pest control, direct contact of sprays with pests is essential. Best applied in early morning or late afternoon when pests tend to be less active. The frequency of application should be adjusted to maintain sufficient pest control. Application frequency may vary depending on crop type, geographical region and local weather conditions. Consult with your state Agricultural Extension Specialist for specific recommendations regarding the crop type and pest control recommendations for your region.

Use the low application rates on the label when pest densities are relatively low. Use the progressively higher application rates on the label for progressively higher pest populations.

TRABON EC has not been tested on all plant varieties under all growing conditions. Test a small number of plants for damage before applying TRABON EC to the entire crop.

INTEGRATED PEST MANAGEMENT AND RESISTANCE MANAGEMENT

Use an effective IPM program. Avoid repeated or sequential use of products in the same mode of pesticidal action. Monitor crop for pest pressure and for loss of efficacy. Contact your extension specialist, certified crop consultant, or manufacturer for the latest resistance management information. Contact the producer to report loss of efficacy.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

APPLICATION EQUIPMENT

TRABON EC may be applied using conventional or low volume ground application equipment including but not limited to backpack sprayers, tractor mounted boom sprayers, and airblast sprayers. Do not apply this product aerially or through any type of irrigation system.

MIXING INSTRUCTIONS

Use sufficient water to ensure thorough and uniform coverage to infested plant material. Add enough water to spray tank to allow agitation. Slowly add TRABON EC while agitator is running and then add remaining water until desired volume is reached. Do not over agitate or excessive foaming may result. Triple rinse and flush spray equipment thoroughly following each use.

TRABON EC (EPA Reg. No. 87749-1)

January 28, 2019 proposed amendments to the 09/18/2012 Accepted Label with 11/15/2016 Notification

Revises "Not for use in California" notations to reflect uses CDPR has agreed to accept.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements set in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water is:

- Coveralls,
- Shoes plus socks, and
- Waterproof gloves.

TANK MIX

TRABON EC has not been tested for use in tank mixes with commonly used pesticides or spray adjuvants. It is advisable to test tank mix combinations for physical compatibility and plant safety. This product cannot be mixed with any product with a label prohibiting mixing. Do not exceed label rates. Application must be made in accordance with the more restrictive label limitations and precautions. Read and follow all label instructions. TRABON EC is most effective in tank mixes with a pH between 6.0 and 8.0. If necessary, buffer the tank mix water to conform to the optimal range.

TRABON EC (EPA Reg. No. 87749-1)

January 28, 2019 proposed amendments to the 09/18/2012 Accepted Label with 11/15/2016 Notification

Revises "Not for use in California" notations to reflect uses CDPR has agreed to accept.

APPLICATION RATE AND FREQUENCY

AVOCADO, BANANA, DATE, FIG, MANGO, PAPAYA, POMEGRANATE, PINEAPPLE †			
INSECT PESTS: Whiteflies, Spider mites, Russet mites, Broad mites, Leafhoppers, Mealybug, Flea beetles, Thrips larvae, Scale insects, Melon aphids, Potato aphid	FUNGAL PESTS: Powdery mildew	FOLIAR APPLICATION RATE: Mix 4.5 to 12.0 fl. oz. TRABON EC with 25 gallons of water. Apply 100 to 400 gallons solution per acre. Ensure thorough penetration and coverage of target pest or disease for maximum effectiveness.	APPLICATION TIMING: Apply when conditions favor pest development or at first signs of damage or infestation. Repeat every 7 to 14 days as necessary. TRABON EC may be applied up to and including the day of harvest.
† Not for use in California.			

BERRY AND SMALL FRUIT GROUP:			
Blackberry (<i>Rubus</i> spp.) (including Andean blackberry, arctic blackberry, bingleberry, black satin berry, boysenberry, brombeere, California blackberry, Chesterberry, Cherokee blackberry, Cheyenne blackberry, common blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, evergreen blackberry, Himalayaberry, hullberry, lavacaberry, loganberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, mora, mures deronce, nectarberry, Northern dewberry, olallieberry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, Southern dewberry, tayberry, youngberry, zarzamora, and cultivars, varieties and/or hybrids of these); Blueberry, highbush (<i>Vaccinium</i> spp.); Blueberry, lowbush (<i>Vaccinium angustifolium</i> Aiton); Grape (<i>Vitis</i> spp.); Kiwifruit, fuzzy (<i>Actinidia deliciosa</i> A. Chev.) (C.F. Liang and A.R. Fergusons, <i>Actinida chinensis</i> Planch.); Kiwifruit, hardy (<i>Actinidia arguta</i> (Siebold and Zucc.) Planch. ex Miq); Raspberry, black and red (<i>Rubus</i> spp.); Strawberry (<i>Fragaria x ananassa</i> Duchesne); and Cultivars, varieties, and/or hybrids of these.			
INSECT PESTS †: Whiteflies, Spider mites, Russet mites, Broad mites, Leafhoppers, Mealybug, Thrips larvae, Scale insects, Melon aphids, Potato aphid	FUNGAL PESTS: Powdery mildew	FOLIAR APPLICATION RATE: Mix 4.5 to 12 fl. oz. TRABON EC with 25 gallons of water. Apply 50 to 200 gallons solution per acre. Ensure thorough penetration and coverage of target pest or disease for maximum effectiveness.	APPLICATION TIMING: Apply when conditions favor pest development or at first signs of damage or infestation. Repeat every 7 to 14 days as necessary. TRABON EC may be applied up to and including the day of harvest.
† Not for use in California.			

BRASSICA (COLE) LEAFY VEGETABLES:

Broccoli (*Brassica oleracea* var. *botrytis*); Broccoli, Chinese (gai lon) (*Brassica alboglabra*); Cabbage (*Brassica oleracea*); Cabbage, Chinese (bok choy) (*Brassica chinensis*); Cabbage, Chinese mustard (gai choy) (*Brassica campestris*); Cauliflower (*Brassica oleracea* var. *botrytis*); Collards (*Brassica oleracea* var. *acephala*); Kale (*Brassica oleracea* var. *acephala*); and Mustard greens (*Brassica juncea*).

BULB VEGETABLES (Allium app.) †:

Chive, fresh leaves (*Allium schoenoprasum* L.); Chive, Chinese, fresh leaves (*Allium tuberosum* Rottler ex Spreng); Garlic, bulb (*Allium sativum* L. var. *sativum*) (*A. sativum* Common Garlic Group); Leek *Allium porrum* L. (syn: *A. ampeloprasum* L. var. *porrum* (L.) J. Gay) (*A. ampeloprasum* Leek Group); Lily, bulb (*Lilium* spp. (*Lilium Leichtlinii* var. *maximowiczii*, *Lilium lancifolium*)); Onion, bulb (*Allium cepa* L. var. *cepa*) (*A. cepa* Common Onion Group); Shallot, fresh leaves (*Allium cepa* var. *aggregatum* G. Don.); and Cultivars, varieties, and/or hybrids of these.

CUCURBIT VEGETABLES:

Chinese waxgourd (Chinese preserving melon) (*Benincasa hispida*); Citron melon (*Citrullus lanatus* var. *citroides*); Cucumber (*Cucumis sativus*); Gourd, edible (*Lagenaria* spp.) (includes hyotan, cucuzza); (*Luffa acutangula*, *L. cylindrica*) (includes hechima, Chinese okra); *Momordica* spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber); Muskmelon (hybrids and/or cultivars of *Cucumis melo*) (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon); Pumpkin (*Cucurbita* spp.); and Squash, summer (*Cucurbita pepo* var. *melopepo*) (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini).

FRUITING VEGETABLES (Except Cucurbits):

Bush tomato, *Solanum centrale* J.M. Black; Currant tomato, *Lycopersicon pimpinellifolium* L.; Eggplant, *Solanum melongena* L.; Pepper, bell, *Capsicum annuum* L. var. *annuum*, *Capsicum* spp.; Pepper, nonbell, *Capsicum chinese* Jacq., *C. annuum* L. var. *annuum*, *C. frutescens* L., *C. baccatum* L., *C. pubescens* Ruiz & Pav., *Capsicum* spp.; Roselle, *Hibiscus sabdariffa* L.; Tomatillo, *Physalis philadelphica* Lam; Tomato, *Solanum lycopersicum* L., *Solanum lycopersicum* L. var. *lycopersicum*; and Cultivars, varieties and/or hybrids of these.

HERBS AND SPICES †:

Basil (*Ocimum basilicum*); Caraway (*Carum carvi*); Catnip (*Nepeta cataria*); Chive, Chinese (*Allium tuberosum*); Coriander (cilantro or Chinese parsley) (leaf) (*Coriandrum sativum*); Coriander (cilantro) (seed) (*Coriandrum sativum*); Curry (leaf) (*Murraya koenigii*); Dill (dillweed) (*Anethum graveolens*); Dill (seed) (*Anethum graveolens*); Lemongrass (*Cymbopogon citratus*); Mustard (seed) (*Brassica juncea*, *B. hirta*, *B. nigra*); Parsley (dried) (*Petroselinum crispum*); Rosemary (*Rosemarinus officinalis*); Sage (*Salvia officinalis*); Tarragon (*Artemisia dracunculus*); and Thyme (*Thymus* spp.).

LEAFY VEGETABLES (Except Brassica Vegetables) †:

Arugula (Roquette) (*Eruca sativa*); Celery (*Apium graveolens* var. *dulce*); Celery, Chinese (*Apium graveolens* var. *secalinum*); Endive (escarole) (*Cichorium endivia*); Fennel, Florence (finocchio) (*Foeniculum vulgare* Azoricum Group); Lettuce, head and leaf (*Lactuca sativa*); Purslane, garden (*Portulaca oleracea*); Radicchio (red chicory) (*Cichorium intybus*); Rhubarb (*Rheum rhabarbarum*); Spinach (*Spinacia oleracea*); and Swiss chard (*Beta vulgaris* var. *cicla*).

LEGUME VEGETABLES (Succulent or Dried) †:

Bean (*Phaseolus* spp.) (includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean); Bean (*Vigna* spp.) (includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean); Broad bean (fava bean) (*Vicia faba*); Pea (*Pisum* spp.) (includes dwarf pea, edible-pod pea, English pea, field pea, garden pea, green pea, snow pea, sugar snap pea); and Soybean (*Glycine max*).

TRABON EC (EPA Reg. No. 87749-1)

January 28, 2019 proposed amendments to the 09/18/2012 Accepted Label with 11/15/2016 Notification

Revises "Not for use in California" notations to reflect uses CDPR has agreed to accept.

ROOT AND TUBER VEGETABLES †:			
Beet, garden (<i>Beta vulgaris</i>); Carrot (<i>Daucus carota</i>); Chervil, turnip-rooted (<i>Chaerophyllum bulbosum</i>); Parsnip (<i>Pastinaca sativa</i>); Radish (<i>Raphanus sativus</i>); Rutabaga (<i>Brassica campestris</i> var. <i>napobrassica</i>); Sweet potato (<i>Ipomoea batatas</i>); and Yam, true (<i>Dioscorea</i> spp.).			
FIELD USE			
INSECT PESTS Whiteflies, Spider mites, Russet mites, Broad mites, Leafhoppers, Mealybug, Flea beetles, Thrips larvae, Scale insects, Melon aphids, Potato aphid	FUNGAL PESTS †: Powdery mildew	FOLIAR APPLICATION RATE: Mix 4.5 to 12.0 fl. oz. TRABON EC with 25 gallons of water. Apply 25 to 100 gallons solution per acre. Ensure thorough penetration and coverage of target pest or disease for maximum effectiveness.	APPLICATION TIMING: Apply when conditions favor pest development or at first signs of damage or infestation. Repeat every 7 to 21 days as necessary. TRABON EC may be applied up to and including the day of harvest.
GREENHOUSE USE			
INSECT PESTS: Whiteflies, Spider mites, Russet mites, Broad mites, Leafhoppers, Mealybug, Flea beetles, Thrips larvae, Scale insects, Melon aphids, Potato aphid	FUNGAL PESTS †: Powdery mildew	FOLIAR APPLICATION RATE: [For WPS use label] Mix 4.5 to 12 fl. oz. TRABON EC with 25 gallons of water. Apply 25 to 100 gallons solution per acre. Ensure thorough penetration and coverage of target pest or disease for maximum effectiveness.	APPLICATION TIMING: Apply when conditions favor pest development or at first signs of damage or infestation. Repeat every 7 to 14 days as necessary. TRABON EC may be applied up to and including the day of harvest.
† Not for use in California.			

CEREAL GRAINS †:			
Corn (<i>Zea mays</i>); Oats (<i>Avena</i> spp.); Rice (<i>Oryza sativa</i>); Rye (<i>Secale cereale</i>); and Sorghum (milo) (<i>Sorghum</i> spp.).			
INSECT PESTS: Whiteflies, Spider mites, Russet mites, Broad mites, Leafhoppers, Mealybug, Flea beetles, Thrips larvae, Melon aphids, Potato aphid	FUNGAL PESTS: Powdery mildew	FOLIAR APPLICATION RATE: Mix 4.5 to 12.0 fl. oz. TRABON EC with 25 gallons of water. Apply 10 to 50 gallons solution per acre. Ensure thorough penetration and coverage of target pest or disease for maximum effectiveness.	APPLICATION TIMING: Apply when conditions favor pest development or at first signs of damage or infestation. Repeat every 7 to 14 days as necessary. TRABON EC may be applied up to and including the day of harvest.

CITRUS FRUITS †:

Citrus spp. *Eremocitrus* spp., *Fortunella* spp., *Microcitrus* spp., and *Poncirus* spp; Grapefruit, *Citrus natsudaoidai* Hayata; Kumquat, *Fortunella* spp; Lemon, *Citrus limon* (L.) Burm. f; Lime, *Citrus aurantiifolia* (Christm.) Swingle; Mediterranean mandarin, *Citrus deliciosa* Ten; Mount White lime, *Microcitrus garrowayae* (F.M. Bailey) *Citrus aurantium* L; Orange, sweet, *Citrus sinensis* (L.); Osbeck Pummelo, *Citrus maxima* (Burm.); *Microcitrus inodora* (F.M. Bailey) Swingle; Satsuma mandarin; Tanaka Tangelo, *Citrus x tangelo* J.W. Ingram & H.E. Moore; Tangerine (Mandarin); and Cultivars, varieties and/or hybrids of these.

POME FRUITS:

Apple, *Malus domestica* Borkh.; Loquat, *Eriobotrya japonica* (Thunb.) Lindl.; Pear, *Pyrus communis* L.; Pear, Asian, *Pyrus pyrifolia* (Burm. f.) Nakai var. *culta* (Makino) Nakai; *Pseudocydonia sinensis* (Thouin) C.K. Schneid.; and Cultivars, varieties and/or hybrids of these.

STONE FRUITS †:

Apricot (*Prunus armeniaca*); Cherry, sweet (*Prunus avium*); Nectarine (*Prunus persica*); Peach (*Prunus persica*); Plum (*Prunus domestica*, *Prunus* spp.); Plum, Japanese (*Prunus salicina*); Plumcot (*Prunus armeniaca* P. *domestica*); and Prune (fresh) (*Prunus domestica*, *Prunus* spp.).

INSECT PESTS †:	FUNGAL PESTS:	FOLIAR APPLICATION RATE:	APPLICATION TIMING:
Whiteflies, Spider mites, Leafhoppers, Mealybug, Thrips larvae, Scale insects, Melon aphids, Potato aphid	Powdery mildew	Mix 4.5 to 12 fl. oz. TRABON EC with 25 gallons of water. Apply 100 to 400 gallons solution per acre. Ensure thorough penetration and coverage of target pest or disease for maximum effectiveness.	Apply when conditions favor pest development or at first signs of damage or infestation. Repeat every 7 to 21 days as necessary. TRABON EC may be applied up to and including the day of harvest.

† Not for use in California.

OILSEED GROUP †:

Castor oil plant, *Ricinus communis* L; Chinese tallowtree, *Triadica sebifera* (L.) Small; Cottonseed, *Gossypium hirsutum* L. *Gossypium* spp; Flax seed, *Linum usitatissimum* L; Jojoba, *Simmondsia chinensis* (Link) C.K. Schneid; Poppy seed, *Papaver somniferum* L. subsp. *Somniferum*; Rapeseed, *Brassica* spp.; *B. napus* L; Rose hip, *Rosa rubiginosa* L; Safflower, *Carthamus tinctorious* L; Sesame, *Sesamum indicum* L., *S. radiatum* Schumach. & honn; Sunflower, *Helianthus annuus* L; and Cultivars, varieties, and/or hybrids of these.

INSECT PESTS:	FUNGAL PESTS:	FOLIAR APPLICATION RATE:	APPLICATION TIMING:
Whiteflies, Spider mites, Russet mites, Broad mites, Leafhoppers, Mealybug, Thrips larvae, Scale insects, Melon aphids, Potato aphid	Powdery mildew	Mix 4.5 to 12.0 fl. Oz. TRABON EC with 25 gallons of water. Apply 75 to 200 gallons solution per acre. Ensure thorough penetration and coverage of target pest or disease for maximum effectiveness.	Apply when conditions favor pest development or at first signs of damage or infestation. Repeat every 7 to 14 days as necessary. TRABON EC may be applied up to and including the day of harvest.

TRABON EC (EPA Reg. No. 87749-1)

January 28, 2019 proposed amendments to the 09/18/2012 Accepted Label with 11/15/2016 Notification

Revises "Not for use in California" notations to reflect uses CDPR has agreed to accept.

FLOWERS AND ORNAMENTAL PLANTS:

Begonia (*Begonia spp.*); Carnation (*Dianthus caryophyllus*); Chrysanthemum (*Chrysanthemum indicum*, *C. morifolium*); Cyclamen (*Cyclamen persicum*); Gerbera (*Gerbera hybrida*); Hydrangea (*Hydrangea macrophylla*); Kalanchoe (*Kalanchoe blossfeldiana*); Orchids (*Arachnis spp.*, *Catteleya spp.*, *Cymbidium spp.*, *Dendrobium spp.*, *Phalaenopsis spp.*); Pansy (*Viola spp.*); Poinsettia (*Euphorbia pulcherrima*); Roses (*Rosa spp.*); and Snapdragon (*Antirrhinum majus*).

NURSERY STOCK †:

Azalea (*Rhododendron spp.*); Camellias (*Camellia spp.*); Cotoneaster (*Cotoneaster spp.*); Fern (*Osmunda spp.*); Grape vines (*Litis spp.*) (non-bearing); Hibiscus (*Hibiscus spp.*); Hydrangea (*Hydrangea spp.*); Juniper (*Juniperus spp.*); Lavender (*Lavandula spp.*); Lilac (*Syringa spp.*); and Rose (*Rosa spp.*).

INSECT PESTS:	FUNGAL PESTS †:	FOLIAR APPLICATION RATE:	APPLICATION TIMING:
Whiteflies, Spider mites, Russet mites, Broad mites, Leafhoppers, Mealybug, Flea beetles, Thrips larvae, Scale insects, Melon aphids, Potato aphid	Powdery mildew	Mix 4.5 to 12 fl. oz. TRABON EC with 25 gallons of water. Apply 50 to 300 gallons solution per 5,000 to 20,000 square feet. Ensure thorough penetration and coverage of target pest or disease for maximum effectiveness.	Apply when conditions favor pest development or at first signs of damage or infestation. Repeat every 7 to 14 days as necessary.

† Not for use in California.

FORESTS, SHADE TREES, AND CONIFERS †:

Ash (*Fraxinus spp.*); Buckeye (*Aesculus spp.*); Cedar (*Cedrus spp.*); Crabapple (*Malus spp.*); Crape myrtle (*Lagerstroemia spp.*); Cypress (*Cupressaceae spp.*); Dogwood (*Cornus spp.*); Elm (*Ulmus spp.*); Fir (*Abies spp.*); Magnolia (*Grandiflora spp.*); Maple (*Aces spp.*); Redbud (*Cercis spp.*); Pine (*Pinus spp.*); Poplar (*Polpulus spp.*); Spruce (*Picae spp.*); and Sycamore (*Platanus spp.*).

INSECT PESTS:	FUNGAL PESTS:	FOLIAR APPLICATION RATE:	APPLICATION TIMING:
Whiteflies, Spider mites, Russet mites, Broad mites, Leafhoppers, Mealybug, Thrips larvae, Scale insects	Powdery mildew	Mix 4.5 to 12.0 fl. oz. TRABON EC with 25 gallons of water. Apply 100 to 300 gallons solution per 10,000 square feet. Ensure thorough penetration and coverage of target pest or disease for maximum effectiveness.	Apply when conditions favor pest development or at first signs of damage or infestation. Repeat every 7 to 14 days as necessary.

† Not for use in California.

TRABON EC (EPA Reg. No. 87749-1)

January 28, 2019 proposed amendments to the 09/18/2012 Accepted Label with 11/15/2016 Notification

Revises "Not for use in California" notations to reflect uses CDPR has agreed to accept.

CONDITIONS OF SALE AND WARRANTY**IMPORTANT: READ BEFORE USE**

Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The Directions for Use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of the Manufacturer and Seller. All such risks shall be assumed by the User or Buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, the Manufacturer and Seller make no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Manufacturer and Seller is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Manufacturer and Seller disclaim any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Manufacturer's or Seller's election, the replacement of product.

PATENT PENDING

EPA Registration No. 87749-1

EPA Establishment No. 87749-MYS-1

MANUFACTURED BY:

Crop Protection Technology, Pte Ltd

71 Bukit Batok Crescent

#07-03 Prestige Centre

Singapore 658071

TRABON™ is a trademark of Crop Protection Technology, Pte Ltd.

[See container for] Batch No. [_____]

Label Version No. _____