



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Biopesticides and Pollution Prevention Division (7511P)
 1200 Pennsylvania Ave., N.W.
 Washington, D.C. 20460

EPA Reg. Number:

87978-9

Date of Issuance:

4/17/2020

NOTICE OF PESTICIDE:

Registration
 Reregistration
 (under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Surtivo Plus

Name and Address of Registrant (include ZIP Code):

AgBiTech Pty Ltd
 8 Rocla Court
 Glenvale, Queensland 4350
 Australia

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Biopesticides and Pollution Prevention Division prior to use of the label in commerce. In any correspondence on this product, always refer to the above EPA Registration Number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA or the Act).

Registration is in no way to be construed as an endorsement or recommendation of this product by the U.S. Environmental Protection Agency (EPA). In order to protect health and the environment, the Administrator, on his or her motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under the Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration or registration review of your product when the EPA requires all registrants of similar products to submit such data.

Signature of Approving Official:

Jeannine Kausch, Product Manager 92
 Microbial Pesticides Branch
 Biopesticides and Pollution Prevention Division (7511P)
 Office of Pesticide Programs

Date:

4/17/2020

2. Submit storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) data as these data requirements are not satisfied. A one-year study is required to satisfy these data requirements. You have 18 months from the date of this registration to provide these data to the EPA.
3. Make the following labeling change before you release this product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 87978-9.”
4. Submit one (1) copy of the final printed labeling for the record before you release this product for shipment.

Should you wish to add/retain a reference to your company’s website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains statements or claims substantially differing from statements or claims made in connection with obtaining a FIFRA section 3 registration, the website will be referred to the EPA’s Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6. A stamped copy of the labeling is enclosed for your records. Please also note that the record for this product currently contains the following acceptable Confidential Statement of Formula (CSF):

- Basic CSF dated 04/07/2020

If you have any questions, please contact Daniel Schoeff by phone at (703) 347-0143 or via email at schoeff.daniel@epa.gov.

Sincerely,



Jeannine Kausch, Product Manager 92
Microbial Pesticides Branch
Biopesticides and Pollution
Prevention Division (7511P)
Office of Pesticide Programs

Enclosure

SURTIVO® PLUS

Biological insecticide for the integrated control of

Helicoverpa zea (corn earworm, cotton bollworm, tomato fruitworm),

Heliothis virescens (tobacco budworm),

Chrysodeixis includens (soybean looper),

Trichoplusia ni (cabbage looper)

Plutella xylostella (diamondback moth)

Anticarsia gemmatalis (velvetbean caterpillar)

Spodoptera frugiperda (fall armyworm)

Spodoptera exigua (beet armyworm)

Spodoptera eridania (southern armyworm)

on labeled food and non-food crops

 FOR ORGANIC PRODUCTION

Active Ingredients*:

<i>Autographa californica</i> Multiple Nucleopolyhedrovirus strain R3:.....	19.5%
<i>Chrysodeixis includens</i> Nucleopolyhedrovirus isolate #460:.....	4.9%
<i>Helicoverpa armigera</i> Nucleopolyhedrovirus strain ABA-NPV-U:.....	4.9%
<i>Spodoptera frugiperda</i> Multiple Nucleopolyhedrovirus strain 3AP2:	4.9%
Other ingredients:	65.8%
Total:	100.0%

* Contains a minimum of 7.5×10^9 occlusion bodies per milliliter of product

KEEP OUT OF REACH OF CHILDREN

CAUTION

See additional precautionary and first aid statements on the back panel

Net Contents: 0.25, 1, 2.5 or 5 gallons

EPA Registration N°: 87978-O

EPA Est. Number:

Manufactured for:

AgBiTech Pty Ltd

8 Rocla Court

Glenvale, Queensland 4350


Australia

Product of USA

Batch code: _____

AgBiTech Pty Ltd

SURTIVO PLUS



FIRST AID	
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment.</p> <p>For general information concerning this product, call the National Pesticide Information Center (NPIC) at 1-800-858-7378 seven days a week, 8 AM to 12 PM PST or at http://npic.orst.edu. For medical emergencies, call the poison control center at 1-800-222-1222.</p>	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through the skin. Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, Loaders, Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Mixers/loaders and applicators must wear a NIOSH-approved particulate respirator with any N, R or P filter with NIOSH approval number prefix TC-84A; or a NIOSH-approved powered air purifying respirator with an HE filter with NIOSH approval number prefix TC-21C. Repeated exposure to high concentrations of microbial proteins can cause allergic sensitization.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions are available, use detergent and hot water for washables. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product may be pathogenic to nontarget Lepidoptera exposed to direct treatment. Do not apply this product while nontarget Lepidoptera are actively visiting the treatment area. Minimize spray drift away from target area to reduce effects to nontarget Lepidoptera.

For terrestrial uses: do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

Do not contaminate water when disposing of equipment wash waters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not use this product until you have read the entire label. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Do not apply by aerial application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farm, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exemptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box apply only to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water), is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

PRODUCT INFORMATION

This product contains a biological insecticide for the control of the following moth larvae on labeled crops:

Helicoverpa zea (corn earworm, cotton bollworm, tomato fruitworm),
Heliothis virescens (tobacco budworm),
Chrysodeixis includens (soybean looper),
Trichoplusia ni (cabbage looper)
Plutella xylostella (diamondback moth)
Anticarsia gemmatalis (velvetbean caterpillar)
Spodoptera frugiperda (fall armyworm)
Spodoptera exigua (beet armyworm)
Spodoptera eridania (southern armyworm)

INSTRUCTIONS FOR USE

Surtivo Plus (*Autographa californica* Multiple Nucleopolyhedrovirus strain R3 (AcMNPV) + *Chrysodeixis includens* Nucleopolyhedrovirus isolate #460 (ChinNPV) + *Helicoverpa armigera* Nucleopolyhedrovirus strain ABA-NPV-U (HearNPV) + *Spodoptera frugiperda* Multiple Nucleopolyhedrovirus strain 3AP2 (SfMNPV)) is a highly specific pathogen of Lepidopteran insects in the following species: *Helicoverpa zea*, *Heliothis virescens*, *Plutella xylostella*, *Trichoplusia ni*, *Chrysodeixis includens*, *Spodoptera frugiperda*, *Spodoptera exigua*, *Spodoptera eridania*, and *Anticarsia gemmatalis*. The effectiveness of Surtivo Plus is dependent on a number of important factors; larval size, environmental conditions, application and the feeding behavior of the pest. Because of the requirement for adequate timing of application, coverage, and weather conditions, the performance of Surtivo Plus may be variable. Once infected, larvae can take up to 8 days to die, although feeding activity is greatly reduced within 3 days post infection, dependent on larvae size. Day time temperatures of 65°F to 95°F are ideal for the infectivity by Surtivo Plus. Infected larvae will amplify the virus, and following death will release large amounts of viral occlusion bodies that can result in ongoing control, particularly under environmental conditions suitable for the virus in Surtivo Plus (warm and humid conditions).

Good coverage of the feeding sites of the larvae is essential, as the product needs to be ingested to be effective. If larvae are feeding low down in a heavy crop canopy, and application of Surtivo Plus does not reach these areas, initial control from the spray will be sub-optimal. However, larvae that die from the Surtivo Plus spray will release large amounts of AcMNPV, ChinNPV, HearNPV and SfMNPV which will spread throughout the crop canopy.

USES

Surtivo Plus should only be used to target small larvae (1st and 2nd). Third instar larvae will take longer to die and cause more significant damage prior to death.

RAIN FASTNESS

The majority of virus uptake by larvae occurs within one (1) hour post-application. For this reason, it is best to avoid applying Surtivo Plus if heavy rain (greater than 0.4 inches per hour) is expected within 1 hour after application. However, do not delay application if only moderate rain is expected, or heavy rain (greater than 0.4 inches per hour) is not imminent.

CROPS, APPLICATION RATES AND CROP SPECIFIC INFORMATION

Crops	Rate of Lepigen per acre	Additional Information
Sorghum	1.7 to 2.8 fl. oz.	Use lower application rates when targeting 1 st instar larvae and in mixtures with sprays for midge control . Use higher application rates when targeting 2 nd instar larvae. Do not use as a stand alone treatment under high pressure situations.
Cereal Grains Group (Crop Group 15 excluding Sweet corn) including: Corn (maize), Popcorn, Rice ¹ Alfalfa (hay and seed) Oilseed Group (Crop Group 20) including: Flax seed, Canola, Safflower, Sunflower Peanut Legume Vegetables (Succulent or Dried) Group (Crop Group 6 excluding Peanut and Soybean) including: Adzuki bean, Broad bean, Chickpeas, Cowpea, Faba bean, Field pea, Kidney bean, Lablab bean, Lentil, Lima bean, Sweet lupin, White lupin, Mung bean, Navy bean, Pigeon pea Vetch	1.7 to 2.8 fl. oz.	Use lower application rates when targeting 1 st instar larvae and in mixtures with other insecticides. Use higher application rates when targeting 2 nd instar larvae. Do not use as a stand alone treatment under high pressure situations.
Soybean (including Edamame)	1.7 to 2.8 fl. oz.	Use lower application rates as a preventive measure in vegetative crop stages. Use the high application rate when the pest population has reached economic threshold. Do not use as a stand alone treatment under high pressure situations.
Sweetcorn	1.7 to 2.8 fl. oz.	Application should be made from the vegetative stages until silking. Applications during silking should employ a high application rate and be in conjunction with other control measures. Application of low rates at regular (3 to 5 day) intervals, particularly via overhead irrigation water, is an effective strategy from vegetative stages, through row tassel to silking. Do not use as a stand alone treatment under high pressure situations.
Root and Tuber Vegetables Group (Crop Group 1) including: Carrot, Sweet potato, Sugar beet, Potato Brassica (Cole) Leafy Vegetables Group (Crop Group 5) including: Broccoli, Brussels sprouts, Cabbages, Cauliflower, Chinese broccoli, Kale, Mustard greens, Mustard spinach, Rape greens Leafy Vegetables (Except Brassica) Group (Crop Group 4) including: Celery, Endive, Lettuce, Roquette, Silver beet, Spinach Fruiting Vegetables Group (crop Group 8 – 10) including: Chili, Eggplant, Okra, Peppers, Tomato Cucurbit Vegetables Group (Crop Group 9) including: Cucumber, Melons Pumpkins Summer and winter Squash, Watermelon, Zucchini Berries Group (Crop Group 13 – 07) including: Blackberry, Blueberry, Boysenberry, Cranberry ² , Currants, Gooseberry, Raspberry, Strawberry Pome Fruit Group (Crop Group 11 – 10) including: Apple, Pear	1.7 to 2.8 fl. oz.	Use the lower application rate to control 1 st instar. Use a higher application rate when flowers, fruit or economic parts of the crop are present, and to control 2 nd instar larvae. Application of low rates at regular (3 to 5 day) intervals, particularly via overhead irrigation water, is an effective strategy in horticultural crops. Do not use as a stand alone treatment under high pressure situations.

¹ Do not apply to flooded fields

² Do not apply to flooded fields

Ornamental flowers and plants Avocado Asparagus		
Cotton	1.7 to 2.8 fl. oz.	High leaf pH in cotton causes rapid NPV deactivation, giving Lepigen very short residual activity and resulting in highly variable performance in this crop. Use lower application rates when targeting 1 st instar larvae and in mixtures with other insecticides. Use higher application rates when targeting 2 nd instar larvae. Do not use as a stand alone treatment under high pressure situations.
Tobacco, Hemp	1.7 to 2.8 fl. oz.	Use lower application rates when targeting 1 st instar larvae and in mixtures with other insecticides. Use higher application rates when targeting 2 nd instar larvae. Do not use as a stand alone treatment under high pressure situations. Using the lower application rate at regular spray intervals is an effective strategy in tobacco.

MIXING INSTRUCTIONS

Shake the container well before use. Spray water pH should be neutral (pH 7.0) – spray water pH above 8 may damage the virus and performance will be reduced. If needed, use a suitable buffer or acidifier. If mixing with other pesticides or foliar fertilizers in water, add Surtivo Plus to the spray tank after the other products are thoroughly diluted. Surtivo Plus should be applied within 10 hours after mixing. Do not let stand overnight.

Compatibility:

In water: Surtivo Plus is highly compatible with the majority of herbicides, insecticides, fungicides and foliar fertilizers when mixed in water. Ensure that the mixture has a pH of 8 or less before adding Surtivo Plus as higher pH levels will damage the virus.

APPLICATION INSTRUCTIONS

Use application parameters (nozzles, swath width, pressure, boom height, speed, etc.) to ensure thorough coverage of the target area.

- I Legume Vegetables (Succulent and Dried) Group (Crop Group 6 excluding Soybean), Root and Tuber Vegetables Group (Crop Group 1) including: Carrot, Sweet potato, Sugar beet, Potato; *Brassica* (Cole) Leafy Vegetables Group (Crop Group 5) including: Broccoli, Brussels sprouts, Cabbage, Cauliflower, Chinese broccoli, Kale, Mustard greens, Mustard spinach, Rape greens); Leafy Vegetables (except *Brassica* Vegetables) Group (Crop Group 4) including: Celery, Endive, Lettuce, Roquette, Spinach; Fruiting Vegetable Group (Crop Group 8 - 10) including: Chili, Eggplant, Okra, Pepper, Tomato; Cucurbit Vegetables Group (Crop Group 9) including: Cucumber, Melon, Pumpkin, Summer and winter squash, Watermelon, Zucchini; Berries Group (Crop Group 13 - 07) including: Blackberry, Blueberry, Boysenberry, Cranberry³, Currant, Gooseberry, Raspberry, Strawberry; Pome Fruit Group (Crop Group 11 - 10) including: Apple, Pear; Ornamental Flowers and Plants; Avocado; Asparagus; Tobacco; Hemp.

Ground Rig

Apply Surtivo Plus by ground rig or hand-held equipment in a minimum of 40 gallons of water per acre.

- II Sorghum, Cereal Grains Group⁴, Alfalfa, Oilseed Group, Peanut, Soybean, Sweet Corn, Cotton

Ground Rig

Apply Surtivo Plus in a minimum of 10 gallons of water per acre.

³ Do not apply to flooded fields

⁴ Rice - Do not apply to flooded fields

Chemigation (via overhead irrigation water):

Surtivo Plus can be effectively applied to crops in overhead irrigation water. The product should be introduced to the irrigation water at the appropriate rate using irrigation equipment. If the product is diluted in water prior to injection into the irrigation water, ensure that the dilution water is clean and not silty with a pH of 7 or less and ensure there is constant agitation. Preferably, rainwater should be used for dilution. Ensure any diluted Surtivo Plus is used within 10 hours of mixing.

For one-pass mobile irrigators (such as: center-pivot, lateral-move, end-tow, side-roll, traveler, big-gun), continuously and evenly introduce the required quantity of Surtivo Plus into the irrigation water over the course of irrigation. It is recommended to apply Surtivo Plus in no more than 0.5 inches of irrigation water. For static irrigators (such as: solid-set or hand-move), introduce the required amount of Surtivo Plus into the irrigation water just prior to completion of the irrigation period, to maximize the concentration of Surtivo Plus applied and the amount that remains on the crop. See the CHEMIGATION section (following) for additional information.

CHEMIGATION**General Requirements:**

- 1) Apply this product only through overhead sprinkler (including center-pivot, lateral-move, end-tow, side-(wheel) roll, traveler, big-gun, solid-set, or hand-move) irrigation systems. Do not apply this product through soil irrigation systems including drip (trickle).
- 2) Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- 3) If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.
- 4) Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- 5) A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make any necessary adjustments should the need arise.

Requirements for Chemigation Systems Connected to Public Water Systems:

- 1) Public water supply means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of 25 individuals daily at least 60 days throughout the year.
- 2) Chemigation systems connected to the public water systems must contain a functional, reduced-pressure zone (RPZ), backflow preventer or the functional equivalent in the water supply upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or the overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- 3) The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- 4) The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system

interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

- 5) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- 6) Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- 7) Do not apply when wind speed favors drift beyond the area intended for treatment.

Requirements for Sprinkler Chemigation:

- 1) The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow.
- 2) The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- 3) The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- 4) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- 5) The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- 6) Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- 7) Do not apply when wind speed favors drift beyond the area intended for treatment.

DAYS TO HARVEST

There are no restrictions on applying Surtivo Plus up to the time of harvest.

STORAGE and DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE

Care must be taken to avoid exposure of Surtivo Plus to high temperatures (above 104°F). Packaged sealed product can be exposed to direct sunlight for brief periods of time (<2 hours), but should be kept out of direct sunlight.

- Surtivo Plus stored in air-conditioned rooms (< 77°F) will be viable for at least 6 months
- Surtivo Plus stored in cool rooms and refrigerators (0°F to 59°F) will be viable for at least 12 months

* Surtivo Plus stored in a freezer (0°F) will remain liquid.

Note: Exposure of Surtivo Plus to temperature from 77°F to 104°F for short periods (<36 hours) will not affect efficacy. Transport time of 36 hours or less in non-refrigerated, covered trucks is acceptable as long as the product does not exceed 104°F.

PESTICIDE DISPOSAL

To avoid wastes, use material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

CONTAINER HANDLING

[Plastic containers with capacities equal to or less than 5 gallons:] Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinance. If burned, stay out of smoke.

WARRANTY

This product is warranted to contain the amount of active ingredients as described in this label and that the product will be as effective as intended if properly transported, used, and applied per the label instructions. The effectiveness of this product may be degraded by improper storage, transportation or handling and may be subject to environmental factors out of AgBiTech Pty Ltd's control. The user must monitor the performance of the product as climatic, geographical or biological variables and/or developed resistance may affect the results obtained. To the extent allowed under the applicable law, AgBiTech Pty Ltd and its subsidiaries makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label or accepts no responsibility in respect of this product. To the extent consistent with applicable law, AgBiTech Pty Ltd and its subsidiaries disclaim any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product