



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

May 3, 2019

Olav Messerschmidt
OMC Ag Consulting, Inc.
Registration Agent for Terramera, Inc.
828 Tanglewood Lane
East Lansing, MI 48823

Subject: Non-PRIA (Pesticide Registration Improvement Act) Labeling Amendment –
Amendment to Add OMRI Logo, Add Alternate Brand Name, Add Tank Size Options,
Improve Descriptive language in the Directions
Product Name: TNO70 Broad Spectrum
EPA Registration Number: 88760-10
Application Date: 11/8/18
OPP Decision Number: 546222

Dear Mr. Messerschmidt:

The amended labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable.

The alternate brand name - Rango – has been added to the registration, and our records have been updated accordingly. This approval does not affect any terms or conditions that were previously imposed on this registration. You continue to be subject to existing terms or conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release this product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. “To distribute or sell” is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR § 152.3.

Should you wish to add/retain a reference to your company’s website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the U.S. Environmental Protection Agency (EPA). If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or

Page 2 of 2
EPA Reg. No. 88760-10
OPP Decision No. 546222

misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA-approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Chris Pfeifer of my team by phone at (703) 308-0031 or via email at pfeifer.chris@epa.gov.

Sincerely,

A handwritten signature in blue ink that reads "Andrew C. Bryceland". The signature is fluid and cursive, with the first name being the most prominent.

Andrew Bryceland, Team Leader
Biochemical Pesticides Branch
Biopesticides and Pollution
Prevention Division (7511P)
Office of Pesticide Programs

Enclosure

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

TNO70 Broad Spectrum (EPA Reg. No. 88760-10)

MASTER LABEL

Sublabel A is for Agricultural Use

Sublabel B is for Commercial Greenhouse and Nurseries

Sublabel C is for Residential Use

TNO70 Broad Spectrum

Biological Insecticide [/Fungicide] [/Nematicide] [/Miticide]

*For control of foliar and soil insects, fungal diseases and nematodes.
For use in vegetables, fruits, tree crops, grapes, row crops, greenhouses,
ornamental plants, nurseries, and other listed plants*



Active Ingredient:

Cold Pressed Neem Oil..... 70.0%

Other Ingredients..... 30.0%

Total 100.0%

KEEP OUT OF REACH OF CHILDREN

READ ALL DIRECTIONS BEFORE USING THIS PRODUCT

Shake Well Before Use

NET CONTENTS: 8 FL. OZ. to 275 GALLONS

EPA Reg. No.: 88760-10
EPA Est. No. 49292-WA-001

Batch No.

Manufactured by:
Terramera, Inc.
6920 Salashan Pkwy D-109
Ferndale, WA 98248

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

[Optional Marketing Claims – Applicable to All Sublabels]

- [Use] The Power of Neem
- [Cold Pressed]] Neem Oil
- Derived From Neem
- Cold-pressed from seeds of the Neem Tree
- [TNO70 Broad Spectrum] contains Cold-Pressed Neem Oil that [provides] [contains] [keeps] the full spectrum of limonoids
- [TNO70 Broad Spectrum] contains the full spectrum of bioactive[s] [compounds] of Neem [Oil]
- Cold pressed neem oil, plant-based [biopesticide] active ingredient
- Cold pressed neem oil, botanical [based] [biopesticide] active ingredient
- Cold pressed neem oil is a botanical active ingredient.
- Cold pressed neem oil is a plant-based active ingredient.
- Protect your plants from insects and diseases
- Broad Spectrum [control] [action]
- Multiple Modes of Action
- Insect Growth Regulator
- [TNO70 Broad Spectrum is a] broad-spectrum bioinsecticide
- [TNO70 Broad Spectrum is a] broad-spectrum biofungicide
- [4-in-1,] [multi-purpose product –] fungicide, insecticide, miticide and nematocide* [for use on listed pest and crops]
- Multi-purpose biopesticide [- 3 in 1] [product]
- One product, triple control
- Insecticide
- Miticide
- Nematicide
- Fungicide
- Botanical Insect Growth Regulator (IGR)
- Kills eggs, larvae and adult insects
- Controls insects and their eggs
- Affects insects at all stages of growth
- [Excellent] for control of spider mites, aphids, whiteflies, leafhoppers, caterpillars
- Controls chewing and sucking insects
- [Also] controls aphids, [beetles,] [stink bugs,] [caterpillars,] [leafhoppers,] [leafminers,] [whiteflies,] [mealy bugs,] [midges,] [spider mites,] [nematodes,] [weevils,] [scales,] [and] [&] [thrips] [other listed insects and nematodes]
- Prevents and controls listed [major] diseases
- Prevents fungal attack of plant tissues
- Tool for prevention and control of powdery mildew and botrytis
- Controls powdery mildew, botrytis, [stem mildew,] [and] [&] [sour rot]
- Use for the prevention and control of powdery mildew, botrytis, stem mildew and others as listed on label.
- [For] Use as part of an Integrated Pest Management [program] [strategy]
- Complement for IPM programs

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

- Alternative within conventional programs
- Peel Here for Directions & Precautions
- Concentrate
- [Only] a 4-hour Re-Entry Interval
- Spray right up to the time of harvest
- Zero-day Pre-Harvest interval
- Can be used up to day of harvest [as a [4 in 1,] multi-purpose product [– fungicide, insecticide, miticide and nematocide *]] [*for use on listed pest and crops]
- For indoor and outdoor use
- For use in hydroponic systems
- Keep plants healthy; keep insects away
- Made in the USA

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

TNO70 Broad Spectrum (EPA Reg. No. 88760-10)

Sublabel A: Agricultural Use

TNO70 Broad Spectrum

[ABN: Rango]

*For control of foliar and soil insects, fungal diseases and nematodes.
For use in vegetables, fruits, tree crops, grapes, row crops, greenhouses,
ornamental plants, nurseries, and other listed plants.*



FOR ORGANIC PRODUCTION

[OMRI Logo]{For Rango only}

Active Ingredient:

Cold Pressed Neem Oil..... 70.0%

Other Ingredients..... 30.0%

Total 100.0%

[This product contains 5.37lbs of cold pressed neem oil per gallon.]

KEEP OUT OF REACH OF CHILDREN

READ ALL DIRECTIONS BEFORE USING THIS PRODUCT

Shake Well Before Use

[See label booklet for additional precautionary statements, directions for use, storage and disposal statements, and warranty.]

NET CONTENTS: 8 FL. OZ. to 275 GALLONS

EPA Reg. No. I: 88760-10
EPA Est. No. 49292-WA-001

Batch No.

Manufactured by:
Terramera, Inc.
6920 Salashan Pkwy D-109
Ferndale, WA 98248

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions are available, wash with detergent and hot water. Keep and store PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment break-down.

ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

USER SAFETY RECOMMENDATIONS

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if contaminated with pesticide. Wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during product application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS).

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Long-sleeved shirt and long pants
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

These requirements apply to uses of this product that are NOT within the scope of the WPS for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. For other uses, including golf courses and other non-agricultural uses, do not enter treated areas without protective clothing until sprays have dried

PRODUCT MODE OF ACTION

TNO70 Broad Spectrum controls target pests on contact or by ingestion. The modes of action on insects are repellence, anti-feedance and interference with the molting process. Diseases are controlled by inhibition of mycelial growth.

GENERAL INFORMATION

Read all directions before using this product.

TNO70 Broad Spectrum is an emulsifiable concentrate containing cold pressed neem oil for the broad spectrum control of listed pests in vegetables, fruits, tree crops, grapes, agronomic crops, ornamental plants, greenhouses, and other listed plants. **TNO70 Broad Spectrum** is exempted from the requirement of a tolerance and may be applied to listed food and non-food crop up to and including the day of harvest.

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

- Thorough coverage is key to providing good insect, mite and disease control.
- For best results maintain constant agitation in spray tank and apply immediately.
- For optimal performance do not mix with cold water (less than 45°F).
- Application in early morning/late evening is recommended to minimize the potential for leaf burn.
- Do not apply under high humidity and temperature conditions >90°F.
- Do not apply to wilted or stressed plants and newly germinated or transplanted crops prior to root establishment.
- Use with care on plants with tender tissue. Test on a small area prior to broader use.
- Do not apply to sensitive plant species such as poinsettias, impatiens, hibiscus flowers, certain carnation and rose flower species, ornamental olive trees and comice pear.
- Weather conditions, intensity, type and physical stages of the pests, and treated crop can influence the degree of product efficacy.
- **DO NOT** apply sulfur or sulfur containing products within 14 days of a **TNO70 Broad Spectrum** application.

RATES

Insecticide and Miticide Foliar Applications:

- Use a concentration of 0.625 -1.25% v/v for normal pest and crop conditions.
- Use a concentration 1.8% v/v **for heavier infestations.**
- Up to a maximum use rate of 3 quarts **TNO70 Broad Spectrum** per acre.

Fungicide Foliar Application:

- Use a concentration of 1.25-1.8% v/v.
- Up to a maximum use rate of 6 quarts **TNO70 Broad Spectrum** per acre.

Soil Applications:

- Use a concentration of 1.25-2.4% v/v.
- Up to maximum use rate of 7.25 quarts **TNO70 Broad Spectrum** per acre.

MIXING INSTRUCTIONS

TNO70 Broad Spectrum Alone:

- **TNO70 Broad Spectrum** is an emulsifiable concentrate and requires only water for the appropriate use dilution. Additional surfactant is not required.
- **Shake the container well before use.**
- Add **TNO70 Broad Spectrum** to a clean spray tank half-filled with water and agitate.
- Next, add additional water to final spray volume, while maintaining continuous agitation.
- Best results are achieved by using a spray water with a temperature of 45°F or warmer.
- If water temperature is below 45°F, achieve a good emulsion by premixing **TNO70 Broad Spectrum** at 1:1 ratio with tepid water, add to half-filled spray tank, agitate, then fill to final spray volume.

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

- Agitate continuously during mixing and application to prevent separation of the emulsion. Inadequate agitation can cause a non-uniform dilution resulting in crop injury and/or reduced efficacy.
- Always use the spray solution promptly after mixing and do not allow mixture to sit for extended periods of time. If allowed to sit, agitate thoroughly before resuming application.
- The recommended pH range of the spray water is between 5.5 – 7 for optimal performance. If needed, adjust by adding a pH modifier.

Neem oil can solidify at temperatures below 60°F. If solidified, thaw the product by setting out in temperatures over 80°F and agitating well before mixing with water. For optimal emulsion, do not use cold water (less than 45°F).

Mixing Order for Tank Mixes:

- Fill clean spray tank with water to 1/3 of the required spray volume.
- Start agitation.
- Add different formulation types in the following order: 1) water dispersible granules, 2) wettable powders.
- Maintain agitation and add water to $\frac{3}{4}$ of final spray volume.
- Next add **TNO70 Broad Spectrum**, other emulsifiable concentrates, water-based solutions, adjuvants, surfacts, oils and/or fertilizers.
- Agitate to achieve complete emulsification. Do not use if a uniform, cloudy emulsion is not formed.
- Continue adding water and agitating to desired final spray volume.
- Always use the spray solution promptly after mixing with water.
- Do not let tank mixture sit for an extended period of time. If tank mixture is allowed to sit, agitate thoroughly again prior to and during application. Sparger line agitators are preferred.
- Tank-mix combinations can alter the pH of the finished spray solution. Adjusting the spray mixture pH to a range between 5.5 and 7.0 will provide optimal performance.

TANK MIX COMPATIBILITY:

To determine the physical compatibility of **TNO70 Broad Spectrum** with other products, test as described below before mixing.

Jar Compatibility Test: Using a quart jar, add the proportionate amounts of products to be tank mixed to 1 quart of water in the following order. Add wettable powders and water-dispersible granular products first, then add liquid flowables, then add emulsifiable concentrates and solutions last. After thoroughly mixing by agitation, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank. All possible tank mixes on all crops have not been tested. Growers must test tank mix combinations for phytotoxicity on a sample of plants prior to use. Do not use mixtures of incompatible products as it may cause phytotoxicity or result in lowered effectiveness.

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

Always read and follow the directions for use, precautions and limitations for use on all product labels used in combination. Applications must follow the precautions and limitations of the most restrictive product label in the mixture. Do not exceed the dosage rates of any product. Check compatibility by using the correct proportion of the products in a small test container.

DO NOT apply sulfur or sulfur containing products within 14 days of a **TNO70 Broad Spectrum** application.

APPLICATION DIRECTIONS

Apply **TNO70 Broad Spectrum** as a foliar spray or as soil treatment (soil drench, in-furrow, drip-applied) using thoroughly clean equipment. Applications can be made with any powered or manual pesticide application equipment including high volume, low volume, ultra-low volume, electrostatic, air blast, and fogging equipment. When applied as a foliar application, ensure complete coverage of the plant surfaces, but avoid pooling or run off. Follow the original equipment manufacturer's instructions.

INSECTICIDE/MITICIDE FOLIAR USE

To control listed insect pests, apply **TNO70 Broad Spectrum** in sufficient amounts of water with adequate spray pressure to achieve thorough coverage of plant surfaces, ensuring that both the top and bottom of leaves are wetted. **TNO70 Broad Spectrum** is most effective when applied before or around the onset of insects, mites or their eggs (see Pests Section) or as soon as they are detected. Apply at a concentration of 0.625-1.8 v/v up to a maximum rate of 3 quarts of **TNO70 Broad Spectrum** per acre. Spray early in the morning or in the evening for best results. Repeat application if it rains within four hours of spraying. Avoid spraying under conditions of high humidity and high temperature (>90°F). For optimal results, repeat the applications at intervals of 7-10 days. Use higher rates and increase spray frequency when pest pressure is high and/or dense crop canopies exist.

Crop	Pest	Concentration	Maximum Rate (TNO70 Broad Spectrum/acre)	Spray Interval
All crops	See Pests Section – Insects and Mites	0.625-1.8% v/v	3 quarts	7-10 days

FUNGICIDE FOLIAR USE

To control listed diseases, apply **TNO70 Broad Spectrum** in sufficient amount of water and with adequate spray pressure to achieve thorough coverage of plant surfaces. **TNO70 Broad Spectrum** is most effective when applied before the onset of disease development. Apply a maximum rate of 6 quarts of **TNO70 Broad Spectrum** per acre at a concentration of 1.25 – 1.8% v/v. Spray early in the morning or in the evening for best results. Repeat application if it rains within four hours of spraying. Avoid spraying under conditions of high humidity and high temperature (>90°F). Do not apply with any sulfur or sulfur containing products within 14 days of a **TNO70 Broad Spectrum** application.

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

Crop	Pest	Concentration	Maximum Rate (TNO70 Broad Spectrum/acre)	Spray Interval
All crops except Grapes	See Pests Section – Diseases	1.25 - 1.8%	6 quarts	10-14 days
Grapes	Powdery Mildew, Stem Mildew, Sour Rot	1.25 - 1.8%	6 quarts	10-14 days from pre-bloom through veraison
	Botrytis			Spray at bloom, pre-bunch closure, veraison and 14 days after veraison

NEMATICIDE AND OTHER SOIL USE

To control listed nematodes, apply as a preventative treatment (see Pests Section for Nematodes) or control treatment after nematodes and other listed pests have been detected. When used as a soil application (soil drench, in-furrow, drip-applied), apply at 1.25 – 2.4% for a maximum rate of 7.25 quarts of **TNO70 Broad Spectrum** per acre to deliver complete and thorough coverage. When applied as a soil drench, avoid excess run off. For best results repeat the applications as necessary.

Crop	Pest	Concentration	Maximum Rate (TNO70 Broad Sepctrum/acre)	Spray Interval
All crops	See Pests Section - Nematodes	1.25 - 2.4%	7.25 quarts	Repeat as Needed

Root-dip Nematicide Use on Strawberries

For bare-root dip applications on strawberries use a concentration of 2.4% (e.g. 2.4 gallons of **TNO70 Broad Spectrum** in 100 gallons of water). If bare-root nursery plants are in cold storage, allow them to thaw to ambient temperature – approximately 20°C (70°F). Submerge the entire plant to be treated in **TNO70 Broad Spectrum** emulsion. Leave the plant completely submerged in for 15-30 minutes. Remove the plants from the treatment solution, shake off excess liquid, and drain for 5-15 min. Plant after treatment, or package the plants in suitable containers and cold store between -2°C and 5 °C (28-48 °F) during shipping and until planting.

CHEMIGATION INSTRUCTIONS

GENERAL CHEMIGATION REQUIREMENTS

Apply this product only through in-furrow or drip (trickle) irrigation & system(s). Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

FURROW CHEMIGATION REQUIREMENTS

Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as a drop structure or weir box to decrease potential for water source contamination from back flow if water flow stops. Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

A supply tank is recommended for this product. If using a supply tank, dilute this product at the rate of 7.25 quarts per 100~200 gallons of water. Frequent agitation is necessary. Apply in the second half of the water application to deliver **TNO70 Broad Spectrum** to the soil pests.

DRIP CHEMIGATION REQUIREMENTS

The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional inter-locking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

to the point where pesticide distribution is adversely affected. Systems must use a metering pump such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

A supply tank is recommended for this product. If using a supply tank, dilute this product at the rate of 6 quarts per 100~200 gallons of water. Frequent agitation is necessary. Apply in the second half of the water application to deliver **TNO70 Broad Spectrum** to the soil pests.

**Dilution Table for Foliar Applications
(10 to 100 gallons per acre)**

Gallons of Water	0.625% v/v	1.25% v/v	1.8% v/v
5	4 fl oz	8 fl oz	12 fl oz
10	8 fl oz	16 fl oz	23 fl oz
20	16 fl oz	32 fl oz	46 fl oz
25	20 fl oz	40 fl oz	58 fl oz
50	40 fl oz	80 fl oz	116 fl oz
75	60 fl oz	120 fl oz	173 fl oz
100	80 fl oz	160 fl oz	230 fl oz

PHYTOTOXICITY: To avoid plant damage, test for crop response by applying the spray solution on a small portion of the area to be treated before applying to the entire area. Make foliar applications in conditions that favor fast drying. Avoid applications during hot temperature conditions >90°F. Make applications early morning/late afternoon to avoid leaf burn. Not all possible mixtures of pesticide sprays, fertilizers, surfactants, and adjuvants have been tested. Therefore, it is the responsibility of the user to test spray mixtures to small areas to ensure crop safety before treating the entire area.

USE SITES

TNO70 Broad Spectrum is exempted from the requirement of a tolerance and may be applied to the following food and non-food crop groups up to and including the day of harvest.

Crop Group 3 - Bulb Vegetable Crops such as

Garlic	Onion
Leek	Shallot

Crop Group 9 - Cucurbit Crops such as:

Cantaloupe	Honeydew Melon	Squash, Summer
------------	----------------	----------------

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

Crenshaw Melon	Persian Melon	Squash, Winter
Cucumber	Pumpkin	Watermelon

Crop Group 8 - Fruiting Vegetable Crops such as:

Eggplant	Tomatillo
Pepper	Tomato

Crop Group 4 & 5 - Leafy & Brassica (Cole) Vegetable Crops such as

Arugula	Cilantro	Mustard green
Broccoli	Collard	Parsley
Brussel sprout	Endive	Radicchio
Cabbage	Greens	Rhubarb
Celery	Kale	Spinach
Chinese cabbage	Kohlrabi	Swiss chard
Cauliflower	Lettuce	

Crop Group 6 - Legume Crops such as:

Bean	Lentil
Chickpea	Pea
Guar	Soybean

Crop Group 1 - Root & Tuber Vegetable Crops such as:

Articoke	Horseradish	Radish
Beet	Parsnip	Sweet Potato
Carrot	Potato	Yam

Crop Group 13 - Small Fruit & Berry Crops such as:

Blackberry	Kiwifruit
Blueberry	Raspberry
Cranberry	Strawberry
Grape	

Crop Group 10, 23 & 24 - Citrus & Tropical Fruit Crops such as:

Avocado	Grapefruit	
Banana	Guava	Olive
Citrus	Lemon	Orange
Date	Lime	Papaya
Fig	Mandarin	Pineapple
	Mango	Pomegranate

Crop Group 11 & 12 - Pome & Stone Fruit Crops such as:

Apple	Nectarine	Peach
-------	-----------	-------

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

Apricot	Pear	Prune
Cherry	Plum	
Crop Group 14 - Tree Nuts such as:		
Almond	Filbert	Pecan
Cashew	Hickory Nut	Pistachio
Chestnut	Macadamia Nut	
Coconut		
Crop Group 19 - Herbs & Spices such as:		
Basil	Dill	Poppy
Chamomile	Fennel	Rosemary
Chive	Mint	Sage
Cinnamon	Mustard	Tarragon
Clove buds	Nutmeg	Wintergreen
Cumin	Pepper	
Curry leaf	Peppermint	
Crop Group 15 - Cereal Grain Crops such as:		
Barley	Oats	Triticale
Corn	Rye	Wheat
Millet	Sorghum (Milo)	Wild Rice
Crop Group 18 - Forage Crops such as:		
Alfalfa	Sainfoin	
Clover	Trefoil	
Lupin	Vetch	
Crop Group 20 & 21 - Miscellaneous Crops such as:		
Canola	Mushroom	Sesame
Coffee	Okra	Sugarcane
Cotton	Peanut	Sunflower
Hops	Safflower	Tobacco
Other Use Sites such as:		
Ornamentals	Greenhouses	Sod Farms
Fencerows	Nurseries	Turf
Mushroom Houses	Shade Houses	

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

PESTS: INSECTS, MITES, NEMATODES, AND DISEASES

(1) Insects and Mites: Use the following rate ranges and spray intervals to control the insects and mites listed below.

Concentration	Maximum Rate (TNO70 Broad Spectrum/acre)	Spray Interval
0.625-1.8% v/v	3 quarts	7-10 days

<p>Aphids such as: Cotton Aphid Cowpea Aphid Pea Aphid Green Peach Aphid</p>	<p>Grasshoppers such as Carolina Grasshopper Rice Grasshopper</p>	<p>Scales such as: California Red Scale Coffee Green Scale San Jose Scale Soft Scale</p>
<p>Beetles such as: Cucumber Beetle Japanese Beetle Spotted Cucumber Beetle</p>	<p>Leafhoppers such as Grape Leafhopper Potato leafhopper</p>	<p>Thrips such as: Flower Thrip Grape Thrip Onion Thrip Western Flower Thrip</p>
<p>Borers such as: Peachtree Borers Peach Twig Borers</p>	<p>Leafminer such as: Citrus Leafminer Tomato Leafminer Vegetable Leafminer</p>	<p>True Plant Bugs such as: Lygus Bug Phylloxera</p>
<p>Caterpillars/Moths/Worms such as: Armyworms Budworms Cutworms Diamondback Moth Gypsy Moth Leafrollers Loopers Navel Orange Worm</p>	<p>Maggots/Grubs such as: Onion Maggots</p>	<p>Spittle Bug Sting Bug Tomato Stink Bug</p>
<p>Flies/Gnats such as: Fruit Fly Fungus Gnat Walnut Husk Fly</p>	<p>Mealy Bugs such as: Citrus Mealy Bug</p>	<p>Weevils such as: Black Vine Weevil Boll weevil Pepper Weevil</p>
	<p>Mites such as: Pacific Spider Mites Red Spider Mite Spider Mites Two Spotted Spider Mite</p>	<p>Wireworms such as: Field/Wheat Wireworm</p>
	<p>Psyllids such as: Asian Citrus Psyllid Pear Psyllid Potato Psyllid</p>	<p>Whiteflies such as: Cotton Whitefly</p>

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

Silverleaf Whitefly
Greenhouse Whitefly

(2) Diseases: Use the following rate ranges and spray intervals to control the diseases listed below.

Crop	Pest	Concentration	Maximum Rate (TNO70 Broad Spectrum/acre)	Spray Interval
All crops except Grapes	See Pests Section – Diseases	1.25 - 1.8%	6 quarts	10-14 days
Grapes	Powdery Mildew, Stem Mildew, Sour Rot	1.25 - 1.8%	6 quarts	10-14 days from pre-bloom through veraison
	Botrytis			Spray at bloom, pre-bunch closure, veraison and 14 days after veraison

Foliar Fungal Diseases

Alternaria	Downey Mildew	Scab
Anthracnose	Molds	Stem Mildew
Blight (early, late, leaf)	Powdery Mildew	Southern Blight
Botrytis	Rust	Sour Rot Grapes

Soil Fungal Diseases

Fusarium Oxysporum
Pythium
Rhizoctonia Solani

(3) Nematodes: Use the following rate ranges and spray intervals to control the nematodes listed below.

Concentration	Maximum Rate (TNO70 Broad Spectrum/acre)	Spray Interval
1.25 - 2.4%	7.25 quarts	Repeat as Needed

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

Nematodes such as:

Dagger Nematode	Reniform Nematode	Soybean Cyst Nematode
Lance Nematode	Root Knot Nematode	Sting Nematode
Lesion Nematode		

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by pesticide storage or disposal.

PESTICIDE STORAGE: Do not store this product above 104°F or below 20°F for extended periods of time. Keep containers tightly closed and in original containers when not in use. Do not store exposed to ultraviolet light (sunlight) or moisture. Neem oil clouds and solidifies at temperatures below 59°F. If oil has solidified, gently thaw by exposing to temperatures over 80°F. Store in such a manner to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out of the reach of children, preferably in a locked storage area. Keep container closed when not in use.

PESTICIDAL DISPOSAL: To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

CONTAINER HANDLING:

{For 5-Gallon or Smaller Containers} Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or dispose of in trash or in a sanitary landfill or by incineration.

{For Containers Larger than 5 Gallons} Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or dispose of in trash or in a sanitary landfill or by incineration.

{For 250 or 275-gallon Refillable Containers} Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

IMPORTANT: READ BEFORE USE

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Seller. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, Terramera, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a purpose or otherwise, that extend beyond the statements made on this label. No agent of Terramera, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Terramera, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Terramera, Inc.'s election, the replacement of product.

[Terramera is a registered trademark of Terramera, Inc.]

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

[Optional Marketing Claims for Sublabel A]

- [Can be used] for organic production

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

TNO70 Broad Spectrum (EPA Reg. No. 88760-10)

Sublabel B: Commercial Greenhouse and Nurseries

TNO70 Broad Spectrum

*For control of foliar and soil insects, fungal diseases and nematodes
in commercial plant production in greenhouses, under shade cloth, in container stock and
field nurseries*



Active Ingredient:

Cold Pressed Neem Oil..... 70.0%

Other Ingredients..... 30.0%

Total 100.0%

[This product contains 5.37lbs of cold pressed neem oil per gallon.]

KEEP OUT OF REACH OF CHILDREN

READ ALL DIRECTIONS BEFORE USING THIS PRODUCT

Shake Well Before Use

[See label booklet for additional precautionary statements, directions for use, storage and disposal statements, and warranty.]

NET CONTENTS: 8 FL. OZ. to 275 GALLONS

EPA Reg. No.: 88760-10
EPA Est. No. 49292-WA-001

Batch No.

Manufactured by:
Terramera, Inc.
6920 Salashan Pkwy D-109
Ferndale, WA 98248

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions are available, wash with detergent and hot water. Keep and store PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment break-down.

ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

USER SAFETY RECOMMENDATIONS

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if contaminated with pesticide. Wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during product application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS).

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Long-sleeved shirt and long pants
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

These requirements apply to uses of this product that are NOT within the scope of the WPS for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. For other uses, including golf courses and other non-agricultural uses, do not enter treated areas without protective clothing until sprays have dried.

PRODUCT MODE OF ACTION

TNO70 Broad Spectrum controls target pests on contact or by ingestion. The modes of action on insects are repellence, anti-feedance and interference with the molting process. Diseases are controlled by inhibition of mycelial growth.

GENERAL INFORMATION

Read all directions before using this product.

TNO70 Broad Spectrum is an emulsifiable concentrate containing 70% cold pressed neem oil for the broad spectrum control of pests in commercial plant production in greenhouses, under shade cloth and in container stock and field nurseries. **TNO70 Broad Spectrum** is exempted from the requirement of a tolerance and may be applied to listed food and non-food crop up to and including the day of harvest.

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

- Thorough coverage is key to providing good insect, mite and disease control.
- For best results maintain constant agitation in spray tank and apply immediately.
- For optimal performance do not mix with cold water (less than 45°F).
- Application in early morning/late evening is recommended to minimize the potential for leaf burn.
- Do not apply under high humidity and temperature conditions >90°F.
- Do not apply to wilted or stressed plants and newly germinated or transplanted crops prior to root establishment.
- Use with care on plants with tender tissue. Test on a small area prior to broader use.
- Do not apply to sensitive plant species such as poinsettias, impatiens, hibiscus flowers, certain carnation and rose flower species, ornamental olive trees and comice pear.
- Weather conditions, intensity, type and physical stages of the pests, and treated crop can influence the degree of product efficacy.
- **DO NOT** apply sulfur or sulfur containing products within 14 days of a **TNO70 Broad Spectrum** application.

RATES

Insecticide and Miticide Foliar Applications:

- Use a concentration of 0.625 - 1.25% v/v for normal pest and crop conditions.
- Use a concentration 1.8% v/v **for heavier infestations.**
- Up to a maximum use rate of 3 quarts **TNO70 Broad Spectrum** per acre.

Fungicide Foliar Applications:

- Use a concentration of 1.25 - 1.8% v/v.
- Up to a maximum use rate of 6 quarts **TNO70 Broad Spectrum** per acre.

Soil Applications:

- Use a concentration of 1.25 - 2.4% v/v.
- Up to maximum use rate of 7.25 quarts **TNO70 Broad Spectrum** per acre.

MIXING INSTRUCTIONS

TNO70 Broad Spectrum Alone:

- **TNO70 Broad Spectrum** is an emulsifiable concentrate and requires only water for the appropriate use dilution. Additional surfactant is not required.
- **Shake the container well before use.**
- Add **TNO70 Broad Spectrum** to a clean spray tank half-filled with water and agitate.
- Next, add additional water to final spray volume, while maintaining continuous agitation.
- Best results are achieved by using a spray water with a temperature of 45°F or warmer.
- If water temperature is below 45°F, achieve a good emulsion by premixing **TNO70 Broad Spectrum** at 1:1 ratio with tepid water, add to half-filled spray tank, agitate, then fill to final spray volume.

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

- Agitate continuously during mixing and application to prevent separation of the emulsion. Inadequate agitation can cause a non-uniform dilution resulting in crop injury and/or reduced efficacy.
- Always use the spray solution promptly after mixing and do not allow mixture to sit for extended periods of time. If allowed to sit, agitate thoroughly before resuming application.
- The recommended pH range of the spray water is between 5.5 – 7 for optimal performance. If needed, adjust by adding a pH modifier.

Neem oil can solidify at temperatures below 60°F. If solidified, thaw the product by setting out in temperatures over 80°F and agitating well before mixing with water. For optimal emulsion, do not use cold water (less than 45°F).

Mixing Order for Tank Mixes:

- Fill clean spray tank with water to 1/3 of the required spray volume.
- Start agitation.
- Add different formulation types in the following order: 1) water dispersible granules, 2) wettable powders.
- Maintain agitation and add water to ¾ of final spray volume.
- Next add **TNO70 Broad Spectrum**, other emulsifiable concentrates, water-based solutions, adjuvants, surfacts, oils and/or fertilizers.
- Agitate to achieve complete emulsification. Do not use if a uniform, cloudy emulsion is not formed.
- Continue adding water and agitating to desired final spray volume.
- Always use the spray solution promptly after mixing with water.
- Do not let tank mixture sit for an extended period of time. If tank mixture is allowed to sit, agitate thoroughly again prior to and during application. Sparger line agitators are preferred.
- Tank-mix combinations can alter the pH of the finished spray solution. Adjusting the spray mixture pH to a range between 5.5 and 7.0 will provide optimal performance.

TANK MIX COMPATIBILITY:

To determine the physical compatibility of **TNO70 Broad Spectrum** with other products, test as described below before mixing.

Jar Compatibility Test: Using a quart jar, add the proportionate amounts of products to be tank mixed to 1 quart of water in the following order. Add wettable powders and water-dispersible granular products first, then add liquid flowables, then add emulsifiable concentrates and solutions last. After thoroughly mixing by agitation, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank. All possible tank mixes on all crops have not been tested. Growers must test tank mix combinations for phytotoxicity on a sample of plants prior to use. Do not use mixtures of incompatible products as it may cause phytotoxicity or result in lowered effectiveness.

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

Always read and follow the directions for use, precautions and limitations for use on all product labels used in combination. Applications must follow the precautions and limitations of the most restrictive product label in the mixture. Do not exceed the dosage rates of any product. Check compatibility by using the correct proportion of the products in a small test container.

DO NOT apply sulfur or sulfur containing products within 14 days of a **TNO70 Broad Spectrum application**.

APPLICATION DIRECTIONS

Apply **TNO70 Broad Spectrum** as a foliar spray or as soil treatment (soil drench, in-furrow, drip-applied) using thoroughly clean equipment. Applications can be made with any powered or manual pesticide application equipment including high volume, low volume, ultra-low volume, electrostatic, air blast, and fogging equipment. When applied as a foliar application, ensure complete coverage of the plant surfaces, but avoid pooling or run off. Follow the original equipment manufacturer’s instructions.

INSECTICIDE/MITICIDE FOLIAR USE

To control listed insect pests, apply **TNO70 Broad Spectrum** in sufficient amounts of water with adequate spray pressure to achieve thorough coverage of plant surfaces, ensuring that both the top and bottom of leaves are wetted. **TNO70 Broad Spectrum** is most effective when applied before or around the onset of insects, mites or their eggs (see Pests Section) or as soon as they are detected. Apply at a concentration of 0.625-1.8 v/v up to a maximum rate of 3 quarts of **TNO70 Broad Spectrum** per acre. Spray early in the morning or in the evening for best results. Repeat application if it rains within four hours of spraying. Avoid spraying under conditions of high humidity and high temperature (>90°F). For optimal results, repeat the applications at intervals of 7-10 days. Use higher rates and increase spray frequency when pest pressure is high and/or dense crop canopies exist.

Crop	Pest	Concentration	Maximum Rate (TNO70 Broad Spectrum/acre)	Spray Interval
All crops	See Pests Section – Insects and Mites	0.625-1.8% v/v	3 quarts	7-10 days

FUNGICIDE FOLIAR USE

To control listed diseases, apply **TNO70 Broad Spectrum** in sufficient amount of water and with adequate spray pressure to achieve thorough coverage of plant surfaces. **TNO70 Broad Spectrum** is most effective when applied before the onset of disease development. Apply a maximum rate of 6 quarts of **TNO70 Broad Spectrum** per acre at a concentration of 1.25 – 1.8% v/v. Spray early in the morning or in the evening for best results. Repeat application if it rains within four hours of spraying. Avoid spraying under conditions of high humidity and high temperature (>90°F). Do not apply with any sulfur or sulfur containing products within 14 days of a **TNO70 Broad Spectrum** application.

Crop	Pest	Concentration	Maximum Rate	Spray Interval
------	------	---------------	--------------	----------------

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

			(TNO70 Broad Spectrum/acre)	
All crops	See Pests Section – Diseases	1.25 - 1.8%	6 quarts	10-14 days

NEMATICIDE AND OTHER SOIL USE

To control listed nematodes, apply as a preventative treatment (see Pests Section for Nematodes) or control treatment after nematodes and other listed pests have been detected. When used as a soil application (soil drench, in-furrow, drip-applied), apply at 1.25 – 2.4% for a maximum rate of 7.25 quarts of **TNO70 Broad Spectrum** per acre to deliver complete and thorough coverage. When applied as a soil drench, avoid excess run off. For best results repeat the applications as necessary.

Crop	Pest	Concentration	Maximum Rate (TNO70 Broad Spectrum/acre)	Spray Interval
All crops	See Pests Section - Nematodes	1.25 - 2.4%	7.25 quarts	Repeat as Needed

Root-dip Nematicide Use on Nursery Plants

For bare-root dip applications on plants use a concentration of 2.4% (e.g. 2.4 gallons of **TNO70 Broad Spectrum** in 100 gallons of water). If bare-root nursery plants are in cold storage, allow them to thaw to ambient temperature – approximately 20°C (70°F). Submerge the entire plant to be treated in **TNO70 Broad Spectrum** emulsion. Leave the plant completely submerged in for 15-30 minutes. Remove the plants from the treatment solution, shake off excess liquid, and drain for 5-15 min. Plant immediately after treatment.

CHEMIGATION INSTRUCTIONS

GENERAL CHEMIGATION REQUIREMENTS

Apply this product only through in-furrow or drip (trickle) irrigation & system(s). Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

FURROW CHEMIGATION REQUIREMENTS

Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as a drop structure or weir box to decrease potential for water source contamination from back

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

flow if water flow stops. Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

A supply tank is recommended for this product. If using a supply tank, dilute this product at the rate of 7.25 quarts per 100~200 gallons of water. Frequent agitation is necessary. Apply in the second half of the water application to deliver **TNO70 Broad Spectrum** to the soil pests.

DRIP CHEMIGATION REQUIREMENTS

The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional inter-locking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

A supply tank is recommended for this product. If using a supply tank, dilute this product at the rate of 7.25 quarts per 100~200 gallons of water. Frequent agitation is necessary. Apply in the second half of the water application to deliver **TNO70 Broad Spectrum** to the soil pests.

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

**Dilution Table for Foliar Applications
(1 to 20 gallons per acre)**

Gallons of Water	0.625% v/v	1.25% v/v	1.8% v/v
1	0.8 fl oz	1.6 fl oz	2.3 fl oz
5	4 fl oz	8 fl oz	12 fl oz
10	8 fl oz	16 fl oz	23 fl oz
20	16 fl oz	32 fl oz	46 fl oz

PHYTOTOXICITY: To avoid plant damage, test for crop response by applying the mixed spray solution on a small portion of the area to be treated before applying to the entire area. Make foliar applications in conditions that favor fast drying. Avoid applications during hot temperature conditions >90F. Make applications early morning/late afternoon to avoid leaf burn. Not all possible mixtures of pesticide sprays, fertilizers, surfactants, and adjuvants have been tested. Therefore, it is the responsibility of the user to test spray mixtures to small areas to ensure crop safety before treating the entire area.

USE SITES

TNO70 Broad Spectrum is exempt from tolerances and may be applied to listed food and non-food crop groups up to and including the day of harvest. Use **TNO70 Broad Spectrum** on the following crop groupings.

Crop Group 3 - Bulb Vegetable Crops such as:

Garlic	Onion
Leek	Shallot

Crop Group 9 - Cucurbit Crops such as:

Cantaloupe	Honeydew Melon	Squash, Summer
Crenshaw Melon	Persian Melon	Squash, Winter
Cucumber	Pumpkin	Watermelon

Crop Group 8 - Fruiting Vegetable Crops such as:

Eggplant	Tomatillo
Pepper	Tomato

Crop Group 4 & 5 - Leafy & Brassica (Cole) Vegetable Crops such as:

Arugula	Cilantro	Mustard green
Broccoli	Collard	Parsley
Brussel sprout	Endive	Radicchio
Cabbage	Greens	Rhubarb
Celery	Kale	Spinach

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

Chinese cabbage	Kohlrabi	Swiss chard
Cauliflower	Lettuce	

Crop Group 6 - Legume Crops such as:

Bean	Lentil
Chickpea	Pea
Guar	Soybean

Crop Group 1 - Root & Tuber Vegetable Crops such as:

Articoke	Horseradish	Radish
Beet	Parsnip	Sweet Potato
Carrot	Potato	Yam

Crop Group 13 - Small Fruit & Berry Crops such as:

Blackberry	Kiwifruit
Blueberry	Raspberry
Cranberry	Strawberry
Grape	

Crop Group 10, 23 & 24 - Citrus & Tropical Fruit Crops such as:

Avocado	Grapefruit	
Banana	Guava	Olive
Citrus	Lemon	Orange
Date	Lime	Papaya
Fig	Mandarin	Pineapple
	Mango	Pomegranate

Crop Group 11 & 12 - Pome & Stone Fruit Crops such as:

Apple	Nectarine	Peach
Apricot	Pear	Prune
Cherry	Plum	

Crop Group 14 - Tree Nuts such as:

Almond	Filbert	Pecan
Cashew	Hickory Nut	Pistachio
Chestnut	Macadamia Nut	
Coconut		

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

Crop Group 19 - Herbs & Spices such as:

Basil	Dill	Poppy
Chamomile	Fennel	Rosemary
Chive	Mint	Sage
Cinnamon	Mustard	Tarragon
Clove buds	Nutmeg	Wintergreen
Cumin	Pepper	
Curry leaf	Peppermint	

Crop Group 15 - Cereal Grain Crops such as:

Barley	Oats	Triticale
Corn	Rye	Wheat
Millet	Sorghum (Milo)	Wild Rice

Crop Group 18 - Forage Crops such as:

Alfalfa	Sainfoin
Clover	Trefoil
Lupin	Vetch

Crop Group 20 & 21 - Miscellaneous Crops such as:

Canola	Mushroom	Sesame
Coffee	Okra	Sugarcane
Cotton	Peanut	Sunflower
Hops	Safflower	Tobacco

Other Use Sites such as:

Ornamentals	Greenhouses	Sod Farms
Fencerows	Nurseries	Turf

PESTS: INSECTS, MITES, NEMATODES, and DISEASES

(1) **Insects and Mites:** Use the following rate ranges and spray intervals to control the insects and mites listed below.

Concentration	Maximum Rate (TNO70 Broad Spectrum/acre)	Spray Interval
0.625-1.8% v/v	3 quarts	7-10 days

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

Aphids such as: Cotton Aphid Cowpea Aphid Pea Aphid Green Peach Aphid	Grasshoppers such as Carolina Grasshopper Rice Grasshopper	Scales such as: California Red Scale Coffee Green Scale San Jose Scale Soft Scale
Beetles such as: Cucumber Beetle Japanese Beetle Spotted Cucumber Beetle	Leafhoppers such as Grape Leafhopper Potato leafhopper	Thrips such as: Flower Thrip Grape Thrip Onion Thrip Western Flower Thrip
Borers such as: Peachtree Borers Peach Twig Borers	Leafminer such as: Citrus Leafminer Tomato Leafminer Vegetable Leafminer	True Plant Bugs such as: Lygus Bug Phylloxera
Caterpillars/Moths/Worms such as: Armyworms Budworms Cutworms Diamondback Moth Gypsy Moth Leafrollers Loopers Navel Orange Worm	Maggots/Grubs such as: Onion Maggots	Spittle Bug Sting Bug Tomato Stink Bug
Flies/Gnats such as: Fruit Fly Fungus Gnat Walnut Husk Fly	Mealy Bugs such as: Citrus Mealy Bug	Weevils such as: Black Vine Weevil Boll weevil Pepper Weevil
	Mites such as: Pacific Spider Mites Red Spider Mite Spider Mites Two Spotted Spider Mite	Wireworms such as: Field/Wheat Wireworm
	Psyllids such as: Asian Citrus Psyllid Pear Psyllid Potato Psyllid	Whiteflies such as: Cotton Whitefly Silverleaf Whiefly Greenhouse Whitefly

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

(2) Diseases: Use the following rate ranges and spray intervals to control the diseases listed below.

Crop	Pest	Concentration	Maximum Rate (TNO70 Broad Spectrum/acre)	Spray Interval
All crops	See Pests Section - Diseases	1.25-1.8%	6 quarts	10-14 days

Foliar Fungal Diseases

Alternaria	Downey Mildew	Scab
Anthrachnose	Molds	Stem Mildew
Blight (early, late, leaf)	Powdery Midlew	Southern Blight
Botrytis	Rust	Sour Rot Grapes

Soil Fungal Diseases

Fusarium Oxyporum
Pythium
Rhizoctonia Solani

(3) Nematodes: Use the following rate ranges and spray intervals to control the nematodes listed below.

Concentration	Maximum Rate (TNO70 Broad Sepctrum/acre)	Spray Interval
1.25 - 2.4%	7.25 quarts	Repeat as Needed

Nematodes such as:

Dagger Nematode	Reniform Nematode	Soybean Cyst Nematode
Lance Nematode	Root Knot Nematode	Sting Nematode
Lesion Nematode		

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by pesticide storage or disposal.

PESTICIDE STORAGE: Do not store this product above 104°F or below 20°F for extended periods of time. Keep containers tightly closed and in original containers when not in use. Do not store exposed to ultraviolet light (sunlight) or moisture. Neem oil clouds and solidifies at temperatures below 59°F. If oil has solidified, gently thaw by exposing to temperatures over 80°F. Store in such a manner to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out of the reach of children, preferably in a locked storage area. Keep container closed when not in use.

PESTICIDAL DISPOSAL: To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

CONTAINER HANDLING:

{For 5-Gallon or Smaller Containers} Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or dispose of in trash or in a sanitary landfill or by incineration.

{For Containers Larger than 5 Gallons} Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or dispose of in trash or in a sanitary landfill or by incineration.

{For 250 or 275-gallon Refillable Containers} Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

IMPORTANT: READ BEFORE USE

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Seller. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, Terramera, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a purpose or otherwise, that extend beyond the statements made on this label. No agent of Terramera, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Terramera, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Terramera, Inc.'s election, the replacement of product.

[Terramera is a registered trademark of Terramera, Inc.]

[Optional Marketing Claims For Sublabel B]

- For use in hydroponic systems
- For greenhouses, shadehouses, interiorscapes, mushroom houses, and nursery uses.
- For indoor and outdoor vegetables, ornamentals, flowers, trees, shrubs, container grown plants, and interiorscapes.
- [Can be used] for organic production

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

TNO70 Broad Spectrum (EPA Reg. No. 88760-10)

Sublabel C: Residential Use

TNO70 Broad Spectrum

[Biological] Insecticide [/Fungicide] [/Nematicide] [/Miticide]

*For Residential Use on Listed Crops
including vegetables, fruits, citrus, nuts, [and] ornamental plants, lawns and other
listed plants*



FOR ORGANIC GARDENING

Active Ingredient:

Cold Pressed Neem Oil.....70.0%

Other Ingredients.....30.0%

Total 100.0%

KEEP OUT OF REACH OF CHILDREN

READ ALL DIRECTIONS BEFORE USING THIS PRODUCT

Shake Well Before Use

[See back label for additional precautionary statements, directions for use, storage and disposal statements, and warranty.]

NET CONTENTS: 3 FL. OZ. to 5 GALLONS

EPA Reg. No.: 88760-10
EPA Est. No. 49292-WA-001

Batch No.

Manufactured by:
Terramera, Inc.
6920 Salashan Pkwy D-109
Ferndale, WA 98248

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION

TNO70 Broad Spectrum controls listed insects on contact or by ingestion. The product acts on insects by way of repellence, anti-feedant action and interference with the molting process. **TNO70 Broad Spectrum** controls listed diseases by inhibiting mycelial fungal growth. The efficacy of this product is dependent on weather conditions, intensity of pest population, area of application, and physical stages of pests and crops.

Do not store this product above 104°F or below 20°F for extended periods of time. Keep containers tightly closed and in original containers when not in use. Do not store exposed to ultraviolet light (sunlight) or moisture.

THAWING (MELTING) INSTRUCTIONS

Neem oil can solidify at temperatures below 60°F. Thaw (melt) solidified product by standing product container in warm water. Make sure cap is tightly closed. Or set out in temperatures over 80°F. Avoid direct exposure to temperatures over 104°F.

MIXING INSTRUCTIONS

TNO70 Broad Spectrum contains cold pressed neem oil and requires only water for the appropriate use dilution. Add **TNO70 Broad Spectrum** to a mixing tank or spray bottle with a little water and agitate thoroughly. For optimal mixing, ensure water temperature is above 45°F. Then add remaining water to the spray tank or spray bottle with continuous agitation. Agitate continuously while spraying. Non-uniform dilution can cause damage to plant or result in lowered effectiveness.

CROP USES: TNO70 Broad Spectrum may be used on bulbs, cucurbits, fruiting vegetables, leafy and brassica (cole) vegetables, legumes, root and tuber vegetables, small fruit and berries, citrus and tropical fruits, pome and stone fruits, tree nuts, herbs, and spices, grass, lawn, and ornamentals.

USE RATES:

- (1) **To Control Insects or Mites:** Apply **TNO70 Broad Spectrum** at 0.625 – 1.8% in sufficient amounts of water to achieve complete coverage in a 7-10 day interval.

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

(2) To Control Diseases: Use at 1.2 – 1.8% in sufficient amounts of water to achieve complete coverage with a 10-14 day interval.

(3) To Control Nematodes and For Soil Use: Use at 1.2 – 2.4% in sufficient amounts of water to achieve complete coverage with a 10-14 day interval.

TNO70 Broad Spectrum is most effective when applied before insects or eggs are present in large numbers.

SPRAY DIRECTIONS

Apply **TNO70 Broad Spectrum** as a foliar spray or as soil drench to control listed pests. When applied as a soil drench, avoid excessive run off. When using as a foliar application, ensure thorough coverage of plant surfaces, but avoid pooling or run off.

Avoid spraying under conditions of high humidity and high temperature (>90°F). To avoid potential leaf burn, apply the product in early morning or late evening. Always test the product on a small portion of plants for compatibility before treating the entire plants.

Instructions for 1-Gallon Sprayer

Pests	Fluid Ounces TNO70 Broad Spectrum per Gallon Water	Tablespoons (Tbsp) TNO70 Broad Spectrum per Gallon Water
Insects – most conditions	0.8-1.5 fl. oz	1.5-3 Tbsp
Insects – heavy infestation	2.3 fl. oz	4.5 Tbsp
Disease control	1.5-2.3 fl. oz	3 – 4.5 Tbsp
Soil Treatments	1.5 - 3 fl. oz	3 – 6 Tbsp

Instructions for 32 oz. (1 Qt.) Spray Bottle

Pests	Teaspoons (tsp) per 32 fl. oz. (1 Qt) Water
Insects – most conditions	1 1/8 - 2 1/4 tsp
Insects – heavy infestation	3 1/2 tsp
Disease control	2 1/4 - 3 1/2 tsp
Soil Treatments	2 1/4 - 4 1/2 tsp

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

PESTS:

TNO70 Broad Spectrum may be used to control a variety of insects and diseases listed below.

Aphids such as: Cotton Aphid Cowpea Aphid Pea Aphid Green Peach Aphid	Grasshoppers such as: Carolina Grasshopper Rice Grasshopper	Scales such as: California Red Scale Coffee Green Scale San Jose Scale Soft Scale
Beetles such as: Cucumber Beetle Japanese Beetle Spotted Cucumber Beetle	Leafhoppers such as: Grape Leafhopper Potato leafhopper	Thrips such as: Flower Thrip Grape Thrip Onion Thrip Western Flower Thrip
Borers such as: Peachtree Borers Peach Twig Borers	Leafminer such as: Citrus Leafminer Tomato Leafminer Vegetable Leafminer	True Plant Bugs such as: Lygus Bug Phylloxera
Caterpillars/Moths/Worms such as: Armyworms Budworms Cutworms Diamondback Moth Gypsy Moth Leafrollers Loopers Navel Orange Worm	Maggots/Grubs such as: Onion Maggots	Weevils such as: Spittle Bug Sting Bug Tomato Stink Bug
Flies/Gnats such as: Fruit Fly Fungus Gnat Walnut Husk Fly	Mealy Bugs such as: Citrus Mealy Bug	Wireworms such as: Field/Wheat Wireworm
	Mites such as: Pacific Spider Mites Red Spider Mite Spider Mites Two Spotted Spider Mite	Whiteflies such as: Cotton Whitefly Silverleaf Whiefly Greenhouse Whitefly
	Psyllids such as: Asian Citrus Psyllid Pear Psyllid Potato Psyllid	

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

Foliar Fungal Diseases

Alternaria	Downey Mildew	Scab
Anthracnose	Molds	Stem Mildew
Blight (early, late, leaf)	Powdery Mildew	Southern Blight
Botrytis	Rust	Sour Rot Grapes

Soil Fungal Diseases

Fusarium Oxysporum
Pythium
Rhizoctonia Solani

Nematodes such as

Dagger Nematode	Reniform Nematode	Soybean Cyst Nematode
Lance Nematode	Root Knot Nematode	Sting Nematode
Lesion Nematode		

STORAGE AND DISPOSAL

Store in a cool dry place out of reach of children and pets and away from direct sunlight, windows, flames and other sources of heat or ignitions. Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE:

Terramera, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Terramera, Inc., and, (2) buyer and user assume the risk of any such use. TO THE EXTENT PERMITTED BY APPLICABLE LAW, TERRAMERA, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

{Note: Texts in [] are optional language. Texts in { } are notes for reviewers.}

[Optional Marketing Claims for Sublabel C]

- TNO70 Broad Spectrum is an effective biological fungicide, insecticide, miticide and nematocidal for organic gardening use on vegetables[, tree fruits] [, nuts] [, vines] [, citrus] [, ornamental flowers] [, shrubs and trees] [, grass lawns] [, and] [houseplants].
- For use on vegetables, [fruits,] [nuts,] [vines,] [citrus,] [ornamental plants,] [shrubs and trees,] [grass lawns,] [houseplants]
- [For] use in and around home and home garden
- [Can be used] for organic gardening
- For greenhouses, shadehouses, interiorscapes and home garden
- For indoor and outdoor vegetables, ornamentals, flowers, trees, shrubs, container grown plants, and interiorscapes.